

Interactive
Comment

Interactive comment on “Chlorophyll *a* specific $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in stream periphyton: implications for aquatic food web studies” by N. F. Ishikawa et al.

Anonymous Referee #3

Received and published: 25 August 2015

This study investigated the chlorophyll-*a* specific isotopic compositions in stream periphyton to examine whether the bulk isotopic compositions of periphyton could be used as representative of aquatic producers. The results showed that periphyton chlorophyll-*a* exhibited ^{13}C and ^{14}C values similar to the bulk tissue, but had higher ^{15}N value than the bulk sample. The difference in ^{15}N value between chlorophyll-*a* and bulk sample was attributed to N isotopic fractionation during chlorophyll-*a* biosynthesis and incorporation of cyanobacteria tissue into periphyton. Because of the novelty of measurement on chlorophyll-*a* specific isotopic compositions of ^{13}C , ^{14}C and ^{15}N of stream periphyton, I would like to recommend this manuscript for Biogeosciences. However, I think that there are some issues to be addressed before final publication.

C4677

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



For example, the authors calculated the relative contribution of algal carbon and terrestrial organic carbon to periphyton based on ^{14}C values of bulk periphyton, chlorophyll-a, and terrestrial plant for each season (April and October). They concluded that the periphyton consisted of 89 – 95 % algal carbon. I wonder if this is a meaningful and reliable calculation. The algal portion of periphyton should consist of both alive and dead (aged) algal tissues. Further, ^{14}C value in periphyton chlorophyll-a changed largely (ca. 60 permil) differed between April and October. Therefore, I suppose that the difference in ^{14}C values of bulk periphyton and chlorophyll-a could be accounted for not only by terrestrial organic carbon incorporation but also by the seasonal variation in ^{14}C of chlorophyll-a. Actually, *Cladophora* sp., the aquatic primary producer, also presented a difference (ca. 10 permil) in ^{14}C between bulk periphyton and chlorophyll-a. The difference is comparable to that in periphyton in October. I think that it would be necessary to consider more carefully about the premise of the calculation.

Additionally, ^{14}C value of chlorophyll-a of terrestrial plant leaves (-10 permil) was much lower than that of bulk ^{14}C (27 permil). The difference was considered to be because of use of old soil CO_2 and soil organic carbon. It should be extremely interesting if the plant can have access to such an old carbon source. The two cited papers (Bloemen et al. and Bruggemann et al.) indeed described the potential importance of these carbon sources for plant production, but these two references did not demonstrate that plants could use such an old carbon for primary production. To my knowledge, most of previous ^{14}C studies have shown that respired soil CO_2 and dissolved soil organic carbon have modern carbon. The recycle of phytol was also used to explain the ^{14}C difference between chlorophyll-a and bulk plant leaves. I like this idea but it is difficult to believe that plant reuse such an old phytol to synthesize chlorophyll-a. Please consider presenting more convincing evidence to support the authors' idea.

Minor comments P11090: Please consider describing the rationale of this study in the first sentence of Abstract. P11090L10, P11098L15: The authors stated that ^{13}C of periphyton do not trace carbon transfer between primary producers and primary con-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

sumers. However, the ^{13}C data clearly indicated that the mayfly larva did not subsist on C of periphyton that was investigated. Please clarify what kind of C flow the authors intended to mention. P11090L15: mixture of only two sources (carbonates and atmospheric CO_2)? What about CO_2 derived from aquatic and terrestrial organic matter? P11091L26: Periphyton ^{14}C is “often” derived P11094L23: washed with H_2O after HCl treatment? P11094L24: when was the periphyton sample collected? P11095L8: Please describe briefly how to confirm that the product was phaeophytin-a. P11096: Please add more explanations about how to transfer the dried chlorophyll-a samples to tin capsules for ^{13}C and ^{15}N and quartz tubes for ^{14}C measurement. P11099L5: proxy for “ ^{13}C ” of bulk algae. P11101L3: It is a great idea. But are there any studies demonstrating that an algae can collect phytol from DOC or POC? Fig.1 and 2.: Please indicate what the error bars stand for.

Interactive comment on Biogeosciences Discuss., 12, 11089, 2015.

BGD

12, C4677–C4679, 2015

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C4679

