

1 Dear Editor,

2 We thank you and the referees for all provided comments, which helped us to further improve the quality  
3 of the manuscript. We carefully considered all of the suggestions and – with the exception of changing the  
4 title - revised the manuscript accordingly in all cases. We hope that it will now be considered adequate for  
5 publication in *Biogeosciences*.

6 Attached please find a point-by-point reply to the issues addressed in the two reviews as well as a revised  
7 clear as well as a marked-up copy of the manuscript and all figures.

8 Sincerely,

9 Alexander Röhl

10

11

12

13 **POINT-BY-POINT REPLY TO THE REVIEWS:**

14  
15

16 **RESPONSE TO REFEREE 1:**

17  
18

Dear Referee,

19  
20

We appreciate your careful reading of our manuscript and the numerous insightful suggestions. Changes to the manuscript detailed below refer to the "markup copy" which is attached as a pdf to this comment. We also attached a clear copy of the manuscript as well as all figures.

21  
22  
23

Sincerely,  
Alexander Röhl

24  
25  
26

27  
28 **General comments**

29  
30

**Referee:** The authors investigated the effect of age and micro-meteorological conditions on transpiration of oil palms in a humid tropical lowland in Indonesia. The authors investigated palms stands varying in age between 2 and 25 years. Medium ages stands had a 12-fold higher transpiration than 2 year old stands. This is a valuable dataset and interesting for the readership of Biogeosciences. The major weak point of this study, however, is that most of the 3-weeks sap flow measurements were not performed simultaneously but were conducted successively and thus under varying weather conditions. To get rid of this methodological problem the authors limited their data evaluation for each stand to the average of three comparably sunny and dry days. Therefore, I wonder how the authors come at the end to the conclusion that the temporal variability of oil palm transpiration is rather low. I do not agree with this conclusion. First of all, the statement itself is misleading. Over the day there is of course a huge temporal variation in transpiration. What the authors probably mean that the diurnal course of transpiration did not vary much among the three days and the stands.

31  
32  
33

**Authors:** We agree with the reviewer that the non-simultaneous measurements in the 15 stands are a weakness of the study; however, it is very complicated under field conditions to conduct such extensive measurements in parallel. After careful exploratory analysis (see exemplary figures in the response to reviewer 2), we are confident that the approach of using three comparably sunny days for the analysis of spatial heterogeneity of transpiration is suitable to eliminate additional variability induced by varying weather conditions.

34  
35

Regarding the low temporal variability of oil palm water use, we do not refer to the analysis of spatial variability among stands on three sunny days, but rather to the low day-to-day variability of oil palm transpiration in all examined stands, which is presented for four stands in this manuscript. We have tried to make this clearer throughout the manuscript.

36  
37  
38

**Referee:** Secondly, to come up with such a conclusion it is not sufficient to evaluate three sunny, dry days. It would require a more sophisticated evaluation of the entire three weeks under contrasting weather conditions and the three plots (BO3, PA, PTPN6) that were monitored over longer periods in parallel. With regard to this aspect it would be very helpful if the authors could present some selected 3-week time series of transpiration.

39  
40  
41

**Authors:** Figure 5 and the according sections in the results/discussion show, that our statement of low temporal variability of oil palm transpiration is not merely based on the analysis of three sunny days, but

42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63

64 rather time series of at least 3 weeks in each stand. In this manuscript, in Figure 5, four such series are  
65 presented and plotted against radiation and VPD, respectively. Both relationships show that water use  
66 seems to ‘level-off’ at relatively low VPD and radiation, respectively, i.e. after a steep initial increase,  
67 further increases in VPD and radiation do not induce substantial increases in water use rates; this lead us  
68 to conclude that the transpirational behavior of oil palms is rather ‘buffered’ to fluctuating environmental  
69 conditions, e.g. in contrast to some of the mentioned studies on other species. We tried to clarify our line  
70 of argument throughout the results and discussion.

71  
72  
73

74 **Referee:** Another point that was somewhat disappointing for me as a reader is that the authors announced  
75 that their study will “shed first light on some of the hydrological consequences of the continuing  
76 expansion of oil palm plantations”. Unfortunately, this very interesting aspect is not lighted at all, and it  
77 would strengthen the manuscript if the authors would add one or two paragraphs in the Discussion about  
78 this issue.

79

80 **Authors:** We agree with the reviewer that the manuscript previously under-delivered on this, and we tried  
81 to work out the main conclusions to be drawn from our study more clearly throughout the discussion and  
82 conclusions, i.e. relatively high (evapo)transpiration from oil palms and rather low day-to-day variability  
83 of transpiration rates.

84  
85  
86

#### 87 **Specific comments**

88

89 **Referee:** p. 9209: The title does not clearly reflect the content of the paper. The title does not reflect the  
90 aspect of micro-meteorological drivers, which is a substantial part of the manuscript.

91

92 **Authors:** While we agree that the title does not reflect the influences of micrometeorological drivers, we  
93 believe that the strong focus on plantation age throughout the manuscript justifies our current, relatively  
94 precise and ‘catchy’ title. After careful consideration, we thus decided to keep the original title.

95  
96  
97

98 **Referee:** p. 9216, line 10: Please add some additional information how the eddy covariance data were  
99 processed. Did you gap fill the data? If yes, how did you do that? Did you use quality flags to filter the  
100 data or did you use all data? What’s about the energy balance closure of the EC flux data. It would help to  
101 assess the quality of the EC flux data if the authors could add some data about the energy balance closure.  
102 Did you apply any method to post-close the energy balance (e.g. Bowen ratio method) or did you use the  
103 raw latent heat flux data?

104

105 **Authors:** We added further information to the method section on eddy covariance measurements.  
106 Generally, no method was applied to post-close the energy balance. Possible methods would be the WPL  
107 correction, as suggested by Liu et al. (2006), or the suggested Bowen ratio method. The first one is a  
108 correct assumption in the case that the energy balance closure is based on an incorrect determination of  
109 the fluxes by the EC method, but this is not always the reason for the missing energy, so we pRefereed  
110 not to use it. The second method might be too simple in some cases, since it is unknown whether scalar  
111 similarity can be assumed for the processes that cause an underestimation of the EC flux under the  
112 assumption that the scalar similarity is fulfilled. Our analysis of sensible and latent heat flux in both sites  
113 showed no similarity between both of them. Therefore we decided not to apply any method to post-close

114 the energy balance (see Ch4. Corrections and Data Quality Control, in Aubinet et al., 2012, Eddy  
115 Covariance, a practical guide to Measurement and Data Analysis SPRINGER ATMOSPHERIC  
116 SCIENCES 2012, DOI: 10.1007/978-94-007-2351-1).

117 **Markup document (page 7):**

118 The eddy covariance technique (Baldocchi, 2003) was used to measure evapotranspiration (ET, mm  
119 day<sup>-1</sup>) in two of the 15 oil palm stands, the 2-year-old (PA) and the 12-year-old (PTPN6) stand (Table  
120 1). Towers of 7 m and 22 m in height, respectively, were equipped with a sonic anemometer (Metek  
121 uSonic-3 Scientific, Elmshorn, Germany) to measure the three components of the wind vector, and an  
122 open path carbon dioxide and water analyzer (Li-7500A, Licor Inc., Lincoln, USA) to derive  
123 evapotranspiration rates (Meijide et al., in preparation). Fluxes were calculated with the software EddyPro  
124 (Licor Inc), planar-fit coordinate rotated, corrected for air density fluctuation and quality controlled.  
125 Thirty-minute flux data were flagged for quality applying the steady state and integral turbulence  
126 characteristic tests (Mauder and Foken, 2006). Data were also filtered according to friction velocity to  
127 avoid the possible underestimation of fluxes in stable atmospheric conditions. Due to the amount of data  
128 gaps created by lack of power and instrument failure, in the two year-old plantation we calculated the  
129 energy balance closure for the selected three sunny days included in the analysis (see Table 1), for which  
130 it was 82%. In the 12 year-old stand, the energy balance closure for the respective full measurement  
131 period (May 2014-February 2015) was 84%. Data used for this analysis were not gap-filled. We selected  
132 three sunny days when most of the thirty-minute measurements during the day were available. When a  
133 single thirty-minute value was missing, the value was filled by linear interpolation between the previous  
134 and the next 30 min value. Measurements were conducted between July 2013 and February 2014 in the 2-  
135 year old and from May 2014 to February 2015 in the 12-year old stand. For the analysis, we used the  
136 average of the same three sunny days that were selected for the sap flux analysis in the respective plots  
137 (see Table 1). Daytime (6am–7pm) evapotranspiration rates were used for the analyses and comparison to  
138 transpiration rates in order to avoid possible measurement errors as a consequence of low turbulent  
139 conditions during nighttime hours.

140  
141  
142

143 *Referee:* p. 9220, line 5: Please introduce the Hill function or give at least a reference to this  
144 function.

145  
146 *Authors:* We provide a reference to the Hill function in the according section.

147  
148 **Markup document (page 9):**

149 Converted to leaf water use, a clear non-linear trend over stand age became apparent ( $R^2_{adj} = 0.61$ ,  $P <$   
150  $0.01$  for the Hill function, see Morgan et al., 1975, fit shown in Appendix Fig. 1b, not shown in Fig. 3b):

151  
152  
153

154 *Referee:* p. 9220, line 16-17: “There was no significant relationship between water use and radiation”  
155 Firstly, this finding is very surprising, because evapotranspiration must be a function of radiation, and

156 secondly this statement contradicts the results that the authors show in Fig. 5b. There, the authors found,  
157 at least for the sites BO3, PTPN6 and HAR\_old, a pronounced linear relationship between leaf water use  
158 and radiation. Please explain!

159  
160 *Authors:* The wording was imprecise here, we did not refer to a general relationship between radiation  
161 and water use, but to the particular relationship between transpiration (on the respective three sunny days)  
162 and the radiation values (on these respective three sunny days), i.e. transpiration differences among sites  
163 could not be explained by differences in radiation during the respective time of measurement. We  
164 adjusted the wording in the according section.

165  
166 **Markup document (page 10):**

167 Potentially, this could be related to differences in radiation on the respective three sunny days that were  
168 chosen for the analysis. However, there was no significant relationship between average water use rates  
169 on the respective three sunny days in the 15 stands and the respective average radiation (or VPD) on those  
170 days (linear regression,  $P > 0.05$ ), i.e. observed spatial variability in transpiration among the 15 stands  
171 could not be explained by differences in weather conditions. A further analysis of the water use rates of  
172 eight medium-aged stands with highly variable transpiration rates also gave no indications of variability  
173 being induced by differences in radiation.

174  
175  
176  
177 *Referee:* p. 9220, line 23-26: I do not agree with the argumentation that the dynamics of leaf water use is  
178 buffered. I think it would help a lot if the authors would discuss their result more in the light of plant  
179 physiological aspect (e.g. light and temperature response curve, stomatal conductance, photosynthesis  
180 etc.). If the light response curve, for example, reaches already at low radiation its maximum than any  
181 further increase in radiation would not increase transpiration but this does not mean that the response of  
182 the water use is buffered.

183  
184 *Authors:* We agree that the use of the word ‘buffered’ might have originally been misleading and have  
185 adjusted the respective section by elaborating further and partly rephrasing.  
186 While we agree that a discussion involving further plant physiological aspects would be highly  
187 interesting, unfortunately the available data basis on oil palm physiology is at this point insufficient to do  
188 so comprehensively. Such issues will certainly have to be addressed in further studies on the water use  
189 characteristics of oil palm.

190  
191 **Markup document (page 18):**

192 At the day-to-day scale, in all 15 oil palm stands, the response of water use rates particularly to changes in  
193 VPD seemed ‘buffered’, i.e. near-maximum daily water use rates were reached at relatively low VPD, but  
194 better environmental conditions for transpiration (i.e. higher VPD) did not induce strong increases in  
195 water use rates (i.e. 1.2-fold increase in water use for a two-fold increase in VPD). Likewise, for both  
196 photosynthesis rates (Dufrene and Saugier, 1993) and water use rates (Niu et al., 2015) of oil palm leaves,  
197 linear increases with increasing VPD were reported at relatively low VPD, until a certain threshold  
198 (1.5–1.8 kPa) was reached, after which no further increases in photosynthesis and water use rates,  
199 respectively, occurred

200

201  
202  
203  
204  
205  
206

**Referee:** Chapter 3.3: Why did you limit your analysis of the environmental drivers to VPD and radiation? Evapotranspiration also depends heavily on wind speed, temperature and atmospheric stability. Did you have also a look on these drivers? Please explain and discuss it in the text!

207  
208  
209  
210  
211  
212  
213  
214

**Authors:** We had recorded a variety of further environmental and micrometeorological parameters (e.g. soil moisture and temperature, air temperature and humidity, air pressure, wind speed, net radiation) and did not limit our analysis to (global) radiation and VPD, but none of the other variables had any significant relationship with water use ( $P > 0.05$  for linear, non-linear and multiple linear regressions), or they had a similar, but weaker relationship as the presented drivers (as e.g. the case for net radiation and global radiation), and we thus did not present them in the manuscript. We included this information in the environmental measurements section of the Methods to make clearer why we focus on VPD and radiation exclusively in this manuscript.

215  
216

**Markup document (page 8):**

217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229

Soil moisture fluctuated only little at the respective locations and during the respective measurement periods and even on a yearly scale, e.g. between  $32 \pm 2\%$  and  $38 \pm 2\%$  between June 2013 and June 2014 (minimum and maximum daily values, mean  $\pm$  SE between the three micrometeorological stations). Soil moisture did e.g. also not fall below 36% during the measurement period in the long-term monitoring (BO3) stand. It was non-limiting for plant water use. As it showed no significant relationship with water use rates, we omitted soil moisture from further analyses of influences of fluctuations in environmental variables on oil palm water use. Likewise, further recorded micrometeorological variables (e.g. air pressure, wind speed) had no significant relationship with water use rates in our study (linear regression,  $P > 0.1$ ) and were thus also omitted. We instead focused on the micrometeorological drivers VPD and global radiation; among an array of micrometeorological variables (e.g. also including temperature, humidity, net radiation) exploratory analysis had shown that they were best suited to explain fluctuations in water use rates. This has also been demonstrated in other studies on plant water use (e.g. Dierick and Hölscher, 2009; Köhler et al., 2009, 2013)

230  
231  
232  
233  
234  
235

**Referee:** p. 9222, line 14-26: This is a Result part, and please describe in the Material and Methods which statistical method you applied to get these numbers.

236  
237  
238  
239

**Authors:** As suggested by the reviewer, we moved the according section to the results and now merely provide a quick summary of these results in the discussion. We included information on the statistical procedure (providing function type, i.e. Hill function, as well as  $R^2$  values, i.e. the percentage of variability that can be explained by the fit) directly into the section.

240  
241

**Markup document (page 10/13):**

242  
243  
244  
245  
246

**Results:** On comparably sunny days, the stand-level transpiration among the 15 oil palm stands varied 12-fold, from 0.2 mm day<sup>-1</sup> in a 2-year old to 2.5 mm day<sup>-1</sup> in a 12-year old stand. A large part of this spatial variability was explained by different stand variables when applying the Hill function. Stand age explained 45% of the observed spatial variability of stand transpiration (i.e.  $R^2_{adj} = 0.45$  at  $P < 0.01$ , Appendix Fig. 1), and variables correlated to stand age, i.e. by average stand trunk height and by stand

247 water conductive area, explained 44% and 43%, respectively (Table 2). Much of the remaining variability  
248 in stand transpiration rates could be explained by varying stand densities (variations of up to 30%  
249 between stands of similar age, see Table 1). Thus, when shifting from the stand level to the palm level, up  
250 to 60% of the spatial variability in palm water use rates could be explained by age and correlated  
251 variables (see Fig. 3c and Table 2). Much of the variability that remains on the palm level is induced by  
252 three stands where palm water use was much higher ( $> 150 \text{ kg day}^{-1}$ ) than in the other 12 stands ( $< 125$   
253  $\text{kg day}^{-1}$ ); excluding these three stands from the analysis, 87% of the spatial variability in palm water use  
254 rates could be explained by age (Table 3).

255  
256 **Discussion:** The observed substantial stand-to-stand variability of transpiration among the 15 stands,  
257 particularly among medium aged plantations, could to 60% be explained by the variables stand age and  
258 density, and up to 87% when excluding three stands with much higher water use. The remaining  
259 unexplained variability as well as the high water use rates in the three mentioned stands could be related  
260 to differences in site and soil characteristics.

261  
262  
263  
264 **Referee:** p. 9223, line 9-15: Please avoid to repeat too many results in the Discussion. Pick up shortly the  
265 main finding and then discuss it.

266  
267 **Authors:** We followed the advice of the reviewer and shortened parts of the discussion that repeated  
268 results in too much detail.

269  
270 **Markup document (page 14):**

271 Our eddy-covariance derived evapotranspiration estimates of 2.8 and 4.7 mm day<sup>-1</sup> (on sunny days, in 2-  
272 and 12-year old stands, respectively) compare very well to the range reported for oil palms in other  
273 studies: For 3–4 year old stands in Malaysia, eddy-covariance derived values of 1.3 mm day<sup>-1</sup> and  
274 3.3–3.6 mm day<sup>-1</sup> were reported for the dry and rainy season, respectively (Henson and Harun, 2005).  
275 For mature stands, a value of 3.8 mm day<sup>-1</sup> was given, derived by the same technique (Henson, 1999).  
276 Micrometeorologically-derived values for 4–5 year old stands in Peninsular India were 2.0–5.5 mm day<sup>-1</sup>  
277 during the dry season (Kallarackal et al., 2004). A catchment-based approach suggested values of 3.3–3.6  
278 mm day<sup>-1</sup> for stands in Malaysia between 2 and 9 years old (Yusop et al., 2008); evapotranspiration rates  
279 derived from the Penman-Monteith equation and published data for various stands were 1.3–2.5 mm  
280 day<sup>-1</sup> in the dry season and 3.3–6.5 mm day<sup>-1</sup> in the rainy season (Radersma and Ridder, 1996). The  
281 values reported in most available studies as well as our values overlap in a corridor from about 3 mm  
282 day<sup>-1</sup> to about 5 mm day<sup>-1</sup>; this range compares to evapotranspiration rates reported for rainforests in  
283 South East Asia (e.g. Tani et al., 2003a; Kumagai et al., 2005). Considering that oil palm stands e.g. have  
284 much lower stand densities and biomass per hectare than natural tropical forests (Kotowska et al., 2015),  
285 this indicates a quite high evapotranspiration from oil palms at both the individual and the stand level.

286  
287  
288  
289 **Referee:** p. 9228: The Conclusions section is in large parts a summary and not a conclusion. Please revise  
290 it and put the focus on your conclusions.

291  
292 **Authors:** We tried to sharpen the conclusions with respect to a stronger focus on the eco-hydrological  
293 implications of the results of our study.

294  
295 **Markup document (page 19):**



296 The study provides first insights into eco-hydrological characteristics of oil palms at varying spatial and  
297 temporal scales and first estimates of oil palm stand transpiration rates across an age gradient. Stand  
298 transpiration rates increased almost 8-fold from an age of two years to a stand age of five years and then  
299 remained constant with further increasing age, but were highly variable among medium-aged plantations.  
300 In some of the studied stands, transpiration was quite high, i.e. higher than values reported for tropical  
301 rainforests. There may be a potential trade-off between water use and management intensity of oil palm  
302 plantations. Total evapotranspirational water fluxes from a two and a 12 year-old oil palm plantation were  
303 also relatively high, i.e. other water fluxes besides transpiration (e.g. from the soil) contributed  
304 substantially and variably to evapotranspiration. This reduced a 12-fold difference in transpiration  
305 between the two stands to a less than two-fold difference in evapotranspiration. In the diurnal course,  
306 most oil palms showed a strong hysteresis between water use and VPD. On the day-to-day basis this  
307 results in a relatively low variability of oil palm water use regardless of fluctuations in VPD and radiation.  
308 In conclusion, oil palm dominated landscapes show some spatial variations in (evapo)transpiration rates,  
309 e.g. due to varying age-structures and stand densities, but the day-to-day variability of oil palm  
310 transpiration is rather low. Under certain site or management conditions, (evapo)transpirational water  
311 fluxes from oil palms can be substantial.

312  
313  
314

315 **Referee:** Figure 3: Please plot the Hill function. That helps to assess the quality of the fit.

316

317 **Authors:** We did not include the Hill function into Figure 3, but now provide an additional figure in the  
318 Appendix that shows that Hill fit for the respective sub-figures.

319

320 **Markup document:** Attached as pdf.

321

322

323

324 **Referee:** Figure 5: It would facilitate the interpretation of the figure if the authors would add the slope of  
325 the regression to the plots.

326

327 **Authors:** We now provide the regression functions in the figure.

328

329 **Markup document:** Attached as pdf.

330

331

332 Technical corrections

333

334 **Referee:** p. 9214, line 17: Please state the manufacturer and give some more information about the probe  
335 type.

336

337 **Authors:** We included manufacturer and a reference for the technical specifications of the sensors.

338

339 **Markup document (page 5):**

340

341 Following a methodological approach for sap flux measurements on oil palms (Niu et al., 2015), we  
342 installed thermal dissipation probe (TDP, Granier, 1985; Uniwerkstätten Universität Kassel, Germany;  
see Niu et al. 2015 for technical specifications) sensors in the leaf petioles of 16 leaves, four each on four



343 different palms, for each of the 15 examined stands. Insulative materials and aluminum foil shielded the  
344 sensors to minimize temperature gradients and reflect radiation.  
345  
346

347 **RESPONSE TO REFEREE 2:**

348

349 Dear Referee,

350

351 We appreciate your careful reading of our manuscript and the numerous insightful suggestions. Changes  
352 to the manuscript detailed below refer to the "markup copy" which is attached as a pdf to this comment.

353 We also attached a clear copy of the manuscript as well as all figures.

354

355 Sincerely,

356 Alexander Röhl

357

358

359

360 **General comments**

361

362 *Referee:* This study presents a study on the transpiration rates in palm oil stands of different ages. With  
363 palm oil plants becoming more and more an important feature of the tropical landscape, and data on  
364 transpiration rates of these sites being rare, I think this manuscript is an important contribution of results  
365 to the scientific community researching tropical landscapes and tropical ecosystem functioning. What is  
366 impressive about this study is the inclusion of 15 different field sites, as well as combining two different  
367 methods for measuring (evapo)transpiration rates. By including this many sites, they were able to show at  
368 what stand age transpiration does not increase anymore. Overall I think this is a well described and  
369 comprehensive study that provides valuable information to the community studying palm oil plant  
370 functioning. There are a few weaknesses to this study as well: the (eddy flux) measurements were not  
371 carried out in parallel, so we will have to assume both periods are comparable (authors could add a table  
372 for example with the meteorological data per site per measuring period). Furthermore, I think including  
373 only 4 trees per site in the sap flux measurements is not so much, although the fact that all trees have the  
374 same age in a plant will reduce the variance between trees of a stand. In addition, I think the authors can  
375 emphasize the urgency and importance of their study and research questions more.

376

377 *Authors:* We thank the reviewer for appreciating the high number of replicates in our study, which we  
378 consider to make our study rather unique. However, we agree that there are weaknesses due to varying  
379 measurement periods, mainly caused by difficulties of carrying out simultaneous measurements in the  
380 field in a tropical environment, e.g. regarding financial and technical aspects. We have tried to adequately  
381 cope with this problem in our study.

382 With regards to the relatively low number of replicates per stand (13 leaves in 4 palms), we followed an  
383 oil palm specific measurement scheme (Niu et al. 2015) that suggests relatively precise estimates of oil  
384 palm transpiration (14% sample-size related uncertainty).

385 During the revision, we consistently tried to sharpen the conclusions to be drawn from the results of our  
386 study, as suggested by the reviewer, and we feel that the manuscript now emphasizes the relevance of our  
387 study and research questions.

388

389

390 **Referee:** As for the presentation, I think some parts of the discussion could be written in a way that they  
391 are less of a repetition of the results, and answer to the research objectives more explicitly. Please find my  
392 more detailed comments below.

393  
394 **Authors:** We agree that parts of the discussion were too repetitive, and we have adjusted the manuscript  
395 accordingly. We also tried to work out conclusions more clearly, and to derive a more overarching  
396 message regarding some of the potential stand-scale eco-hydrological consequences of the continuing oil  
397 palm expansion.

398  
399  
400

#### 401 **ABSTRACT:**

402  
403 **Referee:** P 9210 line 21: “Confronting sap flux and eddy-covariance derived water fluxes” I would use a  
404 different word than ‘confronting’.

405  
406 **Authors:** As suggested, we reworded the sentence.

#### 407 408 **Markup document (page 2):**

409 Comparing sap flux and eddy-covariance derived water fluxes suggests that transpiration contributed 8%  
410 to evapotranspiration in the 2-year old stand and 53% in the 12-year old stand, indicating variable and  
411 substantial additional sources of evaporation, e.g. from the soil, the ground vegetation and from trunk  
412 epiphytes

413  
414  
415

416 **Referee:** P 9211 line 4-6: I do not understand this sentence, it’s too vague.

417  
418 **Authors:** We rephrased the sentence and tried to make it clearer.

#### 419 420 **Markup document (page 2):**

421 The stand transpiration of some of the studied oil palm stands was as high or even higher than values  
422 reported for different tropical forests, indicating a high water use of oil palms under yet to be explained  
423 site or management conditions.

424  
425  
426

#### 427 **INTRODUCTION:**

428  
429 **Referee:** P 9212 line 27: Not clear to what “On the other hand” contrasts with. In line 19 you announce  
430 two possibilities: Water use can increase or decrease with age stand, and you start by listing the reasons  
431 for the latter. Then (line 25) you give reasons for expecting no difference, and in line 27 with a reason to  
432 expect differences. It’s better to already mention in line 19 that there are three (increase, no difference,  
433 decrease in transpiration) rather than two different scenarios to expect. As it reads not, the ‘On the other  
434 hand’ in line 27 threw me off as a reader and I had to reread a couple of times.

435  
436 **Authors:** We rephrased several lines in the respective section to separate the different possibilities more  
437 clearly.

438

439 **Markup document (page 3/4):**

440 Water use patterns over a gradient of plantation age to our knowledge have not yet been studied for oil  
441 palms. Water use could increase or decline with increasing stand age or could remain relatively stable  
442 from a certain age. Reasons for declining water use at a certain age include decreasing functionality of  
443 trunk xylem tissue with increasing age due to the absence of secondary growth in monocot species  
444 (Zimmermann, 1973), a variety of other hydraulic limitations (see review of dicot tree studies in Ryan et  
445 al., 2006) and increased hydraulic resistance due to increased pathway length with increasing trunk height  
446 (Yoder et al., 1994). However, for Mexican fan palms (*Washingtonia robusta* Linden ex André H  
447 Wendl.), no evidence of increasing hydraulic limitations with increasing palm height was found  
448 (Renninger et al., 2009). Reasons for potentially increasing water use in older plantations e.g. include  
449 linearly increasing oil palm trunk height with increasing palm age (Henson and Dolmat, 2003). As trunk  
450 height and thus volume increase, internal water storages probably also increase, possibly enabling larger  
451 (i.e. older) oil palms to transpire at higher rates (Goldstein et al., 1998; Madurapperuma et al., 2009).  
452 Additionally, increased stand canopy height is expected to result in an enhanced turbulent energy  
453 exchange with the atmosphere, i.e. a closer coupling of transpiration to environmental drivers, which can  
454 facilitate higher transpiration rates under optimal environmental conditions (Hollinger et al., 1994;  
455 Vanclay, 2009). The mentioned reasons for possibly increasing and decreasing water use with increasing  
456 plantations age, respectively, could also partly outbalance each other, or could be outbalanced by external  
457 factors (e.g. management related), potentially leading to no clear trend of oil palm transpiration over  
458 plantation age.

459  
460  
461

462 *Referee:* P 9213line 15: Although I think objective 2 is interesting, it's not made clear from the discussion  
463 before why we need to know the ratio between evapo-transpiration and transpiration.

464

465 *Authors:* We added a sentence to the first paragraph to highlight why this knowledge is important.

466

467 **Markup document (page 2/3):**

468 Oil palm (*Elaeis guineensis* Jacq.) has become the most rapidly expanding crop in tropical countries over  
469 the past decades, particularly in South East Asia (FAO, 2014). Besides from losses of biodiversity and  
470 associated ecosystem functioning (e.g. Barnes et al., 2014), potentially negative consequences of the  
471 expansion of oil palm cultivation on components of the hydrological cycle have been reported (e.g.  
472 Banabas et al., 2008). Only few studies have dealt with the water use characteristics of oil palms so far  
473 (Comte et al., 2012). Available evapotranspiration estimates derived from micrometeorological or  
474 catchment-based approaches range from 1.3 to 6.5 mm day<sup>-1</sup> for different tropical locations and climatic  
475 conditions (e.g. Radersma and Ridder, 1996; Henson and Harun, 2005). However, various components of  
476 the water cycle under oil palm yet remain to be studied for a convincing hydrological assessment of the  
477 hydrological consequences of oil palm expansion, e.g. regarding the partitioning of the central water flux  
478 of evapotranspiration into transpirational and evaporative fluxes. Also, to our knowledge, influences of  
479 site or stand characteristics on oil palm water use have not yet been addressed.

480  
481  
482

483 *Referee:* P 9213line 21: "It assesses potential hydrological consequences of large-scale oil palm  
484 expansion on main components of the water cycle." Your results and Discussion underdeliver on this, you  
485 do not scale this to landscape scale or discuss the consequences of expansion of oil palm plants for the

486 region. So better not to promise this in the introduction. Alternatively you could re-write the Discussion  
487 so it can incorporate such an assessment.

488  
489 *Authors:* We both adjusted the sentence as not to over-promise and additionally tried to expand parts of  
490 discussion and conclusions with respect to potential hydrological consequences of oil palm expansion as  
491 not to under-deliver.

492  
493 **Markup document (page 4):**

494 It assesses some of the potential hydrological consequences of oil palm expansion on main components of  
495 the water cycle at the stand level.

496  
497  
498

499 **METHODS:**


500

501 *Referee:* P 9215 line 16: Why use three sunny days and not the average of five days? Would that make a  
502 difference and have you tried comparing how important the inclusion of three or five (or four or six)  
503 sunny days is?

504

505 *Authors:* We used the average of three sunny days rather than just one sunny day in order to make the  
506 results less susceptible to e.g. to extreme values or random events. While the reviewer is right that we  
507 could have also used the average of e.g. five sunny days, data series from some of the 15 sites (as well as  
508 from 24 other, non-oil palm sites in the study region, which will be presented in further publications)  
509 were limited and partly encompassed only relatively few sunny days. Exploratory analyses at the  
510 beginning of the data analysis process showed, that absolute values were very similar when using e.g. 3, 5  
511 or 7 sunny days. Even when using the averages of the complete data series (usually about three weeks per  
512 site), the relative differences among the 15 sites were very similar to when using the three sunny day  
513 approach. Based on our analysis, we are confident that three sunny days constitute a sufficient amount.  
514 The first figure below shows the absolute transpiration values of the 15 stands derived from using three  
515 and five sunny days and all available days, respectively. The second figure shows the very close linear  
516 relationship ( $R^2=0.99$ ,  $P<0.01$ ) between the values derived from three and five sunny days, respectively.

517



518  
519  
520  
521  
522  
523  
524  
525  
526  
527

**Referee:** P 9215 line 22: Are the values behind the \_ standard errors or standard deviations? Please indicate with SD or SE.

**Authors:** We now indicate that this is the standard deviation.

**Markup document (page 6):**

528 We chose days with a daily integrated radiation of more than 17 MJ m<sup>-2</sup> day<sup>-1</sup> and an average daytime  
529 VPD of more than 1.1 kPa; respective averages (mean ± SD) of all days included in the analysis were  
530 20.3 ± 2.6 MJ m<sup>-2</sup> day<sup>-1</sup> and 1.6 ± 0.3 kPa (also see Table 1).

531  
532  
533

**Referee:** P 9215 line 27: How was palm height measured?

535

**Authors:** We included how palm height was measured into the respective sentence, as well as a reference to a more detailed description of stand variable measurements.

538

**Markup document (page 6):**

540 For each sample palm, trunk height to the youngest leaf (m) and diameter at breast height (cm) were  
541 measured (see Kotowska et al., 2015 for detailed methodology) and the number of leaves per palm was  
542 counted.

543  
544  
545

**Referee:** P9216 line 21: This reads like a repetition of the sap flux measurements mentioned under part 2.2?

548

**Authors:** We eliminated the repetitive part from this section.

549

550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598

*Referee:* P9216 line 24: Similarly here, it's like you are describing the measurements again, and therefore repeating what you mentioned in the previous paragraph. I would suggest shortening this part and focusing on what's important: The error in both measurements, and why it gives you confidence that the difference will show the contribution of the soil and other vegetation. The description of this measurement now reads as if it was added to the original paragraph in an afterthought.

*Authors:* We eliminated the repetitive part from the section and now focus more exclusively on the potential measurement errors.

**Markup document (page 7):**

To estimate the contribution of stand transpiration to total evapotranspiration, we confronted sap flux derived transpiration rates with eddy covariance derived evapotranspiration rates. As described in Niu et al. (2015), our methodological approach for estimating sap flux is associated with sample size related measurement errors of about 14%. The eddy covariance measurements were carried out in carefully-chosen and well-suited locations and focused on daytime observations only, when estimation uncertainties are commonly low (< 30%, Richardson et al., 2006). The observed differences between evapotranspiration and transpiration estimates presented in this study are thus likely largely due to natural rather than methodological reasons.

**RESULTS:**

*Referee:* P9219 line16: this non-significant relationship is that per site or with all the data from all the sites together? Can you clarify?

*Authors:* It is using the respective 3-sunny-day averages from all sites. We now explain this more clearly in the respective section to separate this analysis (mainly spatial variability) more clearly from the analysis of the temporal (i.e. day-to-day) variability of oil palm transpiration.

**Markup document (page 10):**

However, three medium-aged stands (PTPN6, BO5, and HO2) that showed increased sap flux densities and leaf and palm water use rates also had higher stand transpiration rates, between 2.0 and 2.5 mm day<sup>-1</sup>. Potentially, this could be related to differences in radiation on the respective three sunny days that were chosen for the analysis. However, there was no significant relationship between average water use rates on the respective three sunny days in the 15 stands and the respective average radiation (or VPD) on those days (linear regression,  $P > 0.05$ ), i.e. observed spatial variability in transpiration among the 15 stands could not be explained by differences in weather conditions.

*Referee:* P9219 line22: 'possibly indicate a slight decline'. That sounds quite uncertain.

*Authors:* We have removed the sentence from the section.

**Markup document (page 10):**



599 As for the leaf- and palm-level water use rates, a Hill function explained the relationship between stand  
600 transpiration and stand age ( $R^2_{adj} = 0.45$ ,  $P < 0.01$ ), but the observed scatter was high, particularly among  
601 medium aged plantations.

602  
603  
604 **Referee:** For the rest of paragraph 3.2: a lot of results are given in the text, why not summarize them in a  
605 table or a figure? That would make it easier to refer to later in the Discussion as well.  
606  
607

608 **Authors:** We agree that a summary table is very helpful and added a table summarizing the main results  
609 for all 15 stands (Table 2). It gives an overview of how leaf and palm water use as well as stand  
610 transpiration could be explained by the variables number of plantation age and stand sapwood area; the  
611 table provides results for both the linear fit and using the frequently mentioned Hill function.

612 We added another table (Table 3), which presents the same results as Table 2, but only for 12 of the 15  
613 stands, i.e. excluding the three stand with much higher water use (PTPN6, BO5, and HO2).

614

615 **Markup document:** Tables 2 and 3 on pages 30 and 31

616

617

## 618 **DISCUSSION**

619

620 **Referee:** P9221 line13: I actually don't think the observed range compares that well with the one you  
621 mention from the Acacia plantation. Yes, for the other studies you refer to, but the Acacia plants seem  
622 quite higher on average. They are in the same order of magnitude, but 3.9 mm a day is a lot higher than  
623 2.5 mm a day. So I would leave the Cienciala study out of the list of comparable rates.

624

625 **Authors:** We removed the value of the 'high density' Acacia plantation from the text and adjusted the  
626 passage accordingly.

627

628 **Markup document (page 12):**

629 Among 13 studied productive oil palm stands (i.e. > 4 years old) stand transpiration rates varied more  
630 than two-fold. The observed range (1.1–2.5 mm day<sup>-1</sup>) compares to transpiration rates derived with  
631 similar techniques in a variety of tree-based tropical land-use systems, e.g. an Acacia mangium plantation  
632 on Borneo (2.3mm day<sup>-1</sup> for stands of relatively low density, Cienciala et al., 2000), cacao monocultures  
633 and agroforests with varying shade tree cover on Sulawesi (0.5–2.2 mm day<sup>-1</sup>, Köhler et al., 2009, 2013)  
634 and reforestation and agroforestry stands on the Philippines and in Panama (0.6–2.5 mm day<sup>-1</sup>, Dierick  
635 and Hölscher, 2009; Dierick et al., 2010).

636

637

638

639 **Referee:** P9222 line1-13: This could be explained more explicit and why it is of interest to your research  
640 objectives. Also, you seem to have more replicates in the medium aged group, how do you know if the

641 variability in this group is not a consequence of having more replicates, rather than the sites being more  
642 variable (Would have more replicates in the older and younger stands not have shown a similar variance  
643 in those age categories?)

644  
645 *Authors:* We agree with the reviewer that this could merely be an issue of higher replication in the  
646 medium aged group, and we adjusted the section accordingly as not to over-interpret our results among  
647 the 20-25 year-old studied plantations.  
648

649 **Markup document (page 10):**

650 As for the leaf- and palm-level water use rates, a Hill function explained the relationship between stand  
651 transpiration and stand age ( $R^2_{adj} = 0.45$ ,  $P < 0.01$ ), but the observed scatter was high, particularly among  
652 medium aged plantations.

653  
654  
655  
656 *Referee:* P9223 line 2-7: It would be good to be more explicit in how you think the management would  
657 influence evapo-transpiration or transpiration. What would be the mechanics behind it? Different soil  
658 structures because of higher maintenance intensity? Would fertilized palms open their stomata more?  
659 Also the trade- off could be highlighted more, I think that is actually an interesting part of the results and  
660 discussion.

661  
662 *Authors:* We agree that the relationship between water use and management intensity is highly interesting  
663 and tried to discuss in more detail how they might be interrelated. However, to our knowledge no hard  
664 data is available yet for oil palms, i.e. the character of this discussion remains partly speculative.

665  
666 **Markup document (page 13):**

667 The remaining unexplained variability as well as the high water use rates in the three mentioned stands  
668 could be related to differences in site and soil characteristics. However, all studied stands were located in  
669 comparable landscape positions (i.e. upland sites of little or medium inclination) and on similar mineral  
670 soils, i.e. loam or clay Acrisols of generally comparable characteristics (Allen et al., 2015; Guillaume et  
671 al., 2015). Differences in management intensity could also contribute to the remaining unexplained  
672 variability of stand transpiration rates over age. E.g., on P-deficient soils such as the Acrisols of our study  
673 region (Allen et al., 2015), fertilization can greatly increase oil palm yield (Breure, 1982) and thus total  
674 primary productivity, which could consequently lead to a higher water use of oil palms. Accordingly, the  
675 highest observed transpiration value in our study came from a stand in an intensively and regularly  
676 fertilized, high yielding commercial plantation. Thus, there may be a trade-off between management  
677 intensity, and hence yield, on the one hand, and water use of oil palms on the other hand. This trade-off is  
678 of particular interest in the light of the continuing expansion of oil palm plantations (FAO, 2014) and  
679 increasing reports of water scarcity in oil palm dominated areas (Obidzinski et al., 2012; Larsen et al.,  
680 2014)

681  
682  
683  
684 *Referee:* P9223 line 9-15: You repeat the results first, which is not bad per se, but I think you can write  
685 the point you are trying to make a bit ‘snappier’.

686  
687 *Authors:* We shortened the respective section and tried to make it less repetitive while putting a stronger  
688 focus on the immediate conclusions to be drawn.

689  
690

**Markup document (page 14):**

691 Our eddy-covariance derived evapotranspiration estimates of 2.8 and 4.7 mm day<sup>-1</sup> (on sunny days, in 2-  
692 and 12-year old stands, respectively) compare very well to the range reported for oil palms in other  
693 studies: For 3–4 year old stands in Malaysia, eddy-covariance derived values of 1.3 mm day<sup>-1</sup> and  
694 3.3–3.6 mm day<sup>-1</sup> were reported for the dry and rainy season, respectively (Henson and Harun, 2005).  
695 For mature stands, a value of 3.8 mm day<sup>-1</sup> was given, derived by the same technique (Henson, 1999).  
696 Micrometeorologically-derived values for 4–5 year old stands in Peninsular India were 2.0–5.5 mm day<sup>-1</sup>  
697 during the dry season (Kallarackal et al., 2004). A catchment-based approach suggested values of 3.3–3.6  
698 mm day<sup>-1</sup> for stands in Malaysia between 2 and 9 years old (Yusop et al., 2008); evapotranspiration rates  
699 derived from the Penman-Monteith equation and published data for various stands were 1.3–2.5 mm  
700 day<sup>-1</sup> in the dry season and 3.3–6.5 mm day<sup>-1</sup> in the rainy season (Radersma and Ridder, 1996). The  
701 values reported in most available studies as well as our values overlap in a corridor from about 3 mm  
702 day<sup>-1</sup> to about 5 mm day<sup>-1</sup>; this range compares to evapotranspiration rates reported for rainforests in  
703 South East Asia (e.g. Tani et al., 2003a; Kumagai et al., 2005). Considering that oil palm stands e.g. have  
704 much lower stand densities and biomass per hectare than natural tropical forests (Kotowska et al., 2015),  
705 this indicates a quite high evapotranspiration from oil palms at both the individual and the stand level.  
706 Additionally to the previously discussed relatively high water use of oil palms under certain site or  
707 management conditions, the high evapotranspiration from oil palm can be explained by substantial  
708 additional water fluxes to the atmosphere. These fluxes (i.e. the differences between evapotranspiration  
709 and transpiration estimates) were substantial in both the 2-year old and the 12-year old oil palm stand, i.e.  
710 2.6 and 2.2 mm day<sup>-1</sup>, respectively.

711  
712  
713

714 *Referee:* Overall, I think that the paragraph 4.2 repeats a lot of results and compares them with other  
715 studies without making a clear statement or conclusion. The Discussion, in my opinion, is the place to put  
716 the results in context. What do these results mean how we think of how these sites function in the tropical  
717 landscape? The answer to that question remains quite implicit like this.

718  
719  
720  
721

719 *Authors:* We tried to consider this suggestion of the reviewer and rewrote the section, shortening the  
720 repetitive parts and trying to derive more clear, over-arching conclusions from the presented results of our  
721 study and the discussed other studies.

722  
723

**Markup document (page 16):**

724 Generally, our comparison of eddy-covariance derived evapotranspiration and sap-flux derived  
725 transpiration suggests significant other water fluxes to the atmosphere than transpiration (e.g. from  
726 evaporation) that are still marginal during the morning hours, reach their peak at the time VPD peaks and  
727 are extremely sensitive to decreasing VPD in the afternoon. In our study, transpiration amounted to only  
728 8% and 53% of evapotranspiration in the two year-old and the 12 year-old oil palm stand, respectively,  
729 which is lower than values reported e.g. for mature coconut stands (68%, Rouspard et al., 2006) and  
730 rainforests in Malaysia (81–86%, Tani et al., 2003b). The low relative contribution of palm transpiration  
731 to total evapotranspiration in oil palm stands could be due to relatively high water fluxes from  
732 evaporation, e.g. after rainfall interception. Interception was reported to be substantially higher in oil palm  
733 stands in the study region (28%, Merten et al., in revision) than e.g. in rainforests in Malaysia (12–16%,  
734 Tani et al., 2003b) and Borneo (18%, Dykes, 1997). The high water losses from interception paired with

735 the relatively high water use of oil palms and the consequent high total evapotranspirational fluxes from  
736 oil palm plantations could contribute to reduced water availability at the landscape level in oil palm  
737 dominated areas, e.g. during pronounced dry periods (Merten et al., in revision).

738  
739  
740 **Referee:** P9226 line 27: I don't think the hysteresis is that unusual, and you give the examples before, that  
741 this actually happens in other vegetation types as well. So I would remove the word 'unusual'.  
742

743 **Authors:** We followed the advice of the reviewer and removed the word.

744

745 **Markup document (page 18):**

746 A contribution of stem water storage to transpiration in the morning could be another potential  
747 explanation (Waring and Running, 1978; Waring et al., 1979, Goldstein et al., 1998). It could explain the  
748 early peak followed by a steady decline of transpiration regardless of VPD and radiation patterns, the  
749 decline being the consequence of eventually depleted trunk water storage reservoirs. Other (palm) species  
750 were reported to have substantial internal trunk water storage capacities (e.g. Holbrook and Sinclair,  
751 1992; Madurapperuma et al., 2009), which can contribute to sustain relatively high transpiration rates  
752 despite limiting environmental conditions (e.g. Vanclay, 2009).

753

754

755 **Referee:** P9228line 1-8: This reads as an afterthought to the previous paragraph, better to integrate  
756 it.

757  
758 **Authors:** As suggested, we integrated the mentioned paragraph into the previous one.  
759

760 **Markup document (page 18/19):**

761 At the day-to-day scale, in all 15 oil palm stands, the response of water use rates particularly to changes in  
762 VPD seemed 'buffered', i.e. near-maximum daily water use rates were reached at relatively low VPD, but  
763 better environmental conditions for transpiration (i.e. higher VPD) did not induce strong increases in  
764 water use rates (i.e. 1.2-fold increase in water use for a two-fold increase in VPD). Likewise, for both  
765 photosynthesis rates (Dufrene and Saugier, 1993) and water use rates (Niu et al., 2015) of oil palm leaves,  
766 linear increases with increasing VPD were reported at relatively low VPD, until a certain threshold  
767 (1.5–1.8 kPa) was reached, after which no further increases in photosynthesis and water use rates,  
768 respectively, occurred. For tropical tree and bamboo species, more sensitive responses to fluctuations in  
769 VPD, i.e. 1.4- to 1.7-fold increases and more than two-fold increases, respectively, have been reported  
770 (e.g. Köhler et al., 2009; Dierick et al., 2010, Komatsu et al., 2010). However, a similar 'levelling-off'  
771 effect of water use rates at higher VPD, as observed for the oil palm stands in our study, has been reported  
772 for Moso bamboo stands in Japan (in contrast to coniferous forests in the same region, where water use  
773 had a linear relationship with VPD, Komatsu et al., 2010). The hydraulic limitations 'buffering' the day-  
774 to-day oil palm water use response to VPD are yet to be explained. As soil moisture was non-limiting,

775 they are likely of micrometeorological or eco-physiological nature. The early peaks of water use rates and  
776 the consequent strong hysteresis to VPD on the intra-daily level, which may point to a depletion of  
777 internal trunk water storage reservoirs early in the day as a possible reason for substantially reduced oil  
778 palm water use rates at the time of diurnally optimal environmental conditions, give some first indications  
779 of the direction that further studies could take.

780  
781  
782

783 **Referee:** For paragraph 4.3 I have the same comments as for 4.2 in general. I like how many studies you  
784 compare your results with, but what is your real message, what does this say about these sites that we  
785 need to know? I would recommend rewriting both these paragraphs in a way that this becomes clearer.

786

787 **Authors:** We tried to consider this suggestion of the reviewer and rewrote both sections, trying to derive  
788 over-arching conclusions from the presented results of our study and the discussed other studies rather  
789 than just enumerating the results.

790

791 **Markup document:** see rewritten sections 4.2 and 4.3 on pages 14-19

792