

1 **Seasonal dynamics of carbon and nutrients from two**
2 **contrasting tropical floodplain systems in the Zambezi**
3 **River Basin**
4

5 **A.L. Zuijggeest^{1,2}, R. Zurbrügg^{1,2,a}, N. Blank^{1,2,b}, R. Fulcri^{1,2,c}, D.B. Senn^{1,2,3}, and B.**
6 **Wehrli^{1,2}**

7
8 ¹Institute of Biogeochemistry and Pollutant Dynamics, ETH Zürich, Universitätstrasse 16, CH-8092
9 Zürich, Switzerland

10 ²Eawag: Swiss Federal Institute of Aquatic Science and Technology, Surface Waters – Research and
11 Management, Seestrasse 79, CH-6047 Kastanienbaum, Switzerland

12 ³San Francisco Estuary Institute, 4911 Central Avenue, Richmond, CA 94804, USA
13

14 ^anow at F. Hoffmann-La Roche Ltd, Basel, Switzerland

15 ^bnow at Departement Bau, Verkehr und Umwelt, Aarau, Switzerland

16 ^cnow at Pro Natura Graubünden, Chur, Switzerland
17

18 Correspondence to Alissa Zuijggeest (alissa.zuijggeest@usys.ethz.ch)
19
20

21 **Abstract**

22 Floodplains are important biogeochemical reactors during fluvial transport of carbon and nutrient
23 species towards the oceans. In the tropics and subtropics pronounced rainfall seasonality results in
24 highly dynamic floodplain biogeochemistry. Massive construction of dams, however, has
25 significantly altered the hydrography and chemical characteristics of many (sub)tropical rivers. In this
26 study, we compare organic matter and nutrient biogeochemistry of two large, contrasting floodplains
27 in the Zambezi River Basin in Southern Africa, the Barotse Plains and the Kafue Flats. Both systems
28 are of comparable size, but differ in anthropogenic influence: while the Barotse Plains are still
29 relatively pristine, the Kafue Flats are bordered by two hydropower dams.

30
31 The two systems exhibit different flooding dynamics, with a larger contribution of floodplain-derived
32 water in the Kafue Flats and a stronger peak flow in the Barotse Plains. Distinct seasonal differences
33 have been observed in carbon and nutrient concentrations, loads, and export and retention behavior in
34 both systems. Simultaneous retention of particulate carbon and nitrogen, and net export of dissolved
35 organic and inorganic carbon and nitrogen suggested that degradation of particulate organic matter
36 was the dominant process influencing the river biogeochemistry during the wet season in the Barotse
37 Plains, and during the dry season in the Kafue Flats. Reverse trends during the dry season indicated
38 that primary production was important in the Barotse Plains, whereas the Kafue Flats seemed to have
39 both primary production and respiration occurring during the wet season, potentially occurring
40 spatially separated in the main channel and on the floodplain, respectively.

41
42 Carbon to nitrogen ratios of particulate organic matter showed that soil-derived material was
43 dominant year round in the Barotse Plains, whereas the Kafue Flats transported particulate organic
44 matter that had been produced in the upstream reservoir during the wet season. Stable carbon isotopes
45 suggested that inputs from the floodplain to the particulate organic matter pool varied throughout the
46 year in both systems, in opposite patterns. In the Kafue Flats, encroachment of woody plants since the
47 construction of the dams could be responsible for the altered pattern. Additionally, the timing of
48 runoff-driven inputs during the wet season has been changed by the presence of the dams. This study
49 revealed effects of dam construction on organic matter and nutrient dynamics on the downstream
50 floodplain that only become visible after longer periods, highlighting the need for continued
51 monitoring after dam construction.

52
53
54 Keywords: Zambezi, tropical floodplain, organic matter, nutrients, yields, dam, Barotse Plains, Kafue
55 Flats
56
57

Alissa Zuijggeest 10/9/15 15:25
Deleted: hydropower

Alissa Zuijggeest 10/9/15 15:53
Deleted: - ... [1]

Alissa Zuijggeest 10/9/15 16:24
Deleted: Differences in the biogeochemistry of the two systems that can be linked to the dams are the timing of the runoff-driven dissolved organic carbon and nitrogen pulses in the wet season and the origin and inputs of particulate organic matter.

Alissa Zuijggeest 10/9/15 16:06
Deleted: s

Alissa Zuijggeest 10/9/15 16:06
Deleted: clear

Alissa Zuijggeest 10/9/15 16:21
Deleted: Man-made reservoirs alter the origin of organic matter, and change the timing of precipitation-driven carbon and nitrogen pulses. Environmental assessments of dam impacts should therefore consider changes in water quality.

75 **1. Introduction**

76 In current global budgets of organic matter and nutrients, large rivers (discharge > 400 km³ yr⁻¹)
77 account for approximately 35% of the total freshwater-related flux to the ocean (Milliman and
78 Farnsworth, 2011). In studies presenting global budgets and models for carbon, nitrogen and
79 phosphorus export via large rivers, tropical systems are often underrepresented (Alvarez-Cobelas et
80 al., 2008; Alvarez-Cobelas et al., 2012; Alvarez-Cobelas et al., 2009). Global extrapolations based on
81 the well-studied temperate and boreal systems are therefore prone to large errors because they neglect
82 the distinct seasonal flooding of extensive tropical floodplain areas (Junk, 1999; Junk et al., 1989).
83 Floodplain systems have been recognized for their potential to alter fluxes of particulate matter,
84 organic carbon, and nutrients transported by rivers (McJannet et al., 2012; Fisher and Acreman, 2004).

85
86 | During transport [from land to sea](#), riverine organic matter is modified by processes in biogeochemical
87 reactors, specifically natural and artificial lakes and wetlands or floodplains. In the past decades,
88 increasing energy demands have resulted in the construction of hydropower dams in most of the
89 world's large river systems (Nilsson et al., 2005). These man-made biogeochemical reactors
90 significantly change the characteristics of river flow. Since water flow is restricted in most lakes,
91 particles have time to settle. The water exiting the lake is therefore depleted in particulate matter and
92 its associated organic carbon and nutrients. Lake stratification favors hypoxia or anoxia in the
93 hypolimnion, which can lead to methane outgassing (Barros et al., 2011; Bastviken et al.,
94 2008; DelSontro et al., 2011), low oxygen concentrations and potentially toxic levels of reduced
95 substances in the outflow from turbines (Kunz et al., 2013). In addition to the direct effects of
96 hydropower reservoirs, energy demands often require flow regimes that deviate from the natural
97 tropical situation, changing the hydrography in the downstream reaches of the river (Lu and Siew,
98 2006; Maingi and Marsh, 2002; Rood et al., 2005).

99
100 Floodplains make up a large fraction of all tropical wetlands (which cover about 2.5-3.5% of the
101 Earth's surface), with areas of >10⁶ km² in South America and >10⁵ km² in Africa (Tockner and
102 Stanford, 2002). Because of the large area, the periodic flooding and corresponding changing redox
103 conditions, the high temperatures, and the intense rates of primary production (Junk and Piedade,
104 1993; Robertson et al., 1999; Ward and Stanford, 1995), the impact of tropical floodplains on riverine-
105 transported organic matter and nutrients can be significant (Hamilton et al., 1997). River-floodplain
106 exchange has been identified as a key process for the ecological and biogeochemical functioning in
107 temperate (Hunsinger et al., 2010; Tockner et al., 2010) and tropical systems [e.g. Melack et al.
108 | (2009)]. Lateral exchange [between a river and its floodplain](#) was shown to affect sediment erosion and
109 transport (Dunne et al., 1998), the composition of the particulate matter (Devol et al., 1995), carbon
110 fluxes (Pettit et al., 2011), and nutrient supply (Villar et al., 1998).

111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141

The type of organic matter transported by tropical rivers co-varies with discharge. Both the Tana River in Kenya (Tamooch et al., 2014) and the Sanaga River in Cameroon (Bird et al., 1998) transported mainly organic matter from the degradation of C₄ plants during the wet season, while dry season runoff is characterized by organic matter originating from C₃ plants. Spencer et al. (2010) also showed that the properties of organic carbon transported by a tributary of the Congo River vary during different hydrographic phases, with highest dissolved organic carbon and lignin concentrations during peak flow. In the same river, the particulate organic carbon concentration increased when the discharge increased (Mariotti et al., 1991).

Following the construction of dams, the hydrological and sediment-related changes have been assessed in several systems, including the Tana River in Kenya (Maingi and Marsh, 2002) and the Lower Mekong River in China (Kummu and Varis, 2007;Fu et al., 2008;Lu and Siew, 2006). However, the impacts of these changes in hydrography on the biogeochemistry of tropical floodplain systems have hardly been studied. Considering the importance of floodplains within the catchment of large tropical rivers, changes in flooding and inundation might have pronounced effects on the biogeochemical behavior of floodplains and can have far-reaching consequences for the downstream catchment.

In this study, we assessed the dynamics and export rates of organic matter and nutrients in two large, understudied floodplains in the Zambezi River Basin, the pristine Barotse Plains and the dam-impacted Kafue Flats, during wet and dry season conditions. This comparative analysis aims at identifying effects of damming on floodplain biogeochemistry and builds on previous studies on river-floodplain interactions in the Kafue Flats (Zurbrügg et al., 2012;Wamulume et al., 2011;Zurbrügg et al., 2013). Based on field campaigns from contrasting seasons, we were able to describe seasonal variability in the two systems. We further quantified the changes in the concentration, speciation, origin, and loads of carbon, nitrogen, and phosphorus along the floodplains in order to assess the implications of river damming and an altered hydrological regime on floodplain biogeochemistry.

Alissa Zuijdgeest 7/9/15 12:54
Deleted: depends on their discharge

Alissa Zuijdgeest 28/9/15 09:24
Deleted: C₄-derived

Alissa Zuijdgeest 28/9/15 09:24
Formatted: Subscript

Alissa Zuijdgeest 7/9/15 13:00
Deleted: Decreased sediment supply from previously flooded areas would affect erosion processes along the river corridor and in the coastal zone, and altered nutrient export could affect riparian and coastal ecosystems (Friedl and Wüest, 2002).

Alissa Zuijdgeest 17/9/15 13:10
Deleted: peak and low-flow

151 **2. Study sites**

152

153 <<Figure 1>>

154

155 At 1.4×10^6 km², the Zambezi River Basin is the fourth largest in Africa, and the only major African
156 river draining into the Indian Ocean. Due to its geographic location, the catchment experiences a
157 pronounced wet season during the passage of the Inter Tropical Convergence zone (Dec-Mar) and a
158 dry season (Apr-Nov) during the remainder of the year.

159

160 <<Figure 2>>

161

162 The Barotse Plains are a near-pristine floodplain area in the upstream part of the Zambezi River in the
163 Western part of Zambia (Fig. 1). The hydrography in the Barotse Plains clearly reflects the climatic
164 conditions, with peak flow around April/May and low flow between July and November (Fig. 2). The
165 total inundatable area is estimated at around 7,700 km² (Hughes and Hughes, 1992). The Kafue Flats
166 are located along the Kafue River, one of the largest tributary of the Zambezi River. Upstream of the
167 Flats, the Itezhi-Tezhi dam (ITT, closed in 1978) stores a significant part of the wet-season runoff in
168 order to allow for a continuous operation of the power station at Kafue Gorge (dam closed in 1972)
169 downstream of the Kafue Flats. [Evaporation from the reservoir changes the water level by 780 mm](#)
170 [year⁻¹, according to](#) Beilfuss (2012). The maximum inundated area of the Kafue Flats is slightly
171 smaller than of the Barotse Plains, at 6,000 km² (Hughes and Hughes, 1992). The hydrography of the
172 Kafue Flats has been significantly altered by the presence of the dams (Fig. 2). Over the last decades,
173 peak flow has been reduced (approximately -50%) and base flow has increased (roughly +50%, [Fig.](#)
174 [2](#)). In consequence, timing and extent of inundation in the Kafue Flats have changed (Mumba and
175 Thompson, 2005). The floodplain area has been reduced by 40% due to permanent inundation, a
176 direct result of elevated base flow sustained by the dam operation.

177

178 The vegetation in the Kafue Flats has been described as a gradient, ranging from open water to
179 floodplain grasslands, water meadows, littoral zones, termitaria grasslands, to woodland areas
180 (Ellenbroek, 1987). After the construction of the dams, the area covered by shrubs has increased
181 (Mumba and Thompson, 2005;Blaser, 2013). For the Barotse Plains a detailed overview of the
182 vegetation zones is lacking, but several sources hint to grasslands, combined with Miombo woodland
183 and deciduous forest patches (Zambezi Society, 2000;Timberlake, 2000).

184

185

186 **3. Methods**

187 **3.1 Sampling**

188 Sampling of the main river channel at multiple locations along the floodplains (Fig. 1) was conducted
189 during peak flow, hereafter called the wet season (April/May; Barotse Plains 2009, 2013; Kafue Flats
190 2008, 2009, 2010) and low flow, referred to as the dry season (October; Barotse Plains 2008, 2013;
191 Kafue Flats 2008). Samples were collected in the middle of the well-mixed channel from surface
192 water (50cm, Barotse Plains), or at mid-depth (Kafue Flats) using a peristaltic pump. The similarity of
193 the results from different years (Zurbrügg et al., 2013; Zurbrügg et al., 2012) allowed combination and
194 averaging of the data sets in order to obtain generalized patterns for the two systems and seasons.
195 Discharge in the main channel was measured using a RiverRay ADCP (for Barotse Plains dry season
196 data from the Zambezi River Authority were used).

198 **3.2 Laboratory analyses**

199 Samples for dissolved nutrient concentrations were filtered through 0.45 µm filters. Dissolved
200 inorganic nitrogen (DIN), phosphate, and the sum of nitrate and nitrite were measured by
201 chemoluminescence detection (Antek 9000). Ammonium was determined by standard colorimetric
202 technique. Total dissolved nitrogen and total phosphorus (TDN and TDP) were determined by
203 chemoluminescence detection (Antek 9000) following persulphate oxidation (Solórzano and Sharp,
204 1980; Bronk et al., 2000). Samples for particulate phosphorus concentrations during the wet season
205 were collected onto 0.7 µm GF/F filters (Whatman) and measured using a sequential phosphorus
206 extraction method (SEDEX, Ruttnerberg (1992) as modified by Slomp et al. (1996)). Samples for
207 dissolved organic carbon (DOC) and dissolved inorganic carbon (DIC) were filtered through 0.7 µm
208 GF/F filters (Whatman) and analyzed on a Shimadzu TOC-L (Barotse Plains) or on a Shimadzu 5050
209 TOC analyzer (Kafue Flats, DOC only). DIC concentrations for the Kafue Flats were calculated from
210 pH and alkalinity measurements (data not shown). Stable oxygen isotopes were determined on filtered
211 water samples (0.45 µm nylon filters) using a Picaro L2120-I Cavity Ringdown Spectrometer
212 (Barotse Plains) or a MultiFlow preparation module connected to a continuous flow IRMS (Isoprime,
213 UK; Kafue Flats) and calibrated against in-house standards ranging from 0 to -22.5‰_{VSMOW}. Riverine
214 suspended matter was collected on pre-weighed GF/F filters (Whatman). After freeze-drying of
215 samples, suspended matter concentrations were determined by weight difference. Particulate organic
216 carbon and nitrogen and their stable isotopic compositions (POC, PN, δ¹³C and δ¹⁵N, respectively)
217 were determined using EA-IRMS (Thermo-Fischer MAT 253 or ThermoFinnigan FlashEA 1112
218 coupled to a DeltaV Advantage Continuous-Flow IRMS), and calibrated against in-house standards
219 (δ¹³C: -15 to -30‰_{VPDB}, precision 0.1‰; δ¹⁵N: -1.1 to +32.7‰_{air}, precision 0.2‰).

Alissa Zuijggeest 4/9/15 16:40
Deleted: in

Alissa Zuijggeest 4/9/15 16:40
Deleted: the

Alissa Zuijggeest 4/9/15 16:40
Deleted: areas

Alissa Zuijggeest 4/9/15 17:02
Deleted:

Alissa Zuijggeest 28/9/15 09:28
Deleted: (Eijkelkamp)

Alissa Zuijggeest 15/9/15 16:13
Deleted: ;

Alissa Zuijggeest 15/9/15 16:13
Deleted:

230 **4. Results**

231

232 **4.1 Hydrology and oxygen isotopes**

233

234 << Figure 3 >>

235

236 During the wet season, the runoff in the main channel of both floodplains was characterized by a
237 discharge minimum roughly in the middle of both systems (Fig. 3). Located around 100 km and 200-
238 300 km downstream in the Barotse Plains and in the Kafue Flats, respectively, constrictions in the
239 river bed were present, which promoted flooding of the surrounding floodplain area (Zurbrügg et al.,
240 2012). During the dry season, the discharge remained rather constant in both systems with a gain due
241 to tributaries along the Barotse Plain and a slight loss in the dry Kafue Flats. Notice, however, that the
242 peak discharge in the wet season was about four times higher in the Zambezi crossing the Barotse
243 Plains compared to the dammed Kafue River (Fig. 2).

244

245 The intense river-floodplain exchange left a distinct $\delta^{18}\text{O}$ signal in both systems: the rivers lost water
246 to the floodplain during the wet season. Downstream of the stretch where discharge in the river
247 decreased, the $\delta^{18}\text{O}$ of the river water sharply turned towards heavier values (Fig. 3). The overall ^{18}O -
248 enrichment of the channel water between the upstream and downstream sampling locations was more
249 pronounced in the Kafue Flats, during the wet season. In the Barotse Plains, the $\delta^{18}\text{O}$ signal showed an
250 overall shift from -2.0‰ in the upstream part to -0.9‰ at the downstream end. In the Kafue Flats a
251 sharp increase towards heavier values was observed downstream of the channel constriction. To
252 correct for different travel distances along the river stretches, the change in $\delta^{18}\text{O}$ per 100 km of river
253 length was estimated: for the Barotse Plains this enrichment was +0.36‰ and for the Kafue +0.56‰
254 per 100 km. During the dry season, no significant increase in isotopic signal of oxygen was observed
255 in the Barotse Plains, while in the Kafue Flats enrichment occurred at +0.17‰ per 100 km.

256

257

258 **4.2 Concentrations and loads**

259

260 <<Figure 4>>

261

262 For comprehensive comparison of the concentrations of carbon, nitrogen, and phosphorus species
263 along the two floodplains during contrasting seasons, all measurements along the floodplain have
264 been considered, irrespective of spatial trends (Fig. 4). The occurrence of large spatial variations

Alissa Zuijdgeest 7/9/15 13:05
Deleted: at a constriction of the river

Alissa Zuijdgeest 7/9/15 13:06
Deleted: These

Alissa Zuijdgeest 7/9/15 13:06
Deleted: located around 100 km and 200-300 km downstream in the Barotse Plains and in the Kafue Flats, respectively, and

Alissa Zuijdgeest 7/9/15 13:15
Deleted: upstream Zambezi

Alissa Zuijdgeest 15/9/15 16:16
Deleted: T

Alissa Zuijdgeest 15/9/15 16:16
Deleted: ,

Alissa Zuijdgeest 15/9/15 16:15
Deleted: water O-isotopic

Alissa Zuijdgeest 17/9/15 13:12
Deleted: At low flow conditions

275 along the floodplain, or differences between measurement methods between the different years,
276 resulted in larger ranges.

277
278 The dissolved inorganic fraction dominated the total carbon concentration in both seasons and both
279 systems (Fig. 4). Dissolved organic nitrogen (DON) was always the main nitrogen species. In the
280 Barotse Plains particulate phosphorus (PP) was the dominant form during the wet season, while
281 dissolved inorganic phosphorus (DIP) was generally the prevailing species during the dry season.
282 Phosphorus concentrations were largely close to detection limit in both systems, and were therefore
283 excluded from the calculation of loads.

284
285 While both systems exhibited very low inorganic nutrient concentrations during the dry season, the
286 Barotse Plains were substantially lower in organic carbon and nitrogen species concentration
287 compared to the Kafue Flats. Differences between the dry season and wet season C and N
288 concentrations within both systems are statistically significant (paired analysis, p-values <0.05) for all
289 species, except for the Kafue Flats DOC (p = 0.23), DON (0.084) and DIN (0.284). The differences in
290 concentrations between the Barotse Plains and Kafue Flats in similar seasons are significant
291 (hypothesis testing, p-values <0.05) for all species, except PP (wet season, p = 0.121) and DIP (dry
292 season, p=0.053).

293
294 <<Figure 5>> <<Table 1>>

295
296 Loads were calculated from the discharge and concentration data for the respective species, as the
297 water column was well mixed (see Supplementary Information for details). Total carbon and nitrogen
298 loads increased along the Barotse Plains during the wet season, mainly due to larger contribution by
299 the dissolved organic form (Fig. 5). The increase in total carbon load in the Kafue Flats during the wet
300 season was mainly attributed to the dissolved inorganic fraction. The magnitude of the wet season
301 carbon loads leaving the floodplain area is comparable between the two systems (roughly 1500 t C d⁻¹,
302 Fig. 5), while the nitrogen loads in the Barotse Plains were almost twice as high as those in the Kafue
303 Flats (44 t N d⁻¹ and 20 t N d⁻¹) During the dry season the loads decrease slightly.

304
305 Net export was determined as the difference between the load at the downstream end of the floodplain
306 and the load at the upstream end of the floodplain (Table 1). During the wet season, the Barotse Plains
307 were a sink for all particulate phases, while the Kafue Flats acted as a source (Table 1). Both systems
308 were sources of DOC and DIC. Dissolved organic nitrogen was exported from both floodplains, but
309 the Barotse Plains retained the small DIN flux, while the Kafue Flats were a minor source. During the
310 dry season, the Barotse Plains acted as source of particulate matter. For the Kafue Flats this could not
311 be determined due to lack of POC and PN measurements in the downstream stretches of the river.

Alissa Zuijdgeest 1/10/15 12:15

Deleted: have

Alissa Zuijdgeest 28/9/15 09:29

Deleted: a

Alissa Zuijdgeest 28/9/15 09:29

Deleted: a lot

Alissa Zuijdgeest 7/9/15 13:17

Deleted: nutrient

Alissa Zuijdgeest 4/9/15 16:22

Deleted: ; Table 1

Alissa Zuijdgeest 4/9/15 16:25

Moved (insertion) [1]

Alissa Zuijdgeest 4/9/15 16:25

Moved (insertion) [2]

Alissa Zuijdgeest 7/9/15 13:59

Deleted: During the wet season, the Barotse Plains showed markedly different concentrations from the dry season and year-round values in the Kafue Flats. Net export for particulate phases from the Kafue Flats could not be determined due to lack of POC and PN measurements at the downstream end of the floodplain.

Alissa Zuijdgeest 7/9/15 14:00

Deleted: The magnitude of the wet season carbon loads leaving the floodplain area is comparable between the two systems (roughly 1500 t C d⁻¹, Fig. 5), while the nitrogen loads in the Barotse Plains were almost twice as high as those in the Kafue Flats (44 t N d⁻¹ and 20 t N d⁻¹)

Alissa Zuijdgeest 4/9/15 16:26

Moved (insertion) [3]

Alissa Zuijdgeest 7/9/15 13:50

Deleted: , most likely caused by aeolian transport of dry floodplain soil material

334 DOC and DIC were retained by both systems. The Barotse Plains were a minor source of dissolved
335 nitrogen, while the Kafue Flats retained both organic and inorganic nitrogen.

Alissa Zuijdggeest 7/9/15 13:51

Deleted: ,

Alissa Zuijdggeest 7/9/15 13:51

Deleted: potentially converted to particulate phases by primary production.

Alissa Zuijdggeest 7/9/15 13:51

Deleted: The general decrease in C loads during the dry season in the Barotse Plains could probably be explained by the slightly higher discharge at the downstream end of the floodplain, due to lower concentrations flowing in from the Luanginga tributary (no data).

338 4.3 C:N ratios and isotopes

340 <<Figure 6>>

342 The C:N ratios of particulate organic matter (Fig. 6) remained fairly constant along the floodplain in
343 the Barotse Plains and Kafue Flats during the wet season (10.8 ± 0.7 and 7.5 ± 0.7 , respectively
344 (statistically significant difference at 95% confidence intervals)). During the dry season the along-
345 floodplain variability within each of the floodplains was larger (10.3 ± 1.5 and 10.3 ± 1.8 ,
346 respectively), but no significant difference was observed between the two systems. On average,
347 particulate carbon was more depleted in ^{13}C in the Kafue Flats compared to the Barotse Plains during
348 the wet season ($\delta^{13}\text{C} = -28.5 \pm 0.9$ and -26.9 ± 1.1 ‰, respectively (statistically significant $p < 0.05$)).

Alissa Zuijdggeest 2/10/15 11:26

Deleted: ^{vpDB}

Alissa Zuijdggeest 2/10/15 11:26

Deleted: ^{vpDB}

Alissa Zuijdggeest 2/10/15 11:26

Deleted: ^{vpDB}

349 During the dry season mean $\delta^{13}\text{C}$ values were -28.5 ± 1.0 ‰ in the Barotse Plains and -26.5 ± 1.9 ‰ in
350 the Kafue Flats, again a significant difference. The organic matter in the Barotse Plains became more
351 enriched in ^{13}C during the wet season compared to the dry season, while in the Kafue Flats lower $\delta^{13}\text{C}$
352 values were observed during the wet season than during the dry season (both significant, $p < 0.05$).

354 The C:N ratio of the dissolved organic phase was more variable: While the wet season values of 17.5
355 ± 1.9 and 23.7 ± 3.4 were fairly similar for the Barotse Plains and Kafue Flats, respectively, they
356 differed widely during the dry season: 166 ± 20 and 22.7 ± 11.3 . The two systems differed
357 significantly ($p < 0.05$) from each other during a given season.

Alissa Zuijdggeest 28/9/15 09:31

Deleted: quite

359 Paired analysis showed that there was no statistically significant ($p < 0.05$) difference in particulate
360 C:N ratios in the Barotse Plains between contrasting seasons, while there was a difference in
361 dissolved C:N ratios and C-isotopic signals. In the Kafue Flats, there was a significant decrease in
362 particulate C:N ratio from the dry to the wet season (as previously reported in Zurbrügg et al. (2013)).

365 5. Discussion

366 5.1 Hydrology and inundation dynamics

367 The discharge patterns (Fig. 3) showed how the bankfull capacity of the Zambezi and Kafue Rivers
368 varied along the floodplain stretch. In both systems water has moved from the main channel onto the
369 floodplain, roughly 600 and $400 \text{ m}^3 \text{ s}^{-1}$ in the Barotse Plains and the Kafue Flats, respectively. Where

384 the capacity of the channel increases again further downstream, water from the floodplains (and
385 potential tributaries) returned to the main channel at higher rates. On the floodplain, flow velocities
386 were extremely low ($< 1 \text{ mm s}^{-1}$ on the Kafue Flats in May 2008, unpublished data), which led to
387 prolonged residence times of the water on the floodplain, during which evaporation might occur,
388 resulting in heavier $\delta^{18}\text{O}$ signatures in floodplain water.

390 Using a mass balance approach based on oxygen isotopic data, Zurbrügg et al. (2012) calculated that
391 $>80\%$ of the water in the Kafue Flats had spent time on the floodplain during the wet season.
392 Logistical constraints prevented the collection of similar remote floodplain samples in the Barotse
393 Plains. Assuming a similar floodplain signal in the Barotse Plains as in the Kafue Flats, a first
394 approximation was made to determine how much water in the Barotse Plains has spent time on the
395 floodplain. This resulted in 50% of the water leaving the pristine floodplain area having spent time
396 outside the channel. This estimate shows that the interaction between river and floodplain was
397 stronger in the Kafue Flats than in the Barotse Plains, and reinforces the observation that a larger
398 fraction of the river discharge in the Kafue Flats was forced onto the floodplain at the constriction
399 location, compared to the Barotse Plains. In the published literature, high contributions of floodplain-
400 derived water are also reported for the Tonle Sap Lake-floodplain system, where water from the
401 Mekong contributed over 50% to the inflows of the lake, and more than 80% of the outflows from the
402 lake returned to the main river channel of the Mekong (Kummu et al., 2014). At peak flow in the
403 Amazon, 97% of the river inflow occurred at overbank flow at the Curuai floodplain, and this water
404 spent on average 19 days on the floodplain, according to the modeling results by Rudorff et al. (2014).

406 During the dry season, the increasing discharge along the Barotse Plains is most likely caused by
407 inflow of the Luanginga tributary. By contrast, the decreasing discharge in the Kafue Flats combined
408 with a calculated 16% of the downstream discharge having spent time on the floodplain (Zurbrügg et
409 al., 2012) indicated that there was still exchange between the river channel and some permanently
410 inundated areas in the downstream reaches of the Kafue Flats. In a regional perspective, the along-
411 floodplain increase in the $\delta^{18}\text{O}$ signal in the Barotse Plains and Kafue Flats during the wet (flooding)
412 season ($+0.21\%_{\text{VSMOV}}$ to $+0.56\%_{\text{VSMOV}}$ per 100 km) was considerably lower than the increase in the
413 Okavango delta during the dry (flooding) season ($+2.04\%$ per 100 km) and during wet season
414 ($+0.74\%_{\text{VSMOV}}$ per 100 km; calculated from Akoko et al. (2013)), indicating that there was
415 significantly less extensive evaporation on the Zambezi catchment floodplains than in the inland
416 Okavango delta.

5.2 Seasonality of C and N export and retention

419 During the wet season, the Barotse Plains were characterized by a net export of dissolved phases and
420 retention of particulate material. Degradation processes or settling of particulate organic matter, either

Alissa Zuijdgeest 17/9/15 11:07

Deleted: Based on the same approach but without a remote sample of floodplain water

Alissa Zuijdgeest 17/9/15 11:08

Deleted: for the Barotse Plains was made using the evaporative isotope value of floodplain water from the Kafue Flats

Alissa Zuijdgeest 18/9/15 10:33

Deleted: , even though the Kafue flood peak was mitigated by Itezhi-Tezhi dam.

Alissa Zuijdgeest 28/9/15 09:31

Deleted:

Alissa Zuijdgeest 7/9/15 14:01

Deleted: could be explained by the

Alissa Zuijdgeest 7/9/15 14:10

Deleted: floodplain contribution of

Alissa Zuijdgeest 7/9/15 14:01

Deleted: [

Alissa Zuijdgeest 7/9/15 14:03

Deleted: here

Alissa Zuijdgeest 28/9/15 09:33

Deleted: In addition, increasing $\delta^{18}\text{O}$ values towards the lower end of in the Kafue Flats during the dry season could also be a consequence of lower flow velocities, prolonged the travel times and stronger evaporation without any floodplain interaction. At present, the data is not sufficient to discriminate between the two mechanisms, but based on the larger ^{18}O enrichment in the Kafue Flats, we can conclude that more evaporation occurs along the Kafue Flats compared to the Barotse Plains.

Alissa Zuijdgeest 17/9/15 11:36

Deleted: -

Alissa Zuijdgeest 15/9/15 14:16

Deleted: Export

Alissa Zuijdgeest 28/9/15 09:33

Deleted: behavior

448 in the main channel or on the floodplain could result in apparent retention of POC and PN. The
449 concurrent export of DOC, DIC, and DON could similarly be a result of degradation, or of leaching of
450 vegetation or soils. During the dry season, the patterns were reversed, indicative of inputs of organic
451 matter from the Plains.

452
453 In contrast, the Kafue Flats were a net source for both particulate and dissolved phases during the wet
454 season, indicating a different balance. The high proportion of DIC to the net dissolved C export
455 suggests that degradation was a dominant process during flooding. While the constant POC:PN ratios
456 contradict large soil inputs, a combination of primary production around the edges of the main
457 channel, and degradation and leaching of soil and vegetation from the inundated floodplain (indicated
458 by low oxygen concentrations of the water from the floodplain (Zurbrügg et al., 2012)) could be
459 responsible for the observed patterns. During the dry season, the retention of DOC, DIC, DON, and
460 DIN pointed towards primary production potentially a minor contribution from sorption of dissolved
461 organic phases onto particulate material.

462
463 The observed net export of particulate organic matter might not have effects beyond the downstream
464 reservoirs of Lake Kariba and Kafue Gorge, respectively (Fig. 1). Both impoundments will trap
465 mobilized particles, and retain 70% and 90% of incoming total N and P within Lake Kariba (Kunz et
466 al., 2011a). Nevertheless, on a catchment scale, mobilization of particulate organic matter from the
467 inundated area of the river-floodplain systems resulted in specific POC and PON yields (net export
468 per inundated area per year; Table 2) from the Barotse Plains, which were close to an order of
469 magnitude higher than previously reported values for the entire Zambezi River (Beusen et al.,
470 2005; Mayorga et al., 2010). Despite the behavior as a sink during the wet season, the normalization to
471 inundated area has resulted in positive annual export from the floodplain. Also, DOC yields from the
472 Barotse Plains were higher than previously estimated for the Zambezi, but comparable to those
473 measured in the Amazon and Orinoco rivers (Table 2; Beusen et al. (2005); Harrison et al. (2005);
474 Lewis and Saunders (1989)). Similarly, DON yields from the pristine floodplain were similar to
475 values measured in the Amazon and Orinoco (Table 2). The Kafue Flats show negative DOC, DON,
476 DIN yields, i.e. are retaining these species. These negative yields show how floodplains can impact
477 the riverine loads in trends opposite to those observed for the whole catchment. Similarly, the high
478 yields from the Barotse Plains underlined the dominant role of floodplains as biogeochemical reactors
479 in riverine transport of organic matter from land to sea.

480
481 <<Table 2>>
482

Alissa Zuijdggeest 4/9/15 16:25

Moved up [1]:

Alissa Zuijdggeest 4/9/15 16:25

Moved up [2]: Dissolved organic nitrogen was exported from both floodplains, but the Barotse Plains retained the small DIN flux, while the Kafue Flats were a minor source. The magnitude of the wet season carbon loads leaving the floodplain area is comparable between the two systems (roughly 1500 t C d⁻¹, Fig. 5), while the nitrogen loads in the Barotse Plains were almost twice as high as those in the Kafue Flats (44 t N d⁻¹ and 20 t N d⁻¹); here the upstream ITT reservoir removed most of the nitrogen (Kunz et al., 2011b).

Alissa Zuijdggeest 4/9/15 16:26

Moved up [3]: During the dry season, the Barotse Plains acted as source of particulate matter, most likely caused by aeolian transport of dry floodplain soil material. For the Kafue Flats this could not be determined due to lack of POC and PN measurements in the downstream stretches of the river. DOC and DIC were retained by both systems, potentially converted to particulate phases by primary production. The Barotse Plains were a minor source of dissolved nitrogen, while the Kafue Flats retained both organic and inorganic nitrogen. The general decrease in C loads during the dry season in the Barotse Plains could probably be explained by the slightly higher discharge at the downstream end of the floodplain, due to lower concentrations flowing in from the Luanginga tributary (no data).

Alissa Zuijdggeest 4/9/15 16:25

Deleted: During the wet season, the Barotse Plains were a sink for all particulate phases, while the Kafue Flats acted as a source (Table 1). Both systems were sources of DOC and DIC. During the wet season, the Barotse ... [2]

Alissa Zuijdggeest 28/9/15 09:47

Deleted: Note that these yields should be seen as conservative estimates, as they are based on maximum inundatable areas.

Alissa Zuijdggeest 9/9/15 08:36

Deleted: are

Alissa Zuijdggeest 4/9/15 17:33

Deleted: (Beusen et al., 2005; Harrison et al., 2005; Lewis and Saunders, 1989)

Alissa Zuijdggeest 1/10/15 14:45

Moved (insertion) [6]

Alissa Zuijdggeest 1/10/15 14:45

Deleted: Similarly, DON yields from the pristine floodplain resemble values measured in the Amazon and Orinoco.

Alissa Zuijdggeest 1/10/15 14:45

Moved up [6]: The Kafue Flats show negative DOC, DON, DIN yields, i.e. are retaining these species.

Alissa Zuijdggeest 16/9/15 11:31

Deleted: ... [3]

567 **5.3 Sources of organic matter**

568 **5.3.1 Dissolved organic matter**

569 Based on the export and retention behavior of the two floodplains, degradation of floodplain-derived
570 organic matter may be a large source of DOC in the Barotse Plains during the wet season. During the
571 dry season, organic matter inputs from the floodplain and sorption of dissolved organic phases to
572 particles may have decreased the DOC concentrations. In the Kafue Flats, degradation of organic
573 matter on the floodplain was contributing to in-stream DOC during the wet season, whereas during
574 the dry season, similarly to the Barotse Plains, primary production and sorption of dissolved phases
575 onto particles were lowering DOC and DON concentrations. The high contribution of DON to TDN
576 further indicates that the Zambezi and Kafue Rivers are still relatively pristine, as anthropogenic
577 activities mainly add N in the form of DIN to aquatic systems (Berman and Bronk, 2003).

578
579 The elevated C:N ratio of the dissolved organic matter was indicative of terrestrial origin of the
580 organic material in both systems. The ITT reservoir did not have a pronounced impact on the
581 dissolved phase (C:N around 23 during both seasons), which has previously been attributed to a
582 mostly refractory dissolved organic matter phase (Zurbrügg et al., 2013). The comparison with the
583 Barotse Plains revealed a much larger variability in C:N of the dissolved matter reaching dry season
584 values of 166 compared to the wet season signatures around 18. While DOC concentrations were
585 fairly similar during both seasons, the large decrease in DON concentrations from the wet to the dry
586 season (Fig. 4) has resulted in this shift in dissolved C:N ratio.

587
588 The increase in DOC and DON concentrations during the wet season in the Barotse Plains compared
589 to the dry season also corresponds to the general observation that DOC export increases with runoff,
590 caused by shallowing of the flow paths through organic-rich upper soils (Mulholland,
591 2003; Aitkenhead-Peterson et al., 2003). This seasonal variability in DOC and DON concentrations
592 has been previously shown in Hawaii (Wiegner et al., 2009) and Congo (Spencer et al., 2010). Runoff
593 from inundated soils, such as found in the Zambezi River Basin during the wet season, also tend to
594 have higher DON concentrations (Aitkenhead-Peterson et al., 2003). This (potentially refractory)
595 source of DON might be responsible for the high DON concentrations found in the Barotse Plains
596 during the wet season. For the Kafue Flats, there was no significant seasonal change in DOC and
597 DON concentrations between the wet and dry seasons. This might be due to the fact that an increase
598 in DOC and DON concentration in the upstream catchment would be diluted and delayed by the
599 presence of the Itzhi-Tezhi dam, showing after peak flow. With a residence time of 0.7 years, large
600 fractions of organic carbon ($\pm 16\%$) and nutrient loads (50% N, 60% P) were trapped in the sediments
601 of the reservoir (Kunz et al., 2011b). Monthly measurements showed that the highest TOC
602 concentrations occurred in the main channel in the floodplain area in May/June, after the peak flow

Alissa Zuijggeest 16/9/15 11:29
Moved (insertion) [4]

Alissa Zuijggeest 16/9/15 11:31
Moved (insertion) [5]

Alissa Zuijggeest 16/9/15 11:31
Deleted: compared to the particulate material

Alissa Zuijggeest 16/9/15 11:31
Deleted: While the particulate matter in the Kafue Flats was heavily influenced by the presence of t

Alissa Zuijggeest 16/9/15 11:31
Deleted: , the reservoir

Alissa Zuijggeest 28/9/15 09:51
Deleted: p

Alissa Zuijggeest 28/9/15 09:51
Deleted: in this undisturbed system with

Alissa Zuijggeest 17/9/15 13:09
Deleted: can explain

Alissa Zuijggeest 17/9/15 13:10
Deleted: peak flow

Alissa Zuijggeest 9/9/15 09:55
Deleted: - ... [4]

Alissa Zuijggeest 4/9/15 17:03
Deleted: peak and lowest flows

616 (Wamulume et al., 2011). This could be a delayed effect of the increased concentrations at higher
617 runoff during the wet season (November-March).

618

619 5.3.2 Particulate organic matter

620 The higher C:N ratio of the suspended matter in the Barotse Plains year-round indicates a soil-derived
621 source in the pristine part of the catchment. In contrast, C:N ratios found in the Kafue Flats during the
622 wet season were indicative of aquatic production (Zurbrügg et al., 2013). This could be attributed to
623 the presence of the ITT reservoir: surface sediments from the reservoir showed an elevated C:N ratio
624 (12.1±0.6, Supplementary information of Zurbrügg et al. (2013)), similar to the numbers found for the
625 suspended matter in the Barotse Plains. Hence, the presence of the dam significantly affected the
626 chemical composition of the suspended matter, and while soil-derived suspended matter settled in the
627 reservoir, mainly photosynthetically produced organic matter from the reservoir surface waters
628 reached the Kafue Flats and eventually the Kafue-Zambezi confluence. The decrease in C:N ratio
629 along the floodplain in the Kafue Flats during the dry season could be indicative of gradual organic
630 matter input from nitrogen-fixating vegetation. As a consequence of nutrient elimination in the ITT
631 reservoir, widespread encroachment of N-fixing woody plants onto the floodplain has been observed
632 (Blaser, 2013).

633

634 While the C:N ratio showed little variation throughout the year in the Barotse Plains, the stable C-
635 isotopic signatures of the particulate matter further suggest different contributors to the POC in the
636 river. During the wet season, the particulate organic matter in the Barotse Plains is ¹³C enriched
637 compared to the dry season (-26.9 and -28.5‰, respectively). Organic matter sources on the
638 floodplain (soils on average -18‰, abundant reeds between -12 and -27‰, unpublished data) had
639 distinctly heavier δ¹³C signatures than the permanent vegetation in the area (average of 6 different tree
640 species -28.3 ± 1.22 ‰, unpublished data). Inputs from permanent vegetation were the dominant
641 source of organic matter during the dry season, whereas inputs from the floodplain during the wet
642 season led to more enriched values. Shifts to isotopically heavier organic matter during the wet season
643 as observed in the Barotse Plains, have been described for the Tana River in Kenya (Tamooh et al.,
644 2014), the Sanaga River in Cameroon (Bird et al., 1998), and the Congo River in Central Africa
645 (Mariotti et al., 1991). These studies clearly showed how the source of organic matter transported by
646 tropical rivers is changing with inundation.

647

648 In contrast, the particulate organic matter in the Kafue Flats was more enriched during the dry season
649 compared to the wet season (-26.5 and -28.5‰, respectively). The average dry season δ¹³C value for
650 the Kafue Flats should be interpreted with caution, since there is a clear spatial pattern: values become
651 more depleted towards the end of the floodplain. This spatial pattern has previously been attributed to

- Alissa Zuijdggeest 16/9/15 11:29
Deleted: Terrestrial or aquatic sources of p
- Alissa Zuijdggeest 16/9/15 09:48
Deleted: more terrestrial, specifically
- Alissa Zuijdggeest 28/9/15 09:51
Deleted: (mostly)
- Alissa Zuijdggeest 2/10/15 11:20
Deleted: (Zurbrügg et al., 2013),
- Alissa Zuijdggeest 2/10/15 11:20
Deleted: which
- Alissa Zuijdggeest 15/9/15 11:04
Deleted: in
- Alissa Zuijdggeest 28/9/15 09:17
Deleted: OM

660 floodplain-derived particulate organic matter, which would consist of phytoplankton and periphyton
661 material in the permanently inundated area in the downstream reaches of this floodplain (Zurbrügg et
662 al., 2013). In the more typical stretch of the floodplain however, the dry season value was even
663 heavier. The encroaching species have resulted in a vegetation pattern with C₄ species occurring close
664 to the river, and C₃ species growing on the higher grounds that are only seasonally flooded (Blaser,
665 2013;Ellenbroek, 1987). The inputs from these encroaching species can be considered as terrestrial
666 inputs of permanent vegetation.

667
668 The difference in composition and origin between dissolved and particulate phases, i.e. DOM from
669 terrestrial sources, POM more aquatic influence, has previously been described for the Amazon
670 (Aufdenkampe et al., 2007;Hedges et al., 1986) and the Fly-Strickland system in Papua New Guinea
671 (Alin et al., 2008). We showed that the interaction of the river with its floodplain is responsible for the
672 changes observed in organic matter characteristics, but that influence of aquatic production in the
673 systems only originated from the reservoir.

675

Alissa Zuijdgeest 16/9/15 11:31

Moved up [5]: The elevated C:N ratio of the dissolved organic matter compared to the particulate material was indicative of terrestrial origin of the organic material in both systems. While the particulate matter in the Kafue Flats was heavily influenced by the presence of the ITT reservoir, the reservoir did not have a pronounced impact on the dissolved phase (C:N around 23 during both seasons), which has previously been attributed to a mostly refractory dissolved organic matter phase (Zurbrügg et al., 2013). The comparison with the Barotse plains revealed a much larger variability in this undisturbed system with C:N of the dissolved matter reaching dry season values of 166 compared to the wet season signatures around 18. While DOC concentrations were fairly similar during both seasons, the large decrease in DON concentrations from the wet to the dry season (Fig. 4) has resulted in this shift in dissolved C:N ratio.

Alissa Zuijdgeest 17/9/15 11:36

Deleted: -

Alissa Zuijdgeest 28/9/15 09:53

Deleted: (

Alissa Zuijdgeest 28/9/15 09:54

Deleted:)

Alissa Zuijdgeest 16/9/15 12:29

Deleted: The constant C:N ratios observed along the floodplain indicate that aquatic production on the inundated floodplains is of limited importance during peak flow.

Alissa Zuijdgeest 16/9/15 11:29

Moved up [4]: The high contribution of DON to TDN further indicates that the Zambezi and Kafue Rivers are still relatively pristine, as anthropogenic activities mainly add N in the form of DIN to aquatic systems (Berman and Bronk, 2003).

Alissa Zuijdgeest 16/9/15 10:00

Deleted: -

... [5]

Alissa Zuijdgeest 16/9/15 10:00

Deleted: Zurbrügg et al. (2013)

Alissa Zuijdgeest 16/9/15 10:00

Deleted:). In contrast, during the dry season the situation is reversed, and the organic matter in the Kafue Flats shows higher C₄ contribution than in the Barotse Plains. The average dry season $\delta^{13}\text{C}$ value for the Kafue Flats should be treated with caution, since there is a clear spatial pattern: values become more depleted towards the end of the floodplain. This has previously been attributed to floodplain-derived particulate organic matter, which would consist of phytoplankton and periphyton material in the permanently inundated area in the downstream reaches of this floodplain (Zurbrügg et al., 2013). ... [6]

729 **6. Conclusions**

730 << Figure 7 >>

731

732 While the pristine Barotse Plains and dam-impacted Kafue Flats seem to have similar properties in
733 terms of timing and dynamics of seasonal flooding, there are several marked differences between the
734 two systems with respect to hydrology, carbon and nutrient dynamics, and sources of the organic
735 matter (Fig. 7). Based on an oxygen isotope mass balance, a larger fraction of water has spent time on
736 the floodplain at the outflow of the Kafue Flats compared to the Barotse Plains. The two floodplains
737 have significantly different concentrations of dissolved carbon and nutrient species during both wet
738 and dry seasons. Over an annual cycle, the Barotse Plains export more carbon and nutrients
739 normalized to the floodplain area (yields) than previously reported for the Zambezi and other tropical
740 ivers. The Kafue Flats are exhibiting negative yields, effectively retaining and accumulating organic
741 matter and nutrients over a full hydrological cycle. Particulate organic carbon $\delta^{13}\text{C}$ values indicated a
742 larger contribution of floodplain-derived organic matter in the Barotse Plains than in the Kafue Flats
743 during the wet season, and the reversed situation during the dry season. The spatial distribution of C_3
744 and C_4 plants in the floodplains disrupts the signal of floodplain inputs during the wet season in the
745 Kafue Flats.

746

747 Differences between the two systems that can be attributed to the presence of the Itezhi-Tezhi
748 reservoir upstream of the Kafue Flats, included a delay of the input of runoff-derived floodplain soil
749 organic matter and altered inputs to the particulate organic matter pool in the Kafue Flats. The
750 difference between sources of organic matter in the two floodplains partly also results from the
751 presence of the Kafue River dams: since the dam construction, woody encroachment onto the Kafue
752 Flats floodplain has increased, contributing to the wet-season signal of permanent vegetation. This is a
753 result of the presence of the dams that only became evident with time, and shows the importance for
754 monitoring after dam construction.

755

756 **Author contributions**

757 A. L. Zuijggeest, R. Zurbrügg, D. B. Senn, and B. Wehrli were responsible for the study design. A. L.
758 Zuijggeest, R. Zurbrügg, N. Blank, and R. Fulcri performed the fieldwork and the laboratory analyses.
759 Data analysis was performed by A. L. Zuijggeest, R. Zurbrügg, and D. B. Senn, and supported by N.
760 Blank, R. Fulcri, and B. Wehrli. The manuscript was prepared by A. L. Zuijggeest with contributions
761 from all co-authors.

762

Alissa Zuijggeest 10/9/15 16:46

Deleted: : while the Barotse Plains retain particles during the flooding season, in the Kafue Flats the load of particulate phases increases along the floodplain. Retention of particles occurs when flow velocities drop on the floodplain, but due to the very low inputs of particulate matter to the Kafue Flats, any material mobilized will easily contribute to a positive yield. Annually

Alissa Zuijggeest 10/9/15 17:17

Deleted: yields

Alissa Zuijggeest 28/9/15 09:54

Deleted: , but in a basin-wide perspective it should be noted that a significant fraction thereof is trapped in the downstream Kariba reservoir

Alissa Zuijggeest 16/9/15 13:01

Deleted: C_4 plants in the Barotse Plains versus a C_3 -dominated vegetation

Alissa Zuijggeest 16/9/15 13:09

Deleted: se differences in plant source of the organic matter owe to the

Alissa Zuijggeest 10/9/15 17:10

Deleted: The difference between vegetation source in the two floodplains partly also results from the presence of the Kafue River dams: since their construction, woody encroachment onto the Kafue Flats floodplain has increased, contributing to the wet-season C_3 signal. Further d

Alissa Zuijggeest 10/9/15 17:10

Deleted: :

Alissa Zuijggeest 10/9/15 17:11

Deleted: origin of

Alissa Zuijggeest 10/9/15 16:50

Deleted: In the Kafue Flats the C:N ratio of particulate organic matter shifts from a terrestrial to an aquatic signal during the wet season, due to a prolonged retention time in the reservoir (Zurbrügg et al., 2013). In contrast, the C:N ratio of the particulate organic matter in the Barotse Plains indicates that soil organic matter is the dominant source year-round. The concentrations of DOC and DON in the Kafue Flats do not change significantly between dry and wet season. Typically, increased precipitation and overland runoff result in higher concentrations of soil-derived organic matter. However, based on previous year-round measurements (Wamulume et al., 2011), the increase in dissolved organic species is likely delayed by the reservoir, and occurs after peak flow. In the Barotse Plains on the other hand, increased export of DOC and DON during the wet season corresponds to the typical trends of increased export with increased precipitation and runoff. .

... [7]

814 **Acknowledgements**

815 The authors thank Wilma Blaser, Griffin Shanungu, Cristian Teodoru, Jason Wamulume, Mongu
816 harbor personnel, and the Zambia Wildlife Authority for fieldwork assistance. Laboratory analyses
817 were supported by Stewart Bishop, Madalina Jaggi, Daniel Montluçon (ETH Zürich) Kate Ashe,
818 Chantal Freymond, Patrick Kathriner, Gijs Nobbe, Ruth Stierli, Stephan Suter, Prosper Zigah
819 (Eawag), Moritz Lehmann, and Mark Rollog (University of Basel). Comments from Tim Kalvelage
820 (ETH Zürich), [and our two reviewers \(Lex Bouwman and Tim Jennerjahn\)](#) improved the manuscript.
821 Institutional support was provided by Imasiku Nyambe (University of Zambia and its Integrated
822 Water Resource Management Center), the Zambia Wildlife Authority, the Zambezi River Authority,
823 and the Zambia Electricity Supply Corporation. Funding for this study came from the Competence
824 Center for Environment and Sustainability (CCES) of the ETH domain, the Swiss National Science
825 Foundation (Grant No. 128707) and Eawag.

826

827

828 **References**

- 829 Aitkenhead-Peterson, J. A., McDowell, W. H., and Neff, J. C.: 2 - Sources, Production, and
830 Regulation of Allochthonous Dissolved Organic Matter Inputs to Surface Waters, in: *Aquatic*
831 *Ecosystems*, edited by: Findlay, S. E. G., and Sinsabaugh, R. L., Academic Press, Burlington, 25-
832 70, 2003.
- 833 Akoko, E., Atekwana, E., Cruse, A., Molwalefhe, L., and Masamba, W. L.: River-wetland interaction
834 and carbon cycling in a semi-arid riverine system: the Okavango Delta, Botswana,
835 *Biogeochemistry*, 114, 359-380, 10.1007/s10533-012-9817-x, 2013.
- 836 Alin, S. R., Aalto, R., Goni, M. A., Richey, J. E., and Dietrich, W. E.: Biogeochemical
837 characterization of carbon sources in the Strickland and Fly rivers, Papua New Guinea, *J Geophys*
838 *Res-Earth*, 113, F01S05, 10.1029/2006jf000625, 2008.
- 839 Alvarez-Cobelas, M., Angeler, D. G., and Sánchez-Carrillo, S.: Export of nitrogen from catchments:
840 A worldwide analysis, *Environmental Pollution*, 156, 261-269, 10.1016/j.envpol.2008.02.016,
841 2008.
- 842 Alvarez-Cobelas, M., Sánchez-Carrillo, S., Angeler, D. G., and Sánchez-Andrés, R.: Phosphorus
843 export from catchments: a global view, *J N Am Benthol Soc*, 28, 805-820, 10.1899/09-073.1,
844 2009.
- 845 Alvarez-Cobelas, M., Angeler, D. G., Sánchez-Carrillo, S., and Almendros, G.: A worldwide view of
846 organic carbon export from catchments, *Biogeochemistry*, 107, 275-293, 10.1007/s10533-010-
847 9553-z, 2012.
- 848 Aufdenkampe, A. K., Mayorga, E., Hedges, J. I., Llerena, C., Quay, P. D., Gudeman, J., Krusche, A.
849 V., and Richey, J. E.: Organic matter in the Peruvian headwaters of the Amazon: Compositional
850 evolution from the Andes to the lowland Amazon mainstem, *Org Geochem*, 38, 337-364,
851 10.1016/j.orggeochem.2006.06.003, 2007.
- 852 Barros, N., Cole, J. J., Tranvik, L. J., Prairie, Y. T., Bastviken, D., Huszar, V. L. M., del Giorgio, P.,
853 and Roland, F.: Carbon emission from hydroelectric reservoirs linked to reservoir age and latitude,
854 *Nat Geosci*, 4, 593-596, 10.1038/ngeo1211, 2011.
- 855 Bastviken, D., Cole, J. J., Pace, M. L., and Van de Bogert, M. C.: Fates of methane from different
856 lake habitats: Connecting whole-lake budgets and CH₄ emissions, *J Geophys Res*, 113, G02024,
857 10.1029/2007JG000608, 2008.
- 858 Beilfuss, R.: A Risky Climate for Southern African Hydro - Assessing hydrological risks and
859 consequences for Zambezi River Basin dams, Technical Report, International Rivers, Berkeley,
860 California, 2012.
- 861 Berman, T., and Bronk, D. A.: Dissolved organic nitrogen: a dynamic participant in aquatic
862 ecosystems, *Aquat Microb Ecol*, 31, 279-305, 10.3354/ame031279, 2003.
- 863 Beusen, A. H. W., Dekkers, A. L. M., Bouwman, A. F., Ludwig, W., and Harrison, J.: Estimation of
864 global river transport of sediments and associated particulate C, N, and P, *Global Biogeochem Cy*,
865 19, GB4S05, 10.1029/2005gb002453, 2005.
- 866 Bird, M. I., Giresse, P., and Ngos, S.: A seasonal cycle in the carbon-isotope composition of organic
867 carbon in the Sanaga River, Cameroon, *Limnol Oceanogr* 43, 143-146,
868 10.4319/lo.1998.43.1.0143, 1998.
- 869 Blaser, W. J.: Impact of woody encroachment on soil-plant-herbivore interactions in the Kafue Flats
870 floodplain ecosystem [Ph.D. thesis], Ph.D., Swiss Federal Institute of Technology, ETH Zürich,
871 Switzerland, 2013.
- 872 Bouillon, S., Yambélé, A., Gillikin, D. P., Teodoru, C., Darchambeau, F., Lambert, T., and Borges, A.
873 V.: Contrasting biogeochemical characteristics of the Oubangui River and tributaries (Congo River
874 basin), *Sci Rep*, 4, 10.1038/srep05402, 2014.
- 875 Bronk, D. A., Lomas, M. W., Glibert, P. M., Schukert, K. J., and Sanderson, M. P.: Total dissolved
876 nitrogen analysis: comparisons between the persulfate, UV and high temperature oxidation
877 methods, *Mar Chem*, 69, 163-178, 10.1016/S0304-4203(99)00103-6, 2000.
- 878 DelSontro, T., Kunz, M. J., Kempster, T., Wüest, A., Wehrli, B., and Senn, D. B.: Spatial
879 Heterogeneity of Methane Ebullition in a Large Tropical Reservoir, *Environ Sci Technol*, 45,
880 9866-9873, 10.1021/es2005545, 2011.

881 Devol, A. H., Forsberg, B. R., Richey, J. E., and Pimentel, T. P.: Seasonal variation in chemical
882 distributions in the Amazon (Solimões) River: A multiyear time series, *Global Biogeochem Cy*, 9,
883 307-328, 10.1029/95GB01145, 1995.

884 Dunne, T., Mertes, L. A. K., Meade, R. H., Richey, J. E., and Forsberg, B. R.: Exchanges of sediment
885 between the flood plain and channel of the Amazon River in Brazil, *Geol Soc Am Bull*, 110, 450-
886 467, 10.1130/0016-7606(1998)110<0450:EOSBTF>2.3.CO;2, 1998.

887 Ellenbroek, G. A.: Ecology and productivity of an African wetland system: The Kafue Flats [Ph.D.
888 thesis], Junk Publisher, Zambia, 1987.

889 Esser, G., and Kohlmaier, G. H.: Modelling terrestrial sources of nitrogen, phosphorus, sulphur, and
890 organic carbon to rivers, in: *Biogeochemistry of major world rivers*, edited by: Degens, E. T.,
891 Kempe, S., and Richey, J. E., Wiley, 297-322, 1991.

892 Fisher, J., and Acreman, M. C.: Wetland nutrient removal: a review of the evidence, *Hydrol Earth
893 Syst Sc*, 8, 673-685, 10.5194/hess-8-673-2004, 2004.

894 Friedl, G., and Wüest, A.: Disrupting biogeochemical cycles - Consequences of damming, *Aquat Sci*,
895 64, 55-65, 10.1007/s00027-002-8054-0, 2002.

896 Fu, K. D., He, D. M., and Lu, X. X.: Sedimentation in the Manwan reservoir in the Upper Mekong
897 and its downstream impacts, *Quatern Int*, 186, 91-99, 10.1016/j.quaint.2007.09.041, 2008.

898 Hall, A., Valente, I. M. C. B. S., and Davies, B. R.: The Zambezi River in Mozambique: The physico-
899 chemical status of the middle and lower Zambesi prior to closure of the Cabora Bassa Dam,
900 *Freshwater Biol*, 7, 187-206, 1977.

901 Hamilton, S. K., Sippel, S. J., Calheiros, D. F., and Melack, J. M.: An anoxic event and other
902 biogeochemical effects of the Pantanal wetland on the Paraguay River, *Limnol Oceanogr* 42, 257-
903 272, 10.4319/lo.1997.42.2.0257, 1997.

904 Harrison, J. A., Caraco, N., and Seitzinger, S. P.: Global patterns and sources of dissolved organic
905 matter export to the coastal zone: Results from a spatially explicit, global model, *Global
906 Biogeochem Cy*, 19, GB4S04, 10.1029/2005gb002480, 2005.

907 Hedges, J. I., Clark, W. A., Quay, P. D., Richey, J. E., Devol, A. H., and Santos, U. D.: Compositions
908 and fluxes of particulate organic material in the Amazon River, *Limnol Oceanogr* 31, 717-738,
909 1986.

910 Hughes, R. H., and Hughes, J. S.: A directory of African wetlands, UNEP, Nairobi & IUCN, Gland
911 Switzerland/WCMC, Cambridge, 1992.

912 Hunsinger, G. B., Mitra, S., Findlay, S. E. G., and Fischer, D. T.: Wetland-driven shifts in suspended
913 particulate organic matter composition of the Hudson River estuary, New York, *Limnol Oceanogr*
914 55, 1653-1667, 10.4319/lo.2010.55.4.1653, 2010.

915 Junk, W. J., Bayley, P. B., and Sparks, R. E.: The flood pulse concept in river-floodplain systems in:
916 *Proceedings of the International Large River Symposium*, edited by: Dodge, D. P., Canadian
917 Special Publication of Fisheries and Aquat Sci 106, 110-127, 1989.

918 Junk, W. J., and Piedade, M. T. F.: Biomass and primary-production of herbaceous plant-communities
919 in the Amazon floodplain, *Hydrobiologia*, 263, 155-162, 10.1007/BF00006266, 1993.

920 Junk, W. J.: The flood pulse concept of large rivers: learning from the tropics, *Large Rivers*, 11, 261-
921 280, 10.1127/lr/11/1999/261, 1999.

922 Kumm, M., and Varis, O.: Sediment-related impacts due to upstream reservoir trapping, the Lower
923 Mekong River, *Geomorphology*, 85, 275-293, 2007.

924 Kumm, M., Tes, S., Yin, S., Adamson, P., Józsa, J., Koponen, J., Richey, J., and Sarkkula, J.: Water
925 balance analysis for the Tonle Sap Lake-floodplain system, *Hydrol Process*, 28, 1722-1733,
926 10.1002/hyp.9718, 2014.

927 Kunz, M. J., Anselmetti, F. S., Wüest, A., Wehrli, B., Vollenweider, A., Thüring, S., and Senn, D. B.:
928 Sediment accumulation and carbon, nitrogen, and phosphorus deposition in the large tropical
929 reservoir Lake Kariba (Zambia/Zimbabwe), *J Geophys Res*, 116, G03003, 10.1029/2010jg001538,
930 2011a.

931 Kunz, M. J., Wüest, A., Wehrli, B., Landert, J., and Senn, D. B.: Impact of a large tropical reservoir
932 on riverine transport of sediment, carbon, and nutrients to downstream wetlands, *Water Resour
933 Res*, 47, W12531, 10.1029/2011wr010996, 2011b.

- 934 Kunz, M. J., Senn, D. B., Wehrli, B., Mwelwa, E. M., and Wüest, A.: Optimizing turbine withdrawal
935 from a tropical reservoir for improved water quality in downstream wetlands, *Water Resour Res*,
936 49, 5570-5584, 10.1002/wrcr.20358, 2013.
- 937 Lewis, W., Jr., and Saunders, J., III: Concentration and transport of dissolved and suspended
938 substances in the Orinoco River, *Biogeochemistry*, 7, 203-240, 10.1007/BF00004218, 1989.
- 939 Lu, X. X., and Siew, R. Y.: Water discharge and sediment flux changes over the past decades in the
940 Lower Mekong River: possible impact of the Chinese dams, *Hydrol Earth Syst Sc*, 10, 181-195,
941 10.5194/hess-10-181-2006, 2006.
- 942 Maingi, J. K., and Marsh, S. E.: Quantifying hydrologic impacts following dam construction along the
943 Tana River, Kenya, *J Arid Environ*, 50, 53-79, 10.1006/jare.2000.0860, 2002.
- 944 Mariotti, A., Gadel, F., Giresse, P., and Kinga, M.: Carbon isotope composition and geochemistry of
945 particulate organic matter in the Congo River (Central Africa): Application to the study of
946 Quaternary sediments off the mouth of the river, *Chem Geol: Isotope Geoscience section*, 86, 345-
947 357, 10.1016/0168-9622(91)90016-P, 1991.
- 948 Mayorga, E., Seitzinger, S. P., Harrison, J. A., Dumont, E., Beusen, A. H. W., Bouwman, A. F.,
949 Fekete, B. M., Kroeze, C., and Van Drecht, G.: Global Nutrient Export from WaterSheds 2
950 (NEWS 2): Model development and implementation, *Environ Modell Softw*, 25, 837-853,
951 10.1016/j.envsoft.2010.01.007, 2010.
- 952 McJannet, D., Wallace, J., Keen, R., Hawdon, A., and Kemei, J.: The filtering capacity of a tropical
953 riverine wetland: II. Sediment and nutrient balances, *Hydrol Process*, 26, 53-72,
954 10.1002/hyp.8111, 2012.
- 955 Melack, J. M., Novo, E. M. L. M., Forsberg, B. R., Piedade, M. T. F., and Maurice, L.: Floodplain
956 Ecosystem Processes, in: *Amazonia and Global Change*, edited by: Keller, M., Bustamente, M.,
957 Gash, J., and Silva Dias, P., *Geophysical Monograph Series*, American Geophysical Union,
958 Washington, DC, 525-541, 2009.
- 959 Milliman, J. D., and Farnsworth, K. L.: *River Discharge to the Coastal Ocean*, Cambridge University
960 Press, 2011.
- 961 Mulholland, P. J.: 6 - Large-Scale Patterns in Dissolved Organic Carbon Concentration, Flux, and
962 Sources, in: *Aquatic Ecosystems*, edited by: Findlay, S. E. G., and Sinsabaugh, R. L., Academic
963 Press, Burlington, 139-159, 2003.
- 964 Mumba, M., and Thompson, J. R.: Hydrological and ecological impacts of dams on the Kafue Flats
965 floodplain system, southern Zambia, *Phys Chem Earth*, 30, 442-447, 10.1016/j.pce.2005.06.009,
966 2005.
- 967 Nilsson, C., Reidy, C. A., Dynesius, M., and Revenga, C.: Fragmentation and Flow Regulation of the
968 World's Large River Systems, *Science*, 308, 405-408, 10.1126/science.1107887, 2005.
- 969 Pettit, N. E., Bayliss, P., Davies, P. M., Hamilton, S. K., Warfe, D. M., Bunn, S. E., and Douglas, M.
970 M.: Seasonal contrasts in carbon resources and ecological processes on a tropical floodplain,
971 *Freshwater Biol*, 56, 1047-1064, 10.1111/j.1365-2427.2010.02544.x, 2011.
- 972 Robertson, A. I., Bunn, S. E., Walker, K. F., and Boon, P. I.: Sources, sinks and transformations of
973 organic carbon in Australian floodplain rivers, *Mar Freshwater Res*, 50, 813-829,
974 10.1071/MF99112 1999.
- 975 Rood, S. B., Samuelson, G. M., Braatne, J. H., Gourley, C. R., Hughes, F. M. R., and Mahoney, J. M.:
976 Managing river flows to restore floodplain forests, *Front Ecol Environ*, 3, 193-201, 10.1890/1540-
977 9295(2005)003[0193:MRFTRF]2.0.CO;2, 2005.
- 978 Rudorff, C. M., Melack, J. M., and Bates, P. D.: Flooding dynamics on the lower Amazon floodplain:
979 2. Seasonal and interannual hydrological variability, *Water Resour Res*, 50, 635-649,
980 10.1002/2013WR014714, 2014.
- 981 Ruttenberg, K. C.: Development of a sequential extraction method for different forms of phosphorus
982 in marine sediments, *Limnol Oceanogr* 37, 1460-1482, 10.4319/lo.1992.37.7.1460, 1992.
- 983 Slomp, C. P., Epping, E. H. G., Helder, W., and Van Raaphorst, W.: A key role for iron-bound
984 phosphorus in authigenic apatite formation in North Atlantic continental platform sediments, *J Mar*
985 *Res*, 54, 1179-1205, 10.1357/0022240963213745, 1996.
- 986 Solórzano, L., and Sharp, J. H.: Determination of total dissolved nitrogen in natural waters, *Limnol*
987 *Oceanogr* 25, 751-754, 10.4319/lo.1980.25.4.0751, 1980.

988 Spencer, R. G. M., Hernes, P. J., Ruf, R., Baker, A., Dyda, R. Y., Stubbins, A., and Six, J.: Temporal
989 controls on dissolved organic matter and lignin biogeochemistry in a pristine tropical river,
990 Democratic Republic of Congo, *J Geophys Res-Bioge*, 115, G03013, G03013
991 10.1029/2009jg001180, 2010.

992 Tamooch, F., Meysman, F. J. R., Borges, A. V., Marwick, T. R., Van Den Meersche, K., Dehairs, F.,
993 Merckx, R., and Bouillon, S.: Sediment and carbon fluxes along a longitudinal gradient in the
994 lower Tana River (Kenya), *J Geophys Res-Bioge*, 119, 2013JG002358, 10.1002/2013JG002358,
995 2014.

996 Tockner, K., and Stanford, J. A.: Riverine flood plains: present state and future trends, *Environ*
997 *Conserv*, 29, 308-330, 10.1017/s037689290200022x, 2002.

998 Tockner, K., Pusch, M., Borchardt, D., and Lorang, M. S.: Multiple stressors in coupled river-
999 floodplain ecosystems, *Freshwater Biol*, 55, 135-151, 10.1111/j.1365-2427.2009.02371.x, 2010.

1000 Villar, C. A., de Cabo, L., Vaithiyathan, P., and Bonetto, C.: River-floodplain interactions: nutrient
1001 concentrations in the Lower Parana River, *Arch Hydrobiol*, 142, 433-450, 1998.

1002 Wamulume, J., Landert, J., Zurbrügg, R., Nyambe, I., Wehrli, B., and Senn, D. B.: Exploring the
1003 hydrology and biogeochemistry of the dam-impacted Kafue River and Kafue Flats (Zambia), *Phys*
1004 *Chem Earth*, 36, 775-788, 10.1016/j.pce.2011.07.049, 2011.

1005 Ward, J. V., and Stanford, J. A.: The serial discontinuity concept: extending the model to floodplain
1006 rivers, *Regul River*, 10, 159-168, 10.1002/rrr.3450100211, 1995.

1007 Wiegner, T. N., Tubal, R. L., and MacKenzie, R. A.: Bioavailability and export of dissolved organic
1008 matter from a tropical river during base- and stormflow conditions, *Limnol Oceanogr* 54, 1233-
1009 1242, 10.4319/lo.2009.54.4.1233, 2009.

1010 Zambezi Society: Biodiversity of the Zambezi Basin Wetlands: Volume III: Land use change and
1011 human impacts, Zambezi Society and Biodiversity Foundation for Africa, 2000.

1012 Zurbrügg, R., Wamulume, J., Kamanga, R., Wehrli, B., and Senn, D. B.: River-floodplain exchange
1013 and its effects on the fluvial oxygen regime in a large tropical river system (Kafue Flats, Zambia),
1014 *J Geophys Res*, 117, G03008, 10.1029/2011jg001853, 2012.

1015 Zurbrügg, R., Suter, S., Lehmann, M. F., Wehrli, B., and Senn, D. B.: Organic carbon and nitrogen
1016 export from a tropical dam-impacted floodplain system, *Biogeosciences*, 10, 23-38, 10.5194/bg-
1017 10-23-2013, 2013.

1018

1019

Tables

Table 1. Net export (in t C d⁻¹ and t N d⁻¹), calculated as the difference between loads at the downstream and upstream ends of the respective floodplain, from the two floodplains during wet and dry seasons. Positive numbers indicate that the floodplain acted as a source (export), negative numbers indicate the floodplain acting as a sink (retention). POC and PN export from the Kafue Flats during the dry season could not be estimated due to lack of measurements at downstream locations.

System		POC	DOC	DIC	PN	DON	DIN
Barotse	<i>Wet season</i>	-38	170	270	-3.7	12	0.0
Plains	<i>Dry season</i>	14	-1.5	-78	1.6	0.0	0.0
Kafue	<i>Wet season</i>	6.5	160	640	0.6	11	0.1
Flats	<i>Dry season</i>	NA	-11	-89	NA	-0.8	-0.2

Alissa Zuijdgeest 4/9/15 17:00

Deleted: peak and base flow

Alissa Zuijdgeest 4/9/15 16:36

Deleted: Units are t C d⁻¹ and t N d⁻¹.

Table 2. Yields of carbon, nitrogen and phosphorus in kg (C or N) km⁻² yr⁻¹ from large river basins and floodplain yields from the Barotse Plains and Kafue Flats. Yields for this study are calculated assuming 6 months of dry-season export, and 6 months of wet-season export. Inundation areas should be considered conservative estimates (see methods section for how inundation areas were estimated).

Dry-season areas were estimated based on river length and width. POC and PN yields from the Kafue Flats during the dry season could not be estimated due to lack of measurements at downstream locations.

Sources: a Beusen et al. (2005), b Bouillon et al. (2014), c Esser and Kohlmaier (1991), d Harrison et al. (2005), e Lewis and Saunders (1989), f Mayorga et al. (2010), g Hall et al. (1977).

River	POC	DOC	PN	DON	DIN
Amazon	2900 ^a	5200 ^d	500 ^a	330 ^d , 180 ^f	170 ^f
Congo	1400 ^a , 400 ^c	3300 ^a , 1600 ^c	200 ^a	92 ^e , 58 ^f	32 ^f
Orinoco	1500 ^e	5600 ^d , 5200 ^e	190 ^e	310 ^d , 190 ^e , 170 ^f	
Oubangui	180-300 ^b	660-1500 ^b	20-29 ^b		
Zambezi	800 ^a	1000-2000 ^f	100 ^a	-	14 ^f , 100-300 ^g
This study:					
Barotse Plains	8000	3000	880	310	0
Kafue Flats	NA	-2700	NA	-200	-110

Alissa Zuijdgeest 4/9/15 17:01
Deleted: a

Alissa Zuijdgeest 17/9/15 10:20
Comment [1]: Deleted: Mayorga et al. (2005)

Figures




Figure 1. Map of the Zambezi catchment, with floodplains (in green) and large dams (red arrows) marked. Inserts show sampling stations during the dry (circles) and wet season (stars) in the Barotse Plains and Kafue Flats. Sampling stations will be further presented in distance along the river (km),

Alissa Zuijdggeest 15/9/15 16:19

Deleted:





Figure 2. River discharge (a) at Senanga, the downstream boundary of the Barotse Plains (1988-2006 average and standard deviation) and (b) at the outflow of Itezhi-Tezhi (ITT) dam, the upstream boundary of the Kafue Flats. Discharge for the Kafue Flats are means of pre-dam (1960-1971), and post-dam construction (Kafue Gorge dam: 1972-1977; ITT dam: 1978-2010) periods. Data from the Department of Water Affairs and Zambezi River Authority, permission for reprint first granted to Blaser (2013).







Figure 3. Discharge and stable oxygen isotope signals in the Barotse Plains and the Kafue Flats during wet and dry seasons. Discharge and $\delta^{18}\text{O}$ data for the Kafue Flats have been published previously in Zurbrugg et al. (2012).






Figure 4. Boxplots of the concentrations of dissolved and particulate carbon, nitrogen and phosphorus species along the Barotse Plains and the Kafue Flats during wet and dry seasons along the floodplain. The boxes represent the first and third quartiles, and the median. No measurements of particulate phosphorus were made on samples from the dry seasons. Carbon and nitrogen data of the Kafue Flats have been previously published in Zurbrügg et al. (2013).

Alissa Zuijdgeest 7/9/15 09:46

Deleted:
Alissa Zuijdgeest 4/9/15 16:49
Deleted: C





Figure 5. Dissolved and particulate carbon and nitrogen loads along the Barotse Plains and the Kafue Flats during wet and dry seasons. The loads of particulate carbon and nitrogen at the two most downstream locations in the Kafue Flats could not be determined for the dry season due to lack of POC and PN measurements.







Figure 6. Carbon to nitrogen (C:N) ratios of particulate and dissolved organic matter as well as carbon isotopic signatures of particulate organic matter during wet (blue triangles) and dry (orange circles) seasons. The Kafue Flats data have been previously published in Zurbrügg et al. (2013).

Alissa Zuijdgeest 7/9/15 09:46



Deleted:

Alissa Zuijdgeest 28/9/15 09:57

Deleted: black squares

Alissa Zuijdgeest 28/9/15 09:57

Deleted: green




Figure 7. Summary of the organic matter characteristics ($\delta^{13}\text{C}$, POC:PN , DOC:DON), oxygen isotopic enrichment along the floodplain, and proportional arrows for net export and removal rates of POM (POC+PN) and DOM (DOC+DON) in the Barotse Plains and Kafue Flats during the wet and dry season.

Alissa Zuijdgeest 16/9/15 14:45

Barotse Plains wet season

Net removal:
 POC 38.4 tC d⁻¹
 PN 3.7 tN d⁻¹

Net export:
 DOC 169.0 tC d⁻¹
 DIC 271.4 tC d⁻¹
 DON 12.5 tN d⁻¹

$\delta^{13}\text{C}$ -26
 C:N_{POM} 1
 C:N_{DOM} 1
 $\delta^{18}\text{O}$ enrichment +0.36 ‰_{VSMOV}

Kafue Flats wet season

$\delta^{13}\text{C}_{\text{POC}}$ -28.5 ‰_{V-PDB} (C
 C:N_{POM} 7.5
 Net POC
 DOC 1
 DIC 6
 PN
 DON 1
 DIN

Deleted:
 Alissa Zuijdgeest 16/9/15 14:44
 Formatted: Normal
 Alissa Zuijdgeest 16/9/15 14:44
 Deleted: C:N_{POM}
 Alissa Zuijdgeest 16/9/15 14:44
 Deleted: C:N_{DOM}

While the Barotse Plains retain particles during the wet season, annual yields of particulate organic carbon and nitrogen are higher than previously reported for the Zambezi and other tropical rivers. Enhanced wet-season runoff adds soil-derived dissolved organic carbon and nitrogen to the Zambezi River, with a corresponding increase in the Barotse Plains. Soil-derived organic matter dominates the particulate phase year-round in the Barotse Plains, and a varying influence of C₃- and C₄-plant vegetation can be observed throughout the year.

In contrast to the Barotse Plains, net export of particulate matter from the Kafue Flats has been observed during the wet season, but over an annual cycle, the Kafue Flats are effectively accumulating dissolved carbon and nutrients. In the Kafue Flats, the runoff-induced increase in dissolved organic carbon and nitrogen concentrations is delayed by the upstream dam operation. The dam reservoir also causes a shift in the source of the particulate organic matter – from soil-derived during the dry season to aquatically produced in the wet season – in the downstream Kafue Flats. Spatial zonation in vegetation and temporal flooding dynamics in the Kafue Flats result in mostly C₃-derived particulate organic matter during wet season, and a dominance of C₄-derived material during dry season. This pattern results from dam-induced changes in vegetation, as dam construction along the Kafue River has led to encroachment of woody plant species onto the Kafue Flats.

During the wet season, the Barotse Plains were a sink for all particulate phases, while the Kafue Flats acted as a source (Table 1). Both systems were sources of DOC and DIC. During the wet season, the Barotse Plains showed markedly different concentrations from the dry season and year-round values in the Kafue Flats. The upstream catchments of both the Zambezi and the Kafue River exhibit the same Late Precambrian geological formations (Geological Map Zambia, Geological Survey Department), after which the Zambezi flows through the Kalahari sands. Therefore, it seems unlikely that erosional processes in the headwaters result in significantly different concentrations during contrasting seasons, and the low concentration of DIC leaving the Barotse Plains was most likely the result of dilution by the high discharge. Dissolved organic nitrogen was exported from both floodplains, but the Barotse Plains retained the small DIN flux, while the Kafue Flats were a minor source. The magnitude of the wet season carbon loads leaving the floodplain area is comparable between the two systems (roughly 1500 t C d⁻¹, Fig. 5), while the nitrogen loads in the Barotse Plains were almost twice as high as those in the Kafue Flats (44 t

N d^{-1} and 20 t N d^{-1}); here the upstream ITT reservoir removed most of the nitrogen (Kunz et al., 2011b).

During the dry season, the Barotse Plains acted as source of particulate matter, most likely caused by aeolian transport of dry floodplain soil material. For the Kafue Flats this could not be determined due to lack of POC and PN measurements in the downstream stretches of the river. DOC and DIC were retained by both systems, potentially converted to particulate phases by primary production. The Barotse Plains were a minor source of dissolved nitrogen, while the Kafue Flats retained both organic and inorganic nitrogen. The general decrease in C loads during the dry season in the Barotse Plains could probably be explained by the slightly higher discharge at the downstream end of the floodplain, due to lower concentrations flowing in from the Luanginga tributary (no data).

Page 11: [3] Deleted Alissa Zuijdgeest 16/09/15 11:31

For downstream ecosystems the different particle organic matter dynamics of both floodplains is of limited consequence: since the pristine Barotse Plains are located in the upstream part of the catchment, it is likely that a lot of the particulate matter mobilized in the floodplain will end up in the sediments of Lake Kariba and the same is true for the export of the Kafue flats which will be deposited in the Cahorra Bassa reservoir (Fig. 1). Mass balance calculations have shown that 70% and 90% of incoming total N and P, respectively, are removed from the water column within Lake Kariba (Kunz et al., 2011a). Hence, while large amounts of organic matter and nutrients are mobilized from (pristine) floodplains, only a small fraction thereof will eventually reach the coastal ocean due to the presence of downstream lakes and reservoirs.

Page 12: [4] Deleted Alissa Zuijdgeest 09/09/15 09:55

Page 14: [5] Deleted Alissa Zuijdgeest 16/09/15 10:00

5.3.3 Vegetation contributions

The stable C-isotopic signatures of the particulate matter further suggest different POC sources. During the wet season, the organic matter in the Barotse Plains is ^{13}C enriched compared to the

Kafue Flats, indicating a C₄-plant-dominated source in the pristine system ($\delta^{13}\text{C}-\text{C}_3$ approx. -26‰_{VPDB}, vs. $\delta^{13}\text{C}-\text{C}_4$ approx. -13‰_{VPDB}, as in

Page 14: [6] Deleted

Alissa Zuijdgheest

16/09/15 10:00

). In contrast, during the dry season the situation is reversed, and the organic matter in the Kafue Flats shows higher C₄ contribution than in the Barotse Plains. The average dry season $\delta^{13}\text{C}$ value for the Kafue Flats should be treated with caution, since there is a clear spatial pattern: values become more depleted towards the end of the floodplain. This has previously been attributed to floodplain-derived particulate organic matter, which would consist of phytoplankton and periphyton material in the permanently inundated area in the downstream reaches of this floodplain (Zurbrügg et al., 2013).

Shifts to more C₄-derived organic matter during the wet season as observed in the Barotse Plains, have been described for the Tana River in Kenya (Tamooch et al., 2014), the Sanaga River in Cameroon (Bird et al., 1998), and the Congo River in Central Africa (Mariotti et al., 1991). These studies clearly showed how the source of organic matter is changing with inundation.

The contrasting trend in vegetation source in the Kafue Flats compared to other tropical rivers could be explained by the spatial distribution of C₃ and C₄ grasses: most of the C₄ species occur close to the river, while C₃ grasses and plants are found on the higher grounds that are only seasonally flooded (Blaser, 2013; Ellenbroek, 1987). This spatial variation could explain why the wet season organic matter in the Kafue Flats shows a higher contribution of C₃-derived organic matter, whereas during low water conditions only C₄ plants are inundated. Since the construction of the dams in the Kafue catchment, encroachment of woody plants onto the floodplain has been observed (Blaser, 2013). The encroaching species most likely contribute to the C₃ signal observed during the wet season. The regulation of water flow through the Kafue Flats has therefore had an indirect effect on the type of the particulate organic matter transported downstream by the river.

Page 15: [7] Deleted

Alissa Zuijdgheest

10/09/15 16:50

In the Kafue Flats the C:N ratio of particulate organic matter shifts from a terrestrial to an aquatic signal during the wet season, due to a prolonged retention time in the reservoir (Zurbrügg et al., 2013). In contrast, the C:N ratio of the particulate organic matter in the Barotse Plains indicates that soil organic matter is the dominant source year-round. The concentrations of DOC and DON in the Kafue Flats do not change significantly between dry and wet season. Typically, increased precipitation and overland runoff result in higher concentrations of soil-derived organic matter.

However, based on previous year-round measurements (Wamulume et al., 2011), the increase in dissolved organic species is likely delayed by the reservoir, and occurs after peak flow. In the Barotse Plains on the other hand, increased export of DOC and DON during the wet season corresponds to the typical trends of increased export with increased precipitation and runoff.

We have shown that dam construction not only affected the hydrological regime but also the sources and concentrations of OM and nutrients within downstream (floodplain) systems, since the presence of a reservoir altered the origin of the particulate organic matter, and changed the timing of a precipitation-driven pulse of dissolved organic carbon and nitrogen. Therefore, environmental assessments of dams should explicitly analyze the potential changes in water quality and their effects on downstream ecosystems.

Barotse Plains	Distance along	Month/Year sampled	Discharge	δ18O	δ18O stdev	POC	DOC	DIC	PN	DON	DIN	PP	DOP	DIP	POC δ13C
<i>Wet season</i>	the channel (km)		(m3/s)	(‰ VSMOW)		(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(‰ VPDB)
Station number															
BP03	0	04/2013	1977	-1.80	0.10	36	206	316	3.2	11.6	0.00	0.25	0.30	0.06	-26.5
BP05	14.4	04/2013		-1.92	0.03	41	197	298	3.3	10.3	0.00	0.28	0.12	0.05	-26.2
BP06	32.8	04/2013		-2.04	0.07	28	210	487	2.7	11.7	0.00	0.27	0.11	0.05	-27.4
BP07	48.6	04/2013	1957	-2.10	0.05	21	198	526	2.0	11.3	0.00	0.32	0.18	0.05	-27.7
BP11	66.7	04/2013		-2.03	0.06	28	197	481	2.6	11.6	0.00	0.24	0.12	0.04	-27.4
BP12	87.5	04/2013		-2.01	0.06	32	196	472	2.9	11.4	0.00	0.29	0.11	0.03	-26.8
BP13	105.7	04/2013	1317	-1.95	0.04	18	180	417	1.7	10.9	0.00	0.19	0.09	0.03	-26.5
BP16	129.1	04/2013		-1.31	0.04	21	213	339	1.8	11.6	0.00	0.09	0.13	0.02	-25.7
BP17	150	04/2013		-1.37	0.03	20	207	359	2.0	13.0	0.00	0.10	0.20	0.02	-25.6
BP01a	140	05/2009		-3.03			191			23.4	1.58		0.14	0.13	
BP02a	143	05/2009		-3.56			169			23.2	0.66		0.25	0.13	
BP03a	160	05/2009		-3.54			173			26.8	0.83		0.21	0.14	
BP18	176.3	04/2013	1717	-1.66	0.06	11	202	450	1.1	12.4	0.00	0.10	0.13	0.02	-27.2
BP04a	180	05/2009		-3.44			180			22.0	0.45		0.17	0.20	
BP20	204.3	04/2013		-1.36	0.05	7	206	351	0.6	12.4	0.00	0.09	0.13	0.02	-26.6
BP05a	200	05/2009		-3.25			175			26.4	0.50		0.26	0.10	
BP06a	210	05/2009		-3.38			179			23.5	0.47		0.24	0.10	
BP21	233	04/2013		-1.39	0.03	15	202	362	1.5	12.4	0.00	0.10	0.13	0.02	-26.1
BP07a	240	05/2009		-3.43			178			24.1	0.47		0.19	0.14	
BP23	251.1	04/2013	1740	-1.22	0.10	13	200	392	1.2	12.5	0.00	0.09	0.08	0.02	-29.3
BP08a	240	05/2009		-3.43			185			24.5	0.48		0.02	0.28	
BP24	278.3	04/2013		-1.19	0.05	16	296	392	1.6	12.4	0.00	0.11	0.08	0.03	-26.1
BP09a	278.4	05/2009		-3.20			181			22.4	0.32		0.27		
BP25	299.5	04/2013		-0.89	0.07	16	222	401	1.4	13.5	0.00	0.10	0.13	0.04	-29.3
BP26	311.8	04/2013	2413	-0.99	0.05	14	236	367	1.3	13.8	0.00	0.15	0.08	0.05	-26.4

Barotse Plains	Distance along	Year sampled	Discharge	δ18O	δ18O stdev	POC	DOC	DIC	PN	DON	DIN	PP	DOP	DIP	POC δ13C
<i>Dry season</i>	the channel (km)		(m3/s)	(‰ VSMOW)		(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(‰ VPDB)
Station number															
BP37	0	10/2013	203	-2.44	0.05	31	134	1641	3.4	0.7	0.07		-0.05	0.06	-29.6
BP39	119.4	10/2013	227	-2.86	0.06	55		1006	4.6	1.5	0.22		-0.02	0.05	-28.1
BP01a	140	10/2008		-2.71			177				0.38		0.27	0.51	
BP02a	143	10/2008		-3.23			124				0.50		0.06	0.55	
BP03a	160	10/2008		-3.06			115				0.53		0.17	0.43	
BP04a	180	10/2008		-3.06			111				0.55		0.02	0.56	
BP05a	200	10/2008		-3.21			120				0.55		0.14	0.52	
BP06a	210	10/2008		-2.95			112				0.48		0.04	0.54	
BP07a	240	10/2008		-3.14			121				0.87		0.08	0.53	
BP08a	240	10/2008		-3.29			111				0.57		0.12	0.54	
BP09a	278.4	10/2008		-2.87			123				0.61		0.14	0.47	
BP38	299.5	10/2013	250	-2.50	0.08	81	103	1030	7.9	0.7	0.05		-0.05	0.05	-27.9

Kafue Flats	Distance along	Year sampled	Discharge	$\delta^{18}O$	$\delta^{18}O$ stdev	POC	DOC	DIC	PN	DON	DIN	PP	DOP	DIP	POC $\delta^{13}C$
<i>Wet season</i>	floodplain (km)		(m³/s)	(‰ VSMOW)		(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(μM)	(‰ VPDB)
Station number															
KFS00	0	05/2009 and 05/2010	483	-4.89	0.05	36	228	1170	4.8	9.6	1.36	0.14	0.06	0.25	-30.5
KFS05	9.3	05/2009 and 05/2010		-4.89	0.03		268	1080		11.1	2.20	0.00	0.03	0.32	-30.4
KFS03	38.9	05/2009 and 05/2010	526	-4.87	0.06	43	280	1080	5.5	11.2	1.04	0.19	0.30	0.17	-29.3
KFS01	64.8	05/2009 and 05/2010	602	-4.88	0.02	38	255	1250	5.1	10.6	1.03	0.17	0.35	0.27	-28.2
KFS27	88.5	05/2009 and 05/2010	575	-4.77	0.04	45	265	1260	5.4	13.9	0.81	0.17	0.14	0.28	-28.3
KFS28	173.3	05/2009 and 05/2010	572	-4.41	0	27	277		3.8	15.0	0.32	0.20	0.00	0.23	-27.5
KFS11	199.1	05/2009 and 05/2010	334	-4.36	0.07	31	298	1390	4.4	11.9	0.36	0.21	0.58	0.19	-28.1
KFS10	230.4	05/2009 and 05/2010		0.00	0		335			10.1	0.81	0.13	0.00	0.29	-28.7
KFS12	231	05/2009 and 05/2010	231	-4.18	0.06	27	282	1640	4.0	15.3	0.64	0.17	0.17	0.36	-27.6
KFS14	254.6	05/2009 and 05/2010	185	-4.05	0.02	17	317	1520	2.7	13.1	0.65	0.11	0.07	0.24	-27.3
KFS16	268.8	05/2009 and 05/2010	218	-3.68	0.09	14	348	1440	2.0	14.4	0.67	0.12	0.05	0.36	-28.3
KFS17	288.3	05/2009 and 05/2010	190	-3.45	0.07	18	363	1470	2.6	15.5	0.70	0.12	0.01	0.33	-28.5
KFS18	321.5	05/2009 and 05/2010	491	-2.80	0.01	21	405	1570	2.8	16.4	0.94	0.13	0.05	0.35	-29.0
KFS21	346.5	05/2009 and 05/2010	344	-2.47		23	359		3.1	13.1	0.70	0.21	0.00	0.19	-28.3
KFS22	359	05/2009 and 05/2010	538	-2.93	0.02	23	418	1710	2.7	17.2	1.21		0.00	0.30	-28.9
KFS23	381.7	05/2009 and 05/2010	528	-2.74	0.04	23	413	1810	3.0	18.2	0.97	0.15	0.00	0.42	-27.8
KFS24	398.2	05/2009 and 05/2010	719	-2.66	0	25	439	1900	2.9	17.6	1.04	0.13	0.00	0.45	-28.3
KFS26	409.6	05/2009 and 05/2010	677	-2.58	0.19	35	393	1750	4.1	19.8	1.11	0.16	0.00	0.36	-27.8

Kafue Flats	Distance along floodplain (km)	Year sampled	Discharge (m3/s)	$\delta^{18}O$ (‰ VSMOW)	$\delta^{18}O$ stdev	POC (μM)	DOC (μM)	DIC (μM)	PN (μM)	DON (μM)	DIN (μM)	PP (μM)	DOP (μM)	DIP (μM)	POC $\delta^{13}C$ (‰ VPDB)
Dry season															
Station number															
KFS00	0	10/2008	214	-3.59	0.19	75	248	1650	6.1	12.8	1.03		0.21	0.16	-25.6
KFS03	38.9	10/2008		-3.48	0.05	74	252	1660	5.8	12.4	0.46				-26.3
KFS01	64.8	10/2008	223	-3.36	0.02	54	324	1590	4.1	14.9	0.33		0.26	0.23	-25.4
KFS27	88.5	10/2008													
KFS28	173.3	10/2008													
KFS11	199.1	10/2008				74	315		7.6						-25.3
KFS10	230.4	10/2008	170				510	2300		24.9	7.16		0.00	1.82	-26.0
KFS12	231	10/2008	170	-3.08	0.25	113		1750	12.8	14.8	0.65		0.36	0.36	
KFS14	254.6	10/2008		-3.40	0.12	129	313	1810	12.2	14.0	0.48		0.41	0.35	-24.1
KFS15	254.7	10/2008													
KFS16	268.8	10/2008		-3.26	0.24		191	1740		14.6	0.60		0.63	0.41	
KFS17	288.3	10/2008		-3.46	0.14	107	274	1780	12.4	15.1	0.71		0.37	0.36	-25.3
KFS18	321.5	10/2008		-3.14	0.13	67	376	1750	6.9	15.1	0.90		0.24	0.41	-27.3
KFS21	346.5	10/2008		-2.95	0.09	72	270		7.6						-29.0
KFS20	348.1	10/2008					759	3910		43.2	0.83		2.78	0.59	
KFS22	359	10/2008		-2.92	0.11		264	1810		12.7	2.21		0.36	0.44	
KFS23	381.7	10/2008		-2.88		68	276	1770	8.2	15.5	0.41		0.22	0.46	-29.8
KFS24	398.2	10/2008	148	-2.64	0.11		285	1810		15.3	0.37		0.21	0.42	
KFS26	409.6	10/2008	148	-2.87	0.18		836			14.2	0.57		0.13	0.49	

Samples collected on the edge of the channel

Barotse Plains	Distance along	Month/Year	Discharge	$\delta^{18}O$	$\delta^{18}O$	POC	DOC	DIC	PN	DON	DIN	PP	DOP	DIP	POC $\delta^{13}C$
<i>Wet season</i>	the channel (km)	sampled	(m ³ /s)	(‰ VSMOW)	stdev	(μ M)	(μ M)	(μ M)	(μ M)	(μ M)	(μ M)	(μ M)	(μ M)	(μ M)	(‰ VPDB)
Station number															
BP03	0	04/2013	1977	-1.82	0.10	34	175	160	2.8	10.4	0.00	0.13	0.22	0.05	-26.8
BP07	48.6	04/2013	1957	-1.82	0.05	26	182	275	2.3	10.0	0.00	0.27	0.08	0.04	-27.6
BP13	105.7	04/2013	1317	-1.91	0.03	25	183	420	2.4	9.7	1.53	0.13	0.08	0.02	-25.7
BP18	176.3	04/2013	1717	-1.31	0.11	21	206	334	1.9	11.8	0.00	0.14	0.12	0.02	-26.1
BP23	251.1	04/2013	1740	-1.44	0.06	15	230	387	1.5	12.9	0.00	0.10	0.10	0.02	-25.2

Samples collected at different depths

Kafue Flats	Water depth (m)	Month/Year	Conductivity	Dissolved oxygen	DOC	TDP	TDN
<i>Wet season</i>		sampled	(μ S/cm)	(mg/L)	(μ M)	(μ M)	(μ M)
KFS06	0.2	05/2008	170	7.66	1212	0.39	21.91
KFS06	0.8	05/2008	-	6.62			
KFS06	1.2	05/2008	170	6.60	1273	0.43	22.48
KFS07	0.2	05/2008	170	8.10	1084	0.36	22.52
KFS07	1.0	05/2008	170	7.79			
KFS07	1.8	05/2008	170	7.53	1034	0.51	22.84
KFS07	2.0	05/2008	170	7.25			
KFS09	0.2	05/2008	7.66	170	1174	0.46	37.00
KFS09	1.0	05/2008	7.65	171			
KFS09	1.5	05/2008	7.62	170	1163	0.58	25.17