

1 **Permafrost coverage, watershed area and season control of dissolved carbon**
2 **and major elements in western Siberia rivers**

3
4 **O.S. Pokrovsky^{1*}, R.M. Manasyov^{2,3}, S. Loiko², L.S. Shirokova^{1,3}, I.A. Krickov²,**
5
6 **B.G. Pokrovsky⁴, L.G. Kolesnichenko², S.G. Kopysov², V.A. Zemtsov²,**
7 **S.P. Kulizhsky², S.N. Vorobyev², S.N. Kirpotin²**

8
9 ¹ GET UMR 5563 CNRS, University of Toulouse, 14 Avenue Edouard Belin, 31400 Toulouse,
10 France, oleg@get.obs-mip.fr

11 ² BIO-GEO-CLIM Laboratory, Tomsk State University, Lenina av., 36, Tomsk, Russia

12 ³ Institute of Ecological Problem of the North, 23 Nab Severnoi Dviny, RAS, Arkhangelsk, Russia

13 ⁴ Geological Institute, Russian Academy of Science, Pyzhevskiy per., 7c1, Moscow, Russia
14
15

16 Submitted to *Biogeosciences*, Special issue

17
18 Keywords: *organic, cations, silica, permafrost, peat, groundwater, soil,*
19
20
21
22
23
24
25
26
27
28
29

ABSTRACT

30
31 Analysis of organic and inorganic carbon (DOC and DIC, respectively), pH, Na, K, Ca, Mg,
32 Cl, SO₄ and Si in ~100 large and small rivers (< 10 km² to ≤ 150,000 km²) of western Siberia
33 sampled in winter, spring, and summer over a more than 1500 km latitudinal gradient allowed
34 establishing main environmental factors controlling the transport of river dissolved components in
35 this environmentally important region, comprising continuous, discontinuous, sporadic and
36 permafrost-free zones. There was significant latitudinal trend consisting in general decrease of
37 DOC, DIC, SO₄, and major cation (Ca, Mg, Na, K) concentration northward, reflecting the interplay
38 between groundwater feeding (detectable mostly in the permafrost-free zone, south of 60°N) and
39 surface flux (in the permafrost-bearing zone). The northward decrease of concentration of inorganic
40 components were strongly pronounced both in winter and spring, whereas for DOC, the trend of
41 concentration decrease with latitude was absent in winter, and less pronounced in spring flood than
42 in summer baseflow. The most significant decrease of K concentration from the southern (< 59°N) to
43 the northern (61-67°N) watersheds occurs in spring, during intense plant litter leaching. The
44 latitudinal trends persisted for all river watershed size, from < 100 to > 10,000 km². Environmental
45 factors are ranked by their increasing effect on DOC, DIC, δ¹³C_{DIC}, and major elements in western
46 Siberian rivers as the following: watershed area < season < latitude. Because the degree of the
47 groundwater feeding is different between large and small rivers, we hypothesize that, in addition to
48 groundwater feeding of the river, there was a significant role of surface and shallow subsurface flow
49 linked to plant litter degradation and peat leaching. We suggest that plant litter- and topsoil derived-
50 DOC adsorbs on clay mineral horizons in the southern, permafrost-free and discontinuous/sporadic
51 permafrost zone but lacks the interaction with minerals in the continuous permafrost zone. It can be
52 anticipated that, under climate warming in western Siberia, the maximal change will occur in small
53 (< 1000 km² watershed) rivers DOC, DIC and ionic composition and this change will be mostly
54 pronounced in summer.

55 **1. Introduction**

56 The Western Siberia Lowland (WSL) can be considered as one of the most vulnerable
57 permafrost-bearing territories with respect to ongoing climate change, due to (i) the dominance of
58 discontinuous, sporadic and intermittent permafrost coverage rather than continuous and
59 discontinuous permafrost of Central and Eastern Siberia and the Canadian High Arctic, (ii) its flat
60 area and high impact of flooding and thermokarst development and most importantly (iii) its high
61 stock of ancient and recent organic carbon in the form of partially frozen peat deposits. Due to the
62 importance of the boreal and sub-arctic continental zones in the Earth's carbon cycle and the high
63 vulnerability of circumpolar zones to the climate warming, the majority of conducted works has been
64 devoted to the biogeochemistry of organic carbon and sediments in large rivers of the Russian boreal
65 circumpolar zone (Gordeev et al., 1996, 2004; Moran and Woods, 1997; Lobbes et al., 2000; Dittmar
66 and Kattner, 2003; Gebhardt et al., 2004; Lobbes et al., 2000; Cooper et al., 2008; Magritsky, 2010,
67 Nikanorov et al., 2010a, b; Holmes et al., 2000, 2001, 2012; Pokrovsky et al., 2010; Feng et al.,
68 2013). While these studies have allowed for the quantification of the carbon and major element
69 delivery fluxes from the continent to the Arctic ocean, the mechanisms responsible for carbon and
70 metals mobilization from the soil/groundwater to the rivers remain very poorly understood. The
71 WSL offers a unique site to test various hypotheses of element sources and to reveal related
72 mechanisms as it presents the full gradient of the permafrost coverage, climate and vegetation over
73 homogeneous sedimentary basement rock, essentially peat soil, flat orography and similar annual
74 precipitation. Taking advantage of these features, in their pioneering studies, Frey et al (2007a, b)
75 and Frey and Smith (2005) provided a first-order assessment of the relative contributions of shallow
76 surface water and deep groundwater to small western Siberia rivers. Their study was conducted
77 during the summer baseflow season presenting the largest contrast between permafrost-free and
78 permafrost-affected rivers. This allowed them to conclude that climate warming should shift the

79 permafrost-affected part of the region from surface-feeding to groundwater feeding while the
80 permafrost-free zone may remain unaffected.

81 However, unlike many regions of the world, the boreal and subarctic river regions exhibit
82 extreme seasonal variations in discharge and chemical elements concentrations (see Voronkov et al.,
83 1966; Gordeev and Sidorov, 1993; Gordeev et al., 1996; Gislason et al., 1996; Gaillardet et al., 2003;
84 Rember and Trefry, 2004; Zakharova et al., 2005, 2007; Bagard et al., 2011, 2013; Prokushkin et al.,
85 2011; Guo et al., 2004b, 2007; Olefeldt and Roulet, 2012; Voss et al., 2015). The quantitative
86 description of these systems, therefore, requires an understanding of how weathering rates and
87 riverine fluxes of major and trace elements as well as their main carrier (organic carbon) vary
88 seasonally. High seasonality implies significant variations in the source of the elements in river flow
89 over the year, which is further accentuated by high variability of the depth of the active layer and
90 relevant contributions of mineral soil weathering and the leaching of the soil organic horizon. As
91 such, the chemistry of fluxes on the seasonal scale depends on the relative role of mineral dissolution
92 vs. plant litter (organic soil) leaching. Although several recent studies have used isotopic techniques
93 in an attempt to resolve the sources of elements in subarctic rivers (Engstrom et al., 2010; Keller et
94 al., 2010; Pokrovsky et al., 2013a; Mavromatis et al., 2014), the relative contributions of mineral and
95 plant litter/organic soil components remain poorly constrained particularly for boreal watersheds.

96 The purpose of the present work is to improve our understanding of western Siberia river
97 transport of organic and inorganic carbon and major elements (Ca, Mg, K, Si) via studying numerous
98 watersheds across the 1500-km latitudinal profile during three main hydrological seasons: winter
99 baseflow, spring flood and summer-autumn period (Zakharova et al., 2014). As a working
100 hypothesis, we assume, following the previous works of Frey et al. (2007 a, b) that the permafrost
101 controls riverine chemical composition via regulating the degree of (i) groundwater feeding and (ii)
102 leaching of elements from unfrozen (active) soil layers. Because groundwaters in the permafrost
103 zone are discharged to the river via unfrozen taliks underneath the river bed (Anisimova, 1981;

104 Bagard et al., 2011, 2013), it can be suggested that the impact of groundwaters via taliks will be
105 mostly visible on large rivers, as it is also known from the geocryological studies of the WSL
106 (Fotiev, 1989, 1991). As a result, the contrast in groundwater-related element concentration between
107 rivers of different latitude is expected to be the largest during winter baseflow. This is especially true
108 in the WSL, exhibiting highly homogeneous, extremely flat topography and similar lithological
109 cover (peat, sand and silt). Therefore, the first objective of this study was to test the effect of river
110 size (watershed area) on inorganic river water components across the permafrost gradient. The
111 second objective was to assess the effect of the permafrost coverage on DOC, DIC and its isotopic
112 composition in rivers during different seasons. Specifically, during spring flood when the majority of
113 soil layer is frozen only surface flux should be important and the concentrations should reflect the
114 degree of DOC and element leaching from the plant litter. The largest contrast between rivers of
115 different size is therefore expected in August, whereas the spring flood should exhibit the lowest
116 differences in terms of DOC transport by rivers of different climate and permafrost zones. Finally,
117 the third objective of this study was first-order assessment of the major river constituent
118 concentration across the 2000-km latitudinal profile. Here, we expect, in accordance with general
119 knowledge of DOC and major cation concentration and export fluxes dependence on temperature,
120 vegetation and permafrost distribution (White and Blum, 1995; Dessert et al., 2003; Gaillardet et al.,
121 2003; Millot et al., 2003; Oliva et al., 2003; Smedberg et al., 2006; Frey and McClelland, 2009;
122 Prokushkin et al., 2011; Beaulieu et al., 2012; Tank et al., 2012a, b; Olefeldt et al., 2014), a gradual
123 or stepwise decrease of all river water constituents northward, from permafrost-free to discontinuous
124 and continuous permafrost zone. Verifying the correctness of these research statements should allow
125 for the quantitative prediction of the degree of river water composition modification in response to
126 changing environmental conditions, notably the increase in the thickness of active (unfrozen) layer,
127 increasing the winter discharge and augmenting plant biomass and productivity.

128

129 **2. Study site and Methods**

130 *2.1. Geographical setting*

131 Western Siberia Lowland (WSL) is the world **second largest flooding territory**, after the
132 Amazon's Varzea. The rivers (mainly the tributaries of the Ob, Pur, and Taz) drain **Pleistocene** sands
133 and clays, covered by thick (1 to 3 m) peat and enclose three main zones of the boreal biome, taiga,
134 forest-tundra and tundra. Quaternary clays, sands, and **silts** ranging in thickness from several meters
135 to 200-250 m have alluvial, lake-alluvial and, rarely, aeolian origin south of 60°N and fluvio-glacial
136 and lake-glacial origin north of 60°N. The older, Paleogene and Neogene, rocks are rarely exposed
137 on the surface and are represented by sands, alevrolites and clays, where carbonate material is
138 present as concretions of individual shells (Geologicheskoe Stroenie, 1958). The climate is humid
139 semi-continental with a mean annual temperature (MAT) ranging from -0.5°C in the south (Tomsk
140 region) to -9.5°C in the north (Yamburg). The annual precipitation **increases from 550 mm at the**
141 **latitude of Tomsk to 650-700 mm at Nojabrsk and further decreases to 600 mm at the lower reaches**
142 **of the Taz River**. The annual river runoff gradually increases northward, from 160-220 mm y⁻¹ in the
143 permafrost-free region to 280-320 mm y⁻¹ in the Pur and Taz river basins located in the
144 discontinuous to continuous permafrost zone (Nikitin and Zemtsov, 1986). A detailed physico-
145 geographical, hydrology, lithology and soil description can be found in earlier works (Botch et al.,
146 1995; Smith et al., 2004; Frey and Smith, 2005, 2007; Frey et al., 2007a, b; Beilman et al., 2009) and
147 in our recent limnological and pedological studies (Shirokova et al., 2013; Manasyrov et al., 2014,
148 2015; Stepanova et al., 2015). A detailed map of studied region together with main permafrost
149 provenances and river runoff in the WSL is given in **Fig. 1** and the list of sampled rivers grouped by
150 watershed size and season is presented in **Table 1**. **The position of the permafrost provenances in the**
151 **WSL shown in this figure is based on extensive geocryological work in this region (Baulin et al.,**
152 **1967; Gruzlov and Trofimov, 1980; Baulin, 1985; Liss et al., 2001).** The hydrological parameters of
153 **the WSL rivers based to calculate a runoff contour line are described in the Supplement 1.**

154 *2.2. Chemical and isotope analyses and statistical treatment*

155 Altogether, 96 river samples were collected in early June 2013 (spring flood), August 2013
156 and 2014 (summer baseflow), October 2013 (autumn) and February 2014 (winter baseflow) along
157 the 1500 km latitudinal gradient (Table 1). All sampled rivers of western Siberian Lowland belong to
158 the Kara Sea basin. The sampling during seasons covered full gradient from south to north, except
159 the month of October which was sampled only in rivers south of 60°N (12 rivers in total). The
160 watershed area of sampled rivers ranged from 2 to 150,000 km², not considering the Ob River in its
161 medium course zone. Collected water samples were immediately filtered in pre-washed 30 mL PP
162 Nalgene® flasks through single-use Minisart filter units (Sartorius, acetate cellulose filter) with a
163 diameter of 25 mm and a pore size of 0.45 µm. The first 20 to 50 mL of filtrate were discarded.
164 Filtered solutions for cation analyses were acidified (pH ~ 2) with ultrapure double-distilled HNO₃
165 and stored in pre-washed HDPE bottles. The preparation of bottles for sample storage was performed
166 in a clean bench room (ISO A 10,000). Filtered samples for DOC, DIC, UV_{280 nm} absorbency and
167 anions were stored in the refrigerator maximum 3 weeks before the analyses. The effect of storage
168 for DOC, DIC and optical measurements in boreal waters was found to be within the uncertainty of
169 analysis (Ilina et al., 2014). Blanks were performed to control the level of pollution induced by
170 sampling and filtration. The DOC blanks of filtrate never exceeded 0.1 mg/L which is quite low for
171 the organic-rich river waters sampled in this study (i.e., 10–60 mg/L DOC). pH was measured in the
172 field using a combined electrode calibrated against NIST buffer solutions (pH of 4.00 and 6.86 at
173 25°C). The accuracy of pH measurements was ±0.02 pH units. DOC and DIC were analyzed using a
174 Carbon Total Analyzer (Shimadzu TOC VSCN) with an uncertainty better than 3%. Special
175 calibration of the instrument for analysis of both form of dissolved carbon in organic-rich, DIC-poor
176 waters was performed as described elsewhere (Prokushkin et al., 2011). Major anions (Cl, SO₄)
177 concentrations were measured by ion chromatography (HPLC, Dionex ICS 2000) with an
178 uncertainty of 2%. The UV-absorbency at 280 nm is used as a proxy for aromatic C and source of

179 DOM in the river water. It was measured using 1 cm quartz cuvette in a CARY-50 UV-VIS
180 spectrophotometer (Bruker, UK). Major cations (Ca, Mg, Na, K) and Si were determined with an
181 ICP-MS Agilent ce 7500 with In and Re as internal standards and 3 various external standards,
182 placed each 10 samples in a series of river water. Approximately 30% of samples were analyzed for
183 Ca, Mg and Na concentration using atomic absorption spectroscopy (flame) with an uncertainty of 2
184 %. Reasonable and non-systematic agreement (between 5 and 10%) with the results of ICP MS
185 analyses was achieved. Aqueous Si concentrations were also determined colorimetrically (molybdate
186 blue method) with an uncertainty of 1% using a Technicon automated analyzer. The SLRS-5
187 (Riverine Water Reference Material for Trace Metals certified by the National Research Council of
188 Canada) was used to check the accuracy and reproducibility of each analysis (Yeghicheyan et al.,
189 2014).

190 The ^{13}C in dissolved inorganic carbon was analyzed in filtered river water sampled in bubble-
191 free sealed glass bottles by gas chromatography and isotope mass spectrometry, using Delta V
192 Advantage and Finnigan Gas-Bench-II in order to determine $\delta^{13}\text{C}_{\text{DIC}}$ (per mil relative to V-PDB;
193 Fritz and Fontes, 1980). For these measurements, 0.1 mg of 100% H_3PO_4 was added to the
194 borosilicate vial and flushed with He (purity of 7.0) for 400 s. Afterwards, 1 mL of the sample was
195 injected into the vial and shaken for 36 h at 24°C. Standard samples of C-O-1 and NBS-19 were
196 routinely analyzed to test the accuracy of our measurements; typically, a disagreement of less than
197 0.3 ‰ between the measured and certified values was observed, with a total estimated measurement
198 uncertainty of ± 0.2 ‰.

199 The concentration of carbon and major elements in rivers were treated using the least squares
200 method, Pearson correlation and one-way ANOVA (SigmaPlot version 11.0/Systat Software, Inc).
201 The ANOVA was used to reveal the differences between different permafrost zones. It was carried
202 out using Dunn's method because each sampling period contained different number of rivers.
203 Regressions and power functions were used to examine the relationships between the dissolved

204 **component** concentrations and the watershed area, river discharge, average latitude of the watershed
205 and seasons. Comparison of **DOC and major element** concentration in rivers sampled in three main
206 permafrost zones (continuous, discontinuous and permafrost-free regions), during all seasons and of
207 different watershed size class was conducted using the non-parametric H-criterion Kraskal-Wallis
208 test. **First, we separated the watershed into four main classes encompassing all studied rivers (except**
209 **Ob): < 100 km², 100 to 1000 km², 1000 to 10,000 km² and > 10,000 km². We considered three main**
210 **season in six different ranges of latitude (56 to 58°N, 58 to 60°N, 60 to 62°N, 62 to 64°N, 64 to 66°N**
211 **and 66 to 68°N). We checked for the variation of measured parameters of each watershed size as a**
212 **function of latitude, separately in each season. In addition, a generalized assessment of the role of**
213 **permafrost type and abundance on river water chemical composition was possible via separating all**
214 **the sampled watersheds on three categories according to the permafrost distribution in WSL:**
215 **permafrost-free, discontinuous and continuous permafrost.**

216

217

218 **3. Results**

219 Results of major element analysis in rivers are listed in **Table S1** of the Supplement **and the**
220 **main results of statistical treatment are listed in Table S2.** Based on the Kraskal-Wallis H statistics,
221 the differences between the seasons and between different latitudes were found to be significantly
222 higher (p-level < 0.0001) for most elements than the difference between watershed size classes,
223 within each seasons and within each latitude range. This is illustrated for pH, DOC, DIC and $\delta^{13}\text{C}_{\text{DIC}}$
224 in **Figs 2, 3, 4 and 5**, respectively which show the measured value as a function of latitude for
225 different watershed classes, individually for each main season. Similar plots for major cations (Ca,
226 Mg, K) and Si are given in supplementary figures **S1, S2, S3 and S4**, respectively. **The latitudinal**
227 **coverage of October was too small to be presented in these figures; however, the October data of 12**
228 **rivers were used for statistical treatment and for assessing the permafrost impact.** There is a clear and

229 significant trend of concentration with latitude; the differences between different latitude ranges are
230 significant at $p < 0.0001$ for all elements, and at $p < 0.05$ for Si. The effect of the watershed size on
231 river water chemical composition in summer, winter and spring is much smaller than that of latitude
232 ($9 < H < 12$, $p < 0.05$ and $20 < H < 50$, $p < 0.001$, respectively). Considering all rivers
233 simultaneously, the effect of the season is clearly seen at $p < 0.001$ for all elements except DOC; the
234 latter, however, is also statistically significant ($H = 10.6$, $p = 0.014$). Considering full data set of all
235 seasons and watershed sizes, we distinguished three geographical zones in terms of the permafrost
236 abundance: continuous, discontinuous and absent. For most river water parameters (pH, DIC, DOC,
237 major anions and cations) the differences between three zones are significant ($30 < H < 95$, p -level $<$
238 0.001). Si concentration exhibited lower but statistically significant differences between different
239 zones ($H = 9.5$, $p = 0.0086$)

240 **Considering all seasons** and watershed sizes revealed a significant decrease of pH, Ca and Mg
241 northward with the largest changes occurring at the beginning of discontinuous permafrost coverage
242 (**Fig. S5 A, B and C**, respectively). The DOC and DIC also decrease in concentration with the
243 increase in the degree of permafrost coverage (**Fig. S6 A, B**, respectively) whereas the isotopic
244 composition of the DIC becomes progressively more negative northward (from ca. -15 ‰ in the
245 permafrost-free zone to -20...-25 ‰ in the continuous permafrost zone, **Fig. S6 C**). In contrast, the
246 effect of the permafrost on Si concentration is not clearly seen; the scatter of the data between
247 different seasons and watersheds does not allow tracing any significant trend (not shown).

248 The optical properties of DOC remain essentially constant throughout the full range of
249 watershed sizes, latitudes and seasons (**Fig. S7**). The largest variation of specific $UV_{280\text{ nm}}$
250 absorbance occurred in winter, when several DOC-rich waters from the southern (permafrost-free)
251 part of WSL demonstrated quite low concentration of aromatic (colored) compounds.

252

253

254 **4. Discussion**

255 *4.1. Effect of latitude (permafrost and vegetation) on major cation, Si and DIC mobilization*
256 *from the soil profile and groundwater to the river*

257 From general knowledge of environmental control on carbon and major element fluxes in
258 rivers of the Russian subarctic (Prokushkin et al., 2011; Pokrovsky et al., 2012) and other boreal and
259 subarctic regions (Laudon et al., 2004; Petrone et al., 2006; Walvoord and Striegl, 2007; Jantze et al.,
260 2013; Giesler et al., 2014), we anticipate a decrease of most element concentration, including DOC,
261 northward regardless of the season and the river size in the WSL due to (1) a decrease of chemical
262 weathering intensity with the temperature, well demonstrated for igneous rocks such as basalts
263 (Dessert et al., 2003) and granites (Oliva et al., 2003); (2) decrease of the thickness of peat deposits
264 in total and the active soil (peat) layer in particular (Liss et al., 2001; Beilman et al., 2009; Stepanova
265 et al., 2015 and references therein); (3) a decrease of plant biomass and related plant litter stock on
266 the surface of the soils (Tyrtikov, 1979; Frey and Smith, 2007); (4) a shortening of the unfrozen
267 period of the year and (5) a decrease of the degree of groundwater feeding (Romanovsky, 1983;
268 Nikitin and Zemtsov, 1986; Fotiev, 1991). The factors capable of enhancing element export fluxes in
269 northern (permafrost-bearing) rivers relative to southern (permafrost-free) rivers of the WSL are (1)
270 the decrease of dissolved organic matter (DOM) respiration by heterotrophs in the water and soil
271 column and thus the increasing removal of allochthonous DOC from the soil to the river (Striegl et
272 al., 2005); (2) the increase of DOC and related element leaching from plant litter and topsoil
273 (Pokrovsky et al., 2005; Giesler et al., 2006; Fraysse et al., 2010) during more pronounced massive
274 freshet event or summer high flow (Michel and Vaneverdingen, 1994; McClelland et al., 2006;
275 White et al., 2007); (3) the decrease of DOM retention (adsorption) on the mineral soil horizon
276 because clay horizon is typically frozen in the north (Kawahigashi et al., 2004); 4) the decrease of
277 authigenic clay and allophane mineral formation in the soil horizons (Targulian, 1971).

278 At the current, rather limited, stage of knowledge of mineral, organic soil horizons and plant
279 biomass chemical composition and reactivity across the WSL, only a few environmental factors can
280 be quantitatively tested based on river water chemical analyses. In the case of the dominance of
281 groundwater feeding of the river, the decrease of element concentrations from water-rock interaction
282 whose transport is not limited by availability of DOM (Ca, Mg, DIC) is expected to be most
283 pronounced in winter, when the groundwater feeding is maximal (see Walvoord and Striegl (2007)
284 for the Yukon River basin example). Moreover, in the permafrost-bearing zone during winter
285 baseflow, one should expect significant difference in element concentration in winter between small
286 rivers (weakly or not affected by taliks) and large rivers (essentially fed by taliks), as it is known
287 from local geocryological conditions (Baulin et al., 1967; Romanovsky, 1983; Fotiev, 1989, 1991;
288 Ivanov and Beshentsev, 2005). In spring, when the active layer is very thin and the majority of the
289 soil column is frozen, the export from the watershed is dominated by surface flow and thus the
290 difference in groundwater-related element concentration between (i) small and large rivers and (ii)
291 north and south should be minimal. However, the abovementioned hypotheses are not supported by
292 DIC, Ca and Mg concentrations observed in rivers (Fig. 4, S1 and S2). First, the DIC concentrations
293 decrease between permafrost-free and discontinuous/continuous permafrost zones is a factor of 15 ± 5
294 in winter (Fig. 4 A) and a factor of 60 ± 10 in spring (Fig. 4 B). Similarly, the decrease of Ca and Mg
295 concentrations between south of 59°N and $62\text{-}66^\circ\text{N}$ zones is $\times 10$ in winter and $\times 20\text{-}30$ in May. In
296 fact, it is the spring period which exhibits the highest contrast in element concentrations between the
297 south and the north. Second, for the latitude concentrations gradient from south to north, the relative
298 DIC, Ca and Mg concentration change between large ($1000\text{-}10,000 \text{ km}^2$ and $> 10,000 \text{ km}^2$) and small
299 ($< 100 \text{ km}^2$) rivers in winter is not statistically significant ($p > 0.05$).

300 However, a systematic decrease of Ca concentration in the WSL rivers northward (Fig. S1,
301 S5 B) is consistent with general decrease of Ca concentration in soil ecosystems as illustrated in Fig.
302 S8. An order of magnitude decrease of Ca concentration in mineral horizons of SWL peat columns

303 occurred between 55 and 66°N (Stepanova et al., 2015). On a smaller scale, a 3 times decrease of
304 exchangeable Ca concentration in alluvial soils of the Ob basin from 56 to 60°N was reported
305 (Izerskaya et al., 2014). These observations confirm strong lithological and soil weathering control
306 on Ca concentration in both deep and surface soil horizons and vegetation, which finally determines
307 the extend of Ca transport via surface flux to the river.

308 North of 66°N, concentrations of Ca, Mg and sulfate increase relatively to their concentration
309 at 62-66°N of discontinuous permafrost zone. This is especially pronounced during the summer
310 period (Fig. S1 C, S2 C). We do not exclude here the influence of marine sedimentary deposits
311 containing salts in the deep part of the mineral soil profile below the peat layer. These deposits are
312 described in the low reaches of Taz and Pur rivers, based on sedimentary cores extracted during
313 extensive drilling of the territory (Liss et al., 2001). This influence, however, cannot be
314 unequivocally evidenced because (i) DIC concentrations also increase in summer, north of 66°N,
315 although DIC is not likely to be affected by marine deposits, and (ii) chloride, an efficient marker of
316 sea salts, is not increasing in the north (not shown).

317 The isotopic composition of DIC confirms the general features of DIC and cation
318 concentration (Fig. 5). The groundwater feeding by taliks in winter is highly uniform over 10 degrees
319 of latitude with the value of $\delta^{13}\text{C}_{\text{DIC}}$ being equal to -15 ± 5 ‰, reflecting both carbonate/silicate
320 weathering and a buildup of CO_2 with a stronger respiratory signal (Finlay, 2003; Striegl et al., 2001;
321 Giesler et al., 2014; Rinta et al., 2015). During this period, the variability of $\delta^{13}\text{C}_{\text{DIC}}$ is the highest in
322 small ($< 100 \text{ km}^2$) watersheds, but no trend of isotopic composition with latitude could be evidenced
323 at $p < 0.05$ (Fig. 5 A). This isotopic signature is preserved in spring for southern ($< 60^\circ\text{N}$) watersheds
324 whereas in permafrost-affected regions, $\delta^{13}\text{C}_{\text{DIC}}$ decreases to c.a. $-25 \dots -20$ ‰ regardless of the river
325 size and the type and the abundance of the permafrost (Fig. 5b). Such low values in the permafrost-
326 affected zone could not anymore represent the influence of carbonate/silicate rock weathering by soil
327 CO_2 and likely reflect direct microbial processing of soil and sedimentary organic matter (Waldron et

328 al., 2007; Giesler et al., 2013), with the DIC isotopic signature similar to that of organic carbon in
329 western Siberia subarctic topsoil ($-26\pm 2\text{‰}$, Gentsch et al., 2015) and the Ob river organic sediments
330 (-25 to -27‰ , Guo et al., 2004a).

331 A plausible explanation for the $\delta^{13}\text{C}_{\text{DIC}}$ seasonal variation being mostly pronounced in the
332 permafrost zone can be that microbial mineralization of dissolved organic carbon occurs most
333 efficiently during the springtime, when significant amounts of fresh organic matter from ground
334 vegetation are leached by melted snow. Higher bioavailability of vegetation leachates relative to
335 more refractory soil humic and fulvic acids is known from studies in other temperate (van Hees et
336 al., 2005) and boreal (Wickland et al., 2007) regions. The lack of $\delta^{13}\text{C}_{\text{DIC}}$ decrease in spring relative
337 to winter in the permafrost-free zone may stem from (i) significant input of the carbonate/silicate
338 rock-hosted groundwaters during full period of the year in the south, or (ii) the different nature of
339 DOM in the south, where the more refractory organic matter originated from peat leaching is less
340 subjected to microbial processing compared to fresh vegetation leachates in the north, where the peat
341 soil in spring is frozen. One has also take into account that the DIC concentrations in spring are a
342 factor of 30 lower in the permafrost-bearing region relative to permafrost-free region (Fig 4 b). As
343 such, a relatively small input of microbially-respired CO_2 will be significantly more visible in the
344 $\delta^{13}\text{C}_{\text{DIC}}$ value of the northern rivers compared to that of the southern rivers.

345 The variation of $\delta^{13}\text{C}_{\text{DIC}}$ along the permafrost/latitude gradient helps to better explain the
346 origin of DIC in rivers in contrasting permafrost zones. Consistent with a progressive decrease of the
347 groundwater feeding of rivers northward (Nikitin and Zemtsov, 1986; Frey et al., 2007b), we observe
348 a distinct trend of $\delta^{13}\text{C}_{\text{DIC}}$ with the latitude during spring period, reflecting the shift of DIC origin
349 from groundwater in the south to plant litter degradation and soil respiration in surface waters north
350 of 62°N (Fig. 5 B). In winter, the $\delta^{13}\text{C}_{\text{DIC}}$ is rather constant within the full latitudinal profile (Fig.
351 5A) confirming the dominant role of carbonate/silicate mineral weathering by atmospheric and soil
352 CO_2 dissolved in the groundwaters. Indeed, hydrological studies in the WSL revealed that the

353 groundwater feeding of small (< 10,000 km² watershed) rivers decreases from 20-30% in the
354 discontinuous and sporadic/isolated part of the WSL to 3-6% in the northern, continuous permafrost
355 zone (Novikov et al., 2009). These numbers agree with estimations based on RHS data of large
356 western Siberian rivers (Nadym, Pur and Taz) and the left tributaries of the Yenisei River (Dubches,
357 Elogyi and Turukhan; Nikitin and Zemtsov, 1986). According to more recent evaluations of Frey et
358 al. (2007b), the groundwater contribution to summertime period river chemical composition ranges
359 between 30 and 80% for the rivers located between 56 and 58°N.

360 **Consistent with these findings, the** pH values of 7 to 7.5 in the southern rivers observed both
361 in winter and spring (Fig. 2 A, B) are indicative of carbonate/silicate rock input. The spring acid
362 pulse, well established in other permafrost-free boreal regions (Buffam et al., 2007), is not at all
363 pronounced in the south of the WSL but becomes clearly visible in the permafrost-affected, northern
364 regions where the spring-time pH decreases to 5...6 (Fig. 2 B). This illustrates the more important
365 role of plant litter and moss leaching in the permafrost-bearing zone on solute export from the
366 watershed. In addition, the dominance of sands north of 62°N (**Liss et al., 2001**) may allow low
367 molecular weight (LMW) organic acids migrate to the river from the soil profile. In the southern,
368 permafrost-free zone, the dominating clays underneath the peat can adsorb acidic LMW organic
369 compounds and thus do not allow the acid pulse to be clearly visible.

370 The increase of pH in summer relative to spring period is again less visible in the south than
371 in the north (Fig. 2 C) and may reflect the persisting role of bedrock dissolution as well as the change
372 of the river feeding regime, from top soil and vegetation **in the north** to the peat soil column leaching
373 **in the south**. The summertime increase of river water pH north of 60°N, in the forest-tundra and
374 tundra zone may be linked to (i) enhanced photosynthesis in rivers of the north **due to better**
375 **insolation and less forest shading** and (ii) mobilization of DOM and other solutes from soil
376 depressions rather than from watershed divides. The depressions are subjected to intense rinsing
377 during the spring seasons, when the majority of soluble acidic compounds are flushed from the litter

378 and O_e horizon. These mechanisms are evidenced from studies of frozen bogs hydrological balance
379 performed in the northern part of studied territory (Novikov et al., 2009). In contrast, the watershed
380 divides contain significant amounts of organic litter and release organic acids only in spring, when
381 they are covered by temporary ponds of melted snow (see Manasypov et al., 2015). This
382 hydrological scheme of river water feeding is based on the seasonal multiannual observations on
383 frozen bogs of the north of WSL (Novikov et al., 2009), although the chemical nature of DOM
384 mobilized from different parts of the watershed remains unknown.

385 The importance of plant litter and ground vegetation leaching as element sources in western
386 Siberian rivers can be assessed from the comparison of K concentrations as a function of latitude
387 during different seasons (Fig. S 3). The most significant decrease of K concentration from the
388 southern (< 59°N) to the northern (61-67°N) watersheds occurs in spring, during intense plant litter
389 leaching. Regardless of latitude, K concentration follows the order spring > winter > summer with
390 the highest concentrations, up to 2500 ppb, recorded in permafrost-free region. Given that the other
391 cations, possibly originating from the water-mineral interaction at some depth, do not exhibit such
392 high concentration in spring, we interpret the spring-time K “pulse” as indicative of plant litter
393 leaching in productive taiga zone. This “pulse” is much less visible in the permafrost zone due to
394 significantly lower biomass and primary productivity of forest-tundra and tundra biomes compared
395 to the taiga of the WSL (Tyrtikov, 1979; Liss et al., 2001).

396 Despite significant variability of Si concentrations among rivers of various sizes across the
397 latitude profile (Fig. S 4), the concentrations in the permafrost zone are not lower than those in the
398 south of the WSL. Results of a previous study of WSL rivers during summer show that Si
399 concentration are weakly dependent on latitude (Frey et al., 2007), as also confirmed in this work for
400 spring flood and winter baseflow period. Given that (i) the dominance of permafrost north of 64°N
401 implies very low groundwater feeding (4 to 6% of the annual discharge, see Nikitin and Zemtsov,
402 1986; Novikov, 2009) and (ii) the upper part of the soil profile including its seasonally frozen and

403 unfrozen parts is mostly peat rather than silicate mineral sediments, the role of groundwater – silicate
404 rock interaction in Si supply to northern rivers should be quite low. Therefore, we hypothesize that
405 elevated concentrations of Si in northern rivers are due to peat leaching and degradation. A depletion
406 of Si in rivers of the southern part of the WSL may be due to Si retaining by abundant bog and forest
407 vegetation. This is consistent with general setting of the WSL, recovering from the last glaciation
408 (Liss et al., 2001), with contemporary peat accumulation in the south and old frozen peat
409 thawing/degradation in the north.

410

411 *4.2. DOC concentration across 1500 km latitude transect of variable permafrost coverage*

412 Results of organic carbon concentration in western Siberia rivers collected over various
413 seasons of the year generally confirm the pioneering findings of Frey and Smith (2005). The strong
414 statistically significant ($p < 0.05$) contrast in DOC concentration between permafrost-free,
415 discontinuous and continuous permafrost zone persists over the course of the year and each season
416 except probably winter (Fig. 3 and Fig. S6 A). This difference is also seen in $\delta^{13}\text{C}_{\text{DIC}}$ value among all
417 three zones (Fig. S6 C) suggesting, on the annual scale, a more significant contribution of microbial
418 processing of plant and soil organic carbon to HCO_3 and CO_2 of the river water in the permafrost-
419 bearing zone compared to the permafrost-free zone.

420 In accordance with the conclusion reached by Frey and Smith (2005), the variation in
421 hydrology may play a limited role in DOC variability and export from the watershed of WSL rivers.
422 The gradient in DOC concentrations along the latitudinal profile remains similar between spring flood
423 and summer baseflow (Fig. 3 b and c). Although the winter period does not exhibit such a clear
424 difference between permafrost-free and permafrost-affected regions (Fig. 3 a), the contribution of the
425 winter discharge to the annual flux of DOC is between 10 and 15% and as such does not significantly
426 affect annual export of DOC from the watersheds.

427 In contrast, the gradient of organic carbon concentration along the latitudinal profile in spring
428 will be mostly controlled by the difference in plant litter stock subjected to leaching by melted snow.
429 As such, one would not expect any significant difference between large and small rivers at otherwise
430 similar runoff, vegetation and bog coverage. This is partially confirmed by the similarity of the
431 $UV_{280\text{ nm}} - \text{DOC}$ slope, corresponding to similar degree of DOM humification, among different
432 seasons and latitudinal positions (Fig. S7). The uniform distribution of UV_{280} absorbance witnesses
433 the main control of DOC by allochthonous (terrestrial) input from peat and/or ground vegetation
434 leachates. The exceptions are the rivers Vasyugan (RF 63), Shegarka (RF 3) and Vatinsky Egan (RF
435 57) exhibiting low $UV_{280\text{nm}}$ at high [DOC] (Fig. S7). These rivers are potentially affected by oil
436 production sites and may contain some uncolored products of hydrocarbon oxidation in the
437 underground waters.

438 Overall, results on western Siberia rivers generally confirm the conclusion of Finlay et al.
439 (2006) on (i) the lack of groundwater contribution to streamflow in arctic watersheds and (ii) that
440 river DOC dynamics are driven essentially by processes occurring at the soil surface. However we
441 doubt the importance of large DOC pool production under very cold conditions as the main reason of
442 sustained high concentration of DOC at snowmelt suggested by Finlay et al. (2006). Indeed, the plant
443 litter degradation in winter, even in the warmest scenario, is minimal and does not contribute
444 significantly to annual litter leaching (Bokhorst et al., 2010, 2013). Instead, we suggest fast plant
445 litter and ground vegetation leaching in spring, at the very beginning of the snow melt. Such a fast
446 enrichment in DOC and colored organic compounds of surface water depressions, on the order of
447 several hours, has been observed in the discontinuous permafrost zone in early June (Manasyrov et
448 al., 2015). Significant release of DOC and nutrients from flooded ground vegetation in the southern
449 part of WSL is also known (Vorovyev et al., 2015).

450 An unexpected result of the study of western Siberian watersheds is the lack of the
451 enrichment in DOC of small headwater streams, in contrast to what was reported for Scandinavian

452 rivers and streams (Agren et al., 2007, 2014 and references therein). In WSL, especially in the
453 northern, permafrost-affected zone, the small (< 100 km²) streams yielded DOC concentration that
454 were not statistically higher ($p > 0.05$) than those of larger rivers, neither in spring flood nor in
455 summer. A number of factors can be responsible for the observed difference between permafrost-free
456 European and permafrost-bearing Siberian watersheds. In the north of western Siberia, the microbial
457 processing of DOM in large rivers may be weakly pronounced. This is confirmed by the observation
458 that the degree of light C isotope enrichment (lowering $\delta^{13}\text{C}_{\text{DIC}}$) in spring is independent ($p > 0.05$)
459 of the size of the river (Fig. 5 B), and, correspondently, of the water residence time on the watershed.
460 According to Kawahigashi et al. (2004), the DOM in northern, permafrost-affected tributaries of the
461 Yenisey River was significantly less biodegradable than that in southern tributaries. This may
462 contribute to better preservation of DOM in the stream yielding its independence of the water travel
463 time. Small watersheds of western Siberia exhibit a runoff and average slope very similar to that of
464 the large rivers, given very flat orographic context of WSL. This contrasts the mountain regions of
465 Sweden and Alaska where the headwater streams may exhibit higher runoff and thus higher export of
466 the dissolved constituents. Finally, the riparian zone, very important for regulation DOC stock and
467 export in small streams draining glacially-formed terrain of NW Europe (Dick et al., 2014;
468 Kuglerova et al., 2014), is much less pronounced in western Siberia, where generally flat, frequently
469 flooded areas dominate the watershed profile.

470 The elevated DOC concentrations in continuous permafrost zone, especially north of 67°N
471 observed in the present study (Fig. 3 B, C) are consistent with previous results showing that, at
472 otherwise similar factors, the permafrost areas are a greater source of DOC than the areas with
473 seasonal frost (Carey et al., 2003). In permafrost areas, meltwater travels through organic-rich layers
474 in the form of so called supra-permafrost flow, as opposed to areas without impermeable permafrost
475 table. In the latter, the infiltration of organic-rich surface waters to deep mineral layer and DOC
476 sorption on clay minerals may occur thus decreasing the overall export of DOC (see Smedberg et al.,

477 2006 for discussion). Given the dominance of peat rather than minerals within the active (unfrozen)
478 layers of soil profile, the difference between permafrost-free and permafrost-affected zones is even
479 more accentuated in western Siberia.

480 A sketch of typical soil profiles of western Siberia in the permafrost-free and permafrost-
481 bearing zone presenting DOC mobilization pathways from the soil to the river in the end of active
482 period is illustrated in **Fig. 6**. The two cross sections shown in this figure are highly representative
483 for two most contrasting cases of soil and watershed flux formation, corresponding to dark
484 coniferous taiga in the permafrost-free zone and dwarf shrubs with green mosses of tundra and
485 forest-tundra in frozen peatlands of continuous permafrost zone; both sites are located at the
486 watershed divide. The detailed position of soil horizons and their attribution to FAO is based on
487 available literature data (Tyrtikov, 1973, 1979; Liss et al., 2001; Pavlov and Moskalenko, 2002) and
488 our recent investigations of the region (Loiko et al., 2015; Stepanova et al., 2015). We hypothesize
489 that plant litter- and topsoil derived-DOC adsorbs on clay mineral horizons in the southern,
490 permafrost-free and discontinuous/sporadic permafrost zone but lacks the interaction with minerals
491 in the continuous permafrost zone. This assumption corroborates results found during another
492 latitudinal river transect of Siberia, along the Yenisey river and its left tributaries draining peatlands
493 of the WSL (Kawahigashi et al., 2004, 2006): the northern tributaries exhibited significantly higher
494 DOC concentrations than the southern tributaries of this river. Specifically, given the significant
495 thickness of the peat even in the northernmost part of the WSL and the active layer thickness of < 50
496 to 80 cm (30 cm on mounds and 80 to 150 m in troughs and depressions, Tyrtikov (1973, 1979),
497 Baulin et al. (1967), Baulin (1985), Khrenov (2011), Novikov et al. (2009)), even in the region of
498 continuous permafrost development, peat soil interstitial solutions might not enter in contact with the
499 mineral soil horizon and thus will not decrease their DOC concentration during migration from the
500 soil column to the river along the permafrost impermeable layer (**Fig. 6**).

501 Therefore, in the permafrost zone, the DOC export is strongly controlled by DOC residence
502 time and water travel pathway through organic topsoil and lichens, moss and litter leaching vs. peat
503 and mineral layer leaching (Fig. 6 B). In this case, it is only the thickness of the unfrozen peat and
504 local permafrost coverage that control the DOC export from the soil to the river. As a result, DOC
505 concentration in the streams will be weakly dependent on the watershed size and seasons. It follows
506 that DOC export from peat soils by medium size ($n \cdot 10,000 - n \cdot 100,000 \text{ km}^2$) rivers located entirely
507 in the permafrost zone may be higher than that of the larger rivers, crossing permafrost-free regions.
508 This hypothesis is supported by available information on DOC yield by rivers of the WSL. Thus,
509 rivers Taz ($s = 150,000 \text{ km}^2$), Pur ($112,000 \text{ km}^2$) and Nadym ($64,000 \text{ km}^2$), entirely located in the
510 discontinuous permafrost zone, exhibit 1.9, 2.1, and $4.4 \text{ t km}^{-2} \text{ y}^{-1}$ DOC yield, respectively (Gordeev
511 et al., 1996 and calculated based on data of the RHS). This is significantly higher than the value
512 suggested for the Ob River ($1.2 \text{ t km}^{-2} \text{ y}^{-1}$, Gordeev et al., 1996).

513

514 *4.3. Possible evolution of western Siberia rivers chemical composition and fluxes under climate* 515 *change scenario.*

516 The most likely scenario of the climate change in western Siberia consists of shifting the
517 permafrost boundary further north and increase of the active layer thickness (Pavlov and
518 Moskalenko, 2002; Frey, 2003; Romanovsky et al., 2010; Vasiliev et al., 2011; Anisimov et al.,
519 2013). The permafrost boundary change, equivalent to the northward shift of the river latitudes, may
520 decrease maximum 2 fold the DOC concentrations of the most northern rivers due to the change of
521 continuous to discontinuous permafrost. The thickness of the active layer is projected to increase
522 more than 30% during this century across the tundra area in the Northern Hemisphere (Anisimov et
523 al., 2002; Stendel and Christensen, 2002; Dankers et al., 2011). In the WSL, this increase will be
524 most dramatic in the north, where the peat deposits are thinner than those in the discontinuous
525 permafrost zone (Botch et al., 1995; Liss et al., 2001; Novikov et al., 2009; Kremenetsky et al.,

2004). Assuming a short-term (hundred years) scenario in the WSL, we hypothesize that the main consequences of this increase will be the involvement of upper clay horizon and sand/silts in water pathways within the soil profile. As a result, the DOC originated from the upper peat layer leaching and plant litter degradation will be retained on mineral surfaces and in the clay interlayers (Kaiser et al., 2007; Oosterwoud et al., 2010; Mergelov and Targulian, 2011; Gentsch et al., 2015). To which degree this change of water pathways in the soil column may affect the other dissolved components cannot be predicted. However, this effect for inorganic solutes is expected to be lower than that for DOC, given much lower affinity of HCO_3^- , major cations and Si to clay surfaces and the lack of unweathered (primary) silicate rocks underneath peat soil column. Nevertheless, the possibility of leaching of inorganic components from the mineral layers should be considered. For example, DOC export exceeded DIC export in a tributary of the Yukon River during high flow, whereas DIC predominated during low flow and the DIC yields increased with decreasing permafrost extend (Dornblaser and Striegl, 2015). Unfortunately, no time series on hydrochemistry of rivers of continuous permafrost development, north of 64°N , are available to test the hypothesis of the impact of climate change on a possible decreasing DOC flux from frozen peatlands and the DOC/DIC change due to ongoing decrease of permafrost protection of mineral layer from adsorbing DOC.

Important modifications linked to the climate change in boreal and subarctic zones concern the change of the hydrological regime (Karlsson et al., 2015), in particular the increase of the winter baseflow (Yang et al., 2004; Ye et al., 2009; Serreze et al., 2000) due to the increase of the groundwater feeding (Frey et al., 2007a,b; Walvoord and Striegl, 2007; Rowland et al., 2010; Walvoord et al., 2012), coupled with the increase of the overall precipitation and, consequently, water runoff (Peterson et al., 2002, McClelland et al., 2006). Here, we argue that 10 to 30% modification in the annual runoff will be within the variation of the DOC and cation concentrations between watersheds of various sizes observed in the present study and as such will not significantly affect the export fluxes of river water constituents.

551 To which degree the ongoing DOC concentration and flux rise in rivers, linked to climate
552 change and/or acidification as reported in Western Europe and Canada (Worrall et al., 2004; Porcal
553 et al., 2009) can be applied to the WSL is unknown. However, we did not observe any significant
554 (i.e., > 30%) change of DOC fluxes over past 30 to 40 years neither in the boreal non-permafrost
555 pristine region of NW Russia (Severnaya Dvina River, Pokrovsky et al., 2010), nor in the Central
556 Siberian, continuous permafrost rivers of the Yenisei basin (Pokrovsky et al., 2005). Moreover, a
557 decrease of DOC fluxes in the Yukon River was reported and suggested to be linked to enhanced
558 mineralization of DOC by biota (Striegl et al., 2005). Note also that the more recent evaluation of
559 the Ob River DOC discharge using flow-weighted concentration of 9.4 mg/L measured in 2003-2007
560 (Cooper et al., 2008) gives a flux of 1.3 t C/km²/y, well comparable with earlier estimate of 1.2 t
561 C/km²/y, based on the RHS data of 1950-1990 (Gordeev et al., 1996).

562 The increase of vegetation productivity reported for Arctic river basins (Sturm et al., 2001,
563 Tape et al., 2006; Kirilyanov et al., 2012) will most likely proportionally increase the spring-time K
564 flux due to its leaching from plant litter but likely decrease the summer –time Si flux, especially in
565 the permafrost-bearing regions. The increase of vegetation density in the next decades to centuries
566 may produce a transient uptake of Si by growing vegetation in the discontinuous permafrost zone
567 during summer period. However, this potential decrease of Si export flux may be outweighed by the
568 increasing release of Si from previously frozen mineral horizons and as such the overall modification
569 of the Si concentration and riverine flux in discontinuous/continuous permafrost zone may be smaller
570 than those projected by simple latitudinal shift.

571

572 *5. Conclusions*

573 An unexpected result of the present study was rather low sensitivity of DOC, DIC, cations
574 and Si concentration and fluxes to the size of the river. The season also played a secondary role in
575 determining element concentration pattern. The most important governing parameter for

576 concentrations of dissolved river water components was the latitude, allowing us to distinguish
577 between permafrost-free, discontinuous and continuous permafrost regions. A northward decrease of
578 DIC and dissolved cation (Ca, Mg) concentrations in the WSL rivers was mostly pronounced during
579 spring flood. It was consistent with general trend of soil cations (such as Ca) concentration decrease
580 from the south to the north, reported for peat, moss and mineral layer.

581 Both seasonal and latitudinal pattern of DOC and DIC concentrations in WSL rivers are
582 consistent with previous observations that in continuous permafrost zone of frozen peat bogs, the
583 underlining mineral layer is not reactive being protected by the permafrost so that the major part of
584 the active layer is located within the organic (peat), not mineral matrix. The variation of $\delta^{13}\text{C}_{\text{DIC}}$
585 along the permafrost/latitude gradient are consistent with a progressive decrease of the groundwater
586 feeding of rivers northward, reflecting the shift of DIC origin from groundwater in the south to plant
587 litter degradation and soil respiration in surface waters north of 62°N. In winter, the $\delta^{13}\text{C}_{\text{DIC}}$ is rather
588 constant within the full latitudinal profile confirming the dominant role of carbonate/silicate mineral
589 weathering by atmospheric and soil CO_2 dissolved in the groundwaters.

590 Because the thickness of the unfrozen peat and local permafrost coverage essentially control
591 the DOC export from the soil to the river, the DOC concentration in the streams is weakly dependent
592 on the watershed size and seasons. It follows that DOC export from peat soils by medium size (<
593 100,000 km²) rivers located entirely in the permafrost zone may be higher than that of the larger
594 rivers, crossing permafrost-free regions. Assuming a short-term (hundred years) climate warming
595 scenario in the WSL, we hypothesize that the increase of the active layer thickness will be bring
596 about the involvement of upper clay horizon and sand/silts in water pathways within the soil profile.
597 As a result, the DOC export in permafrost-affected watersheds may decrease whereas the export of
598 DIC and major cations will increase. Enhanced non-stationary uptake of Si by growing vegetation in
599 the permafrost-bearing zone may attenuate expected increase of its riverine concentration linked to
600 progressive involvement of thawed mineral horizons.

601 **Acknowledgements**

602 We acknowledge support from a BIO-GEO-CLIM grant from the Ministry of Education and Science
603 of the Russian Federation and Tomsk State University (No 14.B25.31.0001) and partial support from
604 RSF grant No 15-17-10009 “Evolution of thermokarst ecosystems” awarded to LS (25%).

605

606 **Authors contribution:** O. Pokrovsky designed the study and wrote the paper; R. Manasyrov, I.
607 Kritzkov, S. Vorobyev performed sampling, analysis of major cations and their interpretation; S.
608 Vorobyev and S. Kirpotin were responsible for the choice of sampling objects and statistical
609 treatment; S. Loiko and S. Kulizhsky provided the background information on soil, peat, and
610 permafrost active layer; L. Shirokova was in charge of DOC, DIC and anion measurements and their
611 interpretation; BP performed ¹³C measurements and their interpretation; L. Kolesnichenko provided
612 GIS-based interpretation, mapping and identification of river watersheds; S. Kopysov and V.
613 Zemtsov performed all primary hydrological data collection, their analysis and interpretation. All 12
614 authors participated in field expeditions. Each co-author have seen and approved the final paper and
615 contributed to writing the manuscript.

616

617 **References**

- 618 Agren, A., Buffam, I., Jansson, M., and Laudon, H.: Importance of seasonality and small streams for
619 the landscape regulation of dissolved organic carbon export, *J. Geophys. Res.*, 112, G03003,
620 doi:10.1029/2006JG000381, 2007.
- 621 Agren, A. M., Buffam, I., Cooper, D. M., Tiwari, T., Evans, C. D., and Laudon, H.: Can the
622 heterogeneity in stream dissolved organic carbon be explained by contributing landscape
623 elements? *Biogeosciences*, 11, 1199–1213, 2014.
- 624 Anisimov, O. A., Shiklomanov, N. I., and Nelson, F. E.: Variability of seasonal thaw depth in
625 permafrost regions: A stochastic modeling approach, *Ecol. Model.*, 153, 217–227, 2002.
- 626 Anisimov, O., Kokorev, V., and Zhil'tsova, Y.: Temporal and spatial patterns of modern climatic
627 warming: Case study of Northern Eurasia, *Climatic Change*, 118, 871–883, 2013.
- 628 Anisimova, N. P.: *Cryohydrochemical Features of Permafrost Zone*, Nauka, 1981 (in Russian).
- 629 Bagard, M. L., Chabaux, F., Pokrovsky, O. S., Prokushkin, A. S., Viers, J., Dupré, B., and Stille, P.
630 Seasonal variability of element fluxes in two Central Siberian rivers draining high latitude
631 permafrost dominated areas, *Geochim. Cosmochim. Ac.*, 75, 3335–3357, 2011.
- 632 Bagard, M. L., Schmitt, A. D., Chabaux, F., Pokrovsky, O. S., Viers, J., Stille, P., Labolle, F., and

- 633 Prokushkin, A. S.: Biogeochemistry of stable Ca and radiogenic Sr isotopes in larch-covered
634 permafrost-dominated watersheds of Central Siberia, *Geochim. Cosmochim. Ac.*, 114, 169–
635 187, 2013.
- 636 Baulin, V. V.: Permanently frozen rocks of oil and gaz producing regions of the USSR, Moscow,
637 Nauka, 176 pp, 1985 (in Russian).
- 638 Baulin, V. V., Belopukhova, E. B., Dubikov, G. I., Shmelev, L. M.: Geocryological conditions of
639 western Siberia Lowland, Moscow, Nauka, 214 pp, 1967 (in Russian).
- 640 Beaulieu, E., Godderis, Y., Donnadieu, Y., Labat, D., and Roelandt, C.: High sensitivity of the
641 continental-weathering carbon dioxide sink to future climate change, *Nature Climate Change*,
642 2, 346–349, 2012.
- 643 Beilman, D. W., MacDonald, G. M., Smith, L. C., and Reimer, P. J.: Carbon accumulation in
644 peatlands of West Siberia over the last 2000 years, *Global Biogeochem. Cy.*, 23, GB1012,
645 doi:10.1029/2007GB003112, 2009.
- 646 Botch, M. S., Kobak, K. I., Vinson, T. S., and Kolchugina, T. P.: Carbon pools and accumulation in
647 peatlands of the former Soviet Union, *Global Biogeochem. Cy.*, 9, 37–46, doi:
648 10.1029/94GB03156, 1995.
- 649 Brown, J., Ferrians Jr., O. J., Heginbottom, J. A., and Melnikov, E. S.: Circum-Arctic Map of
650 Permafrost and Ground Ice Conditions. National Snow and Ice Data Center/World Data Center
651 for Glaciology, Boulder, CO, USA, Digital media, 2002.
- 652 Bokhorst, S., Bjerke, J. W., Melillo, J., Callaghan, T. V., and Phoenix, G. K.: Impacts of extreme
653 winter warming events on litter decomposition in a sub-Arctic heathland, *Soil Biol. Biochem.*,
654 42, 611–617, 2010.
- 655 Bokhorst, S., Metcalfe, D. B., and Wardle, D. A.: Reduction in snow depth negatively affects
656 decomposers but impact on decomposition rates is substrate dependent, *Soil Biol. Biochem.*,
657 62, 157–164, 2013.
- 658 Buffam, I., Laudon, H., Temnerud, J., Mörth, C.-M., and Bishop, K.: Landscape-scale variability of
659 acidity and dissolved organic carbon during spring flood in a boreal stream network, *J.*
660 *Geophys. Res.*, 112, G01022, doi:10.1029/2006JG000218, 2007.
- 661 Carey, S. K.: Dissolved organic carbon fluxes in a discontinuous permafrost subarctic alpine
662 catchment, *Permafrost Periglac.*, 14, 161–171, 2003.
- 663 Cooper, L. W., McClelland, J. W., Holmes, R. M., Raymond, P. A., Gibson, J. J., Guay, C. K., and
664 Peterson, B. J.: Flow-weighted values of runoff tracers ($\delta^{18}\text{O}$, DOC, Ba, alkalinity) from the
665 six largest Arctic rivers, *Geophys. Res. Lett.*, 35, L18606, doi:10.1029/2008GL035007, 2008.

666 Dankers, R., Burke, E. J., and Price, J.: Simulation of permafrost and seasonal thaw depth in the
667 JULES land surface scheme, *Cryosphere*, 5, 773–790, 2011.

668 Dessert, C., Dupré, B., Gaillardet, J., Francois, L. M., and Allégre, C. J.: basalt weathering laws and
669 the impact of basalt weathering on the global carbon cycle, *Chem. Geol.*, 202, 257–273, 2003.

670 Dick, J. J., Tetzlaff, D., Birkel, C., and Soulsby, C.: Modelling landscape controls on dissolved
671 organic carbon sources and fluxes to streams, *Biogeochemistry*, doi: 10.1007/s10533-014-
672 0046-3, 2014.

673 Dittmar, Th. and Kattner, G.: The biogeochemistry of the river and shelf ecosystem of the Arctic
674 Ocean: a review, *Mar. Chem.*, 83, 103–120, 2003.

675 Dornblaser, M. M. and Striegl, R. G.: Switching predominance of organic versus inorganic carbon
676 exports from an intermediate-size subarctic watershed, *Geophys. Res. Lett.*, 42, 386-394, 2015.

677 Engström, E., Rodushkin, I., Ingri, J., Baxter, D., Ecke, F., Österlund, H., and Öhlander, B.:
678 Temporal isotopic variations of dissolved silicon in a pristine boreal river, *Chem. Geol.*, 271,
679 142–152, 2010.

680 FAO, Guidelines for soil description. 4th edition. Rome. FAO, 2006.

681 Feng, X. J., Vonk, J. E., van Dongen, B. E., Gustafsson, O., Semiletov, I. P., Dudarev, O. V., Wang,
682 Z. H., Montlucon, D. B., Wacker, L., and Eglinton, T.I.: Differential mobilization of terrestrial
683 carbon pools in Eurasian Arctic river basins, *P. Natl. Acad. Sci. USA*, 110, 14168–14173,
684 2013.

685 Finlay, J. C.: Controls of streamwater dissolved inorganic carbon dynamics in a forested watershed,
686 *Biogeochemistry*, 62, 231–252, doi:10.1023/A:1021183023963, 2003.

687 Finlay, J., Neff, J., Zimov, S., Davydova, A., and Davydov, S.: Snowmelt dominance of dissolved
688 organic carbon in high-latitude watersheds: Implications for characterization and flux of river
689 DOC, *Geophys. Res. Lett.*, 33, L10401, doi:10.1029/2006GL025754, 2006.

690 Fotiev, C. M. : Taliks and their formations (Taliki and zakonomernosti ix formirovanija), In :
691 *Geocryology of the USSR, Western Siberia*, Ed : E.D. Ershov, Moscow, Nedra, pp. 72-84,
692 1989. (in Russian)

693 Fotiev, C. M. : Formation of taliks of Western Siberia, In : *Permanently frozen rocks and cryogenic*
694 *processes*, Moscow, Nauka, pp. 71-78, 1991. (in Russian).

695 Frayse, F., Pokrovsky, O. S., and Meunier, J.-D.: Experimental study of terrestrial plant litter
696 interaction with aqueous solutions, *Geochim. Cosmochim. Ac.*, 74, 70–84, 2010.

697 Frey, K. E. and Smith, L. C.: Recent temperature and precipitation increases in West Siberia and
698 their association with the Arctic Oscillation, *Polar Res.*, 22, 287–300, 2003.

699 Frey, K. E., McClelland, J. W., Holmes, R. M., and Smith, L. C.: Impacts of climate warming and

700 permafrost thaw on the riverine transport of nitrogen and phosphorus to the Kara Sea, J.
701 Geophys. Res., 112, G04S58, doi: 10.1029/2006JG000369, 2007a.

702 Frey, K. E., Siegel, D. I., and Smith, L. C.: Geochemistry of west Siberian streams and their potential
703 response to permafrost degradation, Water Resour. Res., 43, W03406,
704 doi:10.1029/2006WR004902, 2007b.

705 Frey, K. E. and Smith, L.C.: Amplified carbon release from vast West Siberian peatlands by 2100,
706 Geophys. Res. Lett., 32, L09401, doi:10.1029/2004GL022025, 2005.

707 Frey, K. E. and Smith, L. C.: How well do we know northern land cover? Comparison of four global
708 vegetation and wetland products with a new ground-truth database for West Siberia, Global
709 Biogeochem. Cy., 21, GB1016, doi:10.1029/2006GB002706, 2007.

710 Frey, K. E. and McClelland, J. W.: Impacts of permafrost degradation on arctic river
711 biogeochemistry, Hydrol. Process., 23, 169–182, 2009.

712 Fritz, P. and Fontes, J. C.: Handbook of environmental isotope geochemistry, v. 1, Elsevier, 1980.

713 Gaillardet, J., Millot, R., and Dupré, B.: Chemical denudation rates of the western Canadian orogenic
714 belt: the Stikine terrane, Chem. Geol., 201, 257–279, 2003.

715 Gebhardt, A. C., Gaye-Haake, B., Unger, D., Lahajnar, N., and Ittekkot, V.: Recent particulate
716 organic carbon and total suspended matter fluxes from the Ob and Yenisei Rivers into the Kara
717 Sea (Siberia), Mar. Geol., 207, 225–245, 2004.

718 Gentsch, N., Mikutta, R., Alves, R. J. E., Barta, J., Capek, P., Gitte, A., Hugelius, G., Kuhry, P.,
719 Lashchinskiy, N., Palmtag, J., Richter, A., Santrucková, H., Schneckner, J., Shibistova, O.,
720 Urich, T., Wild, B., and Guggenberger, G.: Storage and transformation of organic matter
721 fractions in cryoturbated permafrost soils across the Siberian Arctic, Biogeosciences Discuss.,
722 12, 2697–2743, 2015.

723 Geological composition of the USSR, v. 1, Stratigraphy, Moscow, Gostoptekhizdat, 588 pp., 1958
724 (in Russian).

725 Giesler, R., Högberg, M. N., Strobel, B. W., Richter, A., Nordgren, A., and Högberg, P.: Production
726 of dissolved organic carbon and low-molecular weight organic acids in soil solution driven by
727 recent tree photosynthate, Biogeochemistry, 84, 1–12, 2006.

728 Giesler, R., Mörth, C.-M., Karlsson, J., Lundin, E. J., Lyon, S. W., and Humborg, C.: Spatiotemporal
729 variations of pCO₂ and δ¹³C-DIC in subarctic streams in northern Sweden, Global
730 Biogeochem. Cy., 27, 176–186, doi:10.1002/gbc.20024, 2013.

731 Giesler, R., Lyon, S. W., Mörth, C.-M., Karlsson, J., Karlsson, E. M., Jantze, E. J., Destouni, G., and
732 Humborg, C.: Catchment-scale dissolved carbon concentrations and export estimates across six

733 subarctic streams in northern Sweden, *Biogeosciences*, 11, 525–537, doi:10.5194/bg-11-525-
734 2014, 2014.

735 Gislason, S. R., Arnorsson, S., and Armannsson, H.: Chemical weathering of basalt as deduced from
736 the composition of precipitation, rivers and rocks in Southwest Iceland: effect of runoff, age of
737 rocks and vegetative/glacial cover, *Am. J. Sci.*, 296, 837–907, 1996.

738 Gordeev, V. V. and Sidorov, I. S.: Concentrations of major elements and their outflow into the
739 Laptev Sea by the Lena River, *Mar. Chem.*, 43, 33–46, 1993.

740 Gordeev, V. V., Martin, J.-M., Sidorov, I. S., and Sidorova, M. V.: A reassessment of the Eurasian
741 river input of water, sediment, major elements, and nutrients to the Arctic Ocean, *Am. J. Sci.*,
742 296, 664–691, 1996.

743 Gordeev, V. V., Rachold, V., and Vlasova, I. E.: Geochemical behavior of major and trace elements
744 in suspended particulate material of the Irtysh river, the main tributary of the Ob river, Siberia,
745 *Appl. Geochem.*, 19, 593–610, 2004.

746 Gruzdov, A. V., Trofimov, V. T.: Abundance and temperature of permanently frozen and seasonally
747 frozen rocks, In: *Cryogenic composition and ice content of permanently frozen rocks of*
748 *Western Siberian Plate*, Moscow, Izd-vo Moscow State University, p. 29-57, 1980. (in
749 Russian)

750 Guo, L., Semiletov, I., Gustafsson, O., Ingri, J., Andersson, P., Dudarev, O., and White, D.:
751 Characterization of Siberian Arctic coastal sediments: Implications for terrestrial organic
752 carbon export, *Global Biogeochem. Cy.*, 18, GB1036, doi:10.1029/2003GB002087, 2004a.

753 Guo, L., Zhang, J. Z., and Guéguen, C.: Speciation and fluxes of nutrients (N, P, Si) from the upper
754 Yukon River, *Global Biogeochem. Cy.*, 18, GB1038, doi:10.1029/2003GB002152, 2004b.

755 Guo, L., Ping, C. L., and MacDonald, R.W.: Mobilization pathways of organic carbon from
756 permafrost to Arctic rivers in a changing climate, *Geophys. Res. Lett.*, 34, L13603,
757 doi:10.1029/2007GL030689, 2007.

758 Holmes, R. M., Peterson, B. J., Gordeev, V. V., Zhulidov, A. V., Meybeck, M., Lammers, R. B., and
759 Vörösmarty, C. J.: Flux of nutrients from Russian rivers to the Arctic Ocean: Can we establish
760 a baseline against which to judge future changes? *Water Resour. Res.*, 36, 2309–2320, 2000.

761 Holmes, R. M., Peterson, B. J., Zhulidov, A. V., Gordeev, V. V., Makkaveev, P. N., Stunzhas, P. A.,
762 Kosmenko, L. S., Köhler, G. H., and Shiklomanov, A. I.: Nutrient chemistry of the Ob' and
763 Yenisey Rivers, Siberia: results from June 2000 expedition and evaluation of long-term data
764 sets, *Mar. Chem.*, 75, 219–227, 2001.

765 Holmes, R. M., McClelland, J. W., Peterson, B. J., Tank, S. E., Bulygina, E., Eglinton, T. I.,
766 Gordeev, V. V., Gurtovaya, T. Y., Raymond, P. A., Repeta, D. J., Staples, R., Striegl, R. G.,

767 Zhulidov, A. V., and Zimov, S. A.: Seasonal and annual fluxes of nutrients and organic matter
768 from large rivers to the Arctic Ocean and surrounding seas, *Estuar. Coast.*, 35, 369–382,
769 doi:10.1007/s12237-011-9386-6, 2012.

770 Ilina, S. M., Drozdova, O. Y., Lapitskiy, S. A., Alekhin, Y. V., Demin, V. V., Zavgorodnyaya, Y. A.,
771 Shirokova, L. S., Viers, J., Pokrovsky, O. S.: Size fractionation and optical properties of dissolved
772 organic matter in the continuum soil solution-bog-river and terminal lake of a boreal watershed,
773 *Org. Geochem.*, 66, 14–24, 2014.

774 Ivanov, Yu. K. and Beshentsev, V. A.: Paleo-geographical aspects of formation of chemical
775 composition of fresh underground waters in Yamalo-Nenets region, *Lithosphere*, No 4, 188–196,
776 2005.

777 Izerskaia, L. A., Vorobyev, S. N, Vorobyeva, T. E., Kolesnichenko, L. G, Zakharchenko, A. V.: The
778 concentration of Mn, Cu, Zn, Co, B, Sr, Cd and Pb in alluvial soils of the Ob River (forest-steppe,
779 southern taiga and central taiga), *Internat. J. Environ. Stud.*, 71(5), 691–697, 2014.

780 Jantze, E. J., Lyon, S. W., and Destouni, G.: Subsurface release and transport of dissolved carbon in a
781 discontinuous permafrost region, *Hydrol. Earth Syst. Sc.*, 17, 3827–3839, doi:10.5194/hess-17-
782 3827-2013, 2013.

783 Kaiser, C., Meyer, H., Biasi, C., Rusalimova, O., Barsukov, P., and Richter, A.: Conservation of soil
784 organic matter through cryoturbation in arctic soils in Siberia, *J. Geophys. Res.*, 112, 9–17, 2007.

785 Karlsson, J. M., Jaramillo, F., and Destouni, G.: Hydro-climatic and lake change patterns in Arctic
786 permafrost and non-permafrost areas, *J. Hydrol.*, in press, 2015.

787 Kawahigashi, M., Kaiser, K., Kalbitz, K., Rodionov, A., and Guggenberger, G.: Dissolved organic
788 matter in small streams along a gradient from discontinuous to continuous permafrost, *Glob.
789 Change Biol.*, 10, 1576–1586, doi:10.1111/j.1365-2486.2004.00827.x, 2004.

790 Kawahigashi, M., Kaiser, K., Rodionov, A., and Guggenberger, G.: Sorption of dissolved organic
791 matter by mineral soils of the Siberian forest tundra. *Global Change Biol.*, 12, 1868–1877,
792 doi:10.1111/j.1365-2486.2006.01203.x, 2006.

793 Keller, K., Blum, J. D., and Kling, G. W.: Stream geochemistry as an indicator of increasing
794 permafrost thaw depth in an arctic watershed, *Chem. Geol.*, 273, 76–81, 2010.

795 Kirdyanov, A. V., Hagedorn, F., Knorre, A. A., Fedotova, E. V., Vaganov, E. A., Naurzbaev, M. M.,
796 Moiseev, P. A., and Rigling, A.: 20th century tree-line advance and vegetation changes along
797 an altitudinal transect in Putorana Mountains, northern Siberia, *Boreas*, 41, 56–67, 2012.

798 Khrenov V. Ya.: Soils of cryolithozone of western Siberia: Morphology, physico-chemical properties
799 and geochemistry, Nauka, Moscow, 214 pp., 2011 (in Russian).

800 Kremenetski, K. V., Velichko, A. A., Borisova, O. K., MacDonald, G. M., Smith, L. C., Frey, K. E.,
801 and Orlova, L. A.: Peatlands of the West Siberian Lowlands: Current knowledge on zonation,
802 carbon content, and Late Quaternary history, *Quaternary Sci. Rev.*, 22, 703–723, 2003.

803 Kuglerová, L., Jansson, R., Ågren, A., Laudon, H., and Malm-Renöfält, B.: Groundwater discharge
804 creates hotspots of riparian plant species richness in a boreal forest stream network, *Ecology*
805 95, 715–725. <http://dx.doi.org/10.1890/13-0363.1>, 2014.

806 Laudon, H., Köhler, S., and Buffam, I.: Seasonal TOC export from seven boreal catchments in
807 northern Sweden, *Aquat. Sci.*, 66, 223–230, 2004.

808 Liss, O. L., Abramova, L. I., Avetov, N. A., Berezina, N. A., Inisheva, L. I., Kurnishnikova, T. V.,
809 Sluka, Z. A., Tolpysheva, T. Yu., and Shvedchikova, N. K.: Wetland systems of West Siberia
810 and their importance for nature conservation, Grif i K publisher, Tula, Russia, 584 pp., 2001,
811 ISBN 5-8125-0150-5 (in Russian).

812 Lobbes, J. M., Fitznar, H. P., and Kattner, G.: Biogeochemical characteristics of dissolved and
813 particulate organic matter in Russian rivers entering the Arctic Ocean, *Geochim. Cosmochim.*
814 *Ac.*, 64, 2973–2983, 2000.

815 Loiko, S. V., Geras'ko, L. I., Kulizhskii, S. P., Amelin, I. I., Istigechev, G. I.: Soil cover patterns in the
816 northern part of the area of aspen-fir taiga in the southeast of Western Siberia, *Eurasian Soil Sci.*,
817 48(4), 359–372, DOI: 10.1134/S1064229315040067, 2015.

818 Manasyrov, R. M., Pokrovsky, O. S., Kirpotin, S. N., and Shirokova, L. S.: Thermokarst lake waters
819 across the permafrost zones of western Siberia, *Cryosphere*, 8, 1177–1193, doi:10.5194/tc-8-
820 1177-2014, 2014.

821 Manasyrov, R. M., Vorobyev, S. N., Loiko, S. V., Kritzkov, I. V., Shirokova, L. S., Shevchenko, V. P.,
822 Kirpotin, S. N., Kulizhsky, S. P., Kolesnichenko, L. G., Zemtsov, V. A., Sinkinov, V. V., and
823 Pokrovsky, O. S.: Seasonal dynamics of organic carbon and metals in thermokarst lakes from the
824 discontinuous permafrost zone of western Siberia, *Biogeosciences*, 12, 3009–3028,
825 doi:10.5194/bg-12-3009-2015, 2015.

826 Mavromatis, V., Prokushkin, A. S., Pokrovsky, O. S., Viers, J., and Korets, M. A.: Magnesium isotopes
827 in permafrost-dominated Central Siberian larch forest watersheds, *Geochim. Cosmochim. Ac.*,
828 147, 76–89, 2014.

829 McClelland, J. W., Déry, S. J., Peterson, B. J., Holmes, R. M., and Wood, E. F.: A pan-Arctic evaluation
830 of changes in river discharge during the latter half of the 20th century, *Geophys. Res. Lett.*, 33,
831 L06715, 2006.

832 Mergelov, N. and Targulian, V.: Accumulation of organic matter in the mineral layers of permafrost-
833 affected soils of coastal lowlands in East Siberia, *Eurasian Soil Sci.*, 44, 249–260, 2011.

834 Methodical recommendations for determination of hydrological parameters under insufficient number
835 of observations, Rota-print GNS AANII, St. Petersburg, 66 pp., 2007 (in Russian).

836 Michel, F. A. and Vaneverdingen, R. O.: Changes in hydrologic regimes in permafrost regions due to
837 climate change, *Permafrost Periglac. Proc.*, 5, 191–195, 1994.

838 Millot, R., Gaillardet, J., Dupré, B., and Allègre, C. J.: Northern latitude chemical weathering rates:
839 Clues from the Mackenzie River Basin, Canada, *Geochim. Cosmochim. Ac.*, 67, 1305–1329,
840 2003.

841 Moran, S. B. and Woods, W. L.: Cd, Cr, Cu, Ni and Pb in the water column and sediments of the Ob-
842 Irtysh Rivers, Russia, *Mar. Pollut. Bull.*, 35, 270–279, 1997.

843 Nikanorov, A. M., Bryzgalo, V. A., Kosmenko, L. S., and Reshetnyak, O. S.: The role of chemical
844 river runoff in the anthropogenic transformation of the state of the aquatic environment in the
845 Enisei mouth area, *Water Resour.*, 37, 471–480, 2010a.

846 Nikanorov, A. M., Smirnov, M. P., and Klimenko, O. A.: Long-term trends in total and
847 anthropogenic discharge of organic and biogenic substances by Russian rivers into the Arctic
848 and Pacific Seas, *Water Resour.*, 37, 361–371, 2010b.

849 Nikitin, S. P. and Zemtsov, V. A.: The variability of hydrological parameters of western Siberia,
850 Nauka, Novosibirsk, 204 pp., 1986 (in Russian).

851 Novikov, S. M., Moskvina, Y. P., Trofimov, S. A., Usova, L. I., Batuev, V. I., Tumanovskaya, S. M.,
852 Smirnova, V. P., Markov, M. L., Korotkevich, A. E., and Potapova, T. M.: Hydrology of bog
853 territories of the permafrost zone of western Siberia, *BBM publ. House*, St. Petersburg, 535
854 pp., 2009 (in Russian).

855 Olefeldt, D., and Roulet, N. T.: Effects of permafrost and hydrology on the composition and
856 transport of dissolved organic carbon in a subarctic peatland complex, *J. Geophys. Res.*, 117,
857 G01005, doi:10.1029/2011JG001819, 2012.

858 Olefeldt, D., Persson, A., and Turetsky, M. R.: Influence of the permafrost boundary on dissolved
859 organic matter characteristics in rivers within the Boreal and Taiga plains of western Canada,
860 *Environ. Res. Lett.*, 9, Art No 035005, doi:10.1088/1748-9326/9/3/035005, 2014.

861 Oliva, P., Viers, J., and Dupré, B.: Chemical weathering in granitic environments, *Chem. Geol.*, 202,
862 225–256, 2003.

863 Oosterwoud, M. R., Temminghoff, E. J. M., and van der Zee, S. E. A. T. M.: Quantification of DOC
864 concentrations in relation with soil properties of soils in tundra and taiga of Northern European
865 Russia, *Biogeosciences Discuss.*, 7, 3189–3226. 2010.

866 Pavlov, A. V. and Moskalenko, N. G.: The thermal regime of soils in the north of Western Siberia,
867 *Permafrost Periglac.*, 13, 43–51, doi:10.1002/ppp.409, 2002.

868 Peterson, B. J., Holmes, R. M., McClelland, J. W., Vorosmarty, C. J., Lammers, R. B., Shiklomanov,
869 A. I., Shiklomanov, I. A., and Rahmstorf, S.: Increasing river discharge to the Arctic Ocean,
870 Science, 298, 2171–2173, 2002.

871 Petrone, K. C., Jones, J. B., Hinzman, L. D., and Boone, R. D.: Seasonal export of carbon, nitrogen,
872 and major solutes from Alaskan catchments with discontinuous permafrost, J. Geophys. Res.,
873 111, G02020, doi:10.1029/2005JG000055, 2006.

874 Pokrovsky, O. S., Schott, J., Kudryavtzev, D. I., and Dupré, B.: Basalts weathering in Central Siberia
875 under permafrost conditions, Geochim. Cosmochim. Ac., 69, 5659–5680, 2005.

876 Pokrovsky, O. S., Schott, J., and Dupre, B.: Trace element fractionation and transport in boreal rivers
877 and soil porewaters of permafrost-dominated basaltic terrain in Central Siberia, Geochim.
878 Cosmochim. Ac., 70, 3239–3260, 2006.

879 Pokrovsky, O. S., Viers, J., Shirokova, L. S., Shevchenko, V. P., Filipov, A. S., and Dupré, B.:
880 Dissolved, suspended, and colloidal fluxes of organic carbon, major and trace elements in
881 Severnaya Dvina River and its tributary, Chem. Geol., 273, 136–149, 2010.

882 Pokrovsky, O. S., Viers, J., Dupré, B., Chabaux, F., Gaillardet, J., Audry, S., Prokushkin, A. S.,
883 Shirokova, L. S., Kirpotin, S. N., Lapitsky, S. A., and Shevchenko, V. P.: Biogeochemistry of
884 carbon, major and trace elements in watersheds of Northern Eurasia drained to the Arctic Ocean:
885 The change of fluxes, sources and mechanisms under the climate warming prospective, C.R.
886 Geosci., 344, 663–677, 2012.

887 Pokrovsky, O. S., Reynolds, B. C., Prokushkin, A. S., Schott, J., and Viers, J.: Silicon isotope variations
888 in Central Siberian rivers during basalt weathering in permafrost-dominated larch forests, Chem.
889 Geol., 355, 103–116, 2013a.

890 Pokrovsky, O. S., Shirokova, L. S., Kirpotin, S. N., Kulizhsky, S. P., and Vorobiev, S. N.: Impact of
891 western Siberia heat wave 2012 on greenhouse gases and trace metal concentration in thaw lakes
892 of discontinuous permafrost zone, Biogeosciences, 10, 5349–5365, 2013b.

893 Porcal, P., Koprivnjak, J.-F., Molot, L. A., and Dillon, P. J.: Humic substaces – part 7: the
894 biogeochemistry of dissolved organic carbon and its interactions with climate change, Environ.
895 Sci. Pollut. R., 16, 714–726, 2009.

896 Prokushkin, A. S., Pokrovsky, O. S., Shirokova, L. S., Korets, M. A., Viers, J., Prokushkin, S. G.,
897 Amon, R., Guggenberger, G., and McDowell, W. H.: Sources and export fluxes of dissolved
898 carbon in rivers draining larch-dominated basins of the Central Siberian Plateau, Environ. Res.
899 Lett., 6, 045212, doi:10.1088/1748-9326/6/4/045212, 2011.

900 Rember, R. D. and Trefry, J. H.: Increased concentrations of dissolved trace metals and organic
901 carbon during snowmelt in rivers of the Alaskan Arctic, *Geochim. Cosmochim. Ac.*, 68, 477–
902 489, 2004.

903 Rinta, P., Bastviken, D., van Hardenbroek, M., Kankaala, P., Leuenberger, M., Schilder, J., Stötter,
904 T., and Heiri, O.: An inter-regional assessment of concentrations and $\delta^{13}\text{C}$ values of methane
905 and dissolved inorganic carbon in small European lakes, *Aquat Sci.*, doi:10.1007/s00027-015-
906 0410-y, 2015.

907 Romanovsky, N.N. : *Underground waters of the cryolithozone*, Moscow, Moscow State University
908 Publ. House, 232 pp, 1983.

909 Romanovsky, V.E., Drozdov, D.S. Oberman, N.G., Malkova G.V., Kholodov A.L., Marchenko, S.S.,
910 Moskalenko, N.G., Sergeev D.O., Ukraintseva, N.G., Abramov A.A., Gilichinsky, D.A., and
911 Vasiliev, A.A.: Thermal State of Permafrost in Russia, *Permafrost Periglacial Proc.*, 21, 136-
912 155, 2010.

913 Serreze, M. C., Walsh, J. E., Chapin, E., Osterkamp, T., Dyugero, M., Romanovsky, V., Oechel, W.
914 C., Morison, J., Zhang, T., and Barry, R. G.: Observation evidence of recent change in the
915 northern high-latitude environment, *Climatic Change*, 46, 159–207, 2000.

916 Shirokova, L. S., Pokrovsky, O. S., Kirpotin, S. N., Desmukh, C., Pokrovsky, B. G., Audry, S., and
917 Viers, J.: Biogeochemistry of organic carbon, CO_2 , CH_4 , and trace elements in thermokarst
918 water bodies in discontinuous permafrost zones of Western Siberia, *Biogeochemistry*, 113,
919 573–593, 2013.

920 Smith, L. C., Macdonald, G. M., Velichko, A. A., Beilman, D. W., Borisova, O. K., Frey, K. E.,
921 Kremenetsky, K. V., and Sheng, Y.: Siberian peatlands as a net carbon sink and global methane
922 source since the early Holocene, *Science*, 303, 353–356, 2004.

923 Smedberg, E., Mörth, C. M., Swaney, D. P., and Humborg, C.: Modeling hydrology and silicon-
924 carbon interactions in taiga and tundra biomes from a landscape perspective: Implications for
925 global warming feedback, *Global Biogeochem. Cy.*, 20, GB2014, doi:10.1029/2005GB002567,
926 2006.

927 Stendel, M. and Christensen, J. H.: Impact of global warming on permafrost conditions in a coupled
928 GCM, *Geophys. Res. Lett.*, 29, 10-1–10-4, 2002.

929 Stepanova, V. M., Pokrovsky, O. S., Viers, J., Mironycheva-Tokareva, N. P. Kosykh, N. P., and
930 Vishnyakova, E. K.: Major and trace elements in peat profiles in Western Siberia: impact of
931 the landscape context, latitude and permafrost coverage, *Appl. Geochem.*, 53, 53–70, 2015.

932 Striegl, R. B., Kortelainen, P., Chanton, J. P., Wickland, K. P., Bugna, G. C., and Rantakari, M.:
933 Carbon dioxide partial pressure and ^{13}C content of north temperate and boreal lakes at spring
934 ice melt, *Limnol. Oceanogr.*, 46, 941–945, 2001.

935 Striegl, R. G., Aiken, G. R., Dornblaser, M. M., Raymond, P. A., and Wickland, K. P.: A decrease in
936 discharge-normalized DOC export by the Yukon River during summer through autumn,
937 *Geophys. Res. Lett.*, 32, L21413, doi:10.1029/2005GL024413, 2005.

938 Sturm, M., Racine, C., and Tape, K.: Increasing shrub abundance in the Arctic, *Nature*, 411, 546–
939 547, 2001.

940 Svod pravil SP 33-101-2003. Determination of basic calculated hydrological parameters. Official
941 release, Gosstroy, Moscow, Russia, 73 pp., 2004.

942 Tank, S. E., Raymond, P. A., Striegl, R. G., McClelland, J. W., Holmes, R. M., Fiske, G. J., and
943 Peterson, B. J.: A land-to-ocean perspective on the magnitude, source and implication of DIC
944 flux from major Arctic rivers to the Arctic Ocean, *Global Biogeochem. Cy.*, 26, GB4018,
945 doi:10.1029/2011GB004192, 2012.

946 Tank, S. E., Frey, K. E., Striegl, R. G., Raymond, P. A., Holmes, R. M., McClelland, J. W., and
947 Peterson, B. J.: Landscape-level controls on dissolved carbon flux from diverse catchments of
948 the circumboreal, *Global Biogeochem. Cy.*, 26, GB0E02, doi:10.1029/2012GB004299, 2012.

949 Tape, K., Sturm, M., and Racine, C.: The evidence for shrub expansion in Northern Alaska and the
950 Pan-Arctic, *Glob. Change Biol.*, 12, 686–702, doi:10.1111/j.1365-2486.2006.01128.x., 2006.

951 Targulian, V. O.: Soil Formation and Weathering in Cold Humid Regions (On Massive-Crystalline
952 and Sandy Polymictic Rocks), Nauka, Moscow, 1971 (in Russian).

953 Tyrtikov, A. P.: Thawing of soils in tundra of western Siberia, in: Popov A.I. (Ed.) Natural
954 environment of western Siberia, Issue 3, Izd-vo MG, Moscow, 160–169, 1973 (in Russian).

955 Tyrtikov, A. P.: Dynamics of vegetation coverage and permafrost development in western Siberia,
956 Moscow, Nauka, 114 pp, 1979 (in Russian).

957 van Hees, P.A.W., Jones, D.L., Finlay, R., Godbold, D.L., Lundstrom, U.S.: 2005. The carbon we do
958 not see—the impact of low molecular weight compounds on carbon dynamics and respiration
959 in forest soils: a review, *Soil Biol Biochem.*, 37, 1–13, 2005.

960 Vasiliev, A. A., Streletskaia, I. D., Shirokov, R. S., and Oblogov, G. E.: Evolution of cryolithozone
961 of coastal zone of western Yamal during climate change, *Kriosfera Zemli*, 2, 56–64, 2011 (in
962 Russian).

963 Vorobyev, S. N., Pokrovsky, O. S., Kirpotin, S. N., Kolesnichenko, L. G., Shirokova, L. S.,
964 Manasyrov, R. M.: Flood zone biogeochemistry of the Ob' River middle course, *Applied*
965 *Geochem.*, 63, 133-145, 2015.

- 966 Voronkov, P. P., Sokolova, O. K., Zubareva, V. I., and Naidenova, V. I.: Hydrochemical features of
967 local discharge during spring flood from the soil coverage of European territory of the USSR,
968 Trudy GGI (Proceedings of State Hydrological Institute), 137, 3–57, 1966 (in Russian).
- 969 Voss, B. M., Peucker-Ehrenbrink, B., Eglinton, T. I., Spencer, R. G. M., Bulygina, E., Galy, V.,
970 Lamborg, C. H., Ganguli, P. M., Montluçon, D. B., Marsh, S., Gillies, S. L., Fanslau, J., Epp,
971 A., and Luymes, R.: Seasonal hydrology drives rapid shifts in the flux and composition of
972 dissolved and particulate organic carbon and major and trace ions in the Fraser River, Canada,
973 *Biogeosciences*, 12, 5597–5618, doi:10.5194/bg-12-5597-2015, 2015.
- 974 Waldron, S., Scott, M. E., and Soulsby, C.: Stable isotope analysis reveals lower-order river
975 dissolved inorganic carbon pools are highly dynamic, *Environ. Sci. Technol.*, 41, 6156–6162,
976 doi:10.1021/es0706089, 2007.
- 977 Walvoord, M. A., Voss, C. I., and Wellman, T. P.: Influence of permafrost distribution on
978 groundwater flow in the context of climate-driven permafrost thaw: Example from Yukon Flats
979 Basin, Alaska, United States, *Water Res. Research*, 48, W07524, doi:
980 10.1029/2011WR011595, 2012.
- 981 Walvoord, M. A. and Striegl, R. G.: Increased groundwater to stream discharge from permafrost
982 thawing in the Yukon River basin: Potential impacts on lateral export of carbon and nitrogen, *J.*
983 *Geophys. Res.*, 34, L12402, doi:10.1029/2007GL030216, 2007.
- 984 White, A. F. and Blum, A. E.: Effects of climate on chemical weathering in watersheds, *Geochim.*
985 *Cosmochim. Ac.*, 59, 1729–1747, 1995.
- 986 White, D., Hinzman, L., Alessa, L., Cassano, J., Chambers, M., Falkner, K., Francis, J., Gutowski Jr.,
987 W. J., Holland, M., Holmes, R. M., Huntington, H., Kane, D., Kliskey, A., Lee, C.,
988 McClelland, J., Peterson, B., Rupp, T. S., Straneo, F., Steele, M., Woodgate, R., Yang, D.,
989 Yoshikawa, K., and Zhang T.: The arctic freshwater system: Changes and impacts, *J. Geophys.*
990 *Res.*, 112, G04S54, doi:10.1029/2006JG000353, 2007.
- 991 Wickland, K. P., Neff, J. C., Aiken, G. R.: Dissolved organic carbon in Alaskan boreal forest:
992 sources, chemical characteristics, and biodegradability, *Ecosystems*, doi: 10.1007/s10021-
993 007-9101-4, 2007.
- 994 Worrall, F., Harriman, R., Evans, C. D., Watts, C. D., Adamson, J., Neal, C., Tipping, E., Burt, T.,
995 Grieve, I., Monteith, D., Naden, P. S., Nisbet, T., Reynolds, B., and Stevens, P.: Trends in
996 dissolved organic carbon in UK rivers and lakes, *Biogeochemistry*, 70, 369–402, 2004.
- 997 Yang, D., Ye, B., and Shiklomanov, A.: Discharge characteristics and changes over the Ob River
998 watershed in Siberia, *J. Hydrometeorol.*, 5, 595–610, 2004.

999 Ye, B., Yang, D., Zhang, Z., and Kane, D. L.: Variation of hydrological regime with permafrost
1000 coverage over Lena basin in Siberia, *J. Geophys. Res.*, 114, D07102, 2009.

1001 Yeghicheyan, D., Bossy, C., Bouhnik Le Coz, M., Douchet, Ch., Granier, G., Heimbürger, A.,
1002 Lacan, F., Lanzanova, A., Rousseau, T. C. C., Seidel, J.-L., Tharaud, M., Candaudap, F.,
1003 Chmeleff, J., Cloquet, C., Delpoux, S., Labatut, M., Losno, R., Pradoux, C., Sivry, Y., and
1004 Sonke, J. E.: A Compilation of Silicon, Rare Earth Element and Twenty-One other Trace
1005 Element Concentrations in the Natural River Water Reference Material SLRS-5 (NRC-CNRC),
1006 *Geostand. Geoanal. Res.*, 37, 449–467, doi:10.1111/j.1751-908X.2013.00232.x, 2013.

1007 Zakharova, E. A., Pokrovsky, O. S., Dupré, B., and Zaslavskaya, M. B.: Chemical weathering of
1008 silicate rocks in Aldan Shield and Baikal Uplift: insights from long-term seasonal
1009 measurements of solute fluxes in rivers, *Chem. Geol.*, 214, 223–248, 2005.

1010 Zakharova, E. A., Pokrovsky, O. S., Dupré, B., Gaillardet, J., and Efimova, L.: Chemical weathering
1011 of silicate rocks in Karelia region and Kola peninsula, NW Russia: Assessing the effect of rock
1012 composition, wetlands and vegetation, *Chem. Geol.*, 242, 255–277, 2007.

1013 Zakharova, E. A., Kouraev, A. V., Rémy, F., Zemtsov, V. A., and Kirpotin, S.N.: Seasonal
1014 variability of the Western Siberia wetlands from satellite radar altimetry, *J Hydrol.*, 512, 366-
1015 378, 2014. DOI: <http://dx.doi.org/10.1016/j.jhydrol.2014.03.002>

1016
1017
1018
1019

Table 1. List of sampled rivers, their watershed area and annual runoff. The codes under the months identify the sampling sites listed in Table S1. The annual runoff was calculated following the approach of Frey et al (2007b) as explained in Supplement 1.

Season				N	E	River	watersheds, km ²	Annual runoff, mm/y	Nuber on the map
June	August	October	February						
RJ-22			RF13	61°29'11.1"	74°09'42.9"	Vach-Yagun	1.79	192	41
RJ-21			RF54	61°29'46.6"	74°15'30.3"	Segut-Yagun	3.37	192	40
	BL-35			60°44'10,9"	77°22'55,9"	Medvedka	7	173	27
RJ-20	RA-18		RF55	61°27'17.3"	74°40'23.3"	Kottym'egan	7.18	192	38
RJ-39			RF31	65°06'48.8"	77°47'58.8"	Tydylyakha	7.46	185	81
RJ-2		R-2		56°43'15.0"	83°55'35.1"	Chybyr'	8.14	44.8	3
	BL-28			61°12'19,5"	75°23'06,5"	Er-Yakh	9.35	173	35
RJ-24			RF52	61°50'28.6"	70°50'28.2"	Vachinguriyagun	9.52	192	42
RJ-56	BL-25		RF48	62°37'08.4"	74°10'15.9"	Petriyagun	9.65	192	47
	RA-6			63°12'43.38"	76°21'27.66"	Goensapur	11	194	60
RJ-40	RY 14-45		RF30	65°12'17.6"	77°43'49.8"	Tydyotta	12.0	309	82
RJ-12		R-12		58°24'38.0"	82°08'46.0"	Istok	12.3	127	14
	RA-5			63°12'45.96"	76°24'1.32"	Denna	15	194	61
RJ-36	RY 14-47		RF33	64°32'07.9"	76°54'21.3"	Seryareyakha	15.2	186	78
RJ-31				64°09'06.4"	75°22'18.1"	Apoku-Yakha	18.8	186	73
	RA-12			63°9'31.38"	75°3'2.58"	Ponto-Yakha	19	194	54
RJ-25			RF14	61°58'05.1"	73°47'03.4"	Lyukh-Yagun	21.6	192	43
RJ-55	BL-23		RF46	62°43'09.9"	74°13'45.9"	Ai-Kirill-Vys'yagun	24.0	192	46
	BL-34			60°45'58,5"	77°26'12,6"	Saim	26	173	28
	BL-31			60°50'43,6"	77°05'03,0"	Kaima	31	173	31
	BL-3			56°54'39,1"	82°33'33,3"	Cherniy Klyuch	32	168	6
	BL-33			60°47'29,3"	77°19'13,5"	Mishkin Saim	32	173	29
	RA-15			63°8'34.02"	74°54'29.1"	Nyudya-Itu-Yakha	32	194	55
RJ-57			RF49	62°33'39.8"	74°00'29.5"	Pintyr'yagun	33.5	192	46
RJ-52;	BL-19		RF39	63°36'48.2"	74°35'28.6"	Khatytayakha	34.6	194	64

Season				N	E	River	watersheds, km ²	Annual runoff, mm/y	Nuber on the map
June	August	October	February						
R-4									
RJ-46			RF26	65°59'05.7"	77°40'52.6"	Tadym-Yakha	39.9	185	88
RJ-45			RF25	65°58'54"	77°34'05"	Yude-Yakha	42.4	185	87
	RA-9			63°13'25.2"	76°5'23.04"	Tlyatsayakha	43	194	51
	BL-32			60°49'32,3"	77°13 46,3"	Alenkin Egan	44	173	30
RJ-35			RF34	64°26'05.2"	76°24'37.0"	Khav' -Yakha	46.4	186	77
	RY 14-48			64°23'30,6"	76°19'50,1"	Khaloku-Yakha	53	186	76
RJ-11			RF10	58°23'16.8"	82°11'39.0"	Tatarkin Istok	58.6	33.4	13
RJ-32				64°12'08.4"	75°24'28.4"	Ngarka-Tyde-Yakha	59.9	186	96
RJ-3		R-3	RF2	56°46'19.5"	83°57'35.7"	Prud	61.5	44.8	2
RJ-30			RF36	64°06'50.7"	75°14'17.3"	Ngarka-Varka-Yakha	67.1	186	72
RJ-33	RY 14-49		RF35	64°17'31.9"	75°44'33.4"	Etu-Yakha	71.6	186	74
RJ-50; R-3	BL-17		RF41	63°49'58,0"	74°39'02,5"	Khanupiyakha	74	194	66
RJ-41	RY 14-44		RF29	65°23'34.1"	77°45'46.7"	Ponie-Yakha	78.9	185	83
	RA-4			63°10'4.68"	76°28'19.08"	Nyudya-Pidya-Yakha	79.5	194	53
	BL-22		RF45	63°11'19,3"	74°36'25,5"	Pyrya-Yakha	82	194	50
	RA-10			63°13'12.06"	75°38'52.26"	Yangayakha	88	194	56
	RA-13			63°10'3.48"	74°45'16.32"	Nekhtyn-Pryn	96	194	52
RJ-34				64°19'10.1"	76°08'26.7"	Varka-Yakha	105	186	75
RJ-43			RF27	65°47'48.6"	78°10'09.0"	Almayakha	106	185	86
RJ-27; Z-86	BL-14; RA-1		RF44	63°47'04.5"	75°37'06.8"	Lymbyd'yakha	115	194	69
RJ-58	BL-5	R-15	RF65	58°40'46.5"	84°27'56.6"	Vyalovka	117	127	20
	RA-8			63°13'3.66"	76°15'24.6"	Chukusamal	121	194	50
	RT2 14-30			66°59'20,9"	79°22'30,5"	Malokha Yakha	157	208	92
	RA-11			63°9'39.84"	75°09'10.86"	Velykh-Pelykh-Yakha	170	194	55

Season				N	E	River	watersheds, km ²	Annual runoff, mm/y	Nuber on the map
June	August	October	February						
RJ-53; R-5	RA-16; BL-21		RF38	63°22'01.6"	74°31'53.2"	Kamgayakha	175	194	63
RJ-8		R-8	RF7	57°52'26.8"	83°11'29.9"	Chemondaevka	177	63.4	10
	RY 14-42			65°46'34,5"	78°08'25,8"	Khiroyakha	183	185	85
			RF62	59°41'01,6"	77°44'33,9"	Kornilovskaya	190	133	22
			RF60	60°08'43"	77°16'53"	Koltogorka	220	155.4	24
	RA-14			63°11'40.68"	74°38'16.92"	Itu-Yakha	250	194	51
			RF61	59°44'09,2"	77°26'06"	Levyi Il'yas	253	133	23
RJ-9		R-9	RF8	57°58'45.7"	82°58'32.2"	Sugotka	275	63.4	11
RJ-51	BL-18		RF40	63°40'41.8"	74°35'20.7"	Pulpuyakha	281	194	65
RJ-6			RF5	57°36'43.3"	83°37'02.1"	Malyi Tatosh	302	63.4	8
RJ-5		R-5	RF4	57°19'20.7"	83°55'53.8"	Brovka	320	63.4	5
RJ-18	RA-19; BL-27			61°19'41.2"	75°04'0.3"	Ur'evskii Egan	359	272	36
	BL-9			58°32'05,8"	80°51'26,8"	Karza	473	148	17
	BL-6			58°37'29,9"	81°06'09,0"	Sochiga	510	148	18
RJ-49	RT2 14-32			65°59'14.7"	78°32'25.2"	Malaya Khadyr-Yakha	512	278	90
RJ-54; R-6	BL-24		RF47	63°38'23,4"	74°10'52"	Kirill-Vys'yagun	598	225	48
	RT2 14-29			67°10'54,8"	78°51'04,5"	Nuny-Yakha	656	312	93
RJ-14		R-14	RF12	58°33'03.1"	81°48'44.3"	Chigas	689	180	16
			RF58	60°30'19"	76°58'57"	Sosninskii Yegan	732	199	25
RJ-29; R-2	BL-16		RF42; RF37	63°51'23.4"	75°08'05.6"	Kharucheyakha	820	292	63
RJ-7		R-7	RF6	57°37'17.3"	83°31'53.3"	Bolshoy Tatosh	1020	74.6	9
RJ-23			RF53	61°34'27.4"	77°46'35.4"	Mokhovaya	1260	192.3	33
	BL-13			63°43'37,9"	75°59'04,1"	Chuchi-Yakha	1396	292	66
			RF51	61°59'39"	73°47'39"	Limpas	1648	320	44
RJ-48	RT2 14-31			66°17'10.8"	79°15'06.1"	Ngarka Khadyta-Yakha	1970	277	91

Season				N	E	River	watersheds, km ²	Annual runoff, mm/y	Nuber on the map
June	August	October	February						
	RA-3; RA-7			63°46'22.92"	76°25'28.86"	Vyngapur	1979	324	59
RJ-17	BL-29; RA-20		RF57	61°11'52.7"	75°25'20.2"	Vatinsky Egan	3190	287	34
	BL-2			57°02'23,75"	82°04'02,44"	Bakchar	3197	96.1	7
RJ-13	RA-22	R-13	RF11	58°26'06.9"	82°05'43.6"	Shudelka	3460	211	15
RJ-42	RY 14-43		RF28	65°41'51.1"	78°01'05.0"	Yamsovey	4030	309	84
RJ-37	RY 14-46		RF32	64°40'14.0"	77°05'27.2"	Purpe	5110	309	79
			RF21	67°24'39"	76°21'12"	Khadutte	5190	346	95
R-1; Z-55; RJ-28	BL-20; RA-2; BL-15		RF43	63°49'54,2"	75°22'47,1"	Pyakupur	9880	324	64
RJ-26; R-7; R-8	RA-17		RF50	62°07'50,0"	73°44'05,6"	Tromyegan	10770	263	45
RJ-4; R-10		R-4	RF3	57°06'39.2"	83°54'41.1"	Shegarka	12000	58.3	4
RJ-15	RA-21; BL-4	R-17	RF64	58°42'34.5"	81°22'22.0"	Parabel	25500	131	19
RJ-38				64°55'55.1"	77°56'08.2 "	Aivasedapur	26100	309	80
RJ-10	RA-23		RF9	58°04'20.8"	82°49'19.7"	Chaya	27200	95.6	12
RJ-19; R-9	BL-26		RF56	61°26'13,6"	74°47'39,7"	Agan	27600	291	37
			RF63	58°59'37"	80°34'00"	Vasyugan	63780	177	21
	BL-30			60°55'41,0"	76°53'49,3"	Vakh	75090	298	32
RJ-44	RY 14-41			65°57'05.5"	78°18'59.1"	Pur	112000	298	89
RJ-47	RT2 14-40			67°22'13.28"	79°00'25,9"	Taz	150000	330	94
RJ-16	BL-36			60°40'28.8"	77°31'29.4"	Ob'	773200	216	26
RJ-1		R-1	RF1	56°31'48"	84°09'44"	Ob'	423100	207	1




Fig.1. Map of the study site with permafrost boundaries (Brown et al., 2002; <http://portal.inter-map.com> (NSIDC)), runoff contour lines (Nikitin and Zemtsov, 1986) and sampling points along the latitudinal transect of rivers Ob, Pur and Taz basin. The numbers of the sampling sites are listed in Table 1.







Fig. 1, continued. Detailed map of the four rectangles in Fig 1 a.




Figure 2. Decrease of river water pH with the increase of the latitude during winter (A), spring (B) and summer (C). The spring acid pulse is seen only in permafrost-affected rivers north of 62°N (B), and the scatter of the values is maximal during summer (C). The variability among different watershed size is smaller than that between the seasons and within the latitude gradient. Diamonds, squares, triangles and circles represent watershed of size < 100 km², 100 to 1000 km², 1000 to 10,000 km², and > 10,000 km², respectively.




Figure 3. Decrease of DOC with latitude during winter (A), spring (B) and summer (C). The latitudinal trend is significant at $p < 0.05$. Considering all seasons together, the differences between different watershed sizes are not statistically significant ($p > 0.05$). The symbols are the same as in Fig. 2.




Figure 4. Significant decrease of DIC with latitude during winter (A), spring (B) and summer (C). Note the logarithmic scale on concentration in all three plots. The symbols represent different size of the watershed, see Fig. 2.




Figure 5. The variation of $\delta^{13}\text{C}_{\text{DIC}}$ with latitude during winter (A) and spring (B) for watershed of different size. The symbols are the same as in Fig. 2. Isotopically-light DIC is observed in permafrost-affected zone during spring, suggesting intensive respiration of soil or plant litter carbon (Ob river sediments are from -25 to -27‰, Guo et al., 2004a).

Forest watershed, 57°N, August
Dark coniferous taiga




Fig. 6. Scheme of DOC pathways within the soil profile and to the river, (A): in forest watershed of the south, permafrost-free zone (57°N). Soil horizons (FAO, 2006): 1, O (Mor, forest litter); 2, Medium-decomposed peat (He) transforming into strongly decomposed peat (Ha) in the bottom layer; 3, Mollic humic horizon (A); 4, ABg surface horizons with stagnic properties; 5, Bg middle stagnic horizon; 6, Cgk carbonate-bearing clays and clay loam.

Frozen bog watershed 67°N, August

B

Dwarf shrubs and green mosses




Fig. 6, continued. (B): DOC pathways in frozen bog peatlands of continuous permafrost (67°N). Soil horizons (FAO, 2006): 7, weakly decomposed peat (Hi); 8, partially decomposed peat (He); 9, humic horizons (AH); 10, cryoturbated frozen stagnic horizon (Bgf@); 11, frozen stagnic horizon (BCgf); 12 sedimentary deposits (Cf).

In the south, DOC is retained by clay and deep in the soil profile, by clay loam with carbonates. In the north, the active layer depth does not exceed the overall thickness of the peat and thus the leachate of ground vegetation and peat layer do not meet mineral horizons during their transit to the river.

Supplement 1. Hydrological parameters of the WSL rivers

The daily, seasonal and annual discharges of some of the studied rivers are available from systematic surveys by the Hydrometeorological State Committee of the former USSR Goskomgidromet and Roskomgidromet (now the Russian Hydrological Survey, RHS). These data are published in the annual issues of the State Water Cadastre (Hydrological Yearbooks) and generalized in the “Resources of surface waters of the USSR, 1964 and 1972”. Given the limited number of observation over the year, the river discharge for each river was averaged for each of the 3 seasons of sampling (May to June, July to October, and November to April). For this, we used available monthly average discharges from the RHS gauging stations in the Kara sea basin from the data base of R-ActicNET (www.r-arcticnet.sr.unh.edu), which is based on mean-multi-annual data of the RHS. The runoff contour lines in Fig 1 A are based on results of Russian Hydrological Survey gauged river monitoring in the region a summarized and compiled in Nikitin and Zemtsov (1986). The southern, permafrost-free part of western Siberia is relatively well covered by RHS stations where monthly discharges are available until 2013-2014. In contrast, the density of stations, especially on small rivers, is much lower in the northern, permafrost – affected part of WSL. However, systematic hydrological study of State Hydrological Institute in 1973-1992 in the northern part of western Siberia allowed reliable evaluation of small and medium rivers discharges (Novikov et al., 2009). In case of the RHS gauging station location which was different from our sampling point of this river, we used an interpolation of the discharge taking into account the watershed area change along the main course of the river (Methodical, 2007; Svod pravil, 2004). In the absence of the gauging station at the river, we used either an analogous river approach or mean values for the area-normalized discharge in the region, given the rather homogeneous geographical setting of WSL (see runoff distribution in Fig. 1). For small and medium rivers of the palsa and polygonal bogs of the permafrost zone, we used empirical formulas accounting for hydrological parameters of these watersheds (Novikov et al., 2009). For southern rivers of the region, in the permafrost-free zone, the annual runoff was taken from available data of the RHS in 2013-2014 and calculated for ungauged rivers using an analogous approach.

References:

Resources of Surface Waters of the USSR (eds. Zhil, I. M. and Alushkinskaya, N. M.). Vol. III, Northern regions. Gidrometeoizdat, Leningrad, 633 pp., 1972.

Resources of Surface Waters of the USSR (ed. Eirikh, G.D.). Vol. XV, Altai and Western Siberia. Issue 3: Low Irtush and Low Ob. Gidrometeoizdat, Leningrad, 432 pp., 1964.




Figure S1. Significant decrease of Ca concentration in western Siberian rivers with latitude during winter (A), spring (B) and summer (C). The symbols represent different size of the watershed, see Fig. 2. Note the logarithmic scale on concentration in all three plots. The latitudinal trend is significant at $p < 0.001$.




Figure S2. Significant decrease of Mg concentration in western Siberian rivers with latitude during winter (A), spring (B) and summer (C). The symbols represent different size of the watershed, see Fig. 2. Note the logarithmic scale on concentration in all three plots. The latitudinal trend is significant at $p < 0.001$.




Figure S3. Evolution of K concentration in western Siberian rivers with latitude during winter (A), spring (B) and summer (C). The symbols represent different size of the watershed, see Fig. 2. The latitudinal trend is significant at $p < 0.001$.




Figure S4. Evolution of Si concentration in western Siberian rivers with latitude during winter (A), spring (B) and summer (C). The symbols represent different size of the watershed, see Fig. 2.




Figure S5. pH (A), Ca (B) and Mg (C) concentration in rivers as a function of latitude representing all seasons and all river watersheds. The difference between three permafrost zone are significant at $p < 0.05$. Note log scale for Ca and Mg concentration as a function of latitude.




Figure S6. DOC (A), DIC (B) and $\delta^{13}\text{C}_{\text{DIC}}$ (C) of all river size as a function of latitude comprising all sampled rivers during all seasons. The differences between 3 groups of sites are significant at $p < 0.05$.




Figure S7. Universal dependence of UV_{280 nm} absorbance of DOC concentration in all rivers during all seasons except winter (black circles) where some hydrocarbon degradation products or oil-field organics may produce significant scatter.





Figure S8. A : Latitudinal trend of Ca concentration across the WSL in peat (diamonds), moss (squares) and mineral horizon (circles) of the peat column (Stepanova et al., 2015). **B:** exchangeable Ca concentration in alluvial soils of the Ob River basin (Slavnina et al., 1981; Afanas'eva et al., 1984 and Izerskaya et al., 2014). Solid lines represent statistically significant trends at $p < 0.05$.

Izerskaia, L. A., Vorobyev, S. N, Vorobyeva, T. E., Kolesnichenko, L. G, Zakharchenko, A. V.: The concentration of Mn, Cu, Zn, Co, B, Sr, Cd and Pb in alluvial soils of the Ob River (forest-steppe, southern taiga and central taiga). *Internat. J. Environ. Stud.*, 71(5), 691–697, 2014.

Slavnina, T.P., Pashneva, G.E., Kakhatkina, M.I., Ivanova, R.G., Abramova, M.D., Seredina, V.P., Izerskaya, L.A.: The soils of the floodplain of the Middle Ob, their meliorative condition and agrochemical characteristics. Tomsk: Tomsk State University, 1981, 226 pp. (In Russian)

Afanas'eva, T.V., Balabko, P.N., Sumerin, M.B., Tereshina, T.B.: The soils of the floodplain of the upper and middle Ob. In: Kovalev R.V. (Ed.) *Problems of use and protection soil in Siberia and the Far East*. Novosibirsk: Nauka. p 62 – 65, 1984. (In Russian)

Table S1. Major physical and hydro-chemical parameters of sampled rivers. See Table 1 and Fig. 1 for localization.

SPRING	Date	S, km ²	D, m3/s	O ₂ , % sat	R, μS cm ⁻¹	pH	DIC, ppm	DOC, ppm	UV _{280nm}	Cl, ppm	SO ₄ , ppm	δ ¹³ C _{DIC} ‰
RJ-22	29.05.2014	1.79	0.017	72.4	106	6.64	5.326	12.22	1.352	15.07	0.070	-12.7
RJ-21	29.05.2014	3.37	0.032	85.9	115	6.85	3.65	12.23	1.384	22.22	0.156	-10.5
RJ-20	29.05.2014	7.18	0.069	70.3	34	6.16	0.785	14.58	1.455	4.403	0.630	-22.8
RJ-39	31.05.2014	7.46	0.044	79	9	5.12	0.309	6.546	1.143	0.128	0.555	-22.8
RJ-2	27.05.2014	8.14	0.062	91	66	6.91	6.037	24.61	1.804	0.161	0.010	-15.8
RJ-24	29.05.2014	9.52	0.091	66.6	20	4.71	0.372	17.62	1.654	0.741	0.273	-25.5
RJ-56	07.06.2014	9.65	0.124	71.6	9	5.16	0.356	8.59	1.264	0.425	0.489	-22
RJ-40	31.05.2014	12.04	0.070	72	10	5.63	0.6474	8.585	1.27	0.150	0.581	-23.5
RJ-12	27.05.2014	12.25	0.400	68.5	84	6.86	8.409	14.54	1.462	0.411	0.957	-14.4
RJ-36	30.05.2014	15.19	0.146	77.5	12	4.83	0.2915	7.684	1.188	0.222	0.381	-21.5
RJ-31	30.05.2014	18.8	0.180	61.3	8	5.43	0.3335	7.346	1.181	0.082	0.267	-21.8
RJ-25	29.05.2014	21.55	0.207	76.3	39	5.91	1.433	17.5	1.677	4.35	0.327	-21.7
RJ-55	07.06.2014	23.95	0.309	60.1	30	4.94	0.3504	21.22	1.876	3.480	0.364	-25.9
RJ-57	07.06.2014	33.48	0.432	75.1	12	5.86	0.638	8.557	1.244	0.466	0.366	-19.7
RJ-52	07.06.2014	34.61	0.447	65.3	19	5.46	0.575	10.18	1.38	2.268	0.370	-23.7
R-4	10.06.2013	34.61	0.447	79	n.d.	6.3	1.461	8.70	0.29	2.366	2.095	n.d.
RJ-46	02.06.2014	39.88	0.233	72.8	10	4.95	0.882	7.57	1.2	0.142	0.626	-23.8
RJ-45	02.06.2014	42.39	0.248	78.2	9	5.61	0.756	6.13	1.155	0.180	0.602	-21.8
RJ-35	30.05.2014	46.4	0.599	67.9	11	5.78	0.483	10.2	1.365	0.0889	0.230	-22.7
RJ-11	27.05.2014	58.56	1.913	59.3	67	7.02	6.611	15.9	1.484	0.251	0.182	-14.4
RJ-32	30.05.2014	59.86	0.772	68.1	39	5.68	0.591	7.63	1.235	8.13	0.276	-22.4
RJ-3	27.05.2014	61.48	0.775	73	76	6.91	7.275	24.4	1.758	0.106	0.0181	-16.5
RJ-30	30.05.2014	67.06	0.865	67.2	10	5.78	0.6641	6.10	1.142	0.076	0.248	-20.8
RJ-33	30.05.2014	71.61	0.924	62.8	28	6.02	1.053	7.99	1.259	3.754	0.329	-22
R-3	10.06.2013	74	0.955	65	n.d.	6.3	3.201	3.20	0.134	7.879	2.439	n.d.
RJ-50	07.06.2014	74	0.54	60.7	29	6.37	1.835	4.57	1.142	1.830	1.367	-23
RJ-41	31.05.2014	78.94	0.461	63.9	15	5.92	1.173	7.93	1.26	0.165	0.455	-21.6
RJ-34	30.05.2014	105.08	1.58	60.7	16	5.99	0.867	8.29	1.288	0.465	0.387	-21.4
RJ-43	31.05.2014	106.16	0.724	67.6	14	5.84	0.951	6.45	1.215	0.704	0.954	-22.9
RJ-27	30.05.2014	115.39	1.86	64.7	10	5.56	0.581	8.64	1.276	0.136	0.353	-21.2
Z-86	07.06.2013	115.39	1.86	82	13	5.22	1.067	29.25	0.2357	0.232	0.328	n.d.
RJ-58	11.06.2014	116.88	6.93	66.2	46	6.59	2.615	30.36	1.988	0.0583	1.168	-20.8
RJ-53	07.06.2014	175.29	2.64	70.7	16	5.86	0.707	11.83	1.41	0.469	0.6014	-23.2
R-5	10.06.2013	175.29	5.33	64	n.d.	6.1	1.774	10.53	0.391	4.71	1.178	n.d.
RJ-8	27.05.2014	177.45	3.16	68.3	286	7.53	35.34	14.82	1.351	0.345	0.1735	n.d.
RJ-9	27.05.2014	275.16	3.52	64.1	258	7.44	30.95	13.01	1.319	0.256	0.2606	-13.7
RJ-51	07.06.2014	280.59	1.32	61.6	13	5.97	0.687	8.69	1.293	0.577	0.4154	-19.8
RJ-6	27.05.2014	302.46	3.81	75.4	179	7.40	19.95	17.97	1.504	0.265	0.5110	-14.3
RJ-5	27.05.2014	320.04	6.15	83.1	256	7.40	30.6	15.82	1.419	0.3019	0.176	-14.2
RJ-18	29.05.2014	359.31	5.17	71.5	118	6.49	2.598	23.8	1.874	22.46	0.4025	-20
RJ-49	04.06.2014	512.33	4.24	78.4	15	5.76	0.6836	13.24	1.447	0.3160	0.667	-25
R-6	10.06.2013	598	11.6	80	n.d.	6.0	1.298	9.948	0.33	1.379	8.258	n.d.
RJ-54	07.06.2014	598	25.8	67.6	156	5.64	0.598	11.17	1.472	40.92	0.123	-20.2
RJ-14	27.05.2014	689	22.5	84.2	112	7.11	11.88	22.31	1.707	0.1145	0.0907	-10.7
R-2	10.06.2013	820.66	9.56	66	n.d.	6.2	2.786	6.606	0.2215	4.815	7.077	n.d.
RJ-29	30.05.2014	820.66	9.56	67.2	32	5.90	0.752	8.247	1.26	5.358	0.4519	-21.7
RJ-7	27.05.2014	1020.47	13.8	57.5	151	7.28	16.47	25.1	1.827	0.2299	0.4997	-16.1
RJ-23	29.05.2014	1260	9.10	72.5	132	6.64	2.976	13.95	1.455	27.31	0.4227	-18
RJ-48	04.06.2014	1970	16.3	77.7	14	5.78	0.675	14.24	1.479	0.1563	0.5026	-25.4
RJ-17	29.05.2014	3190	51.0	109	94	6.56	1.908	19.33	1.611	17.61	0.8858	-19.8
RJ-13	27.05.2014	3460	113	65.5	75	7.25	7.262	19.04	1.591	0.2680	0.5762	-15.7
RJ-42	31.05.2014	4030	44.5	77.5	10	5.70	0.596	6.215	1.17	0.1947	0.6968	-22.3
RJ-37	30.05.2014	5110	68.5	64.7	13	6.72	0.2806	8.776	1.221	0.6556	0.3655	-21.8
Z-55	05.06.2013	9881	134	n.d.	n.d.	n.d.	1.265	11.17	0.374	2.046	0.7693	n.d.
R-1	10.06.2013	9881	134	76	29	6.4	1.498	11.61	0.357	2.915	3.297	n.d.
RJ-28	30.05.2014	9881	134	79.1	17	5.33	0.456	10.55	1.316	1.676	0.3923	-24.3
R-7	10.06.2013	10768	154	79	n.d.	5.7	1.288	10.42	0.331	2.304	1.329	n.d.
R-8	10.06.2013	10768	367	68	31	6.2	1.249	10.82	0.330	2.340	3.904	n.d.
RJ-26	29.05.2014	10768	154	81.1	14	5.27	0.5529	11.39	1.393	0.5485	0.4622	-25.7
RJ-4	27.05.2014	12000	122	58.5	334	7.58	37.25	23.48	1.609	4.799	6.361	-13.7
R-10	12.06.2013	12000	121	67	347	8.1	35.73	26.86	0.709	2.870	7.046	n.d.
RJ-15	27.05.2014	25500	403	72.3	99	6.89	9.555	25.59	1.825	0.6347	0.958	-16.6
RJ-38	31.05.2014	26100	350	83.5	15	5.89	0.8026	10.34	1.362	0.2963	0.703	-24.6
RJ-10	27.05.2014	27200	348	58.1	159	7.11	15.68	32.75	2.024	1.572	1.88	-15.5
R-9	12.06.2013	27622	475	61	42	6.3	2.028	12.86	0.415	1.890	5.60	n.d.
RJ-19	29.05.2014	27622	475	8.04	33	6.37	1.107	13.45	1.445	4.552	0.730	-23.2
RJ-44	31.05.2014	112000	888	79.3	15	5.67	0.8375	7.288	1.231	0.955	0.886	-23.8
RJ-47	04.06.2014	150000	1286	79.6	43	6.87	3.739	11.27	1.37	0.319	1.03	-19.7
RJ-1	27.05.2014	423100	n.d.	181	242	7.90	24.8	5.449	1.056	2.393	13.7	-9.3
RJ-16	29.05.2014	773200	n.d.	95.2	99	7.09	9.034	13.25	1.384	1.167	4.933	-12.7

Table S1, continued.

SUMMER	Date	S, km ²	D, m3/s	O ₂ , % sat	R, μS cm ⁻¹	pH	DIC, ppm	DOC, ppm	UV _{280nm}	Cl, ppm	SO ₄ , ppm
BL-35	22.08.2013	7	0.035	n.d.	n.d.	7.81	23.22	10.44	0.2409	5.98	5.20
RA-18	08.08.2014	7.18	0.061	n.d.	52	6.42	4.253	9.481	1.268	2.90	0.341
BL-28	22.08.2013	9.35	0.080	n.d.	n.d.	7.83	31.51	9.830	0.3644	4.556	1.30
BL-25	22.08.2013	9.65	0.16	n.d.	n.d.	7.08	0.659	10.47	0.1796	1.009	0.283
RA-6	03.08.2014	11	0.10	n.d.	12.6	5.7	n.d.	n.d.	1.21	0.170	0.430
RY 14-45	24.08.2014	12.04	0.11	82.8	47	6.65	2.266	5.851	1.181	0.436	0.855
RA-5	03.08.2014	15	0.14	n.d.	31.9	6.48	4.114	8.410	1.291	0.1725	0.206
RY 14-47	24.08.2014	15.19	0.14	88.7	9	5.66	0.551	61.49	1.225	0.258	0.369
RA-12	04.08.2014	19	0.18	n.d.	29	5.84	n.d.	n.d.	1.678	0.177	0.063
BL-23	21.08.2013	23.95	0.38	n.d.	n.d.	7.25	3.827	7.813	0.277	4.276	3.28
BL-34	22.08.2013	26	0.130	n.d.	n.d.	7.80	27.27	8.842	0.3492	9.43	5.13
BL-31	22.08.2013	31	0.155	n.d.	n.d.	7.42	1.277	25.98	0.769	2.243	39.9
BL-3	12.08.2013	32	0.185	n.d.	n.d.	7.51	6.003	67.53	2.474	3.673	0.1393
BL-33	22.08.2013	32	0.160	n.d.	n.d.	7.42	16.66	19.88	0.594	2.977	0.5419
RA-15	04.08.2014	32	0.30	n.d.	26.3	6.62	3.123	14.47	1.51	0.133	0.208
BL-19	21.08.2013	34.61	0.76	n.d.	n.d.	7.03	3.902	8.519	0.3025	4.9	4.729
RA-9	03.08.2014	43	0.40	n.d.	19.5	6.52	n.d.	n.d.	1.185	0.4350	0.267
BL-32	22.08.2013	44	0.22	n.d.	n.d.	7.37	6.797	33.40	1.1032	2.054	0.250
RY 14-48	24.08.2014	53	0.49	73.5	34	6.24	3.779	4.349	1.073	0.3571	0.638
RY 14-49	24.08.2014	71.61	0.66	68.7	43	6.26	1.657	9.692	1.192	4.7540	0.500
BL-17	21.08.2013	74	0.54	n.d.	n.d.	7.09	3.808	2.828	0.092	8.544	2.679
RY 14-44	24.08.2014	78.94	0.73	65.5	22	6.6	1.689	7.930	1.194	0.627	0.177
RA-4	03.08.2014	79.5	0.74	n.d.	54	6.85	7.893	7.527	1.218	0.1963	0.113
BL-22	21.08.2013	82	0.76	n.d.	n.d.	7.06	9.972	4.130	0.102	11.75	7.22
RA-10	03.08.2014	88	0.82	n.d.	29.9	6.64	4.002	9.859	n.d.	0.136	0.345
RA-13	04.08.2014	96	0.89	n.d.	31.5	6.35	3.831	15.75	1.529	0.330	0.080
BL-14	21.08.2013	115.39	0.60	n.d.	n.d.	6.82	2.273	11.99	0.345	0.644	0.192
RA-1	30.07.2014	115.39	1.07	n.d.	19	6.36	1.186	14.87	1.462	0.143	0.238
BL-5	03.08.2014	116.88	1.04	n.d.	n.d.	7.75	25.81	26.57	0.916	4.314	0.147
RA-8	03.08.2014	121	1.12	n.d.	34.9	6.73	5.046	6.338	1.175	0.115	0.373
RT2 14-30	21.08.2014	157	1.74	89.3	82	7.07	5.965	11.15	1.171	5.49	2.65
RA-11	04.08.2014	170	1.67	n.d.	42.1	6.71	6.062	5.603	1.145	0.658	0.394
BL-21	21.08.2013	175.29	2.87	n.d.	n.d.	7.05	6.102	3.620	0.1015	9.54	3.895
RA-16	08.08.2014	175.29	3.04	n.d.	42	6.49	3.220	11.35	1.387	1.87	0.478
RY 14-42	24.08.2014	183	1.79	77.5	11	5.06	0.499	6.655	1.267	0.321	0.2626
RA-14	04.08.2014	250	2.45	n.d.	32	6.47	2.250	18.95	1.659	3.311	0.2070
BL-18	21.08.2013	280.59	0.81	n.d.	n.d.	7.09	3.003	4.844	0.158	6.973	1.2247
BL-27	22.08.2013	359.31	5.97	n.d.	n.d.	7.88	25.83	14.39	0.5274	4.84	59.2
RA-19	08.08.2014	359.31	4.35	n.d.	293	7.62	24.52	14.14	1.427	23.4	0.164
BL-9	12.08.2013	473	2.32	n.d.	n.d.	7.57	34.65	19.45	0.7005	7.197	0.1424
BL-6	12.08.2013	510	2.50	n.d.	n.d.	7.83	46.62	13.45	0.3092	7.602	0.183
RT2 14-32	21.08.2014	512.33	8.77	110.3	173	7.46	12.17	6.169	1.119	13.69	2.16
BL-24	21.08.2013	598	7.33	n.d.	n.d.	7.22	2.652	10.25	0.3152	1.999	16.97
RT2 14-29	21.08.2014	656	10.4	83	81	7.18	6.506	10.10	1.182	4.34	2.01
BL-16	21.08.2013	820.66	10.2	n.d.	n.d.	7.10	5.348	4.553	0.1486	9.134	7.36
BL-13	21.08.2013	1396	11.5	n.d.	n.d.	6.65	6.012	9.465	0.283	3.928	0.662
RT2 14-31	21.08.2014	1970	6.97	97.4	138	7.26	11.80	15.19	1.322	9.3048	1.371
RA-3	03.08.2014	1979	40.7	n.d.	39	7.22	n.d.	n.d.	1.236	n.d.	n.d.
RA-7	03.08.2014	1979	2.33	n.d.	36.2	6.88	5.446	5.072	1.128	0.105	0.265
BL-29	22.08.2013	3190	45.4	n.d.	n.d.	7.87	29.97	12.87	0.317	4.234	10.54
RA-20	08.08.2014	3190	45.4	n.d.	n.d.	n.d.	22.70	12.80	1.365	107.2	0.296
BL-2	12.08.2013	3197	2.92	n.d.	n.d.	7.65	31.99	51.58	1.591	4.206	2.52
RA-22	08.08.2014	3460	17.0	n.d.	369	7.51	45.23	11.39	1.196	0.328	0.146
RY 14-43	24.08.2014	4030	58	90	32	6.6	2.674	7.196	1.081	0.739	1.096
RY 14-46	24.08.2014	5110	73	90.5	36	6.71	3.190	6.297	1.158	0.779	0.113
RA-2	02.08.2014	9881	142	n.d.	50	6.89	3.179	11.96	1.378	4.568	0.318
BL-20	21.08.2013	9881	60	n.d.	n.d.	7.03	5.835	6.681	0.219	6.46	6.101
BL-15	21.08.2013	9881	142	n.d.	n.d.	6.86	5.930	7.093	0.232	7.02	4.17
RA-17	08.08.2014	10768	120	n.d.	27	6.4	1.498	15.26	1.512	1.70	0.269
BL-4	12.08.2013	25500	73.5	n.d.	n.d.	7.92	43.97	18.59	0.513	8.05	1.84
RA-21	08.08.2014	25500	107	n.d.	406	7.78	49.93	12.09	1.261	1.62	0.811
RA-23	08.08.2014	27200	72.4	n.d.	610	7.96	74.16	12.01	1.237	10.06	1.23
BL-26	22.08.2013	27622	408	n.d.	n.d.	6.86	6.109	10.82	0.287	2.77	2.97
BL-30	22.08.2013	75090	1425	n.d.	n.d.	7.87	8.809	8.062	0.217	3.35	0.310
RY 14-41	24.08.2014	112000	1168	93.3	45	6.69	3.777	71.21	1.111	1.29	0.408
RT2 14-40	16.08.2014	150000	3977	n.d.	80	6.57	8.197	11.45	1.154	0.581	0.743
BL-36	22.08.2013	773200	n.d.	n.d.	n.d.	7.92	19.57	5.974	0.143	2.295	1.51

Table S1, continued.

AUTUMN	Date	S, km ²	D, m3/s	O ₂ , % sat	R, μS cm ⁻¹	pH	DIC, ppm	DOC, ppm	UV _{280 nm}	Cl, ppm	SO ₄ , ppm	δ ¹³ C _{DIC} , ‰
R-2	18.10.2013	8.14	0.0074	93.9	35	6.8	5.951	40.2	1.132	1.462	0.877	n.d.
R-12	18.10.2013	12.25	0.050	88	n.d.	6.9	7.173	26.25	0.888	3.983	0.607	n.d.
R-13	18.10.2013	61.48	0.060	86.9	37	6.5	8.496	36.6	0.9873	2.024	0.458	n.d.
R-15	18.10.2013	116.88	0.19	93.1	25	7.7	7.835	32.94	1.068	4.183	0.591	n.d.
R-8	18.10.2013	177.45	0.23	99.5	199	7.8	67.88	8.775	0.1674	6.083	0.916	n.d.
R-9	18.10.2013	275.16	0.25	93.4	170	7.82	56.04	8.528	0.1585	6.294	0.53	n.d.
R-5	18.10.2013	320.04	0.44	95.3	239	7.2	75.92	7.056	0.1235	7.107	0.640	n.d.
R-14	18.10.2013	689	2.10	106	58	7.06	19.03	24.86	0.790	4.217	0.455	n.d.
R-7	18.10.2013	1020.47	0.99	100	158	7.6	51.2	16.7	0.459	6.20	0.614	n.d.
R-13	18.10.2013	3460	15.4	84.2	87	7	16.34	24.37	0.7399	4.929	0.411	n.d.
R-4	18.10.2013	12000	7.60	98	212	7.8	60.4	22.75	0.521	5.236	8.61	n.d.
R-17	18.10.2013	25500	66.9	84.4	72	7.36	20.09	30.96	0.9709	5.227	1.71	n.d.
R-1	18.10.2013	423100	n.d.	85.4	95	7.8	24.92	3.284	0.0712	n.d.	2.40	n.d.
WINTER												
RF13	22.02.2014	1.79	0.057	n.d.	270	7.48	15.72	14.57	0.554	37.0	0.168	-13.4
RF54	03.03.2014	3.37	0.059	n.d.	239	6.51	15.44	26.51	0.417	29.26	0.616	-12.4
RF55	03.03.2014	7.18	0.034	n.d.	112	6.18	13.36	28.85	0.717	2.77	0.769	-10.2
RF31	26.02.2014	7.46	0.054	n.d.	70	6.33	6.704	5.853	0.463	0.234	0.155	-12.1
RF52	03.03.2014	9.52	0.018	n.d.	79	5.73	n.d.	n.d.	n.d.	n.d.	n.d.	-20.6
RF48	03.03.2014	9.65	0.04	n.d.	28	6.19	2.68	12.65	0.340	0.411	0.438	-11.7
RF30	26.02.2014	12.04	9.80	n.d.	105	6.62	9.603	3.904	0.07	0.499	1.86	-17
RF33	26.02.2014	15.19	0.09	n.d.	32	5.79	1.716	17.57	0.771	0.658	0.947	-18.6
RF14	22.02.2014	21.55	0.103	n.d.	112	6.80	10.73	8.86	0.4158	5.36	0.607	-15.1
RF46	03.03.2014	23.95	0.10	n.d.	91	6.40	7.229	14.14	0.411	7.23	0.295	-14.8
RF49	03.03.2014	33.48	0.06	n.d.	45	6.09	4.433	20.34	0.578	0.469	0.610	-14.5
RF39	27.02.2014	34.61	0.07	n.d.	111	6.37	6.531	7.197	0.5068	4.625	0.261	-13.2
RF26	25.02.2014	39.88	0.094	n.d.	25	5.70	1.848	10.96	0.4798	0.529	0.954	-19.1
RF25	25.02.2014	42.39	0.162	n.d.	33	5.66	1.82	12	0.3275	1.424	1.335	-19.7
RF34	26.02.2014	46.4	0.10	n.d.	56	6.35	6.506	2.709	0.1774	0.406	0.982	-18.7
RF10	19.02.2014	58.56	0.05	n.d.	329	6.84	36.23	18.34	1.6655	0.374	0.205	-10
RF2	19.02.2014	61.48	n.d.	n.d.	144	6.58	33.13	3.185	1.0453	1.201	0.343	-19.3
RF36	26.02.2014	67.06	0.08	n.d.	69	6.28	7.419	4.643	0.095	0.1643	0.105	-17
RF35	26.02.2014	71.61	0.27	n.d.	59	6.38	5.134	8.08	0.1311	3.463	0.725	-18.6
RF41	27.02.2014	74	0.14	n.d.	53	6.40	4.456	2.291	0.0484	2.85	1.43	-19.8
RF29	26.02.2014	78.94	0.34	n.d.	120	6.56	12.47	5.854	0.1625	0.605	0.235	-14.9
RF45	03.03.2014	82	0.15	n.d.	137	6.62	12.18	2.755	0.0485	8.65	0.667	-18.7
RF27	25.02.2014	106.16	0.91	n.d.	68	n.d.	6.79	5.523	0.2241	0.700	0.568	-13.3
RF44	27.02.2014	115.39	0.15	n.d.	240	6.61	18.5	34.51	1.8222	0.628	0.178	-11.3
RF65	05.03.2014	116.88	0.12	n.d.	579	7.08	77.4	14.27	0.2261	0.348	0.057	-11.2
RF38	27.02.2014	175.29	0.78	n.d.	78	6.33	8.03	2.022	0.0761	1.955	0.888	-19.6
RF7	19.02.2014	177.45	0.14	n.d.	665	7.43	83.3	5.25	0.094	0.897	0.427	-14.7
RF62	04.03.2014	190	0.17	n.d.	409	6.86	51.5	9.796	0.2684	0.169	0.0747	-13.9
RF60	04.03.2014	220	0.19	n.d.	258	6.62	34.3	36.25	0.1377	0.158	0.1398	-18
RF61	04.03.2014	253	0.22	n.d.	415	6.86	52.7	10.52	0.266	0.174	0.2331	-10.4
RF8	19.02.2014	275.16	0.16	n.d.	576	7.42	75.05	4.536	0.0763	0.322	0.2176	-14.8
RF40	27.02.2014	280.59	0.16	n.d.	47	6.31	4.66	2.496	0.1074	1.513	0.467	-19.9
RF5	19.02.2014	302.46	0.18	n.d.	621	7.38	79.8	6.377	0.1199	0.516	0.095	-14.3
RF4	19.02.2014	320.04	0.28	n.d.	688	7.45	85.4	4.869	0.0844	1.59	0.739	-13.9
RF37	27.02.2014	350	0.88	n.d.	57	6.48	6.18	1.585	0.0714	1.305	0.680	-18.7
RF47	03.03.2014	598	1.60	n.d.	125	6.15	6.362	14.27	0.6597	16.55	0.483	-12.8
RF12	19.02.2014	689	1.27	n.d.	489	7.17	63.66	12.28	0.2796	0.3438	0.133	-10.2
RF58	04.03.2014	732	1.62	n.d.	258	6.65	30.98	6.234	0.2358	0.267	0.383	-15.6
RF42	27.02.2014	820.66	2.38	n.d.	95	6.41	7.58	2.735	0.0761	6.85	0.469	-18.5
RF6	19.02.2014	1020.47	0.63	n.d.	568	7.29	74.4	7.105	0.1442	0.570	0.186	-14.9
RF53	03.03.2014	1260	1.80	n.d.	237	6.50	16.2	11	0.3805	24.0	0.810	-9.6
RF51	03.03.2014	1648	4.12	n.d.	100	6.24	8.08	15.76	0.2495	8.68	0.525	-12.3
RF57	03.03.2014	3190	n.d.	n.d.	608	6.70	44.22	48.61	0.2349	76.4	0.335	-10.5
RF11	19.02.2014	3460	9.35	n.d.	403	7.06	50.84	10.29	0.3459	0.353	0.163	-12.6
RF28	25.02.2014	4030	11.8	n.d.	95	6.26	8.193	3.954	0.2199	1.455	2.82	-17.2
RF32	26.02.2014	5110	17.1	n.d.	63	6.34	6.143	2.819	0.0552	0.964	0.38	-18.2
RF21	25.02.2014	5190	13.8	n.d.	120	6.34	7.79	11.61	0.3771	4.61	4.34	n.d.
RF43	27.02.2014	9881	33.3	n.d.	132	6.38	9.97	4.347	0.0892	8.36	0.733	-16.3
RF50	03.03.2014	10768	31.9	n.d.	88	6.29	10.02	16.4	0.3752	2.02	0.551	-14
RF3	19.02.2014	12000	n.d.	n.d.	707	7.32	13.78	36.67	0.1549	11.01	5.79	-13.6
RF64	05.03.2014	25500	28.6	n.d.	460	6.92	58.4	8.906	0.1807	1.96	0.843	-16.6
RF9	19.02.2014	27200	21.2	n.d.	558	7.14	68.8	13.44	0.3596	8.78	1.63	-14.4
RF56	03.03.2014	27622	95.3	n.d.	131	6.55	10.9	15.02	0.1124	11.0	0.470	-14.3
RF63	04.03.2014	63780	81.6	n.d.	358	6.96	44.1	52.46	0.3612	4.52	0.735	-12.6
RF1	19.02.2014	423100	n.d.	n.d.	317	7.58	25.7	8.359	0.0538	3.69	13.17	-10.3

Table S2. Results of the non-parametric H-criterion Kraskal-Wallis test. Only Statistically significant values are shown.

S2-A. Comparison of three permafrost zone (all rivers, all seasons)

Component	H (Kryckal-Wallis)	p - level
Specific conductivity	62.654	< 0.0001
pH	91.259	< 0.001
DIC	93.955	< 0.001
DOC	26.559	< 0.0001
Cl	23.982	< 0.0001
Na	27.856	< 0.0001
Mg	91.893	< 0.001
Si	9.5102	0.0086
K	86.669	< 0.001
Ca	116.34	< 0.001

S2-B. Comparison of different seasons (all rivers)

Component	H (Kryckal-Wallis)	p - level
Specific conductivity	45.197	< 0.0001
pH	48.907	< 0.0001
DIC	69.67	< 0.0001
DOC	10.58	0.0142
Cl	29.94	< 0.0001
Na	27.82	< 0.0001
Mg	64.762	< 0.0001
Si	94.877	< 0.001
K	52.302	< 0.0001
Ca	34.145	< 0.0001

S2-C. Comparison of 4 watershed classes (< 100 km², 100 to 1000 km², 1000 to 10,000 km² and > 10,000 km²) in spring

Component	H (Kryckal-Wallis)	p - level
Specific conductivity	7.977	0.0465
pH	8.092	0.0441
DIC	8.140	0.0432
SO4	20.42	0.0001
Si	9.165	0.0272
K	12.57	0.0056

S2-E. Comparison of 4 watershed classes (< 100 km², 100 to 1000 km², 1000 to 10,000 km² and > 10,000 km²) during all seasons.

Component	H (Kryckal-Wallis)	p - level
Specific conductivity	21.282	0.0001
pH	18.908	0.0003
DIC	18.313	0.0004
SO4	25.593	< 0.0001
Na	20.863	0.0001
Mg	20.269	0.0001
K	35.215	< 0.0001
Ca	21.680	0.0001

S2-D. Comparison of 4 watershed classes (< 100 km², 100 to 1000 km², 1000 to 10,000 km² and > 10,000 km²) in winter.

Component	H (Kryckal-Wallis)	p - level
Specific conductivity	11.968	0.0075
pH	11.291	0.0102
DIC	10.349	0.0158
DOC	7.865	0.0489
Cl	7.833	0.0496
SO4	9.033	0.0288
Na	11.37	0.0099
Mg	13.93	0.0030
K	9.795	0.0204
Ca	12.49	0.0059

S2-F. Comparison of 6 latitudinal classes (56-58°, 58-60°, 60-62°, 62-64°, 64-66°, 66-68°) in spring.

Component	H (Kryckal-Wallis)	p - level
Specific conductivity	45.06	< 0.0001
pH	45.05	< 0.0001
DIC	45.50	< 0.0001
DOC	41.60	< 0.0001
Cl	29.65	< 0.0001
Na	29.43	< 0.0001
Mg	51.011	< 0.0001
Si	11.937	0.0357
K	41.129	< 0.0001
Ca	53.051	< 0.0001

S2-G. Comparison of 6 latitudinal classes (56-58°, 58-60°, 60-62°, 62-64°, 64-66°, 66-68°) in winter

Component	H (Kryckal-Wallis)	p - level
Specific conductivity	39.044	< 0.0001
pH	33.483	< 0.0001
DIC	40.452	< 0.0001
DOC, ppm	14.627	0.0055
Cl, ppm	12.766	0.0125
Na	31.139	< 0.0001
Mg	35.491	< 0.0001
Si	16.060	0.0029
K	26.948	< 0.0001
Ca	40.461	< 0.0001