

Interactive comment on “Global analysis of seasonality in the shell flux of extant planktonic foraminifera” by L. Jonkers and M. Kučera

W. Howard (Referee)

will.r.howard@gmail.com

Received and published: 15 March 2015

I concur with the other commenters on the value of this very valuable contribution.

I think the implications of this study could be explored a bit further, perhaps in a future study.

1) For example species with strong peaks in seasonal production might have different patterns in the range of seasonality they reflect in their isotopic or trace-metal variability. This could be another interesting test. Mix (1987) explored this idea in concept, and others have tried to apply it, sometimes coming up with surprising interspecific offsets in isotopic composition due to differences in seasonal production (e.g. King and Howard, 2005)

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



2) It would be interesting to see if, in aggregate, the offsets of individual taxa's peaks from mean temperatures drive biases in multivariate paleotemperature estimates. In principle multivariate biotic approaches can only work well if the taxa have distinct and only partially-overlapping environmental optima (expressed through seasonality or abundance variation against other environmental variable).

3) The authors could stand to add references to (or better yet, in further development, data from) some studies that reinforce their point about strong seasonality.

Eguchi, N. O., H. Ujiie, H. Kawahata, and A. Taira (2003), Seasonal variations in planktonic foraminifera at three sediment traps in the Subarctic, Transition and Subtropical zones of the central North Pacific Ocean, *Mar. Micropaleontol.*, 48(1-2), 149-163

Chapman, M. R. (2010), Seasonal production patterns of planktonic foraminifera in the NE Atlantic Ocean: Implications for paleotemperature and hydrographic reconstructions, *Paleoceanography*, 25, PA1101, doi:10.1029/2008PA001708.

Sagawa, T., A. Kuroyanagi, T. Irino, M. Kuwae, and H. Kawahata (2013), Seasonal variations in planktonic foraminiferal flux and oxygen isotopic composition in the western North Pacific: Implications for paleoceanographic reconstruction, *Mar. Micropaleontol.*, 100, 11-20, doi:10.1016/j.marmicro.2013.03.013.

King, A. L., and W. R. Howard (2001), Seasonality of foraminiferal flux in sediment traps at Chatham Rise, SW Pacific: implications for paleotemperature estimates, *Deep-Sea Research Part I-Oceanographic Research Papers*, 48(7), 1687-1708, doi:10.1016/S0967-0637(00)00106-0.

Pilskaln, C. H., S. J. Manganini, T. W. Trull, L. Armand, W. Howard, V. L. Asper, and R. Massom (2004), Geochemical particle fluxes in the Southern Indian Ocean seasonal ice zone: Prydz Bay region, East Antarctica, *Deep-Sea Research I*, 51, 307-332, doi:10.1016/j.dsr.2003.10.010.

Other references

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

King, A. L., and W. R. Howard (2005), $\delta^{18}\text{O}$ seasonality of planktonic foraminifera from Southern Ocean sediment traps: Latitudinal gradients and implications for paleoclimate reconstructions, *Mar. Micropaleontol.*, 56(1-2), 1-24, doi:10.1016/j.marmicro.2005.02.008.

Mix, A. C. (1987), The oxygen-isotope record of glaciation, in North America and Adjacent Oceans During the Last Deglaciation, edited by W. F. Ruddiman and H. Wright, pp. 111-135, Geological Society of America, Boulder.

Interactive comment on *Biogeosciences Discuss.*, 12, 1327, 2015.

BGD

12, C619–C621, 2015

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

