

1 **Chlorophyll a specific $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in stream**
2 **periphyton: implications for aquatic food web studies**

3

4 **N. F. Ishikawa¹, M. Yamane^{1,2}, H. Suga¹, N. O. Ogawa¹, Y. Yokoyama^{1,2} and N.**
5 **Ohkouchi¹**

6 [1]{Japan Agency for Marine-Earth Science and Technology, 2-15 Natsushima-cho,
7 Yokosuka, Kanagawa 237-0061, Japan}

8 [2]{Atmosphere and Ocean Research Institute, University of Tokyo, 5-1-5 Kashiwanoha,
9 Kashiwa, Chiba, 277-8564, Japan}

10 Correspondence to: N. F. Ishikawa (ishikawan@jamstec.go.jp)

11

12 **Abstract**

13 Periphytic algae attached to a streambed substrate (periphyton) are an important primary
14 producer in stream ecosystems. We determined the isotopic composition of chlorophyll *a* in
15 periphyton collected from a stream flowing on limestone bedrock in the Seri River, central
16 Japan. Stable isotope ratios of carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) and natural radiocarbon
17 abundances ($\Delta^{14}\text{C}$) were measured in chlorophyll *a* ($\delta^{13}\text{C}_{\text{chl}}$, $\delta^{15}\text{N}_{\text{chl}}$ and $\Delta^{14}\text{C}_{\text{chl}}$) and bulk
18 ($\delta^{13}\text{C}_{\text{bulk}}$, $\delta^{15}\text{N}_{\text{bulk}}$ and $\Delta^{14}\text{C}_{\text{bulk}}$) for periphyton, pure aquatic primary producer (*Cladophora*
19 sp.) and terrestrial primary producer (*Quercus glauca*). Periphyton $\delta^{13}\text{C}_{\text{bulk}}$ and $\delta^{13}\text{C}_{\text{chl}}$ values
20 did not necessarily correspond to $\delta^{13}\text{C}_{\text{bulk}}$ for an algal-grazing specialist (*Epeorus latifolium*).
21 Periphyton $\Delta^{14}\text{C}_{\text{chl}}$ values ($-258\text{\textperthousand}$ in April and $-190\text{\textperthousand}$ in October) were slightly lower than
22 $\Delta^{14}\text{C}_{\text{bulk}}$ values ($-228\text{\textperthousand}$ in April and $-179\text{\textperthousand}$ in October), but were close to the $\Delta^{14}\text{C}$ value for
23 dissolved inorganic carbon (DIC) ($-217 \pm 31\text{\textperthousand}$), which is a mixture of weathered carbonates
24 ($\Delta^{14}\text{C} = -1000\text{\textperthousand}$), CO₂ derived from aquatic and terrestrial organic matters (variable $\Delta^{14}\text{C}$)
25 and dissolved atmospheric CO₂ ($\Delta^{14}\text{C}$ approximately $+30\text{\textperthousand}$ in 2013). $\Delta^{14}\text{C}_{\text{chl}}$ values were also
26 close to $\Delta^{14}\text{C}_{\text{bulk}}$ for *E. latifolium* ($-215\text{\textperthousand}$ in April and $-199\text{\textperthousand}$ in October) and *Cladophora* sp.
27 ($-210\text{\textperthousand}$), whereas the $\Delta^{14}\text{C}_{\text{bulk}}$ value for *Q. glauca* ($+27\text{\textperthousand}$) was closer to $\Delta^{14}\text{C}$ for
28 atmospheric CO₂. Although the bulk isotopic composition of periphyton is recognised as a
29 surrogate for the photosynthetic algal community, natural periphyton is a mixture of aquatic

1 and terrestrial organic materials. Our results indicate that the bulk periphyton matrix at the
2 study site consists of 89% to 95% algal carbon (derived from ^{14}C -depleted DIC) and 5% to
3 11% terrestrial organic carbon (derived from ^{14}C -enriched atmospheric CO_2).

4

5 **1 Introduction**

6 The bioavailable energy in a natural ecosystem often originates not only from *in situ*
7 photoautotrophs, but also from resources produced in other ecosystems. In most freshwater
8 ecosystems (e.g., streams), periphytic algae attached to a substrate (periphyton) play an
9 important role as benthic primary producers (Allan and Castillo, 2007). Terrestrial material
10 (e.g., leaf detritus) is another resource for animals, especially in small headwater streams
11 (Vannote et al., 1980). Although the relative importance of aquatic and terrestrial resources
12 for food webs is a major concern in stream ecology (Vannote et al., 1980; Junk et al., 1989;
13 Thorp and DeLong, 1994), the energy flow from periphyton to animal consumers has not yet
14 been adequately assessed, because few studies have traced algal signatures through trophic
15 pathways. In stream food webs, macroinvertebrates are the dominant animal consumers, and
16 observation of their gut contents is a direct measure that can be used to trace energy flow
17 (Winemiller, 1990; Hall et al., 2000). However, the diets of stream macroinvertebrates are
18 sometimes too diverse to identify, and are not necessarily identical to what they actually
19 assimilate (Whitledge and Rabeni, 1997; Finlay, 2001).

20 The stable isotope ratios of carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) have contributed to food web
21 research over the last 40 years (after DeNiro and Epstein, 1978; Minagawa and Wada, 1984).
22 In stream ecosystems, environmental heterogeneity within a small area (e.g., habitat
23 variability in terms of light or flow regimes) is reflected in variations in periphyton $\delta^{13}\text{C}$
24 (Ishikawa et al., 2012a), which often makes it difficult to estimate the relative importance of
25 aquatic (e.g., periphyton) and terrestrial (e.g., leaf detritus) resources for macroinvertebrates
26 (Finlay et al., 1999; Zah et al., 2001; Doi et al., 2007; Dekar et al., 2009).

27 Recently, periphyton and terrestrial leaf detritus have been distinguished using natural
28 radiocarbon abundances ($\Delta^{14}\text{C}$). Periphyton $\Delta^{14}\text{C}$ is often derived from aged carbon reservoirs,
29 such as bedrocks and soils, and is relatively low compared to terrestrial leaf detritus that
30 reflects $\Delta^{14}\text{C}$ value for modern atmospheric CO_2 . Macroinvertebrate and fish $\Delta^{14}\text{C}$ values lie
31 between those for periphyton and leaf detritus, indicating that $\Delta^{14}\text{C}$ can be used to estimate
32 the energy flow in stream food webs (Ishikawa et al., 2014b). Although bulk $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and

$\Delta^{14}\text{C}$ values imply that the periphyton is isotopically identical to periphytic algae, it is actually a mixture of algae, heterotrophic fungi and bacteria, together with the exopolymeric substances exuded by these organisms, protozoa, small metazoa and other non-living particulate organic materials (Cross et al., 2005). All of these components may originate from different sources and have unique $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\Delta^{14}\text{C}$ values (Hladyz et al., 2011; Ishikawa et al., 2012b; Imberger et al., 2014; Fellman et al., 2015). Therefore, the algal and non-algal taxonomic compositions of the periphyton community potentially influence its bulk isotopic composition.

Because the densities of living algae and non-algal materials (e.g., leaf detritus or animal remains) usually differ, algae and other materials in periphyton are sometimes separated by centrifuging slurry washed from stream cobbles or rocks (Hamilton and Lewis, 1992; Small et al., 2011). However, the density-separation method does not often work well when the non-algal fraction contains large amounts of dead algae, and these two components are barely distinguishable even under a microscope (Finlay, 2004). The $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ values for bulk periphyton and its potential carbon sources (e.g., particulate organic carbon: POC, dissolved organic carbon: DOC and dissolved inorganic carbon: DIC) can be used to separate the algal carbon fraction from the non-algal carbon fraction (Fellman et al., 2015), although it is still difficult to quantitatively and directly estimate the relative abundances of the aquatic (i.e., algae) and terrestrial (i.e., leaf detritus) carbon fractions in periphyton based on their bulk isotopic compositions.

To assess the accuracy of using bulk isotopic composition of periphyton to represent that of aquatic primary producers, we used an algal biomarker found in the periphyton matrix. Chlorophylls are the ubiquitous antenna pigments of the photoautotrophs, and the chlorophyll *a* concentration, in particular, has been used as an indicator of *in situ* primary production because it is immediately degraded in the inactive state (Carpenter et al., 1986; Amir-Shapira et al., 1987). Several previous studies have successfully used the $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\Delta^{14}\text{C}$ values for chlorophyll *a* and its derivatives to understand modern environments or reconstruct palaeoenvironments (e.g., Hayes et al., 1987; Sachs et al., 1999; Ohkouchi et al., 2005; 2008; Kusch et al., 2010; Tyler et al., 2010; Higgins et al., 2012).

In this study, differences in the $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\Delta^{14}\text{C}$ values in chlorophyll *a* ($\delta^{13}\text{C}_{\text{chl}}$, $\delta^{15}\text{N}_{\text{chl}}$ and $\Delta^{14}\text{C}_{\text{chl}}$) and bulk ($\delta^{13}\text{C}_{\text{bulk}}$, $\delta^{15}\text{N}_{\text{bulk}}$ and $\Delta^{14}\text{C}_{\text{bulk}}$) for periphyton were compared to distinguish aquatic (i.e., algae) and terrestrial (i.e., leaf detritus) carbon fractions in the

1 periphyton community. Because the $\Delta^{14}\text{C}$ value is internally corrected by its $\delta^{13}\text{C}$ (Stuiver and
2 Polach, 1977), $\Delta^{14}\text{C}_{\text{chl}}$ does not depend on the isotopic fractionation during algal
3 photosynthesis and chlorophyll *a* biosynthesis. Therefore, the $\Delta^{14}\text{C}_{\text{chl}}$ value for periphyton
4 should reflect that for photosynthetic autotrophs (i.e., primary producers) and can be used as a
5 proxy of aquatic carbon for animals at higher trophic levels of the food web. The $\Delta^{14}\text{C}_{\text{chl}}$
6 values for periphyton, DIC and an algal-grazing specialist were compared to identify the
7 trophic transfers of carbon. Pure primary producers (i.e., aquatic algae and terrestrial plants)
8 were used to assess the potential differences in $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\Delta^{14}\text{C}$ values between
9 chlorophyll *a* and bulk cells.

10

11 **2 Materials and methods**

12 **2.1 Study site and sample collection**

13 In April and October 2013, field samplings were undertaken at Kawachi in the upland of the
14 Seri River (watershed area = 30km², 35°15'N, 136°20'E in Shiga Prefecture, central Japan),
15 which flows into Lake Biwa, the largest lake in Japan. The reach of the river studied flows
16 over limestone–basalt bedrock (dominated by cobbles) and contains different light and flow
17 environments. It has a slope of 1% to 2% and was 10 to 15 m wide, 10 to 40 cm deep and 250
18 m in altitude. The dominant riparian trees are from the family Fagaceae and Taxodiaceae
19 (higher plants with C₃ photosynthesis). Further details of this site and the DIC $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$
20 values have been reported in Ishikawa et al. (2012b).

21 We randomly collected several submerged cobbles from various habitats (e.g., open/shaded
22 and riffle/pool), which were rinsed gently with distilled water before the periphyton was
23 removed from the cobble surface with a brush and distilled water. The resulting slurry was
24 placed in a 100 mL polypropylene bottle, which was frozen until further processing. As
25 reference samples of pure aquatic and terrestrial primary producers, a filamentous green alga,
26 *Cladophora* sp., and several fresh leaves from the Japanese blue oak, *Quercus glauca*, were
27 collected in April. Several individuals of the mayfly larva, *Epeorus latifolium*, were collected
28 by hand in both April and October. The larvae of *E. latifolium* have highly specialized mouths
29 for grazing (Takemon, 2005), and their amino acid $\delta^{15}\text{N}$ values indicate that they are algal-
30 grazing specialists (Ishikawa et al., 2014a).

1 **2.2 Laboratory sample processing**

2 All samples were lyophilised with a freeze drier (FDU-1200, Eyela, Tokyo, Japan) in the dark.
3 The gut contents of *E. latifolium* larvae were removed prior to lyophilisation. The periphyton
4 samples were ground to a fine powder with a mortar and pestle, after all large invertebrates
5 (e.g., chironomids) had been manually removed. *Cladophora* sp. and *Q. glauca* were ground
6 with a vibrating mill (TI-100, CMT, Fukushima, Japan). The periphyton, *Cladophora* sp. and
7 *Q. glauca* samples were split into two vials for bulk and compound-specific isotope analyses.
8 The vials for the bulk periphyton and *Cladophora* sp. were treated overnight with 1 M HCl
9 solution to remove any carbonate and were washed and then lyophilised again. The algal
10 community in periphyton previously collected from the same site (November 2008) and the
11 gut contents of *E. latifolium* were observed under a microscope.

12 Chlorophyll *a* was extracted using the modification of the method of Chikaraishi et al. (2005,
13 2007). Briefly, the powdered periphyton, *Cladophora* sp. and *Q. glauca* were sonicated in
14 100% acetone at 0 °C for 15 min, followed by liquid–liquid (water:*n*-hexane = 3:1, v/v)
15 extraction, with NaCl salting out to remove the lipids. The *n*-hexane layer was extracted and
16 dried with a stream of argon, and the precipitate (i.e., pigments) was dissolved in *N,N*-
17 dimethylformamide (DMF) after filtration using a syringe (0.50 × 25 mm; Terumo, Tokyo,
18 Japan) equipped with a filter (4 mm × 0.2 µm PTFE, 100 pk; Grace Dawson Discovery
19 Science, Maryland, USA) to remove any remaining particles. The laboratory standard for
20 chlorophyll *a* was bought commercially (lot DCL2671; Wako Pure Chemical Industries,
21 Osaka, Japan) and the standard for phaeophytin *a* was made by adding 1 M HCl solution to
22 the chlorophyll *a* standard. Absorption spectra of our laboratory standards were consistent
23 with those reported in literatures (Chikaraishi et al., 2007; Tyler et al., 2010).

24 The pigments in DMF were introduced into a high-performance liquid chromatography
25 (HPLC) apparatus (1260 series; Agilent Technologies, California, USA), comprising a
26 G4225A degasser, a G1312B binary pump, a G1367E autosampler, a G1316C column oven, a
27 G1315D diode-array detector and a G1364C fraction collector. All solvents were better than
28 HPLC-grade (Wako Pure Chemical Industries). A Zorbax XDB C18 column (5 µm/4.6 × 250
29 mm; Agilent Technologies) and an XDB C18 guard column (5 µm/4.6 × 12.5 mm) were used
30 in the first purification step. In the first step, the solvent gradient program was as follows:
31 acetonitrile:ethyl acetate:pyridine = 75:25:0.5 (v/v/v) held for 5 min, then gradually changed
32 to 50:50:0.5 (v/v/v) in 55 min. The flow rate of the mobile phase was 1.00 mL min⁻¹. The

1 column oven was set at 30 °C. We identified chlorophyll *a* and phaeophytin *a* based on their
2 retention times and UV/Vis spectral patterns, compared with those of laboratory standards
3 (Fig. B3a, b).

4 The purified chlorophyll *a* and phaeophytin *a* were collected using the fraction collector and
5 were dried with a stream of argon. Because phaeophytin *a* was more abundant than
6 chlorophyll *a* in April, we purified phaeophytin *a* together with chlorophyll *a* and combined
7 them for the isotope measurements. The C and N isotopic compositions of phaeophytin *a* are
8 theoretically identical to those of chlorophyll *a* because phaeophytin *a* is an early degradation
9 product of chlorophyll *a*, and neither a C nor an N atom is replaced in this step. Each fraction
10 was dissolved in DMF and introduced into the HPLC apparatus again. A PAH column (5
11 µm/4.6 × 250 mm, Agilent Technologies) and a PAH guard column (5 µm/4.6 × 12.5 mm)
12 were used in the second purification step. In the second step, the solvent gradient program
13 was as follows: acetonitrile:ethyl acetate:pyridine = 80:20:0.5 (v/v/v) held for 5 min, then
14 gradually changed to 0:100:0.5 (v/v/v) in 35 min. The flow rate of the mobile phase was 1.00
15 mL min⁻¹. The column oven was set at 15 °C. After the second step, the fractions of
16 chlorophyll *a* and phaeophytin *a* were dried and washed with water:*n*-hexane (3:1, v/v). The
17 *n*-hexane layer was carefully extracted, dried again and frozen until the isotope measurements
18 were made. The abundances of chlorophyll *a* and phaeophytin *a* were estimated using
19 conversion formulae between the absorbance at 660 nm and the dry weights of the laboratory
20 standards. The dried chlorophyll *a* and phaeophytin *a* were dissolved in dichloromethane and
21 transferred to tin capsules for δ¹³C and δ¹⁵N measurements or to quartz tubes for Δ¹⁴C
22 measurements. The tin capsules and quartz tubes were dried again prior to measurements.

23 2.3 δ¹³C, δ¹⁵N and Δ¹⁴C measurements

24 The stable isotope ratios of carbon (δ¹³C) and nitrogen (δ¹⁵N) for bulk and chlorophyll *a* from
25 periphyton, *Cladophora* sp. and *Q. glauca* samples and those for bulk *E. latifolium* samples
26 were measured with an elemental analyser (Flash EA1112) coupled to a Delta XP isotope
27 ratio mass spectrometer (Thermo Fisher Scientific, Massachusetts, USA) with a Conflo III
28 interface (Thermo Fisher Scientific) modified for ultra-small-scale isotope measurements
29 (Ogawa et al., 2010). The δ¹³C and δ¹⁵N values are reported relative to those for Vienna Pee
30 Dee belemnite (VPDB) and atmospheric N₂ (AIR), respectively. Data were corrected using
31 two internal standards (tyrosine: δ¹³C_{VPDB} = -20.50‰ ± 0.13‰, δ¹⁵N_{AIR}: 8.44‰ ± 0.05‰;
32 nickel octaethylporphyrin: δ¹³C_{VPDB} = -34.17‰ ± 0.06‰; δ¹⁵N_{AIR}: 0.86 ± 0.03‰), which had

1 been corrected against multiple international standards (Tayasu et al., 2011). The 1σ analytical
2 precision for both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements was within 0.2‰ for bulk and with 0.9‰ for
3 chlorophyll *a*.

4 The samples for $\Delta^{14}\text{C}$ measurements were graphitized, according to the modified methods of
5 Kitagawa et al. (1993) and Yokoyama et al. (2010). Briefly, the bulk samples (approximately
6 1 mg C) and chlorophyll *a* samples (90 to 617 µg C) were combusted in an evacuated quartz
7 tube with copper oxide at 500 °C for 30 min and at 850 °C for 2 h. The CO₂ gas was
8 cryogenically purified in a vacuum line and reduced to graphite with hydrogen and an iron
9 catalyst at 550 °C for 10 h. The $\Delta^{14}\text{C}$ values for the bulk samples and chlorophyll *a* samples
10 were measured with an accelerator mass spectrometer (AMS) at Institute of Accelerator
11 Analysis (Kanagawa, Japan; AMS lab code IAAA) and at Atmosphere and Ocean Research
12 Institute, University of Tokyo (Chiba, Japan; AMS lab code YAUT), respectively. The $\Delta^{14}\text{C}$
13 (‰) value was defined as follows (Stuiver and Polach, 1977):

$$\Delta^{14}\text{C} (\text{\textperthousand}) = \delta^{14}\text{C} - 2(\delta^{13}\text{C} + 25)(1 + \delta^{14}\text{C}/1000)$$

14 (1)

15 $\Delta^{14}\text{C}$ value of the international standard (oxalic acid) took into account the radioactive decay
16 since AD 1950 (Stuiver and Polach, 1977). The 1σ analytical precision of the $\Delta^{14}\text{C}$
17 measurements was within 3‰ for bulk and 8‰ for chlorophyll *a*. The HPLC procedural
18 blank for carbon (e.g., potential contamination by column breeding), assessed with elemental
19 analyser, was below the detection limit (< 0.177 µg C), which was lower than 0.2% carbon in
20 the purified chlorophyll *a* molecules used for the AMS measurement.

21 To determine the carbon transfer pathway in this stream ecosystem, the $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ values
22 for all samples were compared with those for DIC, DOC and POC collected at the same site
23 in the Seri River in 2009 to 2010 (Ishikawa et al., 2012b, 2015).

24

25 **3 Results and discussion**

26 **3.1 Sample observations**

27 Microscopic observations show that diatoms and cyanobacteria are the dominant
28 photoautotrophs in the periphyton community at the study site (Fig. B1). Both the periphyton
29 and gut contents of *E. latifolium* consisted not only of algal cells, but also of amorphous and

unidentified particles (Fig. B2). The exuvium of small invertebrates (approximately 500 µm) was found in the periphyton matrix (Fig. B2a), the isotopic composition of which would have differed from that of pure algae. The UV/Vis spectra show different composition of photosynthetic pigments between April and October. Chlorophyll *a* (Mw, 892.5) and phaeophytin *a* (Mw, 870.6; the Mg atom is replaced by two H atoms in the centre of the tetrapyrrole ring of the chlorophyll *a* molecule) were the dominant pigments in the periphyton matrix in both April and October (Fig. B3). The combined abundance of chlorophyll *a* and phaeophytin *a* per unit dry weight was greater in October than in April, indicating that the algal biomass of the periphyton community was greater in October than in April (Table A2).

3.2 ^{13}C composition

The periphyton $\delta^{13}\text{C}_{\text{bulk}}$ and $\delta^{13}\text{C}_{\text{chl}}$ values were $-20.7\text{\textperthousand}$ and $-20.0\text{\textperthousand}$, respectively, in April, and $-26.2\text{\textperthousand}$ and $-26.0\text{\textperthousand}$, respectively, in October (Fig. 1). The algal-grazer *E. latifolium* $\delta^{13}\text{C}_{\text{bulk}}$ values were $-26.6\text{\textperthousand}$ and $-26.5\text{\textperthousand}$ in April and October (Fig. 1), respectively. In October, the periphyton $\delta^{13}\text{C}_{\text{bulk}}$ and $\delta^{13}\text{C}_{\text{chl}}$ values were close to the *E. latifolium* $\delta^{13}\text{C}_{\text{bulk}}$ value. In contrast, neither the periphyton $\delta^{13}\text{C}_{\text{bulk}}$ nor $\delta^{13}\text{C}_{\text{chl}}$ value was close to the *E. latifolium* $\delta^{13}\text{C}_{\text{bulk}}$ value in April. This is partly because the periphyton $\delta^{13}\text{C}_{\text{bulk}}$ values vary from $-32\text{\textperthousand}$ to $-16\text{\textperthousand}$ among stream habitats (e.g., open/shaded and riffle/pool) in this study site, due to the variable isotopic fractionation between DIC and algae (Ishikawa et al., 2012b). Such a large variation in periphyton $\delta^{13}\text{C}_{\text{bulk}}$ values on a small spatial scale may cause an inconsistency in $\delta^{13}\text{C}$ between periphyton (primary producers) and *E. latifolium* (primary consumers).

A mismatch between the $\delta^{13}\text{C}_{\text{bulk}}$ values for periphyton and grazers is often observed (Dekar et al., 2009), although ^{13}C is not enriched through the trophic levels (Vander Zanden and Rasmussen, 2001). There are four independent scenarios that explain our $\delta^{13}\text{C}$ results. Firstly, *E. latifolium* assimilates the ^{13}C -depleted fraction in periphyton. Secondly, *E. latifolium* assimilates the terrestrial organic matter, which is more ^{13}C -depleted than the periphyton. Thirdly, the periphyton $\delta^{13}\text{C}_{\text{bulk}}$ and $\delta^{13}\text{C}_{\text{chl}}$ values varied by 6\textperthousand , whereas the *E. latifolium* $\delta^{13}\text{C}_{\text{bulk}}$ values did not change greatly between April and October, suggesting that primary consumers integrate temporal fluctuations in $\delta^{13}\text{C}$ values for primary producers. Finally, the $\delta^{13}\text{C}_{\text{chl}}$ value is not a reliable proxy for $\delta^{13}\text{C}$ of bulk algae because the $\delta^{13}\text{C}_{\text{chl}}$ value is affected by the isotopic fractionation that occurs during chlorophyll *a* biosynthesis. To provide a more

1 precise estimate of algal carbon, the $\Delta^{14}\text{C}_{\text{chl}}$ signature is useful because it is corrected for
2 isotopic fractionation by $\delta^{13}\text{C}$ in Eq. (1) (Stuiver and Polach, 1977).

3 $\delta^{13}\text{C}_{\text{bulk}}$ and $\delta^{13}\text{C}_{\text{chl}}$ values were $-23.0\text{\textperthousand}$ and $-24.7\text{\textperthousand}$, respectively, for *Cladophora* sp. and $-$
4 $30.9\text{\textperthousand}$ and $-32.0\text{\textperthousand}$, respectively, for *Q. glauca* (Fig. 1). The $\delta^{13}\text{C}_{\text{chl}}$ value for primary
5 producers is controlled by the $\delta^{13}\text{C}$ value for their carbon source (i.e., DIC for *Cladophora* sp.
6 and atmospheric CO₂ for *Q. glauca*) and by internal isotopic fractionation between bulk cells
7 and chlorophyll *a* molecules. Sachs et al. (1999) reported that $\delta^{13}\text{C}_{\text{chl}}$ values for a cultivated
8 green alga *Dunaliella tertiolecta* were $0.5\text{\textperthousand}$ to $4.0\text{\textperthousand}$ lower than those for their bulk cells,
9 which is consistent with our *Cladophora* sp. data. Chikaraishi et al. (2005) reported the same
10 $\delta^{13}\text{C}_{\text{bulk}}$ value ($-30.9\text{\textperthousand}$) for the fresh leaves of the Mongolian oak *Q. mongolica* as for our *Q.*
11 *glauca* data. In contrast, in this study, the *Q. glauca* $\delta^{13}\text{C}_{\text{chl}}$ value ($-32.0\text{\textperthousand}$) was lower than
12 that for *Q. mongolica* ($-29.2\text{\textperthousand}$) reported in Chikaraishi et al. (2005).

13 **3.3 ^{15}N composition**

14 The periphyton $\delta^{15}\text{N}_{\text{bulk}}$ and $\delta^{15}\text{N}_{\text{chl}}$ values were $-5.7\text{\textperthousand}$ and $-1.5\text{\textperthousand}$, respectively, in April, and
15 $-1.7\text{\textperthousand}$ and $+0.5\text{\textperthousand}$, respectively, in October (Fig. 1). The algal-grazer *E. latifolium* $\delta^{15}\text{N}_{\text{bulk}}$
16 values ($-3.9\text{\textperthousand}$ in April and $+1.4\text{\textperthousand}$ in October) were $1.8\text{\textperthousand}$ to $2.9\text{\textperthousand}$ higher than the
17 periphyton $\delta^{15}\text{N}_{\text{bulk}}$ values. The $\delta^{15}\text{N}_{\text{bulk}}$ and $\delta^{15}\text{N}_{\text{chl}}$ values were $-4.3\text{\textperthousand}$ and $-6.0\text{\textperthousand}$,
18 respectively, for *Cladophora* sp. and $-0.8\text{\textperthousand}$ and $-0.2\text{\textperthousand}$, respectively, for *Q. glauca* (Fig. 1).
19 Sachs et al. (1999) reported that the $\delta^{15}\text{N}_{\text{chl}}$ values were 2\textperthousand to 9\textperthousand lower than the $\delta^{15}\text{N}_{\text{bulk}}$
20 values for phytoplankton because of the isotopic fractionation that occurs during chlorophyll
21 *a* biosynthesis. Kennicutt et al. (1992), on the other hand, reported that the $\delta^{15}\text{N}_{\text{chl}}$ values were
22 relatively close to the $\delta^{15}\text{N}_{\text{bulk}}$ values for terrestrial C₃ plants. Therefore, the relationships
23 between $\delta^{15}\text{N}_{\text{bulk}}$ and $\delta^{15}\text{N}_{\text{chl}}$ values for *Cladophora* sp. and *Q. glauca* are consistent with
24 those for previous studies. In contrast, the periphyton $\delta^{15}\text{N}_{\text{chl}}$ values were $2.2\text{\textperthousand}$ to $4.2\text{\textperthousand}$
25 higher than their $\delta^{15}\text{N}_{\text{bulk}}$ values. This result might be attributable to the presence of
26 cyanobacteria (e.g., *Oscillatoria* sp. or *Homoeothrix* sp., Fig. B1) in the periphyton
27 community, because the $\delta^{15}\text{N}_{\text{bulk}}$ and $\delta^{15}\text{N}_{\text{chl}}$ values for cyanobacteria are usually different
28 from those for algae (Beaumont et al., 2000).

1 3.4 ^{14}C composition

2 The $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ values for DIC at the same study site in the Seri River have been reported
3 previously as $-7.2 \pm 0.2\text{\textperthousand}$ and $-217 \pm 30.7\text{\textperthousand}$, respectively (four-season mean \pm SD, N = 16;
4 Ishikawa et al., 2012b, Figs. 1, 2). These values are balanced by the mixing of weathered
5 carbonates ($\delta^{13}\text{C} = +3.9 \pm 0.3\text{\textperthousand}$ and $\Delta^{14}\text{C} = -1000\text{\textperthousand}$), dissolved atmospheric CO₂ ($\delta^{13}\text{C}$ and
6 $\Delta^{14}\text{C}$ are approximately $-8\text{\textperthousand}$ and $+30\text{\textperthousand}$, respectively, in 2013) and mineralized organic
7 materials (DOC: $\delta^{13}\text{C} = -24.2 \pm 2.9\text{\textperthousand}$, $\Delta^{14}\text{C} = -248 \pm 110\text{\textperthousand}$; POC: $\delta^{13}\text{C} = -25.0 \pm 3.4\text{\textperthousand}$,
8 $\Delta^{14}\text{C} = -109 \pm 52\text{\textperthousand}$) (four-season mean \pm SD, N = 4 for each fraction) at the study site
9 (Ishikawa et al., 2015, Figs. 1, 2).

10 The periphyton $\Delta^{14}\text{C}_{\text{bulk}}$ and $\Delta^{14}\text{C}_{\text{chl}}$ values (mean of the repeated measurements $\pm 1\sigma$
11 analytical precision) were $-228 \pm 2.3\text{\textperthousand}$ and $-258 \pm 4.8\text{\textperthousand}$, respectively, in April, and $-179 \pm$
12 $2.2\text{\textperthousand}$ and $-190 \pm 6.1\text{\textperthousand}$, respectively, in October, showing that chlorophyll *a* is slightly
13 depleted in ^{14}C relative to the bulk of the periphyton (Fig. 1). In particular, the periphyton
14 $\Delta^{14}\text{C}_{\text{chl}}$ value in April was lower than the seasonal range of DIC $\Delta^{14}\text{C}$ (Fig. 1). There are two
15 possible explanations of the periphyton $\Delta^{14}\text{C}_{\text{chl}}$ value in April. Firstly, periphytic algae
16 assimilate CO₂ dissolved from the bedrock limestone at the biofilm–bedrock boundary, in
17 addition to water column DIC. Because respiratory CO₂ and organic acids can mediate
18 carbonate weathering (Berner et al., 1983), ^{14}C -dead (i.e., $\Delta^{14}\text{C} = -1000\text{\textperthousand}$) CO₂ derived from
19 carbonates may enter the algae. Secondly, heterotrophs such as fungi and bacteria in
20 periphyton community consume ambient DOC and release CO₂ during their respiration
21 (Fischer 2003). The CO₂ derived from heterotrophic respiration of DOC may be another ^{14}C -
22 depleted carbon source that is utilized by periphytic algae for photosynthesis.

23 The $\Delta^{14}\text{C}_{\text{bulk}}$ and $\Delta^{14}\text{C}_{\text{chl}}$ values were $-199 \pm 2.7\text{\textperthousand}$ and $-210 \pm 6.8\text{\textperthousand}$, respectively, for
24 *Cladophora* sp. and $+27 \pm 2.3\text{\textperthousand}$ and $-10 \pm 7.3\text{\textperthousand}$, respectively, for *Q. glauca* (Fig. 1). The *Q.*
25 *glauca* $\Delta^{14}\text{C}_{\text{bulk}}$ value was not greatly different from global mean $\Delta^{14}\text{C}$ value for atmospheric
26 CO₂ in 2013 (approximately $+30\text{\textperthousand}$, Levin et al., 2013). Although chlorophyll *a* contains only
27 0.07% of carbon in bulk leaves, *Q. glauca* synthesizes chlorophyll *a* using not only
28 atmospheric CO₂, but also aged (^{14}C -depleted) CO₂ and/or organic matters derived from other
29 carbon sources. A candidate source is soil, as variable $\Delta^{14}\text{C}$ values for soil organic matters
30 have been reported in several previous studies (Trumbore and Zheng 1996; Koarashi et al.,
31 2009). Various terrestrial plants can incorporate soil-derived carbon through their roots
32 (Brüggemann et al., 2011; Bloemen et al., 2013). Although there is no evidence that ^{14}C -

1 depleted organic carbon is transferred from soils to plants, *Q. glauca* and probably other
2 terrestrial plants may be able to make the chlorophyll *a* molecule using recycled phytol, as
3 reported in *Arabidopsis* seedlings (Ischebeck et al., 2006). The chlorophyll *a* biosynthesis has
4 multiple channels to acquire carbon, which is not necessarily originated from atmospheric
5 CO₂. The *Q. glauca* $\Delta^{14}\text{C}_{\text{chl}}$ value will be different from its $\Delta^{14}\text{C}_{\text{bulk}}$ value if *Q. glauca* collects
6 phytol or its precursor from soils. More attentions should be paid in future to plant uptake of
7 soil carbon for understanding carbon allocation in plants and global carbon budget in
8 terrestrial biosphere.

9 To estimate relative abundances of aquatic (e.g., algae) and terrestrial (e.g., leaf detritus)
10 carbon fractions in periphyton bulk matrix, a separate two-source mixing model was applied
11 to each of April and October. We assumed that the periphyton $\Delta^{14}\text{C}_{\text{chl}}$ value (−258‰ in April
12 and −190‰ in October) and the *Q. glauca* $\Delta^{14}\text{C}_{\text{bulk}}$ value (+27‰ in both April and October)
13 represent the aquatic and terrestrial end members, respectively. Therefore, periphyton $\Delta^{14}\text{C}_{\text{bulk}}$
14 values in April (−228‰) and October (−190‰) were explained by both seasonal variation in
15 aquatic end member and relative contributions of the aquatic and terrestrial end members to
16 periphyton bulk matrix. The results of mixing model show that the periphyton bulk matrix
17 consisted of 89% (April) to 95% (October) aquatic carbon and 5% (October) to 11% (April)
18 terrestrial carbon. The *E. latifolium* $\Delta^{14}\text{C}_{\text{bulk}}$ values (−215 ± 2.3‰ in April and −199 ± 2.2‰
19 in October) were within the range of periphyton $\Delta^{14}\text{C}$ values (Fig. 1). The April *E. latifolium*
20 $\Delta^{14}\text{C}_{\text{bulk}}$ value was closer to the periphyton $\Delta^{14}\text{C}_{\text{bulk}}$ value than to its $\Delta^{14}\text{C}_{\text{chl}}$ value, suggesting
21 that *E. latifolium* assimilates not only ¹⁴C-depleted aquatic sources, but also ¹⁴C-enriched
22 terrestrial sources in April. In contrast, the October *E. latifolium* $\Delta^{14}\text{C}_{\text{bulk}}$ value was closer to
23 the periphyton $\Delta^{14}\text{C}_{\text{chl}}$ value than to its $\Delta^{14}\text{C}_{\text{bulk}}$ value, suggesting that *E. latifolium* primarily
24 assimilates aquatic sources in October. This seasonal variation may be attributed to the higher
25 chlorophyll *a* abundance per unit dry weight in October, and/or to the higher terrestrial flux
26 associated with the input of snow melt in April.

27 3.5 Implications of this study

28 Previous studies have assumed that isotopic compositions of bulk periphyton are identical to
29 those of periphytic algae without direct evidence. Regarding identification of aquatic baseline
30 for stream food webs, our $\delta^{13}\text{C}_{\text{chl}}$ and $\Delta^{14}\text{C}_{\text{chl}}$ data indicate that the periphyton $\delta^{13}\text{C}_{\text{bulk}}$ and
31 $\Delta^{14}\text{C}_{\text{bulk}}$ values can be approximated as those for the photosynthetic algal community in
32 periphyton (Fig. 3). However, there remain some uncertainties in our data, such as the results

that the $\delta^{15}\text{N}_{\text{chl}}$ values were higher than $\delta^{15}\text{N}_{\text{bulk}}$ values in periphyton and that the $\Delta^{14}\text{C}_{\text{chl}}$ values were slightly lower than the $\Delta^{14}\text{C}_{\text{bulk}}$ values. These results do not indicate that isotopic compositions of bulk periphyton are completely consistent with those of algae. Bulk isotope analysis may underestimate importance of aquatic production for stream food webs especially in less productive streams, where the terrestrial detritus is more abundant than the algae/cyanobacteria in the periphyton. On the other hand, chlorophyll *a* specific $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values are useful tracers for precisely estimating of the sources of carbon and nitrogen in stream ecosystems, in which heterogeneous resources (e.g., aquatic and terrestrial organic matters) are mixed.

Compound-specific stable isotope and radiocarbon analyses are promising tools for the precise estimation of the sources, dynamics and turnover of various organic molecules (Hayes et al., 1987; Eglinton et al., 1996; Jochmann and Schmidt, 2012; Ohkouchi et al., 2015). Chlorophyll *a* is a unique biomarker of *in situ* photoautotrophs and more accurate than other biochemical compounds (e.g., lipids and amino acids) because it is immediately degraded in the inactive state (Carpenter et al., 1986; Amir-Shapira et al., 1987; Matile et al., 1996). However, a pitfall may exist in chlorophyll *a* recycling system. Some previous studies have suggested that terrestrial plants and cyanobacteria have a salvage pathway of phytol in chlorophyll *a* biosynthesis (Ischebeck et al., 2006; Vavilin and Vermaas 2007). Isotopic composition of chlorophyll *a* is determined by relative contributions of *de novo* synthesis and the recycling system to all chlorophyll *a* molecules. These contributions can be estimated by a separate measurement for isotopic compositions of each of chlorophyll *a* and its bounded phytol (e.g., Chikaraishi et al., 2005).

The isotopic composition of chlorophyll *a* can be used not only in stream ecosystems, but also in coastal ecosystems, where benthic biofilms (i.e., mixtures of algae and other heterotrophs) are important food sources for invertebrates, fish and birds (Kuwae et al., 2008, 2012). Furthermore, primary production in the ocean and lakes is currently estimated using bulk isotopic composition of particulate organic matter, which is a mixture of not only phytoplankton, but also heterotrophs and other organic materials derived from various sources. Chlorophyll *a* specific isotopic compositions can avoid the “mixing effect” on the estimation of *in situ* primary production and provide more precise data for biogeochemical cycling of materials and energy. We conclude that future studies should attempt to test how much $\delta^{13}\text{C}$,

1 $\delta^{15}\text{N}$ and $\Delta^{14}\text{C}$ values differ between bulk and chlorophyll *a* in primary producers collected
2 from multiple ecosystems.

3

4 **Figure captions**

5 Figure 1. The $\Delta^{14}\text{C}_{\text{bulk}}$, $\delta^{13}\text{C}_{\text{bulk}}$ and $\delta^{15}\text{N}_{\text{bulk}}$ values (shaded symbols) for periphyton
6 (diamonds), *Cladophora* sp. (aquatic primary producer; circle), *Q. glauca* (terrestrial primary
7 producer; square) and *E. latifolium* (algal grazer; triangles) and the $\Delta^{14}\text{C}_{\text{chl}}$, $\delta^{13}\text{C}_{\text{chl}}$ and $\delta^{15}\text{N}_{\text{chl}}$
8 values (open symbols) for periphyton, *Cladophora* sp. and *Q. glauca*. DIC: dissolved
9 inorganic carbon; DOC: dissolved organic carbon; POC: particulate organic carbon. *Data
10 from Ishikawa et al. (2012b, 2015).

11 Figure 2. Biplot of $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ data. Carbonate rocks in the Seri River ($\delta^{13}\text{C} = +3.9 \pm 0.3\text{\textperthousand}$
12 and $\Delta^{14}\text{C} = -1000\text{\textperthousand}$) (Ishikawa et al., 2015) and atmospheric CO₂ ($\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ are
13 approximately $-8\text{\textperthousand}$ and $+30\text{\textperthousand}$, respectively, in 2013) are also shown as end members.

14 Figure 3. Schematic view of the carbon cycle at the study site (Seri River) constrained by $\delta^{13}\text{C}$
15 and $\Delta^{14}\text{C}$.

16

17 **Appendix A: Full data set used in this study**

18 Table A1. The $\delta^{13}\text{C}_{\text{bulk}}$, $\delta^{15}\text{N}_{\text{bulk}}$ and $\Delta^{14}\text{C}_{\text{bulk}}$ values (\textperthousand) and C/N ratios (g g⁻¹) of the samples.
19 PP: primary producer. Means and 1σ analytical errors of the repeated measurements are
20 shown.

21 Table A2. The $\delta^{13}\text{C}_{\text{chl}}$, $\delta^{15}\text{N}_{\text{chl}}$ and $\Delta^{14}\text{C}_{\text{chl}}$ values (\textperthousand), C/N ratios of purified chlorophyll *a* (g
22 g⁻¹) (theoretical value: 11.8), chlorophyll *a* abundances per unit dry weight of the samples (μg
23 g⁻¹) and carbon contents of the chlorophyll *a* samples introduced into the AMS (μg C) for
24 periphyton, *Cladophora* sp. and *Q. glauca*. Means and 1σ analytical errors of the repeated
25 measurements are shown. Periphyton in April compiles chlorophyll *a* and phaeophytin *a*. The
26 October periphyton $\delta^{13}\text{C}_{\text{chl}}$ and $\delta^{15}\text{N}_{\text{chl}}$ values were determined based on single measurement.

27

28 **Appendix B: Supplemental information**

1 Figure B1. Illustration of algae and cyanobacteria in the periphyton community observed in
2 November 2008. White scale bars in the bottom right corners indicate 50 µm.
3 Figure B2. Microscopic images of a) periphyton and b) the gut contents of *E. latifolium*
4 collected in April 2013. White scale bars in the bottom right corners indicate 100 µm.
5 Figure B3. Three-dimensional chromatograms of laboratory standards for a) chlorophyll *a*,
6 and b) phaeophytin *a* and periphyton collected from the Seri River in c) April, and d) October
7 2013.

8

9 **Author contribution**

10 N. F. I. conceived the study design and conducted fieldwork. N. F. I. and H. S. conducted
11 pigment purification using HPLC. N. O. O. conducted $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analyses using
12 EA/IRMS. M. Y. and Y. Y. conducted $\Delta^{14}\text{C}$ analysis using AMS. All authors participated
13 discussion. N. F. I. and N. O. wrote the manuscript.

14

15 **Acknowledgements**

16 We thank F. Akamatsu, Y. Kato and T. F. Haraguchi for their help with field and laboratory
17 work, H. Nomaki for microscopic observations, K. Nozaki for the identification of algae and
18 cyanobacteria, Y. Miyairi for technical advice on AMS and Y. Takano and Y. Chikaraishi for
19 discussions about this study. Three anonymous referees provided valuable comments on an
20 early draft of the manuscript. This study was supported by the River Fund (25-1263-017) in
21 charge of the River Foundation, Japan. N.F.I. was supported by a Research Fellowship for
22 Young Scientists (25-1021) from the Japan Society for the Promotion of Science.

23

1 **References**

- 2 Allan, J. D. and Castillo, M. M.: Stream Ecology: Structure and Function of Running Waters,
3 2nd ed., Springer, Dordrecht, Germany, 2007.
- 4 Amir-Shapira, D., Goldschmidt, E. E., and Altman, A.: Chlorophyll catabolism in senescing
5 plant tissues: *In vivo* breakdown intermediates suggest different degradative pathways for
6 citrus fruit and parsley leaves. Proc. Natl Acad. Sci. U. S. A., 84, 1901–1905, 1987.
- 7 Beaumont, V. I., Jahnke, L. L., and Des Marais, D. J.: Nitrogen isotopic fractionation in the
8 synthesis of photosynthetic pigments in *Rhodobacter capsulatus* and *Anabaena cylindrica*.
9 Org. Geochem., 31, 1075–1085, 2000.
- 10 Berner, R. A., Lasaga, A. C., and Garrels, R. M.: The carbonate–silicate geochemical cycle
11 and its effect on atmospheric carbon dioxide over the past 100 million years. Am. J. Sci., 283,
12 641–683, 1983.
- 13 Bloemen, J., McGuire, M. A., Aubrey, D. P., Teskey, R. O., and Steppe, K.: Transport of
14 root-respired CO₂ via the transpiration stream affects aboveground carbon assimilation and
15 CO₂ efflux in trees. New Phytol., 197, 555–565, 2013.
- 16 Brüggemann, N., Gessler, A., Kayler, Z., Keel, S. G., Badeck, F., Barthel, M., Boeckx, P.,
17 Buchmann, N., Brugnoli, E., Esperschütz, J., Gavrichkova, O., Ghashghaie, J., Gomez-
18 Casanovas, N., Keitel, C., Knohl, A., Kuptz, D., Palacio, S., Salmon, Y., Uchida, Y., and
19 Bahn, M.: Carbon allocation and carbon isotope fluxes in the plant–soil–atmosphere
20 continuum: a review. Biogeosciences, 8, 3457–3489, 2011.
- 21 Carpenter, S. R., Elser, M. M., and Elser, J. J.: Chlorophyll production, degradation, and
22 sedimentation: Implications for paleolimnology. Limnol. Oceanogr., 31, 112–124, 1986.
- 23 Chikaraishi, Y., Matsumoto, K., Ogawa, N. O., Suga, H., Kitazato, H., and Ohkouchi, N.:
24 Hydrogen, carbon and nitrogen isotopic fractionations during chlorophyll biosynthesis in C3
25 higher plants. Phytochemistry, 66, 911–920, 2005.
- 26 Chikaraishi, Y., Matsumoto, K., Kitazato, H., and Ohkouchi, N.: Sources and transformation
27 processes of pheopigments: Stable carbon and hydrogen isotopic evidence from Lake Haruna,
28 Japan. Org. Geochem., 38, 985–1001, 2007.

- 1 Cross, W. F., Benstead, J. P., Frost, P. C., and Thomas, S.A.: Ecological stoichiometry in
2 freshwater benthic systems: recent progress and perspectives. *Freshwater Biol.*, 50, 1895–
3 1912, 2005.
- 4 Dekar, M. P., Magoulick, D. D., and Huxel, G. R.: Shifts in the trophic base of intermittent
5 stream food webs. *Hydrobiologia*, 635, 263–277, 2009.
- 6 DeNiro, M. J. and Epstein, S.: Influence of diet on distribution of carbon isotopes in animals.
7 *Geochim. Cosmochim. Acta*, 42, 495–506, 1978.
- 8 Doi, H., Takemon, Y., Ohta, T., Ishida, Y., and Kikuchi, E.: Effects of reach-scale canopy
9 cover on trophic pathways of caddisfly larvae in a Japanese mountain stream. *Mar.*
10 *Freshwater Res.*, 58, 811–817, 2007.
- 11 Eglinton, T. I., Aluwihare, L. I., Bauer, J. E., Druffel, E. R., and McNichol, A.P.: Gas
12 chromatographic isolation of individual compounds from complex matrices for radiocarbon
13 dating. *Anal. Chem.*, 68, 904–912, 1996.
- 14 Fellman, J. B., Hood, E., Raymond, P. A., Hudson, J., Bozeman, M., and Arimitsu, M.:
15 Evidence for the assimilation of ancient glacier organic carbon in a proglacial stream food
16 web. *Limnol. Oceanogr.*, 2015, in press.
- 17 Finlay, J. C.: Stable-carbon-isotope ratios of river biota: Implications for energy flow in lotic
18 food webs. *Ecology*, 82, 1052–1064, 2001.
- 19 Finlay, J. C.: Patterns and controls of lotic algal stable carbon isotope ratios. *Limnol.*
20 *Oceanogr.*, 49, 850–861, 2004.
- 21 Finlay, J. C., Power, M. E., and Cabana, G.: Effects of water velocity on algal carbon isotope
22 ratios: Implications for river food web studies. *Limnol. Oceanogr.*, 44, 1198–1203, 1999.
- 23 Fischer, H.: The role of biofilms in the uptake and transformation of dissolved organic matter,
24 in: Findlay, S. E. G. and Sinsabaugh, R. L. (eds) *Aquatic ecosystems: interactivity of*
25 *dissolved organic matter*. Academic Press, San Diego, 285–313, 2003.
- 26 Hall, R. O. Jr, Wallace, J. B., and Eggert, S. L.: Organic matter flow in stream food webs with
27 reduced detrital resource base. *Ecology*, 81, 3445–3463, 2000.
- 28 Hamilton, S. K. and Lewis, W. M.: Stable carbon and nitrogen isotopes in algae and detritus
29 from the Orinoco River floodplain, Venezuela. *Geochim. Cosmochim. Acta*, 56, 4237–4246,
30 1992.


- 1 Hayes, J. M., Takigiku, R., Ocampo, R., Callot, H. J., and Albrecht, P.: Isotopic compositions
2 and probable origins of organic molecules in the Eocene Messel shale. *Nature*, 329, 48–51,
3 1987.
- 4 Hladyz, S., Cook, R. A., Petrie, R., and Nielsen, D. L.: Influence of substratum on the
5 variability of benthic biofilm stable isotope signatures: implications for energy flow to a
6 primary consumer. *Hydrobiologia*, 664, 135–146, 2011.
- 7 Higgins, M. B., Robinson, R. S., Husson, J. M., Carter, S. J., and Pearson, A.: Dominant
8 eukaryotic export production during ocean anoxic events reflects the importance of recycled
9 NH_4^+ . *Proc. Natl Acad. Sci. U. S. A.*, 109, 2269–2274, 2012.
- 10 Imberger, S. P., Grace, C. M., and Thompson, R.: Tracing carbon sources in small urbanising
11 streams: catchment-scale stormwater drainage overwhelms the effects of reach-scale riparian
12 vegetation. *Freshwater Biol.*, 59, 168–186, 2014.
- 13 Ishikawa, N. F., Doi, H., and Finlay, J. C.: Global meta-analysis for controlling factors on
14 carbon stable isotope ratios of lotic periphyton. *Oecologia*, 170, 541–549, 2012a.
- 15 Ishikawa, N. F., Uchida, M., Shibata, Y., and Tayasu, I.: Natural C-14 provides new data for
16 stream food-web studies: a comparison with C-13 in multiple stream habitats. *Mar.
17 Freshwater Res.*, 63, 210–217, 2012b.
- 18 Ishikawa, N. F., Hyodo, F., and Tayasu, I.: Use of carbon-13 and carbon-14 natural
19 abundances for stream food web studies. *Ecol. Res.*, 28, 759–769, 2013.
- 20 Ishikawa, N. F., Kato, Y., Togashi, H., Yoshimura, M., Yoshimizu, C., Okuda, N., and Tayasu,
21 I.: Stable nitrogen isotopic composition of amino acids reveals food web structure in stream
22 ecosystems. *Oecologia*, 175, 911–922, 2014a.
- 23 Ishikawa, N. F., Uchida, M., Shibata, Y., and Tayasu, I.: Carbon storage reservoirs in
24 watersheds support stream food webs via periphyton production. *Ecology*, 95, 1264–1271,
25 2014b.
- 26 Ishikawa, N. F., Tayasu, I., Yamane, M., Yokoyama, Y., Sakai, S., and Ohkouchi, N.: Sources
27 of dissolved inorganic carbon in two small streams with different bedrock geology: insights
28 from carbon isotopes. *Radiocarbon*, 57, 439–448, 2015.
- 29 Ischebeck, T., Zbierzak, A. M., Kanwischer, M., and Dörmann, P.: A salvage pathway for
30 phytol metabolism in *Arabidopsis*. *J. Biol. Chem.*, 281, 2470–2477, 2006.

- 1 Jochmann, M. A. and Schmidt, T. C.: Compound-specific stable isotope analysis. Roy. Soc.
2 Ch., 376, 2012.
- 3 Junk, W. J., Bayley, P. B., and Sparks, R. E.: The flood pulse concept in river-floodplain
4 system. In: Dodge, D. P. (eds) Proceedings of the International Large River Symposium. Can.
5 Spec. Publ. Fish. Aquat. Sci., 106, 110–127, 1989.
- 6 Kennicutt, M. C. II, Bidigare, R. R., Macko, S. A., and Keeney-Kennicutt, W. L.: The stable
7 isotopic composition of photosynthetic pigments and related biochemicals. Chem. Geol., 101,
8 235–245, 1992.
- 9 Kitagawa, H., Masuzawa, T., Nakamura, T., and Matsumoto, E.: A batch preparation method
10 for graphite targets with low background for AMS ^{14}C measurements. Radiocarbon, 35, 295–
11 300, 1993.
- 12 Koarashi, J., Atarashi-Andoh, M., Ishizuka, S., Miura, S., Saito, T., Hirai, K.: Quantitative
13 aspects of heterogeneity in soil organic matter dynamics in a cool-temperate Japanese beech
14 forest: a radiocarbon-based approach. Glob. Change. Biol., 15, 631–642, 2009.
- 15 Kusch, S., Kashiyama, Y., Ogawa, N. O., Altabet, M., Butzin, M., Friedrich, J., Ohkouchi, N.,
16 and Mollenhauer, G.: Implications for chloro- and pheopigment synthesis and preservation
17 from combined compound-specific $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and $\Delta^{14}\text{C}$ analysis. Biogeosciences, 7, 4105–
18 4118, 2010.
- 19 Kuwae, T., Beninger, P. G., Decottignies, P., Mathot, K. J., Lund, D. R., and Elner, R. W.:
20 Biofilm grazing in a higher vertebrate: the western sandpiper, *Calidris mauri*. Ecology, 89,
21 599–606, 2008.
- 22 Kuwae, T., Miyoshi, E., Hosokawa, S., Ichimi, K., Hosoya, J., Amano, T., Moriya, T.,
23 Kondoh, M., Ydenberg, R. C., and Elner, R. W.: Variable and complex food web structures
24 revealed by exploring missing trophic links between birds and biofilm. Ecol. Lett., 15, 347–
25 356, 2012.
- 26 Levin, I., Kromer, B., and Hammer, S.: Atmospheric $\Delta^{14}\text{CO}_2$ trend in Western European
27 background air from 2000 to 2012. Tellus B, 65, 20092, 2013.
- 28 Matile, P., Hortensteiner, S., Thomas, H., and Krautler, B.: Chlorophyll breakdown in
29 senescent leaves. Plant Physiol., 112, 1403–1409, 1996.

- 1 Minagawa, M. and Wada, E.: Stepwise enrichment of ^{15}N along food chains: further evidence
2 and the relation between $\delta^{15}\text{N}$ and animal age. *Geochim. Cosmochim. Acta*, 48, 1135–1140,
3 1984.
- 4 Ogawa, N. O., Nagata, T., Kitazato, H., and Ohkouchi, N.: Ultra-sensitive elemental
5 analyzer/isotope ratio mass spectrometer for stable nitrogen and carbon isotope analyses, in:
6 Ohkouchi, N., Tayasu, I., and Koba, K. (eds) *Earth, Life, and Isotopes*, Kyoto University
7 Press, Kyoto, Japan, 339–353, 2010.
- 8 Ohkouchi, N., Nakajima, Y., Okada, H., Ogawa, N. O., Suga, H., Oguri, K., and Kitazato, H.:
9 Biogeochemical processes in the saline meromictic Lake Kaiike, Japan: implications from
10 molecular isotopic evidences of photosynthetic pigments. *Environ. Microbiol.*, 7, 1009–1016,
11 2005.
- 12 Ohkouchi, N., Nakajima, Y., Ogawa, N. O., Chikaraishi, Y., Suga, H., Sakai, S., and Kitazato,
13 H.: Carbon isotopic composition of the tetrapyrrole nucleus in chloropigments from a saline
14 meromictic lake: A mechanistic view for interpreting the isotopic signature of alkyl
15 porphyrins in geological samples. *Org. Geochem.*, 39, 521–531, 2008.
- 16 Ohkouchi, N., Ogawa, N. O., Chikaraishi, Y., Tanaka, H., and Wada, E.: Biochemical and
17 physiological bases for the use of carbon and nitrogen isotopes in environmental and
18 ecological studies. *Progr. Earth Planetary Sci.*, 2, 1–17, 2015.
- 19 Sachs, J. P., Repeta, D. J., and Goericke, R.: Nitrogen and carbon isotopic ratios of
20 chlorophyll from marine phytoplankton. *Geochim. Cosmochim. Acta*, 65, 1431–1441, 1999.
- 21 Sachs, J. P. and Repeta, D. J.: The purification of chlorins from marine particles and
22 sediments for nitrogen and carbon isotopic analysis. *Org. Geochem.*, 31, 317–329, 2000.
- 23 Small, G. E., Bixby, R. J., Kazanci, C., and Pringle, C. M.: Partitioning stoichiometric
24 components of epilithic biofilm using mixing models. *Limnol. Oceanogr. Meth.*, 9, 185–193.
25 2011.
- 26 Stuiver, M. and Polach, H. A.: Discussion: Reporting of ^{14}C data. *Radiocarbon*, 19, 355–363,
27 1977.
- 28 Takemon, Y.: Life-type concept and functional feeding groups of benthos communities as
29 indicators of lotic ecosystem conditions. *Jpn. J. Ecol.*, 55, 189–197, 2005 [in Japanese].


- 1 Tayasu, I., Hirasawa, R., Ogawa, N. O., Ohkouchi, N., and Yamada, K.: New organic
2 reference materials for carbon and nitrogen stable isotope ratio measurements provided by
3 Center for Ecological Research, Kyoto University and Institute of Biogeosciences, Japan
4 Agency for Marine-Earth Science and Technology. Limnology, 12, 261–266, 2011.
- 5 Thorp, J. H. and DeLong, M. D.: The riverine productivity model: an heuristic view of carbon
6 sources and organic processing in large river ecosystems. Oikos, 70, 305–308, 1994.
- 7 Tyler, J., Kashiyama, Y., Ohkouchi, N., Ogawa, N. O., Yokoyama, Y., Chikaraishi, Y., Staff,
8 R. A., Ikehara, M., Bronk Ramsey, C., Bryant, C., Brock, F., Gotanda, K., Haraguchi, T.,
9 Yonenobu, H., and Nakagawa, T.: Tracking aquatic change using chlorin-specific carbon and
10 nitrogen isotopes: The last glacial–interglacial transition at Lake Suigetsu, Japan. Geochem.
11 Geophy. Geosy., 11, Q09010, 2010.
- 12 Trumbore, S. and Zheng, S.: Comparison of fractionation methods for soil organic matter ^{14}C
13 analysis. Radiocarbon, 38, 219–229, 1996.
- 14 Vander Zanden, M. J. and Rasmussen, J. B.: Variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ trophic fractionation:
15 implications for aquatic food web studies. Limnol. Oceanogr., 46, 2061–2066, 2001.
- 16 Vannote, R. L., Minshall, G. W., Cummins, K. W., Sedell, J. R., and Cushing, C. E.: The river
17 continuum concept. Can. J. Fish. Aquat. Sci., 37, 130–137, 1980.
- 18 Vavilin, D. and Vermaas, W.: Continuous chlorophyll degradation accompanied by
19 chlorophyllide and phytol reutilization for chlorophyll synthesis in *Synechocystis* sp. PCC
20 6803. Biochim. Biophys. Acta, 1767, 920–929, 2007.
- 21 Winemiller, K. O.: Spatial and temporal variation in tropical fish trophic networks. Ecol.
22 Monogr., 60, 331–367, 1990.
- 23 Whitledge, G. W. and Rabeni, C. F.: Energy sources and ecological role of crayfishes in an
24 Ozark stream: insights from stable isotopes and gut analysis. Can. J. Fish. Aquat. Sci., 54,
25 2555–2563, 1997.
- 26 Yokoyama, Y., Koizumi, M., Matsuzaki, H., Miyairi, Y., and Ohkouchi, N.: Developing ultra
27 small-scale radiocarbon sample measurement at the University of Tokyo. Radiocarbon, 52,
28 310–318, 2010.

1 Zah, R., Burgherr, P., Bernasconi, S. M., and Uehlinger, U.: Stable isotope analysis of
2 macroinvertebrates and their food sources in a glacier stream. Freshwater Biol., 46, 871–882,
3 2001.
4




1 Figure 1. The $\Delta^{14}\text{C}_{\text{bulk}}$, $\delta^{13}\text{C}_{\text{bulk}}$ and $\delta^{15}\text{N}_{\text{bulk}}$ values (shaded symbols) for periphyton
2 (diamonds), *Cladophora* sp. (aquatic primary producer; circle), *Q. glauca* (terrestrial primary
3 producer; square) and *E. latifolium* (algal grazer; triangles) and the $\Delta^{14}\text{C}_{\text{chl}}$, $\delta^{13}\text{C}_{\text{chl}}$ and $\delta^{15}\text{N}_{\text{chl}}$
4 values (open symbols) for periphyton, *Cladophora* sp. and *Q. glauca*. DIC: dissolved
5 inorganic carbon; DOC: dissolved organic carbon; POC: particulate organic carbon. *Data
6 from Ishikawa et al. (2012b, 2015). **Error bars indicate standard deviation (N = 4).**

8



1
2
3
4
5
6
7

Figure 2. Biplot of $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ data. Carbonate rocks in the Seri River ($\delta^{13}\text{C} = +3.9 \pm 0.3\text{\textperthousand}$ and $\Delta^{14}\text{C} = -1000\text{\textperthousand}$) (Ishikawa et al., 2015) and atmospheric CO_2 ($\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ are approximately $-8\text{\textperthousand}$ and $+30\text{\textperthousand}$, respectively, in 2013) are also shown as end members. Error bars indicate standard deviation ($N = 4$).



1
2
3 Figure 3. Schematic view of the carbon cycle at the study site (Seri River) constrained by $\delta^{13}\text{C}$
4 and $\Delta^{14}\text{C}$.
5