

## ***Interactive comment on “Microbial carbon recycling: an underestimated process controlling soil carbon dynamics” by A. Basler et al.***

**A. Basler et al.**

abasler@gwdg.de

Received and published: 14 October 2015

Answer to Anonymous Referee #1

We would like to thank the reviewer for their helpful comments and suggestions, which have greatly improved our manuscript. We hope that our response answers all their concerns.

Interactive comment on “Microbial carbon recycling: an underestimated process controlling soil carbon dynamics” by A. Basler et al.

Anonymous Referee #1

Received and published: 22 July 2015

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



---

[Interactive  
Comment](#)

Basler et al. investigate stabilisation and recycling of soil sugars as processes controlling soil carbon dynamics. This is addressed by  $\delta^{13}\text{C}$  analyses using HPLC/IRMS of soil sugars in density fractions from a natural 30 year old labelling experiment with wheat-maize vegetation change. Overall, this is a well designed and presented study appropriate for publication in BG. The authors clearly state the motivation/relevance of this study for better understanding turnover dynamics of sugars in the introduction, formulate a clear working hypothesis at the end of the introduction (turnover of plant-derived sugars is ruled by stabilization versus turnover of microbial-derived sugars is ruled by recycling) and provide all necessary information where and how the study was performed in the Material and Method section. The authors found that the contribution of maize-derived carbon in the POM fractions is considerably higher in sugars compared to the bulk fractions, equivalent to mean residence times (MRT) being lower for sugars than for bulk C in these fractions. This is interpreted in terms of aggregate formation being fuelled by microbial activity and fresh organic matter. Concerning the working hypothesis, the authors found that the MRT of xylose is considerably lower than the MRT of the other sugars. The authors argue that xylose (plant-derived) dynamic is primarily dominated by stabilization, whereas the dynamic of the other sugars (microbial-derived) is strongly controlled by recycling. Interestingly, this also holds true for arabinose; this is well highlighted and discussed by the authors. However, as alternative interpretation, I would like to suggest (and the authors may want to include this in their discussion) that the dynamic of arabinose, like that of xylose, is primarily controlled by stabilisation (not by recycling). The arabinose/xylose ratio is close to 1 in the soil fractions, possibly because former vegetation contributed relatively high amounts of arabinose to the soil. The addition of wheat/maize sugars with low ara:xyl ratios (1:6 and 1:5, respectively) thus resulted in a low admixture of maize-derived arabinose during the last 30 years, while the admixture with maize-derived xylose was much higher.

Answer: We agree that the proportions of ara and xyl, as well as the other sugars in soil depends on the vegetation. We still do not think that turnover of ara is substantially

C6473

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

influenced by stabilization, as several studies (Basler et al., 2015; Coelho et al., 1988; ) show that arabinose is substantially influenced by the microbial biomass. We do not entirely exclude stabilization, though the results also of our other study show that an influence by recycling to ara seems to be much higher.

Furthermore, if I understand right, the authors have a second MS under review in BGD also focussing on stabilisation versus recycling of soil sugars. Hence, in order to increase the impact of their papers, it may be advantageous to publish both papers as companion papers with a) similar titles, e.g.: 1) Recycling vs. stabilisation of soil sugars –: an underestimated process controlling soil carbon dynamics II – I) a natural 30 yrs old labelling field experiment

2) Recycling vs. stabilisation of soil sugars –: an underestimated process controlling soil carbon dynamics II) a long-term laboratory incubation experiment

b) establishing clear links between these two papers. So far, this is unfortunately not done at all..

Answer: Thanks for this suggestion, we followed your advice and renamed our manuscripts.

1. Microbial carbon recycling: an underestimated process controlling soil carbon dynamics I) a long-term laboratory incubation experiment

2. Microbial carbon recycling: an underestimated process controlling soil carbon dynamics II) a C3-C4 vegetation change field labelling experiment

Minor issues:

- The sugar analyses are performed (in contrast to Amelung et al., 1996) with Serdolit. Is there a reason why you did not use XAD resin as in the original procedure?

Answer: At the time we started our experiment the AMBERLITE® XAD-7 resin was not shippable in the laboratory equipment shops and thus we used an alternative product,

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



the Serdolit PADIV, which was advertised as an alternative resin for the XAD7.

- When emphasizing the importance of recycling dynamics, position-specific d13C differences/methods (co-author M.D. is well known for her excellent expertise on this field) are or will at least soon become of high importance. Hence, the readers will profit from one or two respective sentences and references (maybe in a Conclusion and Outlook chapter)

Answer: We included a brief outlook on the perspectives of position-specific labeling in the manuscript.

- Fig. 1: Please specify what for light/dark grey bars stand for (I guess you mean Ap and E horizons, respectively)

Answer: We apologize for this error, and we have corrected the figure. The light grey presents the Ap horizon and dark grey the E-horizon.

- Table 1: The carbon contents of the POM fractions seem to be very/too low, please check and correct if necessary.

Answer: The given values for the carbon content are correct according to our measurements. The very low numbers result from the fact that the investigated arable soil generally contains only low amounts of POM (see e.g. John et al. (2005) or Helfrich et al. (2006)). Additionally, sampling had taken place early in spring (April) - the season when plant growth had just started and therefore, amounts of POM were generally low.

## References

Basler, A., Dippold, M., Helfrich, M., and Dyckmans, J.: Recycling versus Stabilisation of soil sugars— a long-term laboratory incubation experiment, *Biogeosciences Discuss*, 12, 8819–8847, doi:10.5194/bgd-12-8819-2015, 2015.

Coelho, R. R., Linhares, L. F., and Martin, J. P.: Sugars in hydrolysates of fungal melanins and soil humic acids, *Plant Soil*, 106, 127–133, doi:10.1007/BF02371204,

**BGD**

12, C6472–C6476, 2015

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



1988.

Helfrich, M., Ludwig, B., Buurman, P., and Flessa, H.: Effect of land use on the composition of soil organic matter in density and aggregate fractions as revealed by solid-state C-13 NMR spectroscopy, *Geoderma*, 136, 331–341, 2006.

John, B., Yamashita, T., Ludwig, B., and Flessa, H.: Storage of organic carbon in aggregate and density fractions of silty soils under different types of land use, *Geoderma*, 128, 63–79, doi:10.1016/j.geoderma.2004.12.013, 2005.

---

Interactive comment on *Biogeosciences Discuss.*, 12, 9729, 2015.

**BGD**

12, C6472–C6476, 2015

---

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C6476

