Biogeosciences Discuss., 12, C7612–C7614, 2015 www.biogeosciences-discuss.net/12/C7612/2015/ © Author(s) 2015. This work is distributed under the Creative Commons Attribute 3.0 License.



BGD 12, C7612–C7614, 2015

> Interactive Comment

Interactive comment on "Vanishing coccolith vital effects with alleviated CO₂ limitation" by M. Hermoso et al.

M. Hermoso et al.

michael.hermoso@earth.ox.ac.uk

Received and published: 12 November 2015

Dear Dr Holtz,

Indeed, most of the linear correlations shown in the Discussion Paper between coccolith isotope fractionation and the Bidigare index $[(\mu \times \text{volume}) / ([CO_2 a_q \times \text{surface-area})]$ can be largely explained by changes in ambient CO₂ concentrations (see Figure 1 below) – for *C. pelagicus*, however, cell size and division rate seem important. This important point ought to be emphasised in our manuscript and we thank you very much for drawing our attention of this. The array of CO_{2 aq} concentrations is significant and induces prominent effects on algal physiology (division rates cell geometry) via carbon availability, and most importantly for our study on stable isotope





compositions of coccolith calcite.

Therefore, it may appear more suitable to show the correlations with CO_{2 aq} concentrations in the revised manuscript (combined Figures 2 and 3) as per your suggestion, but we anticipate that colleagues, especially those working on palaeo-CO₂ barometry from alkenone δ^{13} C would like to see how the Bidigare index that they use for organic carbon may work for inorganic carbon (taking into account your previous comments). This could be done as a multi panel figure (as shown below) or as a *Supplement*.

Thanks again for your help with this manuscript!

Michaël Hermoso

BGD

12, C7612-C7614, 2015

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive comment on Biogeosciences Discuss., 12, 15835, 2015.





Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Fig. 1. Raw isotopic composition of cultured coccolith calcite plotted against ambient CO2 concentrations (Top panels) and the "Bidigare Index" (Bottom panels).