

Interactive comment on “Pleistocene sediment offloading and the global sulfur cycle” by S. Markovic et al.

S. Markovic et al.

stefa.markovic@utoronto.ca

Received and published: 30 March 2015

Please see supplement for formatted response.

Comments by B. Mayer

The authors have composed a well-written manuscript describing and interpreting a high-resolution sulfur isotope record of marine sulfate determined on authigenic marine barite crystals from an Eastern Equatorial Pacific ODP site spanning the last 3 million years. The manuscript is written in a logical sequence, is well organized, easy to read and understand. Previous literature is appropriately considered and figures and tables are of good quality. Most conclusions appear well supported by the provided data and by the utilized sulfur cycling model. The analytical approaches are very sophisticated

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



giving confidence in the presented data. The authors describe a $>1.0\%$ decrease in $\delta^{34}\text{S}$ of marine sulfate from circa 22.0% to $<21.0\%$ with the majority of the change occurring between 1.5 and 1.0 Ma. The authors use a sulfur cycle model to conclude that erosion during sea level low-stands was only partly compensated by increased sedimentation during times of sea level high-stands, an interpretation that appears well justified based on the presented data and model runs.

We thank Dr Mayer for his support of the study.

1) The authors suggest that the observed decrease in $\delta^{34}\text{S}$ is related to the Milankovic cycle driven change from 41 kyr (and 23 kyr) interglacial-glacial periodicity earlier in the Pleistocene to 100 kyr by 700,000 years ago. While the end of the reported $\delta^{34}\text{S}$ decrease described in this manuscript is consistent with this interpretation, the question arises why the decrease in $\delta^{34}\text{S}$ values of marine sulfate started as early as 1.5 Ma ago if it is linked to 100 kya cycles? To my best knowledge, oxygen isotope records of benthic foraminifera place this transition somewhere between 1.0 and 0.7 Ma ago (e.g. Ruddiman, W. F. (2008): *Earth's Climate*, Freeman & Co, New York), but your Figure 2 indicates that a 0.7% shift in $\delta^{34}\text{S}$ of marine sulfate had already occurred between 1.5 and 1.2 Ma ago, which appears inconsistent with the larger 100,000 yrs interglacial-glacial periodicity capable of removing much larger sediment loads from the shelf. The authors should add some explanation on the timing of this early onset of the decrease in $\delta^{34}\text{S}$ of marine sulfate reported in their paper, and possibly compare it to the $\delta^{18}\text{O}$ record of benthic foraminifera, which has reportedly been used for age-dating the samples (page 1209, line 3).

This is a very perceptive catch and something that we have overlooked. While the “true” 100kyr cycles did not appear until $\sim 900\text{ka}$ (e.g., Elderfield et al., 2012; Pena and Goldstein, 2014) the early signs of change in amplitude and duration of glaciations probably took place $\sim 1250\text{ka}$ (e.g., Clark et al., 2006; Lisiecki, 2014). The LR04 stack of globally distributed benthic foraminifera $\delta^{18}\text{O}$ records indicates increased severity of glaciations as early as 1.5Ma (Lisiecki and Raymo, 2005; Clark et al. 2006). More

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

pronounced sea level low stands from 1.5Ma to 0.7Ma are reflection of this change (see Figure 1 in discussion paper, data from Miller et al. 2011). It is this lower sea level during glaciations that resulted in increased weathering of shelf pyrite. However, the language we used obscured this main point and suggested that $\delta^{34}\text{S}$ shift is a result of longer glaciations, when in fact it is due to progressively lower sea level and associated aerial exposure of previously inundated continental shelf (Figure 1 in discussion paper). We thank the reviewer for pointing out this ambiguity and we changed the language throughout the text to make it clearer what is the main driver of shift in seawater sulfate $\delta^{34}\text{S}$ value.

2) I am also curious why organic sulfur compounds receive so little attention in the sulfur cycling models, but assume that they play only a very minor role in the reported S fluxes. Nevertheless it should be acknowledged (e.g. on page 1217, line 10) that some sulfur is also buried in organic form.

This is indeed something that we have overlooked. We thank the reviewer for this suggestion and we added the following: “Also note that the burial of organic sulfur is poorly constrained (Werne et al., 2004). Although it is generally considered to be minor compared to pyrite burial (e.g., Goldhaber, 2003; Werne et al., 2004), in some environments (e.g. Peru shelf, the Miocene Monterey Formation, marine sapropels) it is the dominant form of reduced solid sulfur compounds (e.g., Mossman et al., 1991; Canfield et al., 1998; Suits and Arthur, 2000).”

3) And finally, in section 4.2 the authors comment on the link between pyrite oxidation, acidification-enhanced carbonate dissolution/precipitation and associated CO_2 release into the atmosphere. While I agree with the principle geochemical arguments, I am a bit puzzled about the link of the here presented data showing a $\delta^{34}\text{S}$ decrease in marine sulfate starting 1.5 mio years ago and ending 700,000 years ago, and the claim that this is partly related to CO_2 increases 600kyr and 400kyr ago. The authors should provide further arguments on the temporal connections of their S isotope record with the ice core CO_2 record in order to substantiate their argument of a causal relationship.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

We thank you for this comment, which identified the area where we probably promised more than we can deliver at this point. The language we used in discussion might have erroneously suggested that there is a direct link between sulfide oxidation and atmospheric CO₂. However, this link is indirect and complex.

While sulfide weathering increases the dissolved inorganic carbon (DIC) amount in the ocean (Hu and Cai, 2011; Torres et al., 2014,) CO₂ is released only after precipitation of calcium carbonate. Characteristic time scales for this process are in the 100-200 kyr range (Sundquist and Visser, 2003; Zeebe and Wolf-Gladrow, 2009) and therefore, DIC addition and CO₂ release are not synchronized. It is possible that competing effects of biological pump and carbonate precipitation resulted in the increase of atmospheric CO₂ some time after the pulse of DIC produced during sulfide oxidation. However, the temporal changes of these processes are well beyond the scope of this paper. We therefore changed the language in the last section of discussion and dropped the suggestion that there is a direct link between CO₂ concentrations and increased pyrite weathering.

On a technical note, d₃₄S should be followed by “values” (d₃₄S values) rather than compositions” and certainly not “ratios”.

Thank you for the comment. We changed the language throughout the text.

Table A1: add units (‰ after d₃₄S and use same number of significant digits behind the comma for reporting the results;

Done.

Figure 4, y-axis label, unit missing (‰.

Corrected.

References: Canfield, D. E., Boudreau, B. P., Mucci, A. and Gundersen, J. K.: The early diagenetic formation of organic sulfur in the sediments of Mangrove Lake, Bermuda, *Geochim. Cosmochim. Acta*, 62, 767–781, doi:10.1016/S0016-7037(98)00032-5,

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

1998.

Clark, P. U., Archer, D., Pollard, D., Blum, J. D., Rial, J. A., Brovkin, V., Mix, A. C., Pisias, N. G. and Roy, M.: The middle Pleistocene transition: characteristics, mechanisms, and implications for long-term changes in atmospheric pCO₂, *Quat. Sci. Rev.*, 25, 3150–3184, doi:10.1016/j.quascirev.2006.07.008, 2006.

Elderfield, H., Ferretti, P., Greaves, M., Crowhurst, S., McCave, I. N., Hodell, D. and Piotrowski, A. M.: Evolution of Ocean Temperature and Ice Volume Through the Mid-Pleistocene Climate Transition, *Science* (80-.), 337, 704–709, doi:10.1126/science.1221294, 2012.

Goldhaber, M.: Sulfur-rich sediments, *Treatise on Geochemistry*, 254–288, doi:10.1016/B0-08-043751-6/07139-5, 2003.

Hu, X. and Cai, W. J.: An assessment of ocean margin anaerobic processes on oceanic alkalinity budget, *Global Biogeochem. Cycles*, 25, doi:10.1029/2010GB003859, 2011..

Lisiecki, L. E.: Atlantic overturning responses to obliquity and precession over the last 3 Myr, *Paleoceanography*, 29, 71–86, doi:10.1002/2013PA002505, 2014.

Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic d¹⁸O records, *Paleoceanography*, 20, 1–17, doi:10.1029/2004PA001071, 2005.

Miller, K., Mountain, G., Wright, J. and Browning, J.: A 180-Million-Year Record of Sea Level and Ice Volume Variations from Continental Margin and Deep-Sea Isotopic Records, *Oceanography*, 24, 40–53, doi:10.5670/oceanog.2011.26, 2011.

Mossmann, J.-R., Aplin, A. C., Curtis, C. D. and Coleman, M. L.: Geochemistry of inorganic and organic sulphur in organic-rich sediments from the Peru Margin, *Geochim. Cosmochim. Acta*, 55, 3581–3595, doi:10.1016/0016-7037(91)90057-C, 1991.

Pena, L. D. and Goldstein, S. L.: Thermohaline circulation crisis and im-

BGD

12, C938–C943, 2015

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



pacts during the mid-Pleistocene transition., *Science* (80-), 345, 318–322, doi:10.1126/science.1249770, 2014.

Suits, N. S. and Arthur, M. A.: Sulfur diagenesis and partitioning in Holocene Peru shelf and upper slope sediments, *Chem. Geol.*, 163, 219–234, doi:10.1016/S0009-2541(99)00114-X, 2000.

Sundquist, E. T. and Visser, K.: The Geologic History of the Carbon Cycle, in *Treatise on Geochemistry*, vol. 8, edited by W. H. H. D. H. and K. K. T. Schlesinger., pp. 425–472, Elsevier., 2003.

Torres, M. a, West, a J. and Li, G.: Sulphide oxidation and carbonate dissolution as a source of CO₂ over geological timescales., *Nature*, 507, 346–9, doi:10.1038/nature13030, 2014.

Werne, J. P., Hollander, D. J., Lyons, T. W. and Sinninghe Damsté, J. S.: Organic sulfur biogeochemistry: Recent advances and future research directions, *Geol. Soc. Am. Spec. Pap.*, 379, 135–150, doi:10.1130/0-8137-2379-5.135, 2004.

Zeebe, R. E. and Wolf-Gladrow, D. A.: Carbon Dioxide, Dissolved (Ocean), in *Encyclopedia of Paleoclimatology and Ancient Environments*, edited by V. Gornitz, pp. 123–127, Springer Netherlands, Dordrecht., 2009.

Please also note the supplement to this comment:

<http://www.biogeosciences-discuss.net/12/C938/2015/bgd-12-C938-2015-supplement.pdf>

Interactive comment on *Biogeosciences Discuss.*, 12, 1205, 2015.

BGD

12, C938–C943, 2015

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

