

***Interactive comment on “Inhibition of nitrogenase by oxygen in marine cyanobacteria controls the global nitrogen and oxygen cycles” by I. Berman-Frank et al.***

**Anonymous Referee #3**

Received and published: 25 May 2005

This manuscript is essentially a rehash of previously-published discussions about the "oxygen problem" as a constraint on N<sub>2</sub> fixing potential of many diazotrophs, including heterocystous and non-heterocystous cyanobacteria, i.e. Trichodesmium. The explanations for and arguments pertaining to this problem have been made many times before, and I might add, in much more thorough and well-supported ways. There is nothing new here, except for maybe the rewording of somewhat ironic (pardon the pun) paradox that photosynthetic oxygen evolution (which cyanobacteria "invented") has turned out to be a constraint on the extent and magnitude of N<sub>2</sub> fixing potential among the cyanobacteria. This has been more thoroughly articulated by others, including Gallon (1992), Paerl (1990) and Paerl and Zehr (2000). In fact, Paerl (1994) experimentally showed the incompatibility of these processes at high rates of photosynthesis (i.e. O<sub>2</sub>

Full Screen / Esc

Print Version

Interactive Discussion

Discussion Paper

evolution) in *Trichodesmium* 101 well over 10 years ago.

While some interesting molecular response mechanisms to varying ambient O<sub>2</sub> levels were discussed, these have also been presented and discussed far more comprehensively and thoroughly by Zehr et al. (1997). Again, there's nothing new of major significance presented in the current manuscript.

Lastly, all the evolutionary arguments presented here have been made before (see the references cited below).

In summary, this manuscript adds little, if anything, to our understanding of the physiological, ecological, evolutionary and biogeochemical implications of the "oxygen problem". In addition, the authors left out key manuscripts and chapters that previously presented and discussed this issue. I do not recommend publication or for that matter further discussion of this manuscript.

Some specific comments on the manuscript:

P. 262, Abstract. First sentence: Where is the evidence that "N<sub>2</sub> fixation supplies the vast majority of biologically accessible inorganic nitrogen to nutrient-poor aquatic ecosystems"? This seems like a gross and unsupportable generalization. There are certain regions of the world's oceans where N<sub>2</sub> fixation is a significant new N source (i.e. Baltic Sea, parts of the Gulf Stream and Kurishio Current), but these are only a fraction of the Earth's "nutrient-poor aquatic ecosystems".

P. 262, Introduction, line 1. The statement that "Only a small fraction of prokaryotic organisms from the bacterial and archaeal domains can procure and utilize atmospheric nitrogen by reducing it to ammonia" is naïve and incorrect. The process of N<sub>2</sub> fixation is actually fairly broadly distributed among different phylogenetic and physiological groups. In fact, this is probably one reason why the genes encoding for it are so highly conserved in the first place.

P. 263, lines 16 and 17: the part of this sentence "restrictions on nitrogen" makes no

[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)[Discussion Paper](#)

sense. Something appears to be missing here. Also in line 18, it isn't clear what "real-world" means (as opposed to "unreal world"?). Lines 23-24: *Trichodesmium* 101 was initially isolated from North Carolina Atlantic coastal waters (see Prufert-Bebout et al. 1993). It is not known whether this strain "contributes significantly to nitrogen fixation in the tropical and subtropical oceans". This statement is one of numerous examples of gross overgeneralization and unsubstantiated armwaiving that seem to characterize this manuscript. These types of statements are both factually incorrect and add little relevant information.

P. 267, line 6: What does "chronically crippled nitrogenase" mean? Nitrogenase could be functioning at sub-optimal rates due to several (interactive) limiting factors over moderate to long time scales (i.e. days to weeks) and this may not be linked to the O<sub>2</sub> problem. Fe and/or P limitation may be operating over such time scales, or high degrees of mixing and smaller scale turbulence may lead to sub-optimal rates of N<sub>2</sub> fixation.

Line 22: What does "efficiency" mean here? This may not be an "inefficiency" problem at all, if we consider the interactive effects of multiple potential limiting factors (i.e. P, Fe, organic C, turbulence).

Additional references of relevance (not cited in manuscript).

Paerl, H. W. 1990. Physiological ecology and regulation of N<sub>2</sub> fixation in natural waters. *Adv. Microb. Ecol.* 11:305-344.

Paerl, H. W. 1994. Spatial segregation of CO<sub>2</sub> fixation in *Trichodesmium* sp.: Linkage to N<sub>2</sub> fixation potential. *J. Phycol.* 30:790-799.

Paerl, H. W. and J. P. Zehr. 2000. Nitrogen Fixation. Pp. 387-426., In, D. Kirchman [Ed.], *Microbial Ecology of the Oceans*, Academic Press, New York.

Prufert-Bebout, L. E., H. W. Paerl and C. Lassen. 1993. Growth, nitrogen fixation and spectral attenuation in cultivated *Trichodesmium*. *Appl. Environ. Microbiol.* 59:1350-1359.

[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive  
Comment

Full Screen / Esc

Print Version

Interactive Discussion

Discussion Paper