

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

**Metabolic human
CO₂ release**

Y. T. Prairie and
C. M. Duarte

Direct and indirect metabolic CO₂ release by humanity

Y. T. Prairie¹ and C. M. Duarte²

¹Département des sciences biologiques, Université du Québec à Montréal, P.O. Box 8888,
succ. Centre-Ville, Montréal, H3C 3P8, Canada

²IMEDEA (CSIC-UiB), Instituto Mediterráneo de Estudios Avanzados, C/ Miquel Marqués 21,
07190 Esporles (Mallorca), Spain

Received: 26 October 2006 – Accepted: 16 November 2006 – Published: 21 November 2006

Correspondence to: C. M. Duarte (carlosduarte@imedea.uib.es)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The direct CO₂ released by respiration of humans and domesticated animals, as well as the CO₂ derived from the decomposition of their resulting wastes was calculated in order to ascertain the direct and indirect metabolic contribution of humanity to CO₂ release. Human respiration was estimated to release 0.6 Gt C year⁻¹ and that of their associated domestic animals was estimated to release 1.5 Gt C year⁻¹, to which an indirect release of 1.0 Gt C year⁻¹, derived from decomposition of the organic waste and garbage produced by humans and their domestic animals, must be added. These combined direct and indirect metabolic sources, estimated at 3.1 Gt C year⁻¹, has increased 7 fold since pre-industrial times and is forecasted to continue to rise over the 21st century.

1 Introduction

Rapid increase in atmospheric CO₂ has prompted intense research efforts at elucidating and quantifying the sources and sinks of CO₂. The depictions of the global carbon budget consider the role of humans through the CO₂ released from fossil fuel combustion and changes in land use (IPCC, 2001), but does not explicitly estimate the metabolic CO₂ released by respiration by humans, as well as the CO₂ derived from the decomposition of their resulting wastes. In addition, humans maintain a large livestock population to feed themselves as well as a large number of domestic animals for other services, which release CO₂ both via direct respiration as well as via the decomposition of their wastes. We assess here the magnitude of these combined direct and indirect metabolic CO₂ sources and their likely increase since pre-industrial times, to ascertain their role on the global CO₂ budget.

BGD

3, 1781–1789, 2006

Metabolic human CO₂ release

Y. T. Prairie and
C. M. Duarte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

2 Methods

Per capita respiration (R , in watts) and excretion (E , in watts) rates by humans and domestic animals were derived from body weight (W , in kg) using allometric relationships that depict standard metabolic and defecation rates as a power function of body weight as: $R=3.89 W^{0.79}$ and defecation rate $E=3.82 W^{0.63}$ (Peters, 1983). The per capita respiratory CO_2 released by humans was also derived from direct respirometry measurements (Marrieb, 2000). Per capita organic waste production was extrapolated from Organisation for Economic Co-operation and Development figures (OECD, 2002). Human population size was derived from estimates of the past, present and projected estimates (Cohen, 1995; Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat, 2002), and the abundance of livestock was derived from 2004 FAO statistics (<http://faostat.fao.org>).

3 Results and discussion

The estimates of the metabolic CO_2 release from human respiration obtained as the product of present human population size and per capita respiratory CO_2 release estimated either from allometric relationships or from direct respirometric measurements were $0.6 \text{ Gt C year}^{-1}$ and $0.57 \text{ Gt C year}^{-1}$ (Table 1), respectively. We estimated that respiration by decomposers of the excreted organic matter releases an additional $0.29 \text{ Gt C year}^{-1}$ (Table 1). Humans also produce non-metabolic organic waste (as garbage) amounting to about $0.1 \text{ Gt C year}^{-1}$, which must again be largely decomposed to CO_2 by bacterial respiration.

The population of domestic animals maintained by humans exceeds human population by three fold (Table 1). The combined respiratory CO_2 release by domesticated animals is estimated at $1.5 \text{ Gt C year}^{-1}$, and decomposer respiration of their excreted products adds about $0.6 \text{ Gt C year}^{-1}$. The total metabolic release by domesticated animals of $2.1 \text{ Gt C year}^{-1}$, must be considered an indirect human metabolic CO_2 re-

BGD

3, 1781–1789, 2006

Metabolic human CO_2 release

Y. T. Prairie and
C. M. Duarte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

lease, for these animals only exist in such large numbers to provide food and service to humans.

The total direct and indirect metabolic CO₂ release by humans amounts to an estimated 3.1 Gt C year⁻¹ (Table 1). This figure is a conservative estimate of the metabolic CO₂ release associated to human activity, for humans now use about 30–40% of the terrestrial (Vitousek et al., 1986) and 8% of the marine (Pauly and Christensen, 1995) primary production, most of which is eventually decomposed to CO₂.

The calculated direct and indirect metabolic CO₂ release by humans represents about half of the CO₂ released from fossil fuel combustion, and nearly twice that released from changes in land use (IPCC, 2001). Whereas the role of the CO₂ released by fossil fuel combustion has been extensively discussed, that of the anthropogenic metabolic CO₂ release has not. This is likely because anthropogenic metabolic CO₂ release may be considered just an intensification of cycling process between the atmosphere and the biosphere via enhanced crop and pasture production. However, the replacement of natural ecosystems by pasture and crops represents a net human-derived CO₂ emission through a decline in sink capacity, as essentially all of the production of crops and pastures is cropped and eventually decomposed to CO₂. Acknowledging this to be a result of metabolic CO₂ release by humans provides a more direct identification of source components and a better appreciation of the consequences of demographic changes on this source component.

Whereas the increase in atmospheric CO₂ concentration resulting from different scenarios of CO₂ emissions from fossil fuel combustion have been discussed extensively (IPCC, 2001), all of them underestimate the likely increase in atmospheric CO₂ concentration by failing to account for demographics effects on the metabolic CO₂ release. The demographic effects on metabolic CO₂ release were examined by combining estimates of past, present and future human population with the per capita rates of direct and indirect metabolic CO₂ release derived here. Direct and indirect metabolic CO₂ release by humans has been increasing through time from a calculated value of 0.44 Gt C year⁻¹ in 1800 to an expected 4.4 Gt C year⁻¹ by year 2050 (Fig. 1), assuming that

BGD

3, 1781–1789, 2006

Metabolic human CO₂ release

Y. T. Prairie and
C. M. Duarte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

per capita rates are maintained constant. The realized increase may well exceed this estimate, since changes in diet and consumption habits are leading to a rapid increase in human body weight and food ingestion, and therefore per capita metabolic CO₂ release. In addition, the number of domestic animals per capita and per capita waste production are also increasing, and are expected to continue to do so in the future (OECD, 2002; Tilman et al., 2002).

Metabolic CO₂ release can only augment as population size increases in the future. This is a component of the CO₂ flux that must be recognized in future analyses of global CO₂ dynamics and that must be considered to represent a component of the perturbed C cycle, as human population has increased, and will continue to increase, greatly since pre-industrial time. Yet, the direct and indirect metabolic CO₂ emissions by humans is not considered explicitly in the scenarios conducted by the IPCC (2001), and is not incorporated, therefore, into current strategies to mitigate the climatic consequences of greenhouse gas emissions. The absence of explicit consideration of metabolic human CO₂ emissions may be a symptom of a tendency to perceive humans as separate entity to other species, which impacts on the biosphere involve their technology, whereas the results presented here clearly indicate that human demography and metabolism are important factors involved in green house emissions. Whereas metabolic CO₂ release maybe far less amenable to change than emissions derived from deforestation, cement production or fossil fuel use, different human choices can affect human metabolic CO₂ release. The indirect metabolic CO₂ release may be reduced through the promotion of behavioral changes to reduce the per capita consumption of meat and organic waste production, and the direct metabolic CO₂ release may be reduced by adjusting human ingestion to requirements, avoiding the excess food ingestion affecting much of the population in developed societies, and that represents a health hazard as well, responsible for more than 1 in 10 deaths in the EU and USA (Banegas et al., 2003). In contrast, however, per capita food intake is forecasted to increase by about 10% from a present average value of 2803 Kcal year⁻¹ person⁻¹ to 3050 Kcal year⁻¹ person⁻¹ by 2030, increasing to 3500 Kcal year⁻¹ person⁻¹ by

**Metabolic human
CO₂ release**Y. T. Prairie and
C. M. Duarte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2030 in industrialized countries (FAO, 2002). In summary, human metabolic CO₂ release, direct and indirect, is an important and growing component of anthropogenic CO₂ emissions. Explicit consideration of this component may help improve current emission scenarios and mitigation strategies.

5 *Acknowledgements.* This research is a contribution to the “Integrating the aquatic with the terrestrial component of the global carbon budget” Working Group supported by the National Center for Ecological Analysis and Synthesis, a Center supported by NSF (Grant #DEB-94-21535), the University of California at Santa Barbara, and the State of California. Y. T. Prairie and
10 C. M. Duarte were supported by the sabbatical fellowship program from the Spanish Ministry of Education and Science.

References

- Banegas, J. R., López-García, E., Gutiérrez-Fisac, J. L., Guallar-Castillón, P., and Rodríguez-Artalejo, F.: A simple estimate of mortality attributable to excess weight in the European Union, *Eur. J. Clin. Nutr.*, 57, 201–208, 2003.
- 15 Cohen, J. E.: *How Many People Can the Earth Support?*, New York, USA, W. W. Norton, 532 p., 1995.
- IPCC: *Climate change 2001: The Scientific basis*, New York, USA, Cambridge Univ. Press, 892 p., 2001.
- 20 Marrieb, E. N.: *Human Anatomy and Physiology*, 5th edition, New York, USA, Benjamin/Cummings, 1237 p., 2000.
- OECD: *Workshop on Waste Prevention: Toward Performance Indicators*, Paris, France, OECD, 2002.
- FAO: *World agriculture: towards 2015/2030*, Summary report, Food and Agriculture Organization of the United Nations, Rome, Italy, 2002.
- 25 Pauly, D. and Christensen, V.: Primary production required to sustain global fisheries, *Nature*, 374, 255–257, 1995.
- Peters, R. H.: *The ecological implications of body size*, New York, USA, Cambridge University Press, 1983.

Metabolic human CO₂ release

Y. T. Prairie and
C. M. Duarte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat: World Population Prospects: The 2002 Revision, <http://esa.un.org/unpp>, 2002.
- Tilman, D., Cassman, K. G. , Matson, O. A., Naylor, R., and Polasky, S.: Agricultural sustainability and intensive production practices, *Nature*, 418, 671–673, 2002.
- 5 Vitousek, P., Ehrlich, P. R., Ehrlich, A. H., and Matson, P.: Human appropriation of the products of photosynthesis, *Bioscience*, 36, 368–373, 1986.

BGD

3, 1781–1789, 2006

**Metabolic human
CO₂ release**

Y. T. Prairie and
C. M. Duarte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

**Metabolic human
CO₂ release**

Y. T. Prairie and
C. M. Duarte

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Estimates of direct and indirect metabolic CO₂ release by humans and associated animals. Figures correspond to 2003.

		Weight kg	Population (millions)	Respiratory Gt C yr ⁻¹	Excretory Gt C yr ⁻¹	Total Gt C yr ⁻¹
Direct	Humans	70	6100.00	0.57	0.28	0.28
Indirect	Cows	891	1349.00	0.94	0.31	1.26
	Goats	89	740.00	0.08	0.04	0.12
	Sheeps	50	1050.00	0.08	0.04	0.12
	Horses	794	58.00	0.04	0.01	0.05
	Pigs	200	1000.00	0.22	0.09	0.31
	Chickens	2.1	15 090.00	0.09	0.08	0.17
	Buffalos	350	165.90	0.06	0.02	0.08
	Turkeys	11.5	242.90	0.01	0.00	0.01
	Geese	5.2	238.80	0.00	0.00	0.01
	Ducks	2.6	917.70	0.01	0.01	0.01
	Camels	140	19.40	0.00	0.00	0.00
	Cats	2	1100.00	0.01	0.01	0.01
					Total	2.14
Non-metabolic	Waste production					0.10
					Total	3.10

**Metabolic human
CO₂ release**

Y. T. Prairie and
C. M. Duarte




Fig. 1. Estimated direct and indirect metabolic CO₂ release from pre-industrial to year 2050.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion