

## ***Interactive comment on “Modelling an alkenone-like proxy record in the NW African upwelling” by X. Giraud***

### **Anonymous Referee #1**

Received and published: 16 March 2006

The paper by Giraud is an attempt to use a regional ocean model for the Canary upwelling region which includes ecosystems dynamics to forward model the temperature conditions seen by the export of coccolithophores that produce alkenones. This production-weighted temperature is then compared to the alkenone index based on current understandings of their relationship, allowing the model results to be compared directly to the observations of alkenone records in deep sea sediments.

In principle this is a very good use of a regional modelling setup and shows the way forward for a number of frequently used proxies that are analysed in sediments. This study is constrained to assessing the absolute calibration of the alkenone index to mean annual SST, but future studies should be able to assess the sensitivity of the proxy to changes in conditions (which is after all the principle point of the proxy analysis in the first place).

Full Screen / Esc

Print Version

Interactive Discussion

Discussion Paper

## Specific comments:

The full potential of forward modelling is the ability of the forward model to differentiate between the spatial gradients of the climate field and proxies (the relationship between all points at a single time) and the temporal gradient (the relationship at a single point over all times). This approach has been successfully applied to ice core records for instance (Werner et al, 2000, GRL). This goal should be more strongly brought out in the introduction and conclusions.

I cannot comment on the details of the NPZD modelling nor on the regional ocean model set up, but I am satisfied by the sensitivity tests done with the ecosystem model. I am slightly more concerned about the potential sensitivity of the regional results to the climatological boundary conditions. Possibly something could be said about whether boundary conditions derived from specific ocean sections (rather than the highly smoothed Levitus climatology) would significantly alter the results?

In comparison to the data, it might be more useful to average the model output to the grid of the observations to have a cleaner comparison (i.e. it is not clear from fig 4 whether there are really systematic differences). I am a little surprised that no direct comparison to of the seasonal SST is being made (except in a selected fashion in fig 10). Since the main conclusion of the model is based on the differences of the IPT to the annual mean SST, significant biases in the SST (and mixed layer depths) may play a role in the differences between the modelled IPT and core-top alkenone-derived SST.

One major point is that I am not sure that the Levitus SST data are appropriate to be used in Table 3 and fig 10. In particular, I suspect that they are significantly smoothed and do not capture the full temperature gradient near the coast. I would suggest that satellite derived temperatures be used instead since they may have better resolution data. Shipboard analyses may also be helpful.

Long term climate changes is mentioned, but even over the 20th Century this region

[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)[Discussion Paper](#)

may have warmed by about a degree (<http://data.giss.nasa.gov/gistemp/maps/>). It is therefore possible that the core top alkenones may have been produced in slightly colder conditions (making the explanation of the results even more difficult). Although if the mean alkenone ages are in the thousand year range, it may be that mid-Holocene warmer samples are also mixed in (the discussion and Appendix A address this possibility though).

There are some puzzling features in the final results. Firstly, the actual alkenone-derived temperatures are biased high compared to the Levitus annual mean SSTs. This implies that their real IPT (assuming that the Prah et al culture experiments are valid) is warmer than the annual mean (implying a bias towards summer conditions perhaps). (This situation is of course made worse by any recent climate change influence). However, the model produced IPT is biased low compared to the model SST (due to the significant subsurface temperature weighting). I don't have any problem in understanding why the model behaves as it does, but the real world data require explanation. This could lie in the seasonality of the temperatures and in a mis-characterisation of the coccolithophore bloom. Could the Seawifs chlorophyll data help in validating the timing here?

In the conclusions it should be acknowledged that it is conceivable that the bias in IPT found here (due to subsurface production), would be calibrated away in a coretop calibration like Muller et al's since the correlation over large temperature ranges of the IPT and annual mean temperature should still be strong. It is therefore surprising that the Muller calibration is so similar to the Prah values - something that could be explained by an IPT that was warmer than annual mean SST due to seasonality. Thus while this paper is interesting and should be published, there is still more work to do in reconciling the alkenone and modelling results. This paper is a solid first step.

Minor points:

on page 95 it is stated that it is the Muller et al calibration that is used in Table 3, but in

[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)[Discussion Paper](#)

the Table 3 caption it states that Prah1 et al is used. I would suggest that Prah1 is the most suitable, although the differences are very small

p96. line 18: mis-interpret

p 97. line 20. The IPT are 'cooler' (not a cooling).

---

Interactive comment on Biogeosciences Discussions, 3, 71, 2006.

**BGD**

3, S38–S41, 2006

---

Interactive  
Comment

Full Screen / Esc

Print Version

Interactive Discussion

Discussion Paper