

*Biogeosciences Discussions* is the access reviewed discussion forum of *Biogeosciences*

**Input-output of N in  
subtropical forests  
under air pollution**

Y. Fang et al.

# Input and output of dissolved organic and inorganic nitrogen in subtropical forests of South China under high air pollution

Y. T. Fang<sup>1</sup>, P. Gundersen<sup>2</sup>, J. M. Mo<sup>1</sup>, and W. X. Zhu<sup>3</sup>

<sup>1</sup>South China Botanical Garden, the Chinese Academy of Sciences, Guangzhou, 510650, China

<sup>2</sup>Forest and Landscape Denmark, Faculty of Life Sciences, Univ. of Copenhagen, Hørsholm Kongevej 11, 2970 Hørsholm, Denmark

<sup>3</sup>Depart. of Biological Sciences, State Univ. of New York – Binghamton, Binghamton, NY 13902, USA

Received: 30 August 2007 – Accepted: 29 October 2007 – Published: 12 November 2007

Correspondence to: Y. T. Fang (fangyt@scbg.ac.cn)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

The nitrogen (N) emissions to the atmosphere and are thereby N deposition to forest ecosystems increasing rapidly in Southeast Asia, but little is known about the fate and effects elevated N deposition in warm and humid forests. Here we report the concentrations and fluxes of dissolved inorganic (DIN) and organic N (DON) in precipitation, throughfall, surface runoff and soil solution for three subtropical forests in a region of South China under high air pollution, to investigate how deposited N is processed and examined the importance of DON in N budget. The precipitation DIN input was 32–34 kg N ha<sup>-1</sup> yr<sup>-1</sup>. An additional input of 18 kg N ha<sup>-1</sup> yr<sup>-1</sup> as DON was measured in 2005, which to our knowledge is the highest DON flux ever measured in precipitation. Dry deposition was of minor importance at the site. A canopy uptake of DIN was indicated in two young conifer dominated forests (72–85% of DIN input reached the floor in throughfall), whereas no uptake occurred in an old-growth broadleaf forest. The DON fluxes in throughfall of all forest were similar to that of precipitation. In the young forests DIN was further retained in the soil, but 41–63% of precipitation DIN was still leached. Additionally, about half of the DON input was retained in these forests. The N retention in the two young aggrading forests (21–28 kg N ha<sup>-1</sup> yr<sup>-1</sup>) was in accordance with estimates of N accumulation in biomass and litter accretion. In the old-growth forest, no N retention occurred, but rather a net loss of 8–16 kg N ha<sup>-1</sup> yr<sup>-1</sup> from the soil was estimated. In total up to 60 kg N ha<sup>-1</sup> yr<sup>-1</sup> was leached, indicating that this forest was completely N saturated and could not retain additional anthropogenic N inputs. We found that the majority of DIN deposition and DIN leaching simultaneously occurred in the rainy season (March to August) and monthly DIN concentrations and fluxes in leaching were positively related to those in throughfall in all three forests, implying that part of the N leaching was hydrological driven by the abundant precipitation in the monsoon climate at the site. Our results suggest that long-term high N deposition has caused elevated N leaching in all studied forest types although most pronounced in the old-growth forest where wood increment was negligible or even negative. N availability

**BGD**

4, 4135–4171, 2007

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

even exceeded the biotic N demand in the young aggrading forests, albeit intensive rain in the growing season is likely to enhance N leaching in these forests.

## 1 Introduction

Increases in the deposition of atmospheric nitrogen (N) influence N cycling in forest ecosystems and can result in several negative consequences including acidification and leaching of nitrate into groundwater (Aber et al., 1989). A large body of research to assess the risk and consequence of N saturation has been carried out in temperate regions, where industrial development occurred earliest (e.g. Gundersen et al., 2006). Forest ecosystems have been shown to vary in their responses to increased N deposition. The timing and magnitude of response are thought to depend largely on the nutrient status of the forest and how close it is to N saturation (Gundersen et al., 1998; Aber et al., 2003). At the large scale, different climate regime (temperature and precipitation) is considered to affect forest N cycling rate and subsequently the response to anthropogenic N inputs (Hall and Matson, 2003; Lohse and Matson, 2005). At more local scale, differences in soil N pool size, species composition, stand age and land-use history may be major factors controlling the response pattern, because they influence the balance between N availability and demand (Fenn et al., 1998; Lovett et al., 2002; Kirstensen et al., 2004; Magill et al., 2004).

Atmospheric N deposition increases in densely populated areas of tropical and subtropical Asia due to the intensification of fossil fuel use and expansion of industrial and agricultural activities. Several authors have raised their concerns over the consequences of N enrichment of forest ecosystems in such warm and humid regions (Matson et al., 1999; Galloway et al., 2002; Chen and Mulder, 2007a, b). Elevated deposition of N in precipitation greater than  $25 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ , the threshold above which elevated N leaching always occurs in temperate forests (e.g. MacDonald et al., 2002), has already been reported for areas in southern China with rapid economic growth (Fan and Hong, 2001; Zhang, 2006; Chen and Mulder, 2007a; Luo et al., 2007), and it

**BGD**

4, 4135–4171, 2007

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

is expected to increase further in the coming decades (Zheng et al., 2002). However, little is known about how precipitation N interacts with forest canopies in warm and humid climates. Also the current N status and N process rates are unknown for forest ecosystems in these regions (Chen and Mulder, 2007a, b).

In temperate forests the response to elevated N deposition has been described in stages of decreasing biological control over the N cycle (Stoddard, 1994). Nitrate leaching first appears in the dormant season where biological demand is small, and it gradually appears also in the growing season as plant and microbial demand for N become saturated. The responses of subtropical forest ecosystems in China may differ from those in temperate zone because of different climate, species composition and soil properties (Chen and Mulder, 2007). Due to its position near Pacific Ocean in the east and the Indian Ocean in the south, south China has a monsoonal climate with a high abundance of heat, light, and water throughout the rainy season where a major fraction of the N deposition also occurs (Zhou and Yan, 2001). Elevated N deposition thus coincide with the most productive season and may be retained by biological processes, on the other hand high water fluxes occur during rain events which may favor leaching of deposition N. The balance between biological uptake, contact time and flow rate determine the fate of deposition N in the rainy season, whereas deposition N will most likely be retained throughout the dry season where plants are still productive and flows are minimal (or flow rates are low).

In the present study we have measured N input and N leaching at three subtropical forest types in the Dinghushan Biosphere Reserve (DHSBR) in southern China over two years to improve our understanding of N cycling in warm humid forest ecosystems. Short-term measurements of bulk precipitation in periods throughout the 1990's revealed atmospheric N deposition of 20–38 kg N ha<sup>-1</sup> yr<sup>-1</sup> in precipitation to this reserve (Huang et al., 1994; Zhou and Yan, 2001; Mo et al., 2002). We recently reported similar dissolved inorganic N (DIN) deposition in bulk precipitation in 2004 (Fang et al., 2007). Here we have continued these measurements but including also analysis of dissolved organic N (DON) input in bulk and wet-only precipitation.

**Input-output of N in subtropical forests under air pollution**

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

---

**Input-output of N in  
subtropical forests  
under air pollution**Y. Fang et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The forests types used, one mature monsoon evergreen broadleaf forest (old-growth) and two young forests (a pine forest and a pine-broadleaf mixed forest), are included in an ongoing N addition experiments (Fang et al., 2006). The old-growth forest is a regional climax type and has been protected for more than 400 years by monks in the nearby temples (Wang et al., 1982). The two young forests both originated from the 1930's clear-cut and subsequent pine plantation but developed differently later due to different patterns of human disturbance (Wang et al., 1982). We expect that these forest types would respond differently to the elevated N deposition. The old-growth forest is likely to be N-saturated but the two young ones may be still N-limited (Fang et al., 2006). However the canopy differences (old-growth broadleaf vs. two conifer canopies) may modify the response due to higher pollution interception in the conifer canopies than the broadleaf ones. Here we present two years of data from control plots in these forests to explore how ecosystem N retention has been affected by at least 15 years of high atmospheric N deposition. The importance of DON in the N budgets were examined in the second year, since this form of N has been commonly ignored in studies of warm humid ecosystems, although studies from other forest ecosystems have indicated its potential importance (Perakis and Hedin, 2002; Cornell et al., 2003).

## 2 Materials and methods

### 2.1 Site description

The study site is located in Dinghushan Biosphere Reserve (DHSBR) in the middle part of Guangdong province, South China ( $112^{\circ}10' E$  and  $23^{\circ}10' N$ ). This reserve is 20 km east of the relatively small city Zhaoqing (330 thousand inhabitants), about 90 km west of the metropolitan Guangzhou (10 millions inhabitants), and 180 km northwest of Hong Kong (7 millions inhabitants). The climate is warm and humid. The mean annual rainfall of 1927 mm has a distinct seasonal pattern, with 75% falling from March to August and only 6% from December to February (Huang and Fan, 1982). Mean annual relative

humidity is 80% and mean annual temperature is 21.0°C, with average temperatures in the coolest month (January) and the hottest month (July) of 12.6°C and 28.0°C, respectively (Huang and Fan, 1982).

A survey conducted in 2003 showed that in the old-growth evergreen broadleaf forest the major species were *Castanopsis chinensis*, *Machilus chinensis*, *Schima superba*, *Cryptocarya chinensis*, *Syzygium rehderianum* in the canopy and sub-canopy layers, which represented up to 80% of total basal area. Both young forests were originated from the 1930's clear-cut and subsequent pine plantation (Fang et al., 2006). The colonization from natural dispersal of regional broadleaf species has changed plant composition in the mixed forest (main species were *Pinus massoniana*, *Schima superba*, and *Castanopsis chinensis*), while the pine forest is dominated by *Pinus massoniana* under continuous human disturbances (generally the harvesting of understory and litter) (Mo et al., 2003). The old-growth forest had a basal area of almost twice ( $26.2 \text{ m}^2 \text{ ha}^{-1}$ ) those in the pine and mixed forests ( $14.0$  and  $13.8 \text{ m}^2 \text{ ha}^{-1}$ ), but less litter accumulation in forest floor ( $8.9$ ,  $23$  and  $20 \text{ Mg ha}^{-1}$  in the old-growth, pine and mixed forests, respectively; Fang et al., 2006).

The topography is highly heterogeneous, with slopes ranging from 15° to 35°. The soil is lateritic red earth formed from sandstone (He et al., 1982). The soil depths vary with forests. In the old-growth forest the soil depth ranges from 30 cm to 70 cm. The soil is about 40 cm deep in the mixed forest, and generally less than 40 cm in the pine forest. The old-growth forest had significant higher concentrations of total C, N and P, and extractable  $\text{NO}_3^-$ , but lower soil pH, C/N ratio, soil bulk density and extractable  $\text{NH}_4^+$  concentration than the pine and mixed forests (Table 1). Soil condition in the pine and mixed forests did not differ significantly (Table 1).

## 2.2 Sampling protocol

We sampled both bulk and wet-only precipitation on an open area in the reserve. Bulk precipitation was collected using two open glass funnels (15 cm in diameter), each connected to a 2.5 L sampling bottle with black polypropylene tubes. Wet-only precipitation

**BGD**

4, 4135–4171, 2007

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

was taken from a standard automatic wet-only collector (a 300 mm in diameter stainless steel container for wet deposition and a 150 mm in diameter glass container for dry precipitation, APS-3, Hunan Xianglan Ltd. China) located near the bulk collectors. To collect throughfall five collectors made of split longitudinally PVC pipes (intercept area 0.8 m<sup>2</sup> for each collector) were laid out randomly about 1.3 m above the ground in each forest. Each collector was connected to two 50 L sealed buckets (avoiding overflow) with black polypropylene tubes. The contribution from stemflow was negligible (<4% of throughfall N; Fang et al., 2007).

Since the plots are situated on steep slopes, one of the control plots in both the pine and old-growth forests had been delimited hydrologically, by plastic and concrete barriers to sample and quantify surface runoff. Surface runoff was not collected from the mixed forest due to its similarities of floor litter amount and slope degree with those in the pine forest. Soil solutions from 20 cm below the surface were sampled from each of the three control plots (10 m×20 m) in all three forest types. Two zero tension tray lysimeters (755 cm<sup>2</sup> per tray) was installed in each plot in the spring of 2003 to collect soil solutions. Each lysimeter was connected to a 5 L bottle using the steep slope of the sites to facilitate sampling. Previous study showed that more than 70% of the fine root (<5 mm) was distributed in the upper 20 cm soil in the mixed and old-growth forests (Wen et al., 1999). Thus, the solution N leaching estimated in this study was probably greater than that actually leached from the systems, since a small fraction of N was likely to be retained in the deeper soil. Soil solution at 40 cm depth obtained with ceramic suction cups were collected in 2004, but were terminated due to technical difficulties. The measurement revealed that annual volume-weight DIN concentrations at 40 cm were slightly lower than at 20 cm soil depth in the pine forest (2.5 vs. 3.0 mg N L<sup>-1</sup>), and the difference was somewhat more pronounced in the old-growth forest (5.2 vs. 6.8 mg N L<sup>-1</sup>; Fang et al., 2007).

Water samples were taken from January 2004 to December 2005. Precipitation samples were collected generally following a rainy day or series of rainy days, according to the weather forecast. We took soil solution samples every two weeks, with one sam-

**Input-output of N in  
subtropical forests  
under air pollution**

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

pling date around the middle and the other around the end of each month. Throughfall and surface runoff samples were taken in the first and third week of each month, and additional samples were taken again when we collected soil solution (if precipitation had occurred), which were pooled (volume weighted) with their respective regular samples before chemical analysis. For all the throughfalls and soil solutions, we recorded the water volume before sampling. All collectors were washed with distilled water immediately after each collection.

### 2.3 Sampling processing and analysis

Samples were filtrated within 24–48 h of collection through 0.45  $\mu\text{m}$  filters in the laboratory, and then stored in plastic bottles at 4°C until chemical analysis. Concentrations of  $\text{NH}_4^+\text{-N}$  and  $\text{NO}_3^-\text{-N}$  was determined for all samples. Total dissolved N (TDN) concentration was determined for samples collected in 2005. Concentration of  $\text{NH}_4^+\text{-N}$  was analyzed by the indophenol blue method followed by colorimetry, and  $\text{NO}_3^-\text{-N}$  was analyzed after cadmium reduction to  $\text{NO}_2^-\text{-N}$ , followed by sulfanilamide-NAD reaction (Liu et al., 1996). Total N was determined using persulphate oxidation to  $\text{NO}_3^-\text{-N}$  followed by colorimetric determination (Liu et al., 1996). Dissolved organic N (DON) concentration was calculated as the differences between TDN and DIN.

### 2.4 Calculations and statistics

The precipitation and air temperature used in this study was from the weather station in the reserve (Fig. 1). The recorded volumes of precipitation, throughfall, and solution were multiplied by their concentrations for the same period to determine N fluxes in  $\text{kg ha}^{-1}$ , which then was summed to get monthly and annual fluxes for each collector. We used estimated surface runoff volumes according to the observed relationship between precipitation and surface runoff to calculate its N fluxes (Fang et al., 2007). Monthly mean concentrations and fluxes were used to explore the relationships among precipitation, throughfall and solution for each forest type, using correlation analysis.

**BGD**

4, 4135–4171, 2007

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**



One-way ANOVA with Tukey's-b post hoc was used to identify the effects of forest type and year on annual fluxes of  $\text{NH}_4^+$ -N and  $\text{NO}_3^-$ -N in throughfall and soil solution. One-way ANOVA was also carried out to examine the effect of forest type on annual DON fluxes in 2005. All analyses were conducted using SPSS 10.0 for Windows. Statistical significant differences were set with P-values  $<0.05$  unless otherwise stated.

### 3 Results

#### 3.1 Precipitation

There were no obvious differences in the concentrations of either DIN or DON between bulk and wet-only precipitations (Fig. 2–4), suggesting a very little contribution of dry deposition. Thus, the means of the three collectors in the open area were used in the statistical analysis thereafter. Ammonium and  $\text{NO}_3^-$  concentrations showed a similar seasonal pattern during the course of our study (Fig. 2, 3), as confirmed by their highly significant linear correlation ( $r^2=0.68$ ,  $n=34$ ,  $P<0.001$ ). Their concentrations were highest in March/April, and then decreased considerably after the rainy season started, and reached the lowest levels in May or June (Fig. 2, 3). Annual volume-weighted concentration of  $\text{NH}_4^+$  was markedly higher in 2004 ( $1.8 \text{ mg NL}^{-1}$ ) than that in 2005 ( $1.1 \text{ mg NL}^{-1}$ ). For  $\text{NO}_3^-$  annual volume-weighted concentration was similar in the two years (both around  $0.8 \text{ mg NL}^{-1}$ ).

Despite 330 mm more precipitation (25%) in the second year, annual DIN input in precipitation was 5% less in the second than in the first year (Table 2). Ammonium was the dominant form of DIN input in precipitation with a contribution of 68% and 56% in the first and second year, respectively (Table 2). In both years, 82–83% of the precipitation fell in the rainy season (March to August, Fig. 1). Correspondingly, from 80% to 92% of DIN input in precipitation occurred in this period of the year (Fig. 5). Monthly DIN inputs were positively correlated to the mean DIN concentrations ( $r^2=0.56$ ,  $P=0.001$ ,  $n=16$ ) but were weakly correlated to the monthly precipitation amount.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Precipitation DON concentration (measured in 2005 only) showed a different seasonal fluctuation from DIN (Fig. 2–4). As a result, it did not correlate with either  $\text{NH}_4^+$  or  $\text{NO}_3^-$  ( $P > 0.05$ ), which indicated that DIN and DON might have different sources at our site. Dissolved organic N input amounted to  $17.8 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ , accounting for 36% of total dissolved N (TDN) input ( $47.6 \text{ kg N ha}^{-1}$  in 2005, Table 2). Annual volume-weighted concentration of DON was  $1.1 \text{ mg N L}^{-1}$ . Monthly DON inputs correlated positively with the concentrations ( $r^2 = 0.85$ ,  $P = 0.003$ ,  $n = 7$ ) but were independent of monthly precipitation amount.

### 3.2 Throughfall

In throughfall, the seasonal patterns of both  $\text{NH}_4^+$  and  $\text{NO}_3^-$  concentrations were similar to those in precipitation (Fig. 2, 3). The concentrations of  $\text{NO}_3^-$  in small samples collected in December 2004 were particularly high (Fig. 3), which might be due to dry deposition in a two months drought period before the sampling (Fig. 1), but could also be due to mineralisation/nitrification of litter in the canopy or the collector. Thus the concentration in precipitation at the same sampling date was used to calculate the N input in throughfall. This led to an underestimation of the total N input, but throughfall in that month accounted for only 16–24 mm, namely 0.9–2% of annual water fluxes in throughfall. Monthly mean DIN concentrations in throughfall showed significant correlations with that in precipitation for all three forests ( $P < 0.001$ ).

Annual volume weighed concentrations of  $\text{NH}_4^+$  in throughfall were  $1.1$  to  $1.5 \text{ mg N L}^{-1}$  in 2004 and  $0.9$ – $1.1 \text{ mg N L}^{-1}$  in 2005, which were lower than those observed in precipitation. The tree canopy of all forest types was thereby a sink for  $\text{NH}_4^+$ , removing  $4.7$ – $10.4 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  in 2004 and  $3.6$ – $6.6 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  in 2005, respectively (Table 2). Annual volume weighed concentrations of  $\text{NO}_3^-$  were always higher in throughfall ( $1.0$ – $1.7 \text{ mg N L}^{-1}$  in 2004 and  $0.9$ – $1.3 \text{ mg N L}^{-1}$  in 2005) than in precipitation, and annual throughfall  $\text{NO}_3^-$  input was close to that of precipitation input (in both the pine and mixed forest) or increased (in the old-growth forest) after the interaction

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

with tree canopies (Table 2). Consequently, a DIN uptake in the tree canopy on the order of 5 to 9 kg N ha<sup>-1</sup> yr<sup>-1</sup> must have occurred in the two young forests, which represented from 15% to 28% of their total DIN input in precipitation. But, in the old-growth forest, which had a broadleaf canopy, N fluxes remained unchanged (Table 2).

Throughfall DON concentrations exhibited a similar seasonality with that in precipitation (Fig. 4), indicating that DON in precipitation might be a main source of throughfall DON flux. Like in precipitation, there were no significant relationships between DON and NH<sub>4</sub><sup>+</sup> or between DON and NO<sub>3</sub><sup>-</sup> in throughfall in any forest. Forest throughfall DON inputs varied from 14.6 to 20.1 kg N ha<sup>-1</sup> yr<sup>-1</sup>, but all accounted for about 40% of their TDN inputs (Table 2). Annual volume weighed DON concentration in throughfall was 1.2–1.5 mg N L<sup>-1</sup>.

### 3.3 Surface runoff

Surface runoff was collected from the pine and mature forests, but not from the mixed forest (see Sect. 2.2). Concentrations of NH<sub>4</sub><sup>+</sup>-N were slightly lower than those of NO<sub>3</sub><sup>-</sup>-N and DON in both forests (data not shown). Losses via surface runoff were lower than those via seepage leaching and the difference between forests was minor relative to the difference of seepage leaching (Table 2). From 3.6 to 5.1 kg N ha<sup>-1</sup> yr<sup>-1</sup> as DIN and 1.1–2.1 kg N ha<sup>-1</sup> yr<sup>-1</sup> as DON (2005) were lost via surface runoff (Table 2).

### 3.4 Soil solution

Both NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> concentrations in soil solution at 20 cm depth had seasonal patterns comparable to those in throughfall (Figs. 2, 3). But the monthly mean DIN concentration relationship between throughfall and solution was forest-specific. In the pine and mixed forests, the seasonal changes were close to those in the throughfall concentrations; the correlations were highly significant ( $P < 0.001$  and  $P = 0.003$ , respectively). The old-growth forest had a larger seasonal variability. Furthermore, a marked increase in NO<sub>3</sub><sup>-</sup> concentration was observed in some months with high precipitation

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

amount, and most pronounced in the second year (for example, April, May, August and September of 2005, Fig. 3). Consequently, in this forest monthly mean concentrations in throughfall and solution was only marginally correlated ( $P=0.06$ ).

We also found that solution  $\text{NO}_3^-$  concentration was generally higher in 2005 than in 2004, particularly in the old-growth and pine forests (Fig. 3). Increased concentration in the second year might be caused by 330 mm more precipitation (Fig. 1), which might favor the soil nitrification. Active nitrification had been observed in the mineral soil in the pine and old-growth forests, but not in the mixed forest (author unpublished data). Annual volume weighted concentrations of  $\text{NO}_3^-$  ranged from 2.0 to 7.1  $\text{mg NL}^{-1}$ , whereas  $\text{NH}_4^+$  concentrations were 0.1 to 0.5  $\text{mg NL}^{-1}$ . The seasonal pattern of monthly DIN leaching followed those of throughfall input (Figs. 6, 7), as indicated by the significant relationships between leaching losses and throughfall inputs (Fig. 8). The leaching rate of DIN was higher in the old-growth forest than in the other two, as shown by a steeper slope of the regression line in Fig. 8a.

In the pine and mixed forests, from 10.0 to 16.6  $\text{kg N ha}^{-1} \text{yr}^{-1}$  were leached as DIN, which accounted for 37–62% of N deposition in throughfall. However, the N-rich old-growth forest lost significantly more N due to its higher solution  $\text{NO}_3^-$  level than the two young forests (Table 2); DIN and DON leached in this forest was 2.6–4.4 and 2.2–2.6 times higher than the two young forests, respectively (Table 2). Annual DIN leaching loss was measured to 37.8  $\text{kg N ha}^{-1} \text{yr}^{-1}$  in 2004, similar to the throughfall input (Table 2). A higher DIN loss (43.0  $\text{kg N ha}^{-1} \text{yr}^{-1}$ ) was observed in 2005, partly due to more precipitation amount, and it was 11  $\text{kg N ha}^{-1} \text{yr}^{-1}$  in excess of its throughfall input.

DON leached in 2005 was 8.4 to 16.9  $\text{kg N ha}^{-1} \text{yr}^{-1}$ , and accounted for 28–38% of the TDN leached and 42–84% of its input in throughfall, respectively (Table 2). Annual volume weighed DON concentration was 1.7 to 3.2  $\text{mg NL}^{-1}$ , which also was higher than those in throughfall. Seasonal variation of DON concentration generally followed those for DIN, and a significant but weak correlation of DIN and DON concentration in solution was found across all samplings in all three forests ( $r^2=0.094$ ,  $P=0.048$ ,

**Input-output of N in  
subtropical forests  
under air pollution**

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$n=126$ ), but this relationship was not significant when analyzed for each individual forest. Monthly DON leaching was positively correlated with throughfall input in the mixed forest (Fig. 8b)

### 3.5 Total N leaching and N retention

5 Total N leaching losses (surface runoff and seepage leaching) from the upper 20 cm soil was 14–20 kg N ha<sup>-1</sup> yr<sup>-1</sup> as DIN in the two young forests, and was 42–48 kg N ha<sup>-1</sup> yr<sup>-1</sup> in the old-growth forest (Table 3). Total DON leaching losses varied from 8 to 19 kg N ha<sup>-1</sup> yr<sup>-1</sup>, with the old-growth being doubled than the other two (Table 3). In the two young forests, 22 and 28 kg N ha<sup>-1</sup> yr<sup>-1</sup> or 41% and 55% of precipitation TDN input in 2005 was retained in the upper 20 cm soil, while no retention, but a net loss was found in the old-growth forest (Table 3). These retention estimates based on input-output budgets also accounts for the potential gaseous loss of N by denitrification. Gaseous losses as N<sub>2</sub>O were measured at nearby plot and estimated to 3.2±1.2 kg N ha<sup>-1</sup> yr<sup>-1</sup> for the three forest types (Tang et al. 2006).

## 15 4 Discussion

### 4.1 Wet N deposition

We did not observe marked differences in N concentrations between wet-only and bulk precipitation, indicating that dry deposition (and potential contamination from bird and insects) was negligible on the bulk collectors. No difference in concentrations of NH<sub>4</sub><sup>+</sup>-N and NO<sub>3</sub><sup>-</sup>-N between wet-only and bulk precipitation was also observed at another site in Guangzhou area (Aas et al., 2007). At the high humidity and frequent rainfalls in the wet season of this region wet deposition is likely to dominate the total atmospheric deposition.

Precipitation DIN input was estimated at 31.6 and 34.2 kg N ha<sup>-1</sup> yr<sup>-1</sup> in the two years

---

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

studied (Table 2), which are comparable to the highest deposition levels observed in Europe (MacDonald et al., 2002; Kristensen et al., 2004), but higher than those observed in most forests in North America (Fenn et al., 1998; Campbell et al., 2004) and Japan (Ohte et al., 2001). Precipitation DIN deposition at our site is higher in those  
5 in some other parts of southern China as well (Gan et al., 1995; Chen et al., 1997; Sha et al., 2002; Chen and Mulder, 2007a), except in Zhangzhou of Fujiang Province and Shanghai where averaged 37 and 58 kg N ha<sup>-1</sup> yr<sup>-1</sup> in precipitation were reported, respectively (Xiao et al., 2005; Zhang, 2006).

Consistent with our result, NH<sub>4</sub><sup>+</sup> is the main form of inorganic N in precipitation in most reports from China (Xiao et al., 2005; Zhang, 2006; Chen and Mulder, 2007a).  
10 The source of the high ammonium deposition is mainly thought to be the intensive agriculture covering much of the landscape between our site and the major cities. The warm climate may increase the emission rates above those observed in temperate regions, but emission inventories for the region are lacking. The oxidized N emissions  
15 from the industrialized areas in China are better documented. For instance, Richter et al. (2005) observed a highly significant increase of about 50% in the tropospheric column amount of NO<sub>2</sub> over south China using the satellite instruments.

In addition to the DIN input, a surprisingly high DON input of 17.8 N ha<sup>-1</sup> yr<sup>-1</sup> in precipitation was measured at our site. This is higher than any fluxes in precipitation reported in a global synthesis on organic N deposition (0.6–10.9 kg N ha<sup>-1</sup> yr<sup>-1</sup> with a median of 2.2 kg N ha<sup>-1</sup> yr<sup>-1</sup>; Neff et al., 2002). We therefore checked the quality of the measurements, but found no reason to doubt the results since both event samples and the wet-only sampler revealed the same high concentrations (mean 1.1 mg N L<sup>-1</sup>).  
20 Further, we identified a study from China with a similar high DON input (on average 15.8 kg N ha<sup>-1</sup> yr<sup>-1</sup>) for 11 sites in Zhangzhou city of Fujian province, with total mean DON concentration of 1.1 mg N L<sup>-1</sup> and mean total N input of 53 kg N ha<sup>-1</sup> yr<sup>-1</sup> (Xiao, 2005). A very recent study from Japan also measured 10.1 kg N ha<sup>-1</sup> yr<sup>-1</sup> of DON in precipitation in an intensive agricultural area (Ham and Tamiya, 2006). Our results  
25 combined with the observation in Zhangzhou suggest that high DON deposition does

---

**Input-output of N in  
subtropical forests  
under air pollution**Y. Fang et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

exist in the heavily air polluted regions in China, and so we need to consider this additional input in environmental studies. Future studies should identify the sources and investigate the formation of DON in precipitation.

Our data showed that DON concentration had a different seasonal fluctuation than DIN (Fig. 2–4), and it did not correlated with either  $\text{NH}_4^+$  or  $\text{NO}_3^-$  concentration, which indicate that DIN and DON might have different sources at our site. The synthesis by Neff et al. (2002) also revealed that there were no strong correlations between DON and either  $\text{NH}_4^+$  or  $\text{NO}_3^-$ . Conceptually, atmospheric organic N (AON), which we observe as DON in precipitation and throughfall, can be divided into three types of nitrogen: organic nitrate, reduced AON and biological/terrestrial AON (Neff et al., 2002). The biological/terrestrial AON refers to biological and particulate forms of organic N including bacteria, dust particles and pollen (Neff et al., 2002). While we find only negligible dry deposition of DON (Fig. 4) at our study site, these particulate forms are probably a minor source at least in dry form. Reduced AON compounds (e.g. from intensive agricultural activities) may form a significant contribution to DON in deposition at our site, as suggested for a site in Japan (Ham and Tamiya, 2006). Organic nitrates are oxidized end products of reactions of hydrocarbons with  $\text{NO}_x$  ( $\text{NO} + \text{NO}_2$ ) in polluted air masses (Neff et al., 2002) such as those occurring over southern China (Richter et al., 2005). Thus organic nitrates are also a likely contributor to the high DON deposition observed at our site.

## 4.2 Throughfall N

In temperate forests at high levels of N deposition, throughfall N is generally found to be higher than bulk precipitation N due to the filtering effect of the forest canopy on dry particles and gases, and conifers tend to have higher throughfall N input than broadleaves forests (Kristensen et al., 2004). However, this pattern was different from our results presented here, where a considerable DIN uptake (5 to  $9 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) were observed for the conifer-dominant canopies in the pine and mixed forests representing from 15% to 28% of their DIN input in precipitation (Table 2). Our results are,

### Input-output of N in subtropical forests under air pollution

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

however, in agreement with the observation in two fir forests in Fujian of China, where a decrease in DIN from precipitation to throughfall was reported after passing through the canopy (Fan and Hong, 2001). The old-growth forest with a broadleaf canopy on the contrary exhibited a slight increase or no change in N fluxes after the interaction with canopy (Table 2). A previous study also showed that the total N input in throughfall of this old-growth forest ( $39.2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) was slightly greater than that in precipitation ( $35.7 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ , Huang et al., 1994). Adding the DON fluxes to the canopy balance did not change the pattern of canopy uptake and release among the three forests, since the DON flux in throughfall was close to that of the precipitation and the minor differences followed those observed for DIN (Table 2).

The climatologic data from the weather station showed that total annual precipitation amount were 1327 and 1657 mm in 2004 and 2005, respectively (Table 2), which both were below the long-term average precipitation of 1997 mm. Furthermore, the rain in both years fell almost exclusively in the period from March to September (83% of the annual precipitation in 2004 and 100% in 2005, Fig. 1). This precipitation pattern in the study period may have led to an underestimation of the total DIN deposition to the canopies because there might have been some dry deposition in the period from October 2004 to February 2005, which was not included in our estimate. Unfortunately the very low amounts of rain made it difficult to obtain uncontaminated throughfall samples for the dry season in these particular years. Although we might slightly underestimate the total throughfall N deposition it was still higher ( $22\text{--}35 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) than most observations of throughfall N fluxes in Europe and North America (Kristensen et al., 2004; Campbell et al., 2004).

### 4.3 Impacts of N deposition on N leaching

In forest ecosystems, throughfall fluxes of N are often used as a first approximation of the total deposition (Gundersen et al., 2006). Several studies show that increased  $\text{NO}_3^-$ -N leaching start to occur in temperate and boreal forest when the throughfall N

---

**Input-output of N in  
subtropical forests  
under air pollution**

Y. Fang et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



input exceed  $10 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  and always occur above  $25 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  (Aber et al., 2003; Gundersen et al., 2006). With the N deposition well above these thresholds in our study forests, one might expect high N leaching. The old-growth forest leached up to  $37\text{--}42 \text{ kg NO}_3\text{-N ha}^{-1} \text{ yr}^{-1}$  below the surface 20 cm soil layer (Table 2), which indeed is substantially higher than those in almost all natural forests investigated in China (Gan et al., 1995; Chen et al., 1997; Sha et al., 2002). For example, leaching losses were found to be  $1.4 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  at the 25 cm depth from a mountain evergreen broadleaf forest in Ailaoshan of Yunnan (Gan et al., 1995),  $5.9 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  in a seasonal rain forest in Xishuanbanna of Yunan (Sha et al., 2002), and  $6.1 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  in a mountain rain forest in Jiangfengling of Hainan (Chen et al., 1997), respectively. These three forests received relatively low N deposition, ranging from 8.9 to  $14.2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ . The DIN losses in leaching in our study sites were also higher than most reported data in Europe and North America forests (Gundersen et al., 2006). In addition, we found that the N leaching loss from the old growth forest was higher than that observed 15 years ago ( $27.5 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ), where  $8 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  was still retained in the ecosystem (Huang et al., 1994).

Total N leaching loss mounted up  $67 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  in the old-growth forest, which exceeded the TDN inputs in throughfall ( $52 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) or precipitation ( $50 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) by 15 or  $17 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  (Table 2, 3). The results above indicate that the upper 20 cm soil is well saturated with N, and that “mining” of the pre-existing organic N may be occurring. Compared to the old-growth forest, the two young forests lost less TDN ( $22\text{--}29 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) and thereby retained more N from the precipitation (approximately  $21\text{--}28 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ , Table 3) in the plants or soils. These two stands have more N-poor soils (higher C/N ratios, Table 1) than the old-growth forest, which may suggest that the capacity to retain N depended largely on ecosystem N status as proposed by Gundersen et al. (1998).

The N retained in forest ecosystems mainly ends up in the plant biomass and the soil organic matter (SOM) (Nadelhoffer et al., 1999). In the aggrading pine forest where human disturbance had ceased recent estimate suggests that the canopy tree, the un-

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

derstory plants and standing floor litter accumulated 9.1, 6.0 and 6.5 kg N ha<sup>-1</sup> yr<sup>-1</sup>, respectively, during the period from 1990 to 2000 (Mo et al., 2004). Slightly higher amount of N may have been accumulated in the mixed forest, since it has higher litter production and higher foliar N concentration (Mo et al., 2007) than the pine forest.

5 In the pine forest, in total 22 kg N ha<sup>-1</sup> yr<sup>-1</sup> were sequestered in the three measured aboveground pools, which is sufficient to explain the observed 21 kg N ha<sup>-1</sup> yr<sup>-1</sup> that was retained above the upper 20 cm soil (Table 3). Also in the mixed forest N accumulation in the plant biomass and in the increasing litter layer appears to be sufficient to account for the retained 28 kg N ha<sup>-1</sup> yr<sup>-1</sup> (Table 3), especially if we also account

10 for the potential gaseous loss of N by denitrification that are include in the retention estimated (Sect. 2.5). The accumulation of plant biomass and plant derived litter is thus the dominant sink for N in these young forest, whereas the N sink in the mineral soil SOM is minor. In the old-growth forest, high N leaching losses may in contrast be a function of low increment that would diminish the ability of the vegetation to retain N.

15 Recent monitoring suggested that this forest might be experiencing a decline in tree productivity (Zhang et al., 2002; Guan et al., 2004). The biomass of woody plants (>1 cm in DBH) declined by 15.2% from 1994 to 1999 (Zhang et al., 2002). A decreasing trend of litter production was also found over the last two decades (Guan et al., 2004). From our present study we cannot conclude if the declining growth is the

20 reason for reduced N retention or if the high N load and the N saturation of this natural stand is the cause of the reduced growth. A mean NO<sub>3</sub><sup>-</sup>-N leaching for the two years of observation of 39 kg N ha<sup>-1</sup> yr<sup>-1</sup> imposed a strong acidification of the upper 20 cm soil equivalent to 2.8 kmol of H<sup>+</sup> ha<sup>-1</sup> yr<sup>-1</sup> (Gundersen et al., 2006) that may potentially affect plant growth and nutrition through base cation leaching.

25 The pattern of N retention (plant N demand being the major N sink for increased N deposition) demonstrated in the young forest seems to be different from that in the temperate and boreal zone, where the forests had a high retention efficiency for throughfall (or added) N (Aber et al., 1998; Gundersen et al., 1998; MacDonald et al., 2002). A major fraction of the input was incorporated into the large organic matter pool of the

---

**Input-output of N in  
subtropical forests  
under air pollution**Y. Fang et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

forest floor (Nadelhoffer et al., 1999). We suspect that the thinner or lack forest floor horizon in our forests (including the old growth) may contribute to the reduced retention capacity in soils compared to the temperate sites.

Further, reduced tree growth, as forests are aging or possibly as a consequence of N saturation as indicated in the old-growth forest of our study, may with time increase the fraction of N loss more generally in the region. Additional research is, however, needed to determine the partition of anthropogenic N in soil and plant in other air N polluted forests of southern China, with differing disturbance histories, floristic compositions, and soils, so that these suggested patterns can be confirmed.

#### 4.4 Seasonality of DIN deposition and leaching

Temperate forests are generally N-limited and cycle DIN tightly, and leaching loss is thought to be under biologically control. Therefore, the concentration and seasonality of leached  $\text{NO}_3^-$  in stream water has been taken as a primary indicator for the N excess in forested catchments in Europe and North America (Stoddard, 1994). The early stages of N excess are characterized by apparent seasonality and increases in the severity and frequency of  $\text{NO}_3^-$  episodes in the growing season. The later stages of N excess are marked by elevated concentrations of  $\text{NO}_3^-$  even at base flow (Stoddard, 1994). In this study, a considerable amount of DIN ( $14\text{--}48 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) was leached from all three forests in despite of the aggrading condition of the two young forest (Table 3), indicating that current N deposition seems sufficient for the biotic demand for N by plants and microbes. Moreover, abundant precipitation in the rainy season is likely to aggravate N leaching loss. We found a good correlation between throughfall and leaching for monthly DIN concentration as well as flux (Fig. 8a) indicating that part of leaching losses could be hydrological driven in these forest.

In the old-growth forest, a somewhat different seasonal variation in soil solution  $\text{NO}_3^-$  was observed with markedly increased  $\text{NO}_3^-$  concentrations in some months with high precipitation, whereas no such increase was found in the young forests (Fig. 3). This

---

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

increase in  $\text{NO}_3^-$  concentrations may be caused by increased net mineralization and nitrification stimulated by abundant precipitation in combination with high temperature (Fig. 1). Such climatic conditions may also have stimulated net mineralization rate in the two young forests, but a concurrent increase in plant N uptake most probably limited the loss from soil solution. This result indicates that episodic climatic events are likely to contribute more to DIN leaching losses in N-rich forest than in less N-rich forest.

#### 4.5 Importance of DON

The importance of DON in ecosystem nutrient budgets has been largely ignored probably due to poor analytical methods (Campbell et al., 2000). However, a number of recent studies in temperate climates have shown that DON can comprise about 50% or more of the TDN flux in throughfall, soil leachate, and streamwater in both low and moderate N deposition areas (Perakis and Hedin, 2002; McDowell et al., 2004). In this study, DON is also shown to be a very important component in the N cycle under high N deposition (Table 2). In 2005,  $17.8 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  was deposited to our study forests as DON, comprising 36% of TDN input in precipitation. Comparable DON fluxes ( $14.6$  to  $20.1 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) were found in throughfall. In soil solution, from  $6.5$  to  $16.9 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  leaved systems as DON, and made up 28–38% of TDN output. The absolute amount and percentage of DON in N fluxes in this study suggested that inclusion of DON is critical in assessing N dynamics.

We note that the concentrations and fluxes of DON in precipitation, throughfall, as well as solution leaching from the forests that we investigated are markedly higher than those reported in most other forests (Michalzik et al., 2001; Perakis and Hedin, 2002). For instance, in pristine temperate forest watersheds, DON is the dominant form of N loss, but usually DON concentrations are below  $0.6 \text{ mg N L}^{-1}$  and the loss in the order of  $1\text{--}3 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  (Gundersen et al., 2006). A recent study from the humid tropics in North Thailand estimated the DON leaching loss in tree forest types to be only  $0.5\text{--}0.8 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  (Möller et al., 2005), which was comparable to the input of DON with precipitation ( $1.1 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) in that area. High fluxes of DON were found in only a

### Input-output of N in subtropical forests under air pollution

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

few sites in the compilation of Michalzik et al. (2001) for temperate forests. However, despite the high concentrations and fluxes in our study, we found the proportions of DON in TDN (28–40%) in precipitation, throughfall and solution were well within the range for the previous reports (Michalzik et al., 2001; Neff et al., 2002; Cornell et al., 2003; Ham et al., 2007).

Throughfall DON may be a source of soil solution DON at our sites. But a significant relationship between monthly DON leaching and throughfall DON input was only found in one of three forests (Fig. 8b). This suggests that soil solution DON was influenced either by soil sources of DON (decomposition of litter and soil organic matter, plant exudates, fine-root and mycorrhizal turnover, and the waste products of macro and microorganisms), or by DON sinks (e.g. adsorption in organic and mineral soil horizons), or both.

## 5 Conclusions

Our measurements of N deposition in precipitation and forest throughfall confirm that atmospheric N input (up to  $50 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) in this region of south China is among the highest in world and previous data suggests that it has persisted for more than a decade. The contribution of DON ( $18 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) is the highest ever reported in precipitation and this flux needs more research attention to detail sources, pathways as well as potential effects. Measurements in throughfall indicate a minor importance of dry deposition to the forest canopies at the site.

Considerable amount of DIN ( $14\text{--}48 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ ) regardless of forest age, vegetation composition and land-use history was leached out of the upper 20 cm soil where a majority of roots grows, indicating that present atmospheric N loading may have already exceeded the plant and microbial N demand in the study region. A higher N retention in two young forests could be explained by the tree growth and litter accumulation, which on the other hand, suggests a limited retention capacity of the mineral soil. The N-excess was aggravated in the old-growth forest, where no retention but

**BGD**

4, 4135–4171, 2007

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

rather a net loss of  $17 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  (including DON) from the soil pool was observed.

The consequences of the elevated N deposition for forests in this region still need to be further investigated. Nevertheless, the nitrate leaching we observed have caused soil acidification that may impair forest nutrition. Forests of the region have been used heavily in the past and thus younger stands, like the pine and mixed forests in our study may still be able to retain a fraction of the atmospheric N. However, hydrologically driven N leaching will occur in the growing season regardless of the forest N status due to the abundant rainy season precipitation in the region.

*Acknowledgements.* This study was funded by the National Natural Science Foundation of China (No. 30670392), postdoctoral fellowship of South China Botanical Garden, CAS (200611), Key Project of Chinese Academy of Sciences Knowledge Innovation Program (KZCX2-YW-432-2, KSCX2-SW-133), and the Provincial Natural Science Foundation of Guangdong (No. 7006915). We thank L. Dejun, F. Hua, L. Xiankai, F. Xiaoming, L. Jurong and others for assistance in the field and laboratory. We especially thank M. Yoh, S. Brown and G. Y. Zhou for their advices throughout the study and thank L. O. Nilsson for his improvement of manuscript.

## References

- Aas, W., Shao, M., Jin, L., Larssen, T., Zhao, D. W., Xiang, R. J., Zhang, J. H., Xiao, J. S., and Duan, L.: Air concentrations and wet deposition of major inorganic ions at five non-urban sites in China, 2001–2003, *Atmos. Environ.*, 41, 1706–1716, 2007.
- Aber, J. D., Goodale, C. L., Ollinger, S. V., Smith, M. L., Magill, A. H., Martin, M. E., and Stoddard, J. L.: Is nitrogen altering the nitrogen status of northeastern forests?, *BioScience*, 53, 375–390, 2003.
- Aber, J. D., McDowell W., Nadelhoffer K., Magill A., Berntsen G., Kamakea, M., McNulty S., Currie W., Rustad, L., and Fernandez, I.: Nitrogen saturation in temperate forest ecosystems – hypothesis revisited, *BioScience*, 48, 921–934, 1998.
- Aber, J. D., Nadelhoffer, K. J., Steudler, P., and Melillo, J. M.: Nitrogen saturation in northern forest ecosystems – hypotheses and implications, *BioScience*, 39, 378–386, 1989.

**BGD**

4, 4135–4171, 2007

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

**Input-output of N in  
subtropical forests  
under air pollution**

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Campbell, J. L., Hornbeck, J. W., Mitchell, M. J., Adams, M. B., Castro, M. S., Driscoll, C. T., Kahl, J. S., Kochenderfer, J. N., Likens, G. E., Lynch, J. A., Murdoch, P. S., Nelson, S. J., and Shanley, J. B.: Input-output budgets of inorganic nitrogen for 24 forest watersheds in the northeastern United States: a review, *Water Air Soil Pollut.*, 151, 373–396, 2004.
- 5 Campbell, J. L., Hornbeck, J. W., McDowell, W. H., Buso, D. C., Shanley, J. B., and Likens, G. E.: Dissolved organic nitrogen budgets for upland, forested ecosystems in New England, *Biogeochemistry*, 49, 123–142, 2000.
- Chen, B. F., Zhou, G. Y., Zeng, Q. B., Li, Y. D., and Wu, Z. M.: Study on hydrochemical cycling in tropical mountain rain forest ecosystem, *Forest Res.*, 10, 111–117, 1997.
- 10 Chen, X. Y. and Mulder, J.: Atmospheric deposition of nitrogen at five subtropical forested sites in South China, *Sci. Total Environ.*, 378, 317–330, 2007a.
- Chen, X. Y. and Mulder, J.: Indicators for nitrogen status and leaching in subtropical forest ecosystems, South China, *Biogeochemistry*, 82, 165–180, 2007b.
- Cornell, S. E., Jickell, T. D., Cape, J. N., Rowland, A. P., and Duce, R. A.: Organic nitrogen de-  
position on land and coastal environments: a review of methods and data, *Atmos. Environ.*,  
15 37, 2173–2191, 2003.
- Fan, H. B. and Hong, W.: Estimation of dry deposition and canopy exchange in Chinese fir plantation, *Forest Ecol. Manag.*, 147, 99–107, 2001.
- Fang, Y. T., Yoh, M., Mo, J. M., Gundersen, P., and Zhou, G. Y.: Response of nitrogen leaching to simulated nitrogen deposition in a disturbed and a mature forest in southern China, *Pedosphere*, in press, 2007.
- 20 Fang, Y. T., Zhu, W. X., Mo, J. M., Zhou, G. Y., and Gundersen, P.: Dynamics of soil inorganic nitrogen and their responses to nitrogen additions in three subtropical forests in south China, *J. Environ. Sci. China*, 18, 752–759, 2006.
- 25 Fenn, M. E., Poth, M. A., Aber, J. D., Baron, J. S., Bormann, B. T., Johnson, D. W., Lemly, A. D., McNulty, S. G., Ryan, D. F., and Stottlemyer, R.: Nitrogen excess in North American ecosystems in: Predisposing factors, ecosystem responses and management strategies, *Ecol. Appl.*, 8, 706–733, 1998.
- Galloway, J., Cowling, E., and Kessler, E.: Reactive nitrogen, *Ambio*, 31, 59, 2002.
- 30 Gan, J. M., Xue, J. Y., and Zhao, H. K.: A preliminary study on changes of nutrient import and export in the process of rainfall in the Ailao Mountains region of Yunnan Province, *J. Natural Res.*, 10, 43–50, 1995.
- Guan, L. L., Zhou, G. Y., Zhang, D. Q., Liu, J. X., and Zhang, Q. M.: Twenty years of litterfall



- dynamics in subtropical evergreen broad-leaved forests at the Dinghushan forest ecosystem research station, *Acta Phytoecol. Sin.*, 28, 449–456, 2004.
- Gundersen, P., Emmett, B. A., Kjønnaas, O. J., Koopmans, C., and Tietema, A.: Impact of nitrogen deposition on nitrogen cycling in forest: a synthesis of NITREX data, *Forest Ecol. Manag.*, 101, 37–55, 1998.
- Gundersen, P., Schmidt, I. K., and Raulund-Rasmussen, K.: Leaching of nitrate from temperate forests – effects of air pollution and forest management, *Environ. Rev.*, 14, 1–57, 2006.
- Hall, S. J. and Matson, P. A.: Nutrient status of tropical rain forests influences soil N dynamics after N additions, *Ecol. Monogr.*, 73, 107–129, 2003.
- Ham, Y. S., Tamiya, S., and Choi I. S.: Contribution of dissolved organic nitrogen deposition to nitrogen saturation in a forested mountainous watershed in Tsukui, Central Japan, *Water Air Soil Pollut.*, 178, 113–120, 2006.
- Ham, Y. S. and Tamiya, S.: Contribution of dissolved organic nitrogen deposition to total dissolved nitrogen deposition under intensive agricultural activities, *Water Air Soil Pollut.*, 178, 5–13, 2006.
- He, J. H., Chen, Z. Q., and Liang, Y. A.: The soil of Ding Hu Shan, *Trop. Subtrop. For. Ecosys.*, 1, 25–38, 1982.
- Huang, Z. F. and Fan, Z. G.: The climate of Ding Hu Shan, *Trop. Subtrop. For. Ecosys.*, 1, 11–23, 1982.
- Huang, Z. L., Ding, M. M., Zhang, Z. P., and Yi, W. M.: The hydrological processes and nitrogen dynamics in a monsoon evergreen broad-leaved forest of Dinghu shan, *Acta Phytoecol. Sinic.*, 18, 194–199, 1994.
- Kristensen, H. L., Gundersen, P., Callesen, I., and Reinds, G.: Throughfall nitrogen deposition has different impacts on soil solution nitrate concentration in European coniferous and deciduous forests, *Ecosystems*, 7, 180–192, 2004.
- Liu, G. S., Jiang, N. H., and Zhang, L. D.: Soil physical and chemical analysis and description of soil profiles, Standards Press of China, Beijing, 1996.
- Lohse, A. L. and Matson, P.: Consequences of nitrogen additions for soil processes and solution losses from wet tropical forests, *Ecol. Appl.*, 15, 1629–1648, 2005.
- Lovett, G. M., Weather, K. C., and Arthur, M. A.: Control of nitrogen loss from forested watersheds by soil carbon: nitrogen ratio and tree species composition, *Ecosystems*, 5, 712–718, 2002.
- Luo, L. C., Qin, B. Q., Song, Y. Z., and Yang, L. Y.: Seasonal and regional variations in precipi-

**BGD**

4, 4135–4171, 2007

---

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**



- tation chemistry in the Lake Taihu Basin, China, *Atmos. Environ.*, 41, 2674–2679, 2007.
- MacDonald, J. A., Dise, N. B., Matzner, E., Armbruster, M., Gundersen, P., and Forsius, M.: Nitrogen input together with ecosystem nitrogen enrichment predict nitrate leaching from European forests, *Global Change Biol.*, 8, 1028–1033, 2002.
- 5 Magill, A. H., Aber, J. D., Currie, W. S., Nadelhoffer, K. J., Martin, M. E., McDowell, W. H., Melillo, J. M., and Steudler, P.: Ecosystem response to 15 years of chronic nitrogen additions at the Harvard Forest LTER, Massachusetts, USA, *Forest Ecol. Manag.*, 196, 7–28, 2004.
- Matson, P. A., McDowell, W. H., Townsend, A. R., and Vitousek, P. M.: The globalization of N deposition: ecosystem consequences in tropical environments, *Biogeochemistry*, 46, 67–83, 10 1999.
- McDowell, W. H., Magill, A. H., Aitkenhead-Peterson, J. A., Aber, J. D., Merriam, J. L., and Kaushal, S. S.: Effects of chronic nitrogen amendment on dissolved organic matter and inorganic nitrogen in soil solution, *Forest Ecol. Manag.*, 196, 29–41, 2004.
- Michalzik, B., Kalbitz, K., Park, J. H., Solinger, S., and Matzner, E.: Fluxes and concentrations 15 of dissolved organic carbon and nitrogen – a synthesis for temperate forests, *Biogeochemistry*, 52, 173–205, 2001.
- Mo, J. M., Brown, S., Peng, S. L., and Kong, G. H.: Nitrogen availability in disturbed, rehabilitated and mature forests of tropical China, *Forest Ecol. Manag.*, 175, 573–583, 2003.
- Mo, J. M., Brown, S., Xue, J. H., Fang, Y. T., Li, Z. A., Li, D. J., and Dong, S. F.: Response of nutrient dynamics of decomposing pine (*Pinus massoniana*) needles to simulated N deposition 20 in a disturbed and a rehabilitated forest in tropical China, *Ecol. Res.*, 22, 649–658, 2007.
- Mo, J. M., Fang, Y. T., Zhang, D. Q., Kong, G. H., and Fen, Z. N.: Effects of rainfall reallocation on nutrient dynamic of a pine forest in Dinghushan, Guihaia, 22, 529–533, 2002.
- Mo, J. M., Peng, S. L., Brown, S., Kong, G. H., and Fang, Y. T.: Nutrient dynamics in response to 25 harvesting practices in a pine forest of subtropical China, *Acta. Phytocool. Sin.*, 28, 810–822, 2004.
- Möller, A., Kaiser, K., and Guggenberger, G.: Dissolved organic carbon and nitrogen in precipitation, throughfall, soil solution, and stream water of the tropical highlands in northern Thailand, *J. Plant. Nutr. Soil. Sci.*, 168, 649–659, 2005.
- 30 Nadelhoffer, K. J., Emmett, B. A., Gundersen, P., Kjønaas, O. J., Koopmans, C. J., Schleppi, P., Tietema, A., and Wright R. F.: Nitrogen deposition makes a minor contribution to carbon sequestration in temperate forests, *Nature*, 398, 145–148, 1999.
- Neff, J. C., Holland, E. A., Dentener, F. J., McDowell, W. H., and Russell, K. M.: The origin,

**BGD**

4, 4135–4171, 2007

---

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

composition and rates of organic nitrogen deposition: A missing piece of the nitrogen cycle?, *Biogeochemistry*, 57/58, 99–136, 2002.

Ohte, N., Mitchell, M. J., Shibata, H., Tokuchi, N., Toda, H., and Iwatsubo, G.: Comparative evaluation on nitrogen saturation of forest catchements in Japan and Northeastern United States, *Water Air Soil Pollut.*, 130, 649–654, 2001.

Perakis, S. S. and Hedin, L. O.: Nitrogen loss from unpolluted South American forests mainly via dissolved organic compounds, *Nature*, 415, 416–419, 2002.

Richter, A., Burrows, J. P., Nub, H., Granier, C., and Niemeier, U.: Increase in tropospheric nitrogen dioxide over China observed from space, *Nature*, 437, 129–132, 2005.

Sha, L. Q., Zheng, Z., Feng, Z. L., Liu, Y. H., Liu, W. J., Meng, Y., and Li, M. R.: Biogeochemical cycling of nitrogen at a tropical seasonal rain forest in Xishuanbanna, SW China, *Acta. Phytoecol. Sin.*, 26, 689–694, 2002.

Stoddard, J. L.: Long-term changes in watershed retention of nitrogen-its causes and aquatic consequences, *Environ. Chem. Lakes Reservoirs*, 237, 223–284, 1994.

Wang, Z., He, D., Song, S., Chen, S., Chen, D., and Tu, M.: The vegetation of Dinghushan Biosphere Reserve, *Trop. Subtrop. For. Ecosys.*, 1, 77–141, 1982.

Wen, D. Z., We, I. P., Kong, G. H., and Ye W. H.: Production and turnover rate of fine roots in two lower subtropical forest sites at Dinghushan, *Acta. Phytoecol. Sin.*, 23, 361–369, 1999.

Xiao, J.: The formation and harm of N wet N deposition in Zhuangzhou city, *Energy Environ.*, 59–61, 2005.

Zhang, X. F.: Atmospheric nitrogen wet deposition and its effects on wetland water environment of Shanghai area, *Chinese J. Appl. Ecol.*, 17, 1099–1102, 2006.

Zhang, Y. M., Zhou, G. Y., Wen, D. Z., Zhang, D. Q., and Zhang Q. M.: Biomass dynamics of the *Castanopsis chinensis-Schima superba-Cryptocarya concinna* community of monsoon evergreen broad-leaved forest in Dinghushan Reserve, *Trop. Subtrop. For. Ecosys.*, 9, 10–17, 2002.

Zheng, X. H., Fu, C. B., Xu, X. K., Yan, X. D., Huang, Y., Han, S. H., Hu, F., and Chen, G. X.: The Asian nitrogen cycle case study, *Ambio*, 31, 79–87, 2002.

Zhou, G. Y. and Yan, J. H.: The influence of region atmospheric precipitation characteristics and its element inputs on the existence and development of Dinghushan forest ecosystems, *Acta. Ecol. Sin.*, 21, 2002–2012, 2001.

**BGD**

4, 4135–4171, 2007

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

**Table 1.** Characteristic of the mineral soil (0–10 cm) in the pine, mixed and old-growth forests at DHSBR in southern China.

| Parameter   | Pine forest     | Mixed forest    | Old-growth forest | P      |
|---|-----------------|-----------------|-------------------|--------|
| Bulk density ( $\text{g cm}^{-3}$ )                   | 1.16 (0.03) ab  | 1.22 (0.03) a   | 0.98 (0.06) b     | 0.026  |
| pH  | 4.04 (0.04) a   | 3.95 (0.01) a   | 3.83 (0.02) b     | 0.003  |
| Total C (%)   | 2.8 (0.3) b     | 2.6 (0.3) b     | 4.6 (0.23) a      | 0.004  |
| Total N (%)   | 0.11 (0.01) b   | 0.10 (0.01) b   | 0.19 (0.01) a     | 0.001  |
| C/N ratio   | 25 (1.1) ab     | 28 (0.7) a      | 22.1 (1.3) b      | 0.05   |
| Total P (%)   | 0.043 (0.003) b | 0.044 (0.004) b | 0.059 (0.003) a   | 0.024  |
| Extractable $\text{NH}_4^+$ ( $\text{mg N kg}^{-1}$ ) | 3.20 (0.35) a   | 3.03 (0.22) a   | 2.04 (0.04) a     | 0.027  |
| Extractable $\text{NO}_3^-$ ( $\text{mg N kg}^{-1}$ ) | 2.48 (0.43) b   | 2.80 (0.40) b   | 11.70 (1.00) a    | <0.001 |

SE in parentheses,  $n=3$ . Significant differences between forests were indicated by different letters. Data from Fang et al. (2006).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Input-output of N in subtropical forests under air pollution

Y. Fang et al.

**Table 2.** Annual fluxes of water (mm) and nitrogen ( $\text{kg N ha}^{-1} \text{ yr}^{-1}$ ) in precipitation, throughfall, surface runoff and the soil solution below the surface 20 cm soil layer in the pine, mixed and old-growth forests at DHSBR in southern China.

|                      | 2004     |                          |                          |             | 2005     |                          |                          |              |           |              |
|----------------------|----------|--------------------------|--------------------------|-------------|----------|--------------------------|--------------------------|--------------|-----------|--------------|
|                      | Water    | $\text{NH}_4^+\text{-N}$ | $\text{NO}_3^-\text{-N}$ | DIN         | Water    | $\text{NH}_4^+\text{-N}$ | $\text{NO}_3^-\text{-N}$ | DIN          | DON       | TDN          |
| Precipitation        | 1327     | 23.2                     | 10.9                     | 34.2        | 1657     | 17.8                     | 13.8                     | 31.6         | 17.8      | 49.5         |
| Throughfall (n=5)    |          |                          |                          |             |          |                          |                          |              |           |              |
| Pine forest          | 1120(27) | 12.8(0.6)                | 12.6(1.0) b              | 24.6(1.4) b | 1315(15) | 12.9(0.7)                | 13.9(0.6) b              | 26.8(1.3) b  | 18.2(2.1) | 44.9(3.2) ab |
| Mixed forest         | 1135(8)  | 15.2(1.0)                | 11.6(0.9) b              | 26.1(1.8) b | 1229(12) | 11.2(0.7)                | 11.5(0.4) c              | 22.8(1.0) b  | 14.6(2.2) | 37.3(2.9) b  |
| Old-growth forest    | 1126(22) | 16.5(1.5)                | 18.9(2.1) a              | 35.4(3.4) a | 1333(57) | 14.2(1.1)                | 17.4(0.9) a              | 31.6(1.5) a  | 20.1(0.8) | 51.7(2.1) a  |
| Surface runoff (n=1) |          |                          |                          |             |          |                          |                          |              |           |              |
| Pine forest          | 106      | 1.6                      | 2.7                      | 4.3         | 133      | 1.1                      | 2.4                      | 3.6          | 1.1       | 4.7          |
| Old-growth forest    | 226      | 1.6                      | 2.2                      | 3.8         | 283      | 1.3                      | 3.8                      | 5.1          | 2.1       | 7.2          |
| Solution (n=3)       |          |                          |                          |             |          |                          |                          |              |           |              |
| Pine forest          | 397(58)  | 1.6(0.5)                 | 9.0(3.0) b               | 10.6(3.5) b | 515(49)  | 1.2(0.2)                 | 15.5(6.3) b              | 16.6(6.3) b  | 8.4(3.4)  | 25.0(9.7) b  |
| Mixed forest         | 377(31)  | 1.5(0.2)                 | 8.5(1.4) b               | 10.0(1.6) b | 469(137) | 1.1(0.5)                 | 9.5(2.8) b               | 10.6 (3.3) b | 6.5(2.0)  | 17.1(5.2) b  |
| Old-growth forest    | 527(78)  | 0.6(0.1)                 | 37.2(7.7) a              | 37.8(7.7) a | 601(28)  | 1.2(0.1)                 | 41.7(7.1) a              | 43.0(7.1) a  | 16.9(3.9) | 59.8(10.7) a |

SE in parentheses. Different letters were used to indicate differences among treatments within the same forest when the effect was significant,  $P < 0.05$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Input-output of N in subtropical forests under air pollution**

Y. Fang et al.

**Table 3.** Total leaching loss and retention of nitrogen ( $\text{kg N ha}^{-1} \text{yr}^{-1}$ ) of precipitation inputs in the pine, mixed and old-growth forests at DHSBR in southern China.

|   | 2004 |     | 2005 |     |
|---|------|-----|------|-----|
|   | DIN  | DIN | DON  | TDN |
| Total N leaching losses (surface runoff plus leaching solution) |      |     |      |     |
| Pine forest   | 15   | 20  | 9    | 29  |
| Mixed forest  | 14   | 14  | 8    | 22  |
| Old-growth forest   | 42   | 48  | 19   | 67  |
| Ecosystem N retention to precipitation input                    |      |     |      |     |
| Pine forest   | 19   | 12  | 9    | 21  |
| Mixed forest  | 20   | 18  | 10   | 28  |
| Old-growth forest   | -8   | -16 | -1   | -17 |

Note: surface runoff N fluxes in the pine forest was used to estimate those in the mixed forest, due to their similar amount of floor litter and similar landscape. The runoff N fluxes were a minor component comparing to leaching in the younger forests, and a tiny fraction in the old-growth forest (see Table 2).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Input-output of N in subtropical forests under air pollution**

Y. Fang et al.



**Fig. 1.** Monthly precipitation and monthly mean air temperature at DHSBR in southern China.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Input-output of N in subtropical forests under air pollution

Y. Fang et al.



**Fig. 2.** Concentrations of  $\text{NH}_4^+$  in precipitation (a), throughfall (b) and soil solution (c) from three subtropical forests at DHSBR in southern China.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Input-output of N in subtropical forests under air pollution

Y. Fang et al.



**Fig. 3.** Concentrations of  $\text{NO}_3^-$  in precipitation (a), throughfall (b) and soil solution (c) from three subtropical forests at DHSBR in southern China.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 4.** Concentrations of DON in precipitation (a), throughfall (b) and soil solution (c) from three subtropical forests at DHSBR in southern China. DON concentration was measured in 2005 only.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Input-output of N in subtropical forests under air pollution

Y. Fang et al.




Fig. 5. Monthly fluxes of NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup> and DON in precipitation at DHSBR in southern China. DON flux was measured in 2005 only.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶


◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Fig. 6.** Monthly fluxes of NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup> and DON in throughfall in the pine, mixed and old-growth forests from DHSBR in southern China. DON flux was measured in 2005 only. \* Concentration of DIN in precipitation was used to calculated for throughfall N input in that month (see the text).

|                          |              |
|--------------------------|--------------|
| Title Page               |              |
| Abstract                 | Introduction |
| Conclusions              | References   |
| Tables                   | Figures      |
| ◀                        | ▶            |
| ◀                        | ▶            |
| Back                     | Close        |
| Full Screen / Esc        |              |
| Printer-friendly Version |              |
| Interactive Discussion   |              |

Input-output of N in subtropical forests under air pollution

Y. Fang et al.



**Fig. 7.** Monthly fluxes of NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup> and DON in soil solution from the pine, mixed and old-growth forests from DHSBR in southern China. DON flux was measured in 2005 only.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Input-output of N in subtropical forests under air pollution

Y. Fang et al.



**Fig. 8.** Relationships between monthly solution leaching and monthly throughfall input for DIN (a) and DON (b) in the pine, mixed and old-growth forests in DHSBR of southern China. (a) Pine forest,  $y=0.51x$ ,  $r^2=0.24$ ,  $P=0.062$ ,  $n=15$ ; Mixed forest,  $y=0.41x$ ,  $r^2=0.63$ ,  $P<0.001$ ,  $n=15$ ; Old-growth forest,  $y=1.16x$ ,  $r^2=0.33$ ,  $P=0.021$ ,  $n=15$ . (b) Mixed forest,  $y=0.44x$ ,  $r^2=0.54$ ,  $P=0.037$ ,  $n=7$ . In the pine and old-growth forest, the relationships were not significant. DON concentration was measured in 2005 only.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion