

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

Stable isotopes as ecological tracers: an efficient method for assessing the contribution of multiple sources to mixtures

M. N. Bugalho¹, P. Barcia⁵, M. C. Caldeira^{2,3}, and J. O. Cerdeira^{2,4}

¹Centro de Ecologia Aplicada “Baeta Neves”, Instituto Superior de Agronomia, Universidade Técnica de Lisboa, Lisboa, Portugal

²Centro de Estudos Florestais, Instituto Superior de Agronomia, Universidade Técnica de Lisboa, Lisboa, Portugal

³Departamento Florestal, Instituto Superior de Agronomia, Universidade Técnica de Lisboa, Lisboa, Portugal

⁴Departamento de Matemática, Instituto Superior de Agronomia, Universidade Técnica de Lisboa, Lisboa, Portugal

⁵Faculdade de Economia, Universidade Nova de Lisboa, Lisboa, Portugal

Received: 7 April 2008 – Accepted: 7 April 2008 – Published: 9 June 2008

Correspondence to: M. N. Bugalho (migbugalho@isa.utl.pt)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Stable isotopes and multiple sources

M. N. Bugalho et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Stable isotopes are increasingly being used as tracers of ecological processes potentially providing relevant information to environmental management issues. An application of the methodology consists in relating the stable isotopic composition of a sample mixture to that of sources. The number of stable isotopes, however, is usually lower than that of potential sources existing in an ecosystem, which creates mathematical difficulties in correctly tracing sources. We discuss a linear programming model which efficiently derives information on the contribution of sources to mixtures for any number of stable isotopes and any number of sources by addressing multiple sources simultaneously. The model identifies which sources are present in all, present in a subset of the samples or absent from all samples simultaneously and calculates minimum and maximum values of each source in the mixtures. We illustrate the model using a data set consisting on the isotopic signatures of different plant sources ingested by primary consumers in tropical riverine habitat in Asia. The model discussed may contribute to extend the scope of stable isotopes methodology to a range of new problems dealing with multiple sources and multiple tracers. For instance, in food web studies, if particular organic matter sources disappear or decrease in availability (e.g. climate change scenarios) the model allows simulation of alternative diets of the consumers providing potentially relevant information for managers and decision makers.

1 Introduction

Stable isotopes are naturally occurring chemical forms of the same element that differ in their molecular weights. Contrary to radioactive isotopes, stable isotopes do not undergo radioactive decay (West et al., 2006). Due to differential weights, stable isotopes are discriminated in physical, chemical, or biological reactions: the lighter and more abundant isotopes (e.g. ^{12}C) are more reactive than heavier ones (e.g. ^{13}C). As a consequence reaction products tend to be depleted in lighter isotopes and un-reacted

BGD

5, 2425–2444, 2008

Stable isotopes and multiple sources

M. N. Bugalho et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Stable isotopes and
multiple sources**M. N. Bugalho et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

products tend to be enriched in heavier isotopes (McKechnie, 2004). This property makes stable isotopes useful as natural integrators and tracers of ecological processes (Dawson et al., 2002) explaining their increasing use in ecological research and environmental management (West et al., 2006). Stable isotopes have provided insight into plant-environment and plant-plant relationships (Caldeira et al., 2001; Dawson et al., 2002), animal migrations (Kelly et al., 2005; Rubenstein and Hobson, 2004), nutritional ecology (Cherel et al., 2005; Inger et al., 2006), paleoecology (Passey et al., 2002; Zazzo et al., 2000) and ecosystem ecology (Jackson et al., 2002).

Stable isotope abundances are quantified through mass spectrometry and expressed by comparison with internationally accepted reference standards (West et al., 2006). The most commonly used stable isotopes in ecological research are from H, C, N, O and S elements. An important aspect of stable isotopes is their use as tracers, as they allow following the fate and transformation of a resource. This requires that the different potential sources have distinct isotopic values and that stable isotopes do not undergo significant fractionation (Dawson et al., 2002). In many fields of science, stable isotopes are used as tracers to determine the proportional contributions of several sources to a mixture (Phillips and Gregg, 2003). Examples range from plant ecology, where natural abundance of H and O isotopes have been used to determine the importance of different sources of water for plant use (Querejeta et al., 2007), to archaeological science, where stable isotopes of human remains and associated archaeofauna provided an estimate of human diets (Schoeninger and Moore, 1992; Newsome et al., 2004). Animal ecology is one of the fields that often uses tracing of stable isotopes to study animal migrations and diets. Stable isotope measurements in animal tissues reflect those isotope values in food webs, and may give information on the animal diet or location of feeding provided that isotope signatures vary among potential dietary components and locations of feeding (Cerling et al., 2006). Quite often, particularly in diet studies (e.g. Ben-David et al., 1997), the number of potential food sources is much higher than the number of stable isotopes which creates mathematical difficulties in identifying and quantifying the contribution of each source to a particular mixture or

sample of the animal tissue analysed, because infinitively many multiple solutions are possible.

Initial models for tracing stable isotopes were limited to consider one tracer at a time (e.g. van Dover et al., 1992). More recently, other models allowed for a more complex set of potential sources to be considered (Phillips and Gregg, 2003; Lubetkin and Simenstad, 2004).

IsoSource (Phillips and Gregg, 2003), the most commonly used algorithm, iteratively tests within a specified tolerance, all possible source contributions to a mixture. Specifically, a source increment s is user defined (e.g., entering a value of $s=1\%$) and every combination of multiples of s of each source adding up to 100% is tested as a feasible solution (within the given tolerance) to a particular mixture. This approach has been used successfully in a number of studies (Inger et al., 2006; Benstead et al., 2006) but becomes computationally intense with an increasing number of sources. For instance, an example of seven sources and two isotopes would require examining more than 1.7 billion partitions at 1% increment or over 32 million partitions at 2% increments (Lubtekin and Simenstad, 2004).

Lubtekin and Simenstad (2004) proposed a model specifically addressed to food webs which only enumerates the (basic) solutions corresponding to mixtures where the number n of sources does not exceed the number of tracers i plus 1. This gives a set S of $\binom{n}{i+1} = \frac{n(n-1)\dots(n-i)}{(i+1)i\dots 1}$ mixtures, from which some mathematically possible “nonsensical mixtures” (i.e. those not having all fractions of sources combinations comprised between 0 and 1) may have to be excluded. The solutions in S are those that define the limits of the region (of the n -dimensional Euclidean space) where the remaining feasible solutions are, which considerably reduces the size of the output relatively to that of IsoSource. For instance, for the same example of seven sources and two stable isotopes, the approach suggested by Lubetkin and Simenstad (2004) would take 35 corner points to delineate the outer boundaries of the solution range. This approach also considers the arithmetic mean of the solutions in S as a representative mixture.

Stable isotopes and multiple sources

M. N. Bugalho et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Stable isotopes and multiple sources

M. N. Bugalho et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Instead of an iterative model we use a simple linear programming algorithm which efficiently derives relevant information on source contributions to mixtures by addressing multiple samples simultaneously. The model provides the minimum and maximum contributions of each source among all the (infinitely many) possible mixtures and efficiently generates qualitative information on the simultaneous presence of sources in the mixtures. Addressing multiple samples simultaneously allow questions such as: Which sources are present in *all* mixtures? Which sources are absent from *all* mixtures? Or which sources are certainly present in particular mixtures?

We illustrated this approach with an example of diet composition, by applying the model to a data set used by Benstead et al. (2006), consisting of the isotopic signatures of different organic matter sources ingested by primary consumers in a tropical riverine habitat in Asia.

2 Material and methods

2.1 The data base

For computational experiments we used the data base provided in the Ecological Archives E087-018-A2 for the Ecology paper by Benstead et al. (2006). The database consists of: 1) the isotopic signatures of sources of organic matter (mangrove litter, freshwater swamp-forest litter, seagrasses, seagrass epiphytes, and marine particulate organic matter) from four estuaries in Kosrae, Federated States of Micronesia, that are consumed by primary consumers (fishes and crabs) in a tropical riverine habitat; and 2) the isotopic signatures of samples of tissue collected from primary consumers (the mixtures). Data were collected independently in 4 different study areas. Primary consumers include fish species from mangrove and reef flat areas as well as mangrove crabs. We used stable isotopes $\delta^{13}\text{C}$ and $\delta^{34}\text{S}$ to run our model, as the latter authors did. We did not consider the marine particulate organic matter source as its $\delta^{13}\text{C}$ and $\delta^{34}\text{S}$ isotopic signatures were taken from the literature and not directly measured and

also because this source was irrelevant to the diet of consumers analysed (Benstead et al., 2006).

The latter authors grouped individual organic matter sources into 5 different groups: mangrove leaf litter, freshwater swamp-forest leaf litter, seagrasses, seagrass epiphytes and marine particulate organic matter. We did not group organic matter sources as 1) the model can handle any number of sources, 2) to avoid any assumptions of a priori grouping of sources (this allows leaving this decision for posterior analysis after interpretation of the model output) and 3) because we wanted to explicitly address all individual samples simultaneously (in spite of magnitude of differences of their isotope signatures).

The model was run for each of the 4 study areas. We used information on the isotopic signatures of 19 primary consumers and 14 organic matter sources in study area Yela; 13 primary consumers and 12 organic matter sources in study area Okat; 9 primary consumers and 19 organic matter sources in study area Finkol; 14 primary consumers and 17 sources of organic matter in study area Yeseng.

2.2 The model

The model, a variant of the method of Barcia et al. (2007) for estimating diet composition of herbivores using natural markers in plant cuticles, uses linear programming to classify (see below) and determine the maximum and minimum proportions of each source in each sample. (An implementation of the model is available at http://www.isa.utl.pt/~orestes/LP_Tracer/.) More precisely, if S_i is the column vector of the isotope signatures of source i ($i=1, \dots, n$), M the column vector containing the isotope signatures of a sample and x_i the variable indicating the contribution of source i to the sample, the maximum and minimum values of each x_i subject to:

$$\sum_i S_i x_i = M, \quad (1)$$

Stable isotopes and multiple sources

M. N. Bugalho et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$\sum_i x_i = 1, \quad (2)$$

$$x_i \geq 0, \text{ for } i = 1, \dots, n \quad (3)$$

are determined. This can be easily achieved using any linear programming solver.

For each sample mixture (e.g. consumer), the $2 \times n$ maximum and minimum solutions obtained this way are highly meaningful for data interpretation (e.g. Demopoulos et al., 2007), among an infinite list of feasible solutions. Combined for all samples these solutions allow the classification of sources. If the maximum value of the proportion of a given source is zero for all the samples then we can conclude that that source is *absent* from every mixture, and it can be omitted from further consideration. If the minimum value is greater than zero we can conclude that the source is necessary to explain that sample. A source is *mandatory* if that source is necessary to explain all samples simultaneously. A source is *conditional mandatory* if that source is necessary to explain at least one, but not every sample. A (non absent) source is *optional* if the minimum value is zero for every sample. Unless the maximum value is zero, it is not possible to conclude that an optional source is indeed part of the mixture.

Beyond this qualitative classification, the solutions obtained allow deriving relevant information. Relatively high minimum values indicate significant mandatory or conditional mandatory sources, whilst low maximum values indicate relatively low importance sources. When minimum and maximum values coincide a reliable estimate of the source contribution is achieved.

3 Results

A graphical solution is presented for each estuary (Fig. 1). The polygon for each study area is the convex hull of the sources (i.e. the smallest convex set containing all the sources). Samples falling outside the polygon bounds give rise to “nonsensical mixtures” and were excluded from the analyses: *Epinephelus merra*, *Caranx melampygus*,

Stable isotopes and multiple sources

M. N. Bugalho et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Stable isotopes and
multiple sources**M. N. Bugalho et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aphareus furca, *Lutjanus monostigma* and *Kyphosus cinerascens* in study area Yela; *Liza vaigiensis*, *Caranx ignobilis*, *Lutjanus semicinctus* and *Moolgarda engeli* in study area Okat; *Moolgarda seheli*, *Cephalopholis leopardus*, *E. merra*, *Lethrinus harak* in study area Finkol; and *Kyphosus vaigiensis* and *Sargocentron punctatissimum* in study area Yeseng. The user, however, can use his/her ecological sensibility to decide if some of the sources should be included in the analysis by changing the tolerance value of the model (see the help menu of model implementation). Tables 1 to 4 summarize main aspects of the solutions produced by our model, identifying minimum and maximum source contributions to consumer diets in each estuary. No sources were classified as absent in any of the 4 study areas. (Indeed, when using 2 markers only, a source would only be absent if, and only if, all mixtures were aligned along a segment to which that source does not belong.) Five organic matter sources were classified as conditional mandatory (study areas Yela, Yeseng and Okat) and 1 as mandatory (study area Okat). All remaining sources were classified as optional.

In study area Yela, *Xylocarpus granatum*-R was included in the diet of *Siganus vermiculatus* with a value between 3% and 26% (Table 1). This justifies its classification as conditional mandatory (Table 1). The other sources were classified as optional, and we can not conclude if they were part of the diet of any consumer. However, low positive maximum values of an optional source imply it has low importance in the diet. For instance, none of the mangrove trees occurs in the diet of *L. vaigiensis* with more than 10% (Table 1). Similarly, in the diet of *S. vermiculatus*, the sources *Rhizophora apiculata*-I, *Nypa fruticans*-R and *N. fruticans*-I did not exceed 7%, 6% and 7%, respectively. Conversely, the seagrasses *Enhalus acoroides* or *Thalassia hemiprichii* entered, with high proportions, in the diet of several consumers. Indeed, for these consumers, when minimizing in the model the value of one of the seagrasses the other takes a high positive value. This can be perceived from the position of the seagrasses (S*) in Fig. 1. In other words, if the two seagrasses were considered a single one, i.e., a functional group, it would be conditional mandatory.

In study area Okat, *E. acoroides* was classified as mandatory source species, i.e.,

**Stable isotopes and
multiple sources**M. N. Bugalho et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

it was present in the diet of all consumers with a minimum contribution of, at least, 28% in the diet of *Scylla serrata*, and exceeding half of the diets of *E. merra* (92%), *L. equulus* (56%), *Lutjanus fulvus* (58%), *Parupeneus indicus* (57%), *U. vittatus* (63%) and *Siganus randalli* (55%) (Table 2). *Barringtonia racemosa*, a conditional mandatory source, completes the diet of *E. merra* with the remaining 8%. *E. acoroides* epiphytes (Ep.) also classified as conditional mandatory, was present in the diet of *M. seheli*, *L. equulus*, *L. harak* and *U. vittatus*. The diets of *L. equulus* and *U. vittatus* are fully explained by *E. acoroides* Ep. (44% and 37%) combined with *E. acoroides* (56% and 63%). Additionally, the diet of *M. seheli* consists almost entirely of *E. acoroides* and its epiphytes (52% minimum of *E. acoroides* Ep. and 40% minimum of *E. acoroides*). *E. acoroides* Ep. is also present in more than a quarter of the diet of *L. harak* (Table 2).

In study area Finkol, although there were only optional species, it is possible to conclude that the seagrasses *E. acoroides* or *T. hemprichii* entered, with relatively high proportions, in the diet of *Caranx* sp., *L. equulus* and *L. fulvus* (Table 3). Note that, similarly to study area Yela, if the two seagrasses were considered a single one it would be conditional mandatory as it can also be perceived from the position of the seagrasses (S*) in Fig. 1. From remaining sources *X. granatum*-R and *X. granatum*-F had relatively lower importance in the diet of all consumers, as compared to the other sources (Table 3).

Finally, in study area Yeseng, *E. acoroides* and *Lumnitzera littorea*-F were part of the diet of several consumers and were thus classified as conditional mandatory. *E. acoroides* was present in the diet of *Myripristis woodsi* (values between 26% and 62%), *L. harak* (7% and 72%), *Abudedefduf septemfasciatus* (77% and 98%) and *Acanthurus triostegus* (27% and 82%); and *L. littorea*-F occurred in the diet of *M. woodsi* (35% and 39%), *Caranx* sp. (25% and 56%), *L. monostigma* (12% and 26%), *Parupeneus bifasciatus* (9% and 19%) and *A. triostegus* (13% and 19%). *E. acoroides* and *L. littorea* explained, at least, 61% and 40% of the diet of *M. woodsi* and *A. triostegus*, respectively. Remaining source species were optional. Note that *T. hemprichii* and *T. hemprichii* epiphytes (Ep.) are the only optional sources that may be significantly

present in the diet of *M. woodsi* with a joint contribution that may be up to 39% (given that minimum contributions of the 2 mandatory sources in the diet of *M. woodsi* are 35% and 26%) (Table 4). Note also that *T. hemprichii* and *T. hemprichii* Ep. are the optional species with relatively high maximum values in almost all cases (Table 4). This may imply the relevance of these sources in the diet of several consumers. Also in this case, differences between *E. acoroides* and *T. hemprichii* in the isotopic signature of $\delta^{13}\text{C}$ were sufficient to allow discrimination between the two seagrasses and conclude that *E. acoroides* was present in the diet of some consumers (conditional mandatory).

4 Discussion

Our results confirmed the importance of seagrasses and their epiphytes as food sources in a riverine tropical fishery. Other researchers have also shown that food chains in tropical streams are largely dominated by algae (e.g. Brito et al., 2006; Benstead et al., 2006) and suggested the importance of epiphytes (e.g. Moncreiff and Sullivan, 2001). The model presented here, also showed the most relevant sources in the diet of consumers. For instance, we could conclude that *E. acoroides* was present in the diet of all (e.g. study area Okat) or several consumers (e.g. study area Yeseng) and that epiphytes of *E. acoroides* were part of the diet of several consumers (e.g. study area Okat). The majority of species were classified as optional. This result is not surprising and could be expected in unconstrained mixing models with a number of sources much higher than that of markers, and when several different samples (consumers) are considered simultaneously. However, because the model indicated minimum and maximum proportions of sources in the consumers, it also generated relevant information from optional sources that considerably improved the interpretation of results.

Contributions from mangrove-derived organic matter were variable and relatively low among estuaries and species. This is consistent with other author findings (e.g. Nagelkerken and van der Velde, 2004) and with the results obtained by Ben-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



stead et al. (2006) for the same estuaries. In addition, we also showed that there were two main mangrove litter source species (*X. granatum* and *L. littorea*) contributing to the consumer diets and we were able to identify those consumers for which mangrove litter sources were not dietary relevant.

5 Contribution of litter from freshwater swamp-forest was relatively low. Benstead et al. (2006) showed limited incorporation of *Terminalia carolinensis* into the food webs of the estuaries analysed, which is in agreement with our results. Adding to this, however, we have also shown that litter of *B. racemosa*, another fresh-water swamp tree, was certainly included (although in low quantities) in the diet of *E. merra* in study area Okat.

10 Our model has the computational ability to analyse any number of individual sources but the value of the output is likely to be higher if discrimination of stable isotope signatures among sources increases. For instance, in area Yeseng, stable isotope discrimination along the $\delta^{13}\text{C}$ was sufficient to allow discrimination between seagrasses leading us to conclude that *E. acoroides* was the dominant seagrass in the diet of different consumers. However, in study areas Yela and Finkol, it was not possible to assess which seagrass was mainly contributing to the consumer diets. In this case, *E. acoroides* and *T. hemprichii* had similar tracer signatures and function roles in the consumer diets and should be considered as a group.

20 Graphical interpretation may help decisions on grouping of sources with similar isotopic signatures, but for more than 3 markers methods such as the Nearest Neighbourhood Distance (Lubetkin and Simenstad, 2004) or other multivariate statistical criteria must be used. Grouping, however, must be considered and interpreted carefully. For instance, statistical meaningful groups may not correspond to ecologically functional groups (e.g. Dove and Mayes, 2000). Also, a priori grouping may lead to exclusion of samples from the convex polygon as it happened with *Megalops cyprinoides* in study areas Yeseng and Finkol (Benstead et al., 2006).

25 Some of the samples of consumers fell outside the minimum convex polygon defined by the sources. This is something that one must frequently deal with. Trophic fractionation of stable isotopes (e.g. Layman, 2007), non-sampling of important organic matter

Stable isotopes and multiple sourcesM. N. Bugalho et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

sources, or a priori grouping of sources may contribute to explain this and need to be considered when analysing the results.

The model presented here suggests a computationally efficient way of classify and generate minimum and maximum contributions of stable isotopes to multiple sources simultaneously. The model identified those sources that were important, unimportant or composed a substantial proportion of the consumer diets, but was also able to explain most of the diet of some consumers (e.g. *M. woodsi* in study area Yeseng). This contrasts with Benstead et al. (2006) contention that natural-abundance isotope surveys are powerful in disproving the importance of certain sources, but usually weak in showing the importance of a source of interest.

Contrary to other methods the model presented here addresses multiple samples simultaneously. In the present case we have dealt with multiple samples of different identities (i.e. primary consumers) but the approach is also suited for cases where several replicates of a population are present.

The importance of using stable isotopes as tracers in ecological research is increasing rapidly. Strong limitation of stable isotopes arises when dealing with numbers of sources higher than those of isotope tracers. In such cases, an approach such as the one shown in the present paper provides qualitative and quantitative information contributing to data interpretation. The model may contribute to data interpretation in other kind of problems. For instance, within the context of climate change scenarios, particular sets of organic matter sources may disappear or become less available within the ecosystem. In such cases, the model would allow simulation of alternative diets and assessment of the relatively importance of substitute sets of organic matter sources to consumers, a potentially relevant information for managers and decision makers. Additionally the model can handle any number of markers which may also contribute for extending the scope of stable isotopes to a range of new questions dealing with multiple sources and multiple tracers.

Acknowledgements. We thank John Benstead for permission to use Kosrae data. M. N. Bugalho was funded by the Portuguese Foundation for Science and Technology (FCT)

Stable isotopes and multiple sourcesM. N. Bugalho et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

grant SFRH/BPD/27216/2006 and M. C. Caldeira by SFRH/BPD 38710/2007 and the projects POCTI/AGG/48704/2002 and POCTI/AGR/63322/2004. J. O. Cerdeira was supported by program POCTI of FCT.

References

- 5 Barcia, P., Bugalho, M. N., Campagnolo, M. L., and Cerdeira, J. O.: Using N-alkanes to estimate the diet composition of herbivores, *Animal (Farming Systems and Environment)*, 1, 141–149, 2007.
- Ben-David, M., Flynn, R. W., and Schell, D. M.: Annual and seasonal changes in diets of martens: evidence from stable isotope analysis, *Oecologia*, 111, 280–291, 1997.
- 10 Benstead, J. P., March, J. G., Fey, B., Ewel, K. C., and Pringle, C. M.: Testing IsoSource: Stable isotope analysis of a tropical fishery with diverse organic matter sources, *Ecology*, 87, 327–333, 2006.
- Brito, E. F., Moulton, T. P., Souza, M. L., and Bunn, S. E.: Stable isotope analysis indicates microalgae as the predominant food source of fauna in a coastal forest stream, south-east Brazil, *Austral Ecol.*, 31, 623–633, 2006.
- 15 Caldeira, M. C., Ryel, R. J., Lawton, J. H., and Pereira, J. S.: Mechanisms of positive biodiversity-production relationships: insights provided by $\delta^{13}\text{C}$ analysis in experimental Mediterranean grassland plots, *Ecol. Lett.*, 4, 439–443, 2001.
- Cerling, T. E., Wittemyer, G., Rasmussen, H. B., Vollrath, F., Cerling, C. E., Robinson, T. J., and Douglas-Hamilton, I.: Stable isotopes in elephant hair document migration patterns and diet changes, *P. Natl. Acad. Sci. USA*, 103, 371–373, 2006.
- 20 Cherel, Y., Hobson, K. A., Bailleul, F. R., and Groscolas, R.: Nutrition, physiology, and stable isotopes: New information from fasting and molting penguins, *Ecology*, 86, 2881–2888, 2005.
- 25 Dawson, T. E., Mambelli, S., Plamboeck, A. H., Templer, P. H., and Tu, K. P.: Stable isotopes in plant ecology, *Ann. Rev. Ecol. Syst.*, 33, 507–559, 2002.
- Demopoulos, A. W. J., Fry, B., and Smith, C. R.: Food-web structure in exotic and native mangroves: a. Hawaii-Puerto Rico comparison, *Oecologia*, 153, 675–686, 2007.
- 30 Dove, H. and Mayes, R. W.: Measurement of dietary nutrient intake in free-ranging mammalian herbivores, *Nutr. Res. Rev.*, 13, 107–138, 2000.

BGD

5, 2425–2444, 2008

Stable isotopes and multiple sources

M. N. Bugalho et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Inger, R., Ruxton, G. D., Newton, J., Colhoun, K., Robinson, J. A., Jackson, A. L., and Bearhop, S.: Temporal and intrapopulation variation in prey choice of wintering geese determined by stable isotope analysis, *J. Anim. Ecol.*, 75, 1190–1200, 2006.
- Jackson, R. B., Banner, J. L., Jobbagy, E. G., Pockman, W. T., and Wall, D. H.: Ecosystem carbon loss with woody plant invasion of grasslands, *Nature*, 418, 593–594, 2002.
- Kelly, J. F., Ruegg, K. C., and Smith, T. B.: Combining isotopic and genetic markers to identify breeding origins of migrant birds, *Ecol. Appl.*, 15, 1487–1494, 2005.
- Layman, C. A.: What can stable isotopes ratios reveal about mangrove as fish habitats, *B. Mar. Sci.*, 80, 513–527, 2007.
- Lubetkin, S. C. and Simenstad, C. A.: Multi-source mixing models to quantify food web sources and pathways, *J. Appl. Ecol.*, 41, 996–1008, 2004.
- Mayes, R. W. and Dove, H.: Measurement of dietary nutrient intake in free-ranging mammalian herbivores, *Nutr. Res. Rev.*, 13, 107–138, 2000.
- McKechnie, A. E.: Stable isotopes: powerful new tools for animal ecologists, *S. Afr. J. Sci.*, 100, 131–134, 2004.
- Moncreiff, C. A. and Sullivan, M. J.: Trophic importance of epiphytic algae in subtropical sea-grass beds: evidence from multiple stable isotope analyses, *Mar. Ecol.-Progr. Ser.*, 215, 93–106, 2001.
- Nagelkerken, I. and van der Velde, G.: Relative importance of interlinked mangroves and sea-grass beds as feeding habitats for juvenile reef fish on a Caribbean island, *Mar. Ecol.-Progr. Ser.*, 274, 153–159, 2004.
- Newsome, S. D., Phillips, D. L., Culleton, B. J., Guilderson, T. P., and Koch, P. L.: Dietary reconstruction of an early to middle Holocene human population from the central California coast: insights from advanced stable isotope mixing models, *J. Arch. Sci.*, 31, 1101–1115, 2004.
- Passey, B. H., Cerling, T. E., Perkins, M. E., Voorhies, M. R., Harris, J. M., and Tucker, S. T.: Environmental Change in the Great Plains: An Isotopic Record from Fossil Horses, *J. Geol.*, 110, 123–140, 2002.
- Phillips, D. L. and Gregg, J. W.: Source partitioning using stable isotopes: coping with too many sources, *Oecologia*, 136, 261–269, 2003.
- Phillips, D. L., Newsome, S. D., and Gregg, J. W.: Combining sources in stable isotope models: alternative methods, *Oecologia*, 144, 520–527, 2005.
- Querejeta, J. I., Estrada-Medina, H., Allen, M. F., and Jiménez-Osornio, J. J.: Water source

Stable isotopes and multiple sources

M. N. Bugalho et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

partitioning among trees growing on shallow karst soils in a seasonally dry tropical climate, *Oecologia*, 152, 26–36, 2007.

Rubenstein, S. R. and Hobson, K. A.: From birds to butterflies: animal movement patterns and stable isotopes, *Trends Ecol. Evol.*, 19, 256–263, 2004.

5 Schoeninger, M. J. and Moore, K.: Bone stable isotope studies in archaeology, *J. World Prehist.*, 6, 247–296, 1992.

van Dover, C. L., Grassler, J. F., Fry, B., Garritt, R. H., and Starczak, V. R.: Stable isotope evidence for entry of sewage-derived organic material into a deep-sea food web, *Nature*, 360, 153–155, 1992.

10 West, J. B., Bowen, G. J., Cerling, T. E., and Ehleringer, J. R.: Stable isotopes as one of nature's ecological recorders, *Trends Ecol. Evol.*, 21, 408–414, 2006.

Zazzo, A., Bocherens, H., Brunet, M., Beauvilain, A., Billiou, D., Mackaye, H. T., Vignaud, P., and Mariotti, A.: Herbivore paleodiet and paleoenvironmental changes in Chad during the Pliocene using stable isotope ratios of tooth enamel carbonate, *Paleobiology*, 26, 294–309, 2000.

BGD

5, 2425–2444, 2008

Stable isotopes and multiple sources

M. N. Bugalho et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Source category and minimum and maximum proportions of organic matter source species in the diet of consumers at study area Yela as estimated by stable isotopes $\delta^{13}\text{C}$ and $\delta^{14}\text{S}$. Letters mean: O – Optional, C – Conditional mandatory, Ep – Epiphytes, R – Riverine, I – Interior, F – fringe.

Source Species	Category	Consumer Species													
		L. vaigiensis	S. prasin	C. ignobilis	L. equulus	L. fulvus	L. harak	M. flavolineatus	P. indicus	U. arge	Oxyeleotris sp.	S. vermiculatus	S. barracuda	S. qenie	S. serrata
Mangrove Trees															
<i>B. gymnorhiza</i> -R	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.08	0.37	0.41	0.29	0.33	0.22	0.33	0.39	0.28	0.66	0.14	0.34	0.24	0.60
<i>B. gymnorhiza</i> -I	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.09	0.40	0.44	0.31	0.36	0.24	0.35	0.41	0.30	0.70	0.14	0.36	0.24	0.64
<i>R. apiculata</i> -R	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.09	0.36	0.46	0.32	0.37	0.25	0.37	0.43	0.31	0.70	0.12	0.38	0.22	0.66
<i>R. apiculata</i> -I	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.10	0.21	0.49	0.34	0.27	0.22	0.39	0.41	0.33	0.41	0.07	0.33	0.13	0.52
<i>R. apiculata</i> -F	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.10	0.39	0.50	0.35	0.41	0.27	0.40	0.47	0.34	0.76	0.13	0.42	0.23	0.73
<i>X. granatum</i> -R	C	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.00
		0.10	0.46	0.33	0.21	0.41	0.27	0.40	0.42	0.34	0.57	0.26	0.42	0.31	0.45
<i>N. fruticans</i> -R	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.09	0.17	0.43	0.32	0.21	0.18	0.32	0.33	0.29	0.33	0.06	0.26	0.10	0.41
<i>N. fruticans</i> -I	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.09	0.21	0.47	0.33	0.26	0.22	0.38	0.41	0.32	0.40	0.07	0.33	0.12	0.51
<i>H. nunu</i>	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.08	0.38	0.42	0.27	0.34	0.23	0.34	0.40	0.29	0.67	0.19	0.35	0.26	0.58
Freshwater Swamp Trees															
<i>B. racemosa</i>	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.09	0.41	0.43	0.28	0.37	0.25	0.36	0.43	0.31	0.72	0.20	0.38	0.27	0.59
Seagrasses and epiphytes															
<i>E. acoroides</i>	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.81	0.63	0.55	0.38	0.65	0.76	0.64	0.59	0.69	0.34	0.78	0.64	0.75	0.38
<i>T. hemprichii</i>	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.88	0.64	0.57	0.40	0.68	0.79	0.67	0.61	0.72	0.35	0.78	0.66	0.76	0.39
<i>E. acoroides</i> Ep.	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.38	0.45	0.62	0.74	0.56	0.47	0.69	0.63	0.72	0.35	0.15	0.64	0.27	0.43
<i>T. hemprichii</i> Ep.	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.27	0.32	0.58	0.71	0.40	0.33	0.60	0.55	0.54	0.31	0.11	0.49	0.19	0.39

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Stable isotopes and multiple sources

M. N. Bugalho et al.

Table 2. Source category and minimum and maximum proportions of organic matter source species in the diet of consumers at study area Okat as estimated by stable isotopes $\delta^{13}\text{C}$ and $\delta^{14}\text{S}$. Letters mean: O – Optional, C – Conditional mandatory, M – Mandatory, Ep – Epiphytes, R – Riverine, I – Interior, F – fringe.

Source species	Category	Consumer species								
		<i>M. seheli</i>	<i>E. merra</i>	<i>C. ignobilis</i>	<i>L. fulvus</i>	<i>L. semicinctus</i>	<i>L. harak</i>	<i>P. indicus</i>	<i>S. randalli</i>	<i>S. serrata</i>
Mangrove trees										
<i>B. gymnorhiza</i> -R	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.03	0.00	0.00	0.26	0.00	0.14	0.19	0.24	0.28
<i>B. gymnorhiza</i> -I	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.04	0.00	0.00	0.27	0.00	0.17	0.22	0.28	0.32
<i>R. apiculata</i> -R	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.04	0.00	0.00	0.29	0.00	0.20	0.26	0.33	0.38
<i>R. apiculata</i> -I	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.05	0.00	0.00	0.22	0.00	0.22	0.27	0.26	0.42
<i>R. apiculata</i> -F	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.05	0.00	0.00	0.32	0.00	0.24	0.32	0.38	0.46
<i>S. alba</i> -R	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.04	0.00	0.00	0.31	0.00	0.17	0.23	0.29	0.34
<i>N. fruticans</i> -R	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.05	0.00	0.00	0.24	0.00	0.22	0.29	0.29	0.43
<i>N. fruticans</i> -I	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.06	0.00	0.00	0.22	0.00	0.26	0.27	0.26	0.48
<i>H. nunu</i>	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.02	0.00	0.00	0.20	0.00	0.11	0.15	0.18	0.21
Freshwater Swamp Trees										
<i>B. racemosa</i>	C	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.02	0.08	0.00	0.18	0.00	0.10	0.13	0.17	0.20
Seagrasses and epiphytes										
<i>E. acoroides</i>	M	0.40	0.92	0.76	0.58	0.92	0.34	0.57	0.55	0.28
		0.43	0.92	0.76	0.71	0.92	0.46	0.72	0.70	0.52
<i>E. acoroides</i> Ep.	C	0.52	0.00	0.24	0.00	0.08	0.28	0.00	0.00	0.00
		0.58	0.00	0.24	0.24	0.08	0.56	0.29	0.29	0.52

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Source category and minimum and maximum proportions of organic matter source species in the diet of consumers at study area Finkol as estimated by stable isotopes $\delta^{13}\text{C}$ and $\delta^{14}\text{S}$. Letters mean: O – Optional, Ep – Epiphytes, R – Riverine, I – Interior, F – fringe.

Source species	Category	Consumer species				
		<i>M. cyprinoides</i>	<i>Caranx</i> sp.	<i>L. equulus</i>	<i>L. fulvus</i>	<i>S. serrata</i>
Mangrove trees						
<i>B. gymnorhiza</i> -R	O	0.00	0.00	0.00	0.00	0.00
		0.41	0.32	0.31	0.42	0.29
<i>B. gymnorhiza</i> -I	O	0.00	0.00	0.00	0.00	0.00
		0.36	0.28	0.28	0.37	0.26
<i>B. gymnorhiza</i> -F	O	0.00	0.00	0.00	0.00	0.00
		0.53	0.40	0.41	0.50	0.38
<i>R. apiculata</i> -R	O	0.00	0.00	0.00	0.00	0.00
		0.57	0.32	0.43	0.40	0.42
<i>R. apiculata</i> -I	O	0.00	0.00	0.00	0.00	0.00
		0.57	0.33	0.44	0.42	0.41
<i>R. apiculata</i> -F	O	0.00	0.00	0.00	0.00	0.00
		0.74	0.34	0.45	0.43	0.54
<i>X. granatum</i> -R	O	0.00	0.00	0.00	0.00	0.00
		0.25	0.24	0.23	0.31	0.22
<i>X. granatum</i> -F	O	0.00	0.00	0.00	0.00	0.00
		0.25	0.25	0.25	0.33	0.23
<i>S. alba</i> -F	O	0.00	0.00	0.00	0.00	0.00
		0.43	0.34	0.33	0.45	0.31
<i>N. fruticans</i> -R	O	0.00	0.00	0.00	0.00	0.00
		0.72	0.28	0.38	0.36	0.58
<i>N. fruticans</i> -I	O	0.00	0.00	0.00	0.00	0.00
		0.49	0.38	0.37	0.50	0.35
Freshwater swamp forest trees						
<i>T. carolinensis</i>	O	0.00	0.00	0.00	0.00	0.00
		0.37	0.29	0.29	0.39	0.27
<i>H. nunu</i>	O	0.00	0.00	0.00	0.00	0.00
		0.35	0.29	0.29	0.39	0.27
<i>B. racemosa</i>	O	0.00	0.00	0.00	0.00	0.00
		0.34	0.28	0.27	0.37	0.26
<i>Mixed-species leaf litter</i>	O	0.00	0.00	0.00	0.00	0.00
		0.37	0.31	0.31	0.41	0.29
Seagrasses and epiphytes						
<i>E. acoroides</i>	O	0.00	0.00	0.00	0.00	0.00
		0.26	0.64	0.57	0.53	0.28
<i>T. hemprichii</i>	O	0.00	0.00	0.00	0.00	0.00
		0.26	0.65	0.58	0.54	0.29
<i>E. acoroides</i> Ep.	O	0.00	0.00	0.00	0.00	0.00
		0.68	0.30	0.40	0.37	0.74
<i>T. hemprichii</i> Ep.	O	0.00	0.00	0.00	0.00	0.00
		0.46	0.52	0.69	0.66	0.57

Stable isotopes and multiple sources

M. N. Bugalho et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4. Source category and minimum and maximum proportions of organic matter source species in the diet of consumers at study area Yeseng as estimated by stable isotopes $\delta^{13}\text{C}$ and $\delta^{14}\text{S}$. Letters mean: O – Optional, C – Conditional mandatory, Ep – Epiphytes, R – Riverine, I – Interior, F – fringe.

Source Species	Category	Consumer Species											
		M. cyprinoides	O. labiosus	M. woodsii	Caranx sp.	L. fulvus	L. monostigma	L. harak	P. bifasciatus	A. septemfasciatus	Oxyeleotris sp.	S. spinus	A. tristegus
Mangrove Trees													
<i>B. gymnorhiza</i> -R	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.44	0.05	0.01	0.09	0.13	0.04	0.02	0.03	0.01	0.62	0.09	0.02
<i>B. gymnorhiza</i> -I	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.32	0.04	0.01	0.06	0.09	0.03	0.02	0.02	0.00	0.46	0.07	0.01
<i>B. gymnorhiza</i> -F	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.43	0.06	0.01	0.11	0.16	0.05	0.03	0.03	0.01	0.79	0.11	0.02
<i>R. apiculata</i> -R	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.25	0.04	0.01	0.06	0.09	0.03	0.02	0.02	0.00	0.46	0.07	0.01
<i>R. apiculata</i> -I	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.26	0.03	0.01	0.06	0.08	0.02	0.02	0.02	0.00	0.41	0.06	0.01
<i>R. apiculata</i> -F	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.21	0.04	0.01	0.06	0.09	0.03	0.03	0.02	0.00	0.46	0.07	0.01
<i>L. littorea</i> -F	C	0.00	0.00	0.35	0.25	0.00	0.12	0.00	0.09	0.00	0.00	0.00	0.13
		0.71	0.12	0.39	0.56	0.40	0.26	0.05	0.19	0.03	0.45	0.16	0.19
<i>S. alba</i> -I	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.32	0.05	0.01	0.09	0.13	0.04	0.03	0.03	0.01	0.64	0.09	0.02
<i>N. fruticans</i> -R	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.30	0.04	0.01	0.07	0.10	0.03	0.02	0.02	0.01	0.50	0.07	0.01
<i>N. fruticans</i> -I	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.20	0.03	0.01	0.05	0.07	0.02	0.02	0.01	0.00	0.35	0.05	0.01
Freshwater Swamp trees													
<i>T. carolinensis</i>	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.48	0.06	0.01	0.11	0.15	0.04	0.03	0.03	0.01	0.75	0.10	0.02
<i>B. racemosa</i>	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.85	0.08	0.03	0.24	0.32	0.10	0.03	0.07	0.02	0.49	0.12	0.04
Mixed-species leaf litter	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.59	0.08	0.02	0.14	0.20	0.06	0.03	0.04	0.01	0.68	0.12	0.03
Seagrasses and epiphytes													
<i>E. acoroides</i>	C	0.00	0.00	0.26	0.00	0.00	0.00	0.07	0.00	0.77	0.00	0.00	0.27
		0.12	0.93	0.62	0.51	0.68	0.78	0.72	0.83	0.98	0.38	0.92	0.82
<i>T. hemprichii</i>	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.13	0.92	0.40	0.55	0.74	0.85	0.42	0.91	0.22	0.41	0.87	0.60
<i>E. acoroides</i> Ep.	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.11	0.20	0.04	0.37	0.52	0.15	0.32	0.10	0.03	0.36	0.38	0.07
<i>T. hemprichii</i> Ep.	O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
		0.11	0.38	0.08	0.44	0.60	0.27	0.60	0.19	0.05	0.36	0.70	0.12

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Stable isotopes and
multiple sources

M. N. Bugalho et al.




Fig. 1. Graphical solution (convex hull) produced by the model. Stable isotope ratios, $\delta^{34}\text{S}$ and $\delta^{13}\text{C}$, of organic matter sources (solid squares) and consumer samples (open circles) for the four estuaries of Kosrae (Yela, Okat, Finkol and Yeseng). M refers to mandatory and C to conditional mandatory organic matter sources. S* refers to seagrasses.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

