

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

Carbon and oxygen isotope analysis of leaf biomass reveals contrasting photosynthetic responses to elevated CO₂ near geologic vents in Yellowstone National Park

S. Sharma¹ and D. G. Williams^{1,*}

¹Department of Renewable Resources, University of Wyoming, Laramie, WY, USA

*also at: Department of Botany, University of Wyoming, Laramie, WY, USA

Received: 1 August 2008 – Accepted: 8 August 2008 – Published: 17 September 2008

Correspondence to: S. Sharma (shikha@uwyo.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

In this study we explore the use of natural CO₂ emissions in Yellowstone National Park (YNP) in Wyoming, USA to study responses of natural vegetation to elevated CO₂ levels. Radiocarbon (¹⁴C) analysis of leaf biomass from a conifer (*Pinus contortus*; lodgepole pine) and an invasive, non-native herb (*Linaria dalmatica*; Dalmation toadflax) was used to trace the inputs of vent CO₂ and quantify assimilation-weighted CO₂ concentrations experienced by individual plants near vents and in comparable locations with no geologic CO₂ exposure. The carbon and oxygen isotopic composition and nitrogen percent of leaf biomass from the same plants was used to investigate photosynthetic responses of these plants to naturally elevated atmospheric CO₂ concentrations. The coupled shifts in carbon and oxygen isotope values suggest that dalmation toadflax responded to elevated CO₂ exposure by increasing stomatal conductance with no change in photosynthetic capacity and lodgepole pine apparently responded by decreasing stomatal conductance and photosynthetic capacity. Lodgepole pine saplings exposed to elevated levels of CO₂ likewise had reduced leaf nitrogen concentrations compared to plants with no enhanced CO₂ exposure, further suggesting that this widespread and dominant conifer down-regulated photosynthetic capacity under elevated CO₂ levels near geologic vents.

1 Introduction

Effects of elevated CO₂ on plant photosynthesis and growth have been intensively studied because of the prominent role terrestrial vegetation plays in the global carbon cycle. Unresolved is whether and to what degree photosynthesis will be stimulated and how stomatal conductance and carboxylation and photochemical capacities are likely to react to globally increased CO₂ levels (Ainsworth and Rogers, 2007). Free-Air CO₂ Enrichment (FACE) is the preferred approach for studying effects of elevated CO₂ on natural ecosystems under realistic soil conditions and climate. However, cost con-

BGD

5, 3825–3843, 2008

Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



straints and technical difficulties have severely limited application of FACE technology across different ecosystems and climate. Further due to cost constraints most FACE experiments are short term providing only limited opportunities to study feedbacks on community structure and soil biogeochemistry. Geologic CO₂ emissions at the terrestrial surface offer an alternative low cost approach for studying elevated CO₂ effects on plant communities growing in natural environments (Bettaraini et al., 1995; Miglietta et al., 1998; Tognetti et al., 2000). Although natural CO₂ gradients established near geologic CO₂ vents can be confounded with other environmental factors, observations from such systems provide an opportunity to examine effects of long-term exposure to elevated CO₂ to compare against results from short-term Free-Air CO₂ Enrichment (FACE) experiments.

In this study we explore the use of natural CO₂ emissions in Yellowstone National Park (YNP) in Wyoming, USA to study responses of a native conifer (*Pinus contortus*; Lodgepole pine) and an invasive, non-native herb (*Linaria dalmatica*; Dalmation toadflax) to elevated CO₂ levels. There are numerous cold geologic features in YNP emitting large quantities of CO₂ at very high concentrations (>99%) with very minimal traces of H₂S and CH₄ and the native vegetation has been exposed to elevated levels of CO₂ at these sites for generations. The CO₂ emitted from geologic vents in YNP has no ¹⁴C, whereas background air has distinguishable and stable levels of ¹⁴CO₂ incorporated into plant biomass during photosynthesis. Hence, radiocarbon (¹⁴C) analysis of leaf biomass was used to trace the inputs of vent CO₂ and quantify assimilation-weighted CO₂ concentrations experienced by individual plants near vents and in comparable locations with no geologic CO₂ exposure.

Changes in plant photosynthesis associated with increased atmospheric CO₂ concentration can be detected in the ¹³C and ¹⁸O isotope composition of leaves (Farquhar et al., 1989; Ehleringer et al., 1993; Scheidegger et al., 2000; Barbour, 2007). The combined ¹³C and ¹⁸O isotopic composition of leaves can be used as long-term integrators of plant response to elevated CO₂ conditions (Grams et al., 2007). Increases in leaf carbon isotope discrimination can be a result of either

BGD

5, 3825–3843, 2008

Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1. a proportional increases in stomatal conductance relative to photosynthetic CO₂ demand by chloroplasts or
2. a proportional decrease in photosynthetic CO₂ demand by chloroplasts relative to stomatal conductance (Scheidegger et al., 2000).

5 Measurements of the carbon and oxygen isotopic composition of leaf organic matter can be used to investigate independent effects of photosynthetic capacity and stomatal conductance on the ratio of internal to ambient CO₂ composition (c_i/c_a) and hence $\Delta^{13}\text{C}$ (Scheidegger et al., 2000; Barbour, 2007; Grams et al., 2007). The combined
10 investigate photosynthetic responses of both plant species to naturally elevated atmospheric CO₂ concentrations.

2 Methods

2.1 Leaf sampling and determination of CO₂ exposure

15 Foliage samples were collected from lodgepole pine saplings at Ochre Springs and from adult plants of the invasive, non-native forb dalmation toadflax at Mammoth Upper Terraces (Fig. 1). Plants were sampled from near low-temperature CO₂ vents (“elevated” CO₂ sites) and from nearby sites where CO₂ exposure from vents was minimal (“control” CO₂ sites). Sampling locations were selected based on observed atmospheric CO₂ concentration values measured with a portable infra-red gas analyzer (IRGA; LI 6200, LiCor Inc., Lincoln NE, USA) and from site surveys (Tereck, 2004¹). Instantaneous CO₂ concentrations within plant canopies measured with the
20 IRGA were elevated (400–600 $\mu\text{L/L}$) above background levels near the CO₂ vents and

¹Tereck, M.: Plant communities growing under elevated CO₂ concentrations in Yellowstone National Park, Field survey, unpublished report, 2004.

BGD

5, 3825–3843, 2008

Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Effect of elevated
CO₂ on plants in YNP**S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were near background levels (378 $\mu\text{L/L}$) at comparable locations distant from vents. The control CO₂ site at Ochre Springs (44°37'1''N; -110°26'12''W) was located approximately 1600 m from the elevated CO₂ site (44°36'45''N; -110°24'52''W). At Mammoth Upper Terraces the control site (44°57'42''N; -110°42'51''W) and elevated CO₂ site (44°57'45''N; -110°42'54''W) were at distance of ≈ 200 m from each other. Soil temperature variations can potentially effect the photosynthesis and water relations in different plant communities (Schwarz et al., 1997; Dang and Cheng, 2004; Zhang and Dang, 2005). Therefore, soil temperatures were measured at all sampling sites and we noted that soil temperatures at 10 cm depth were similar (between 20 and 25°C at midday) at both control and elevated CO₂ sites.

Five individuals of lodgepole pine were haphazardly selected for isotope sampling at each of the control and elevated CO₂ sites in the Ochre Springs area. Isotopic composition of leaves might be effected by increased hydraulic resistance resulting from increased xylem path lengths in taller trees and longer branches (Warren and Adams, 2000) and the effective CO₂ exposure could also vary considerably with canopy height. To reduce this variability in this study needles were sampled (from the 2006 cohort) from branches of same length at the middle crown position from 0.5 to 1 m tall trees. The light environment of leaves can also potentially affect their carbon isotope signature (Lockheart et al., 1997; Grams et al., 2007). Therefore, we sampled needles only from tips of south facing branches of trees exposed to similar microhabitat conditions within the forest stand. Leaves of dalmation toadflax at Mammoth Upper Terraces were collected from 5 plants growing near a point CO₂ emission source and from 4 toadflax plants growing in a nearby site where atmospheric CO₂ levels were similar to background levels. At both the elevated and control site leaves were sampled from 4 to 5 plants growing at a distance of 3–4 m from each other along a 15m transect. Leaves (10–15) were collected from 4 different stems on each plant avoiding the small leaves subtending flowers.

Quantitative exposure to CO₂ from the geologic emissions integrated over the growth period of leaf development and maturation is difficult to assess solely from short-term

Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



IRGA measurements of air in the plant canopy. Therefore, we used the ¹⁴C content of leaves to estimate the assimilation-weighted CO₂ concentration exposure. The CO₂ at our sampling sites can only come from 2 sources i.e. atmosphere or the vent and therefore the sum of fractions of CO₂ fixed by plant coming from the vent and normal atmosphere should be equal to unity. Using mass balance approach, the measured ¹⁴C abundance in plant material (M_p) will be equal to the sum of fractions coming from vent and atmosphere weighted by the abundance of ¹⁴C in each of these gases (van Gardingen et al., 1995). CO₂ emitted from the vents originates from dissolution of carbonate sediments and therefore is highly depleted in ¹⁴C (so called “dead CO₂”) and ¹⁴C contribution from vent can be taken as 0 (Bettarini et al., 1995; van Gardingen et al., 1995). Simple mass-balance approach was then used to calculate the average CO₂ concentration surrounding individual lodgepole pine and dalmation toadflax plants:

$$[\text{CO}_2]_p = [\text{CO}_2]_b (M_b / M_p) \quad (1)$$

where $[\text{CO}_2]_b$ is the mean CO₂ concentration in the normal background atmosphere (taken to be 380 ppm). M_b and M_p are the ¹⁴C abundances (reported as fraction of modern standard) in leaves growing at ambient CO₂ concentrations and higher CO₂ concentrations, respectively.

Carbon isotope discrimination was determined from $\delta^{13}\text{C}$ of leaf biomass ($\delta^{13}\text{C}_l$) and $\delta^{13}\text{C}$ of air surrounding the plant ($\delta^{13}\text{C}_p$) from (Farquhar et al., 1989):

$$\Delta^{13}\text{C} = (\delta^{13}\text{C}_p - \delta^{13}\text{C}_l) / (1 + \delta^{13}\text{C}_l / 1000) \quad (2)$$

The $\delta^{13}\text{C}$ of CO₂ in air surrounding the plant ($\delta^{13}\text{C}_p$) will represent a mixture of $\delta^{13}\text{C}$ of CO₂ from the vent ($\delta^{13}\text{C}_v$) and background air ($\delta^{13}\text{C}_b$). CO₂ emitted from geologic vents at Mammoth Upper Terraces and Ochre Spring sites has $\delta^{13}\text{C}$ values of -3.9 and -3.2‰ , respectively (Werner and Brantley, 2003). The $\delta^{13}\text{C}$ of background atmospheric CO₂ was assumed to be -8‰ . The assimilation-weighted $\delta^{13}\text{C}$ value of CO₂

in the air surrounding individual plants was calculated using mass-balance:

$$\delta^{13}C_p = \delta^{13}C_b + f(\delta^{13}C_v - \delta^{13}C_b) \quad (3)$$

where f is the fraction of CO_2 contribution from the vent. The fraction of CO_2 contributed by vent is calculated using mass balance approach:

$$f = (\Delta^{14}C_l - \Delta^{14}C_b) / (\Delta^{14}C_v - \Delta^{14}C_b) \quad (4)$$

where $\Delta^{14}C$ is the relative deviation of ^{14}C activity of the sample from the oxalic Acid I standard in per mil values (after accounting for ^{14}C fractionation with a $\delta^{13}C$ correction). $\Delta^{14}C_l$ is the measured value of leaf samples, $\Delta^{14}C_b$ is the measured value for 2006 atmosphere at Pt. Barrow Alaska (50‰) and $\Delta^{14}C_v$ is the assumed value for CO_2 from vent (-1000‰).

The calculated carbon isotope discrimination was then used to estimate the influence of elevated CO_2 concentration on the ratio of leaf intercellular to ambient CO_2 concentration (c_i/c_a) using the simplified model of Farquhar et al. (1989) as:

$$c_i/c_a = (\Delta^{13}C - a) / (b - a) \quad (5)$$

where a is the carbon isotope discrimination associated with CO_2 diffusion through air (-4.4‰), and b is the net discrimination associated with CO_2 fixation in C_3 plants (-27‰).

2.2 Leaf isotopic and N content analyses

Whole leaf samples of dalmation toadflax and lodgepole pine needles from the 2006 growth year were dried and ground in a mill to a consistent particle size. Stable isotope analysis was carried out at the University of Wyoming Stable Isotope Facility (<http://www.uwyo.edu/sif/>). The $\delta^{13}C$ and N % values were determined by continuous flow EA-IRMS using a Costech 4010 element analyzer online with a Finnigan Delta plus XP isotope ratio mass spectrometer. The $\delta^{18}O$ composition of bulk leaf material was

BGD

5, 3825–3843, 2008

Effect of elevated CO_2 on plants in YNP

S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



measured by online pyrolysis to CO using a Finnigan TC/EA coupled to the Finnigan Delta Plus XP isotope ratio mass spectrometer. Measurement precision was assessed by replicate analysis of samples and internal lab standards and was better than $\pm 0.1\%$ for $\delta^{13}\text{C}$ and $\pm 0.25\%$ for $\delta^{18}\text{O}$. The $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values are reported in per mil (‰) relative to V-PDB and V-SMOW, respectively. Radiocarbon analysis was performed at the W. M. Keck AMS Laboratory at the University of California, Irvine, CA, USA. Radiocarbon concentrations are reported as fractions of the Modern Standard and $\Delta^{14}\text{C}$ is reported following conventions of Stuiver and Polach (1977).

3 Results and discussion

The assimilation-weighted CO_2 exposure of a leaf calculated from radiocarbon values of leaves Eq. (1) ranged from 379–381 (control site) to 401–607 ppm (elevated site) in dalmation toadflax from the Mammoth Upper Terraces site and from 378–383 (control site) to 419–482 ppm (elevated site) in lodgepole pine at the Ochre Springs site. Changes in leaf $\delta^{13}\text{C}$ values in response to variations in environmental conditions integrate changes in the relative demand of CO_2 by chloroplasts during photosynthesis and in the supply of CO_2 associated with stomatal conductance in plants (Farquhar et al., 1989; Ehleringer et al., 1993). We found that $\Delta^{13}\text{C}$ was significantly positively correlated with assimilation-weighted CO_2 exposure levels (Fig. 2) for both dalmation toadflax and lodgepole pine (dalmation toadflax, $r^2=0.77$, $P<0.002$; lodgepole pine, $r^2=0.85$, $P<0.001$) indicating that elevated CO_2 levels caused an increase in leaf c_i/c_a . The observed increase in $\Delta^{13}\text{C}$ values reflect a 13% and 17.5% increase in leaf c_i/c_a in the case of toadflax and lodgepole pine samples, respectively.

We observed a 2.4‰ decrease in $\delta^{18}\text{O}$ of leaf biomass with increasing CO_2 exposure in dalmation toadflax and a 2.5‰ increase in $\delta^{18}\text{O}$ of leaf biomass along the gradient of increasing CO_2 exposure in lodgepole pine (Fig. 3a and b). Because of close proximity, we assumed that the oxygen isotopic composition of soil water and

BGD

5, 3825–3843, 2008

Effect of elevated CO_2 on plants in YNP

S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



atmospheric vapor was similar for plants growing within control or elevated CO₂ sites, and the conceptual model proposed by Grams et al. (2007) can be used to understand components of the photosynthetic response of the two species. This model infers the changes in stomatal conductance, c_i/c_a and photosynthetic capacity of plants using a dual carbon and oxygen isotope approach. This model is an expansion of the conceptual model proposed by Scheidegger et al. (2000) which estimates changes in leaf-to-air vapor pressure deficit from the $\delta^{18}\text{O}$ of leaf biomass for prediction of changes that would have likely occurred in stomatal conductance. The model of Grams et al. (2007) infers stomatal conductance changes directly from $\delta^{18}\text{O}$ of leaf cellulose bypassing the previously used relative humidity/ $\delta^{18}\text{O}$ relationship in the model of Scheidegger et al. (2000). Grams et al. (2007) did not find a significant correlation between $\delta^{18}\text{O}$ of leaf biomass and leaf cellulose in their study and therefore used the $\delta^{18}\text{O}$ of leaf cellulose as the input parameter in their model. However, a reliable correlation between $\delta^{18}\text{O}$ of leaf biomass and cellulose has been observed in several herbaceous and woody plants (Barbour and Farquhar, 2000; Barbour et al., 2001). We feel justified in using the $\delta^{18}\text{O}$ of leaf biomass instead of $\delta^{18}\text{O}$ of leaf cellulose as an input parameter in the conceptual model of Grams et al. (2007).

In case of dalmation toadflax there is an increase in $\Delta^{13}\text{C}$ accompanied by a decrease in $\delta^{18}\text{O}$ of leaf biomass associated with increasing CO₂ exposure (Fig. 3a). The decrease in $\delta^{18}\text{O}$ of leaf biomass is likely a result of an increase in stomatal conductance. This is because higher stomatal conductance results in a higher rate of transpiration which cools the leaf surface and reduces the leaf intercellular vapor pressure resulting in lower $\delta^{18}\text{O}$ enrichment at evaporating sites. Further, due to increased transpiration, the convective flux of unenriched source water to evaporating sites increases, thereby reducing the enrichment of bulk leaf water (Farquhar and Lloyd, 1993). This $\delta^{18}\text{O}$ enrichment signal of leaf water is transferred to the bulk leaf organic matter by isotope exchange reactions. Using the dual isotope model of Grams et al. (2007) the increase in c_i/c_a can then be inferred to be a result of increased stomatal conductance accompanied by no response of photosynthetic capacity. Our results do not fit predic-

**Effect of elevated
CO₂ on plants in YNP**S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tions from FACE sites which generally suggest increasing exposure to CO₂ results in an average 22% decrease in stomatal conductance across all plant species (reviewed in Ainsworth and Rogers, 2007). However, preliminary studies in a CO₂ enrichment experiment (<http://www.phace.us>) near Cheyenne, Wyoming USA (Williams, unpublished data) also confirm that dalmation toadflax shows about a 6‰ ($n=2-3$) decline in leaf biomass $\delta^{18}\text{O}$ values under elevated CO₂ conditions.

Several studies have shown that soil water savings induced by decreased stomatal conductance under elevated CO₂ conditions is primarily observed when moisture is limiting (Field et al., 1997; Wullschlegler et al., 2002). However, when soil water availability is high several plant species show higher stomatal conductance and are insensitive to elevated CO₂ conditions (Mooney et al., 1983; Knapp et al., 1996; Schomp, 2007). We hypothesize that since dalmation toadflax maintains deeper maximal rooting depth and greater root mass density in deep soil layers compared to other co-existing native species it has access to surplus moisture stored in deeper soil layers under elevated CO₂ conditions (Schomp, 2007). Due to this more favorable plant water status it shows higher stomatal conductance under elevated CO₂ conditions.

A strikingly different pattern was observed in lodgepole pine. Here an increase in $\Delta^{13}\text{C}$ was accompanied by an increase in $\delta^{18}\text{O}$ of leaf biomass with increasing CO₂ exposure (Fig. 3b). The increase in $\delta^{18}\text{O}$ of leaf biomass is likely a result of decreased stomatal conductance. The decreased stomatal conductance should result in a decrease in c_i/c_a if photosynthetic CO₂ demand is unchanged. However, we observed an increase in c_i/c_a in lodgepole pine exposed to elevated CO₂ levels likely due to decreased photosynthetic capacity. The N concentration of plant leaves is positively correlated with photosynthetic capacity (Field and Mooney, 1986), as most of the N in leaves is used for synthesis of photosynthetic enzymes, and in particular Rubisco (Shangguan et al., 2000). The N concentration of lodgepole pine with an increase in atmospheric CO₂ concentration ($r^2=0.47$; $P<0.03$) (Fig. 4). This decrease in leaf N% at elevated CO₂ might be an additional indicator of decreased photosynthetic capacity in lodgepole pine at elevated CO₂ levels. No correlation was observed between leaf

**Effect of elevated
CO₂ on plants in YNP**S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



N and $\Delta^{13}\text{C}$ in dalmation toadflax (Fig. 4). Therefore we can infer that photosynthetic capacity of lodgepole pine decreases under higher CO_2 concentrations. Our result again contradicts the results obtained from FACE sites which frequently show that elevated CO_2 stimulates photosynthesis in C_3 plants, with trees showing the largest net increase in photosynthesis with increasing CO_2 exposure (reviewed in Nowak et al., 2004 ; Ainsworth and Long, 2005).

Acclimation or down regulation of photosynthesis is likely to occur in ecological situations where nutrient conditions are low or nitrogen is limiting (Stitt and Krapp, 1999; Sinclair et al., 2000). Nonetheless, even if down regulation is observed photosynthesis is often stimulated in long-term experiments even after 7–10 years of FACE (Nowak et al., 2004). The sampled lodgepole pine stand in this study most likely has been exposed to elevated CO_2 levels since centuries. We hypothesize that decreased photosynthetic capacity observed in our case could be due to severe reduction in plant available nitrogen in this forest stand due to phenomenon known as progressive nitrogen limitation (PNL). PNL develops because elevated CO_2 leads to N immobilization by plants and microbes which deplete soils of N, causing slower rates of N mineralization progressively reducing the mineral N available for plant uptake in the long term (Gill et al., 2002; Zak et al., 2000; Luo et al., 2004).

In this study, the combined use of ^{14}C , $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values of leaf biomass provided useful insight into the photosynthetic response of a native conifer and a non-native forb to elevated CO_2 conditions in Yellowstone National Park. The non-native forb, dalmation toadflax, responded by increasing stomatal conductance with no change in photosynthetic capacity, and the native lodgepole pine by decreasing stomatal conductance and photosynthetic capacity under elevated atmospheric CO_2 . The results from our study are in contrast to those from studies conducted in various FACE sites, emphasizing the importance of long-term, high resolution ecosystem-scale CO_2 experiments to understand responses of natural ecosystems to globally increasing CO_2 levels.

**Effect of elevated
 CO_2 on plants in YNP**S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. We thank J. Cable and P. Koenig for field assistance. The project was funded and supported by a Faculty Development grant from the University of Wyoming Research Office and the National Science Foundation – EPSCoR grant EPS-0447681. Radiocarbon analyses were performed at W. M. Keck AMS Lab at the University of California, Irvine, CA, USA.

References

- Ainsworth, E. A. and Long, S. P.: What have we learned from fifteen years of Free Air Carbon Dioxide Enrichment (FACE)? A Meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂, *New Phytol.*, 165, 351–372, 2005.
- Ainsworth, E. A. and Rogers, A.: The response of photosynthesis and stomatal conductance to rising [CO₂]: mechanisms and environmental interactions, *Plant Cell Environ.*, 30, 258–270, 2007.
- Barbour, M. M. and Farquhar, G. D.: Relative humidity- and ABA-induced variation in carbon and oxygen isotope ratios of cotton leaves, *Plant Cell Environ.*, 23, 473–485, 2000.
- Barbour, M. M., Andrews, T. J., and Farquhar, G. D.: Correlations between oxygen isotope ratios of wood constituents of *Quercus* and *Pinus* samples from around the world, *Aust. J. Plant Physiol.*, 28, 335–348, 2001.
- Barbour, M. M.: Stable oxygen isotope composition of plant tissue: a review, *Funct. Plant Biol.*, 34, 83–94, 2007.
- Bettarini, I., Calderoni, G., Miglietta, F., Raschi, A., and Ehleringer, J.: Isotopic carbon discrimination and leaf nitrogen content of *Erica arborea* L. along a CO₂ concentration gradient in a CO₂ spring in Italy, *Tree Physiol.*, 15, 327–332, 1995.
- Dang, Q. L. and Cheng, S.: Effects of soil temperature on ecophysiological traits in seedlings of four boreal tree species, *Forest Ecol. Manag.*, 194, 379–387, 2004.
- Ehleringer, J. R., Hall, A. E., and Farquhar, G. D.: *Stable Isotopes and Plant Carbon-Water Relations*, Elsevier, New York, USA, 1993.
- Farquhar, G. D., Ehleringer, J. R., and Hubick, K. T.: Carbon isotope discrimination and photosynthesis, *Annu. Rev. Plant Phys.*, 40, 503–537, 1989.
- Farquhar, G. D. and Lloyd, J.: Carbon and oxygen isotope effects in the exchange of carbon dioxide between terrestrial plants and the atmosphere, in: *Stable Isotopes and Plant Carbon-*

BGD

5, 3825–3843, 2008

Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Water Relations, edited by: Ehleringer, J. R., Hall, A. E., and Farquhar, G. D., Academic Press, San Diego, CA, USA, 47–70, 1993.

Field, C. and Mooney, H. A.: The photosynthesis - nitrogen relationship in wild plants, in: On the Economy of Plant Form and Function, edited by: Givnish, T. J., Cambridge University Press, New York, NY, USA, 25–55, 1986.

Field, C. B., Lund, C. P., Chiariello, N. R., and Mortimer, B. E.: CO₂ effects on the water budget of grassland microcosm communities, *Glob. Change Biol.*, 3, 197–206, 1997.

Gill, R. A., Polley, H. W., Johnson, H. B., Anderson, L. J., Maherali, H., and Jackson, R. B.: Nonlinear grassland responses to past and future atmospheric CO₂, *Nature*, 417, 279–282, 2002.

Grams, T. E. E., Kozovitis, A. R., Haberele, K. H., Matyssek, R., and Dawson, T. E.: Combining $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ analyses to unravel competition, CO₂ and O₃ effects on the physiological performance of different-aged trees, *Plant Cell Environ.*, 30, 1023–1034, 2007.

Knapp, A. K., Hamerlynck, E. P., Ham, J. M., and Owensby, C. E.: Responses in stomatal conductance to elevated CO₂ in 12 grassland species that differ in growth form, *Plant Ecol.*, 125, 31–41, 1996.

Lockheart, M. J., Van Bergen, P. F., and Evershed, R. P.: Variations in the stable carbon isotope compositions of individual lipids from the leaves of modern angiosperms: implications for the study of higher land plant-derived sedimentary organic matter, *Org. Geochem.*, 26, 137–153, 1997.

Luo, Y., Currie, W., Dukes, J., Finzi, A., Hartwig, U., Hungate, B., McMurtrie, R., Oren, R., Parton, W., Pataki, D., Shaw, M., Zak, D., and Field, C.: Progressive nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide, *BioScience*, 54, 731–739, 2004.

Miglietta, F., Bettarini, I., Raschi, A., Körner, C., and Vaccari, F. P.: Isotope discrimination and photosynthesis of vegetation growing in the Bossoleto CO₂ spring, *Chemosphere*, 36, 771–776, 1998.

Morgan, J. A., Pataki, D. E., Körner, C., Clark, H., Del Grosso, S. J., Grunzweig, J. M., Knapp, A. K., Knippert, J. B., Nowak, R. S., Parton, W. J., Polley, H. W., and Shaw, M. R.: Water relations in grassland and desert ecosystems exposed to elevated atmospheric CO₂, *Oecologia*, 140, 11–25, 2004.

Mooney, H. A., Field, C., VasquezYanes, C., and Chu, C.: Environmental controls on stomatal conductance in a shrub of the humid tropics, *P. Natl. Acad. Sci. USA*, 80, 1295–1297, 1983.

Nowak, R. S., Ellsworth, D. S., and Smith, S. D.: Functional responses of plants to elevated

BGD

5, 3825–3843, 2008

Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Effect of elevated
CO₂ on plants in YNP**S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



atmospheric – do photosynthetic and productivity data from FACE experiments support early predictions?, *New Phytol.*, 162, 253–280, 2004.

Pataki, D. E., Ellsworth, D. W., King, J. S., Leavitt, S. W., Lin, G., Pendall, E., Siegwolf, R., van Kessel, C., and Ehleringer, J. R.: Tracing changes in ecosystem function under elevated CO₂, *Bioscience*, 53, 805–818, 2003.

Scheidegger, Y., Saurer, M., Bahn, M., and Siegwolf, R.: Linking stable oxygen and carbon isotopes with stomatal conductance and photosynthetic capacity: a conceptual model, *Oecologia*, 125, 350–357, 2000.

Schomp, J. M.: Rising atmospheric CO₂ concentrations and plant invasion in the northern mixed-grass prairie: an ecophysiological perspective, M.S. thesis, University of Wyoming, Laramie, WY, USA, 2007

Schwarz, P. A., Fahey, T. J., and Dawson, T. E.: Seasonal soil temperature effects on photosynthesis in red spruce (*Picea rubens*) saplings, *Tree Physiol.*, 17, 187–194, 1997.

Shangguan, Z. P., Shao, M. A., and Dyckmans, J.: Nitrogen nutrition and water stress effects on leaf photosynthetic gas exchange and water use efficiency in winter wheat, *Environ. Exp. Bot.*, 44, 141–149, 2000.

Sinclair, T. R., Pinter Jr., P. J., Kimball, B. A., Adamsen, F. J., LaMorte, R. L., Wall, G. W., Hunsaker, D. J., Adam, N., Brooks, T. J., Garcia, R. L., Thompson, T., Leavitt, S., and Matthias, A.: Leaf nitrogen concentration of wheat subjected to elevated [CO₂] and either water or N deficits, *Agr. Ecosyst. Environ.*, 79, 53–60, 2000.

Stitt, M. and Krapp, A.: The interaction between elevated carbon dioxide and nitrogen nutrition: the physiological and molecular background, *Plant, Cell Environ.*, 22, 583–621, 1999.

Stuiver, M. and Polach, H. A.: Reporting of ¹⁴C data, *Radiocarbon*, 19, 355–363, 1977.

Sullivan, P. F. and Welker, J. M.: Variation in leaf physiology of *Salix arctica* within and across ecosystems in the High Arctic: test of a dual isotope (Δ¹³C and Δ¹⁸O) conceptual model, *Oecologia*, 151, 372–386, 2007.

Tognetti, R., Minnocci, A., Peñuelas, J., Raschi, A., and Jones, M. B.: Comparative field water relations of three Mediterranean shrub species co-occurring at a natural CO₂ vent, *J. Exp. Bot.*, 51, 1135–1146, 2000.

van Gardingen, P. R., Grace, J., Harkness, D. D., Miglietta, F., and Raschi, A.: Carbon dioxide emissions at an Italian mineral spring: measurements of average CO₂ concentration and air temperature, *Agr. Forest Meteorol.*, 73, 17–27, 1995.

Warren, C. R. and Adams, M. A.: Water availability and branch length determine δ¹³C in foliage

of *Pinus pinaster*, *Tree Physiol.*, 20, 637–643, 2000.

Werner, C. and Brantley, S.: CO₂ emissions from the Yellowstone volcanic systems, *Geochem. Geophys. Geosys.*, 4, 1–27, 2003.

Wullschleger, S. D., Tschaplinski, T. J., and Norby, R. J.: Plant water relations at elevated CO₂ – implications for water-limited environments, *Plant Cell Environ.*, 25, 319–331, 2002.

Zak, D. R., Pregitzer, K. S., King, J. S., and Holmes, W. E.: Elevated atmospheric CO₂, fine roots and the response of soil microorganisms: a review and hypothesis, *New Phytol.*, 147, 201–222, 2000.

Zhang, S. and Dang, Q. L.: Effects of soil temperature and elevated atmospheric CO₂ concentration on gas exchange, in vivo carboxylation and chlorophyll fluorescence in jack pine and white birch seedlings, *Tree Physiol.*, 25, 523–531, 2005.

BGD

5, 3825–3843, 2008

Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Effect of elevated
CO₂ on plants in YNP**S. Sharma and
D. G. Williams

Fig. 1. Map of Yellowstone National Park showing the Ochre Spring and Mammoth Hot Springs sampling locations.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams




Fig. 2. Correlation between carbon isotope discrimination ($\Delta^{13}\text{C}$) and average flux-weighted CO_2 in the leaves of dalmation toadflax (open and filled triangles) and lodgepole pine (open and filled circles) were exposed to. The flux-weighted CO_2 exposure was calculated from the radiocarbon or ^{14}C content of leaves using Eq. (1).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams




Fig. 3. Correlation between ¹³C discrimination ($\Delta^{13}\text{C}$) and the $\delta^{18}\text{O}$ of bulk leaf material in dalmation toadflax and lodgepole pine. The filled and open symbols represent data from the control and elevated sites respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Effect of elevated CO₂ on plants in YNP

S. Sharma and
D. G. Williams




Fig. 4. Nitrogen concentrations of leaf biomass show different relationships over gradients of CO₂ exposure between dalmation toadflax (open and filled triangles) lodgepole pine (open and filled circles).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

