

*Biogeosciences Discussions* is the access reviewed discussion forum of *Biogeosciences*

**Modeling  
biogeochemical  
responses to storms**

M. Fujii and Y. Yamanaka

# Effects of storms on primary productivity and air-sea CO<sub>2</sub> exchange in the subarctic western North Pacific: a modeling study

M. Fujii<sup>1</sup> and Y. Yamanaka<sup>2,3</sup>

<sup>1</sup>Sustainability Governance Project, Creative Research Initiative, Hokkaido University, N9W8, Kita-ku, Sapporo, Hokkaido 060-0809, Japan

<sup>2</sup>Graduate School of Environmental Earth Science, Hokkaido University, N10W5, Kita-ku, Sapporo, Hokkaido 060-0810, Japan

<sup>3</sup>Frontier Research System for Global Change, 3173-25 Showa-machi, Kanazawa-ku, Yokohama, Kanagawa 236-0001, Japan

Received: 13 November 2007 – Accepted: 29 November 2007 – Published: 4 January 2008

Correspondence to: M. Fujii (mfujii@sgp.hokudai.ac.jp)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

Biogeochemical responses of the open ocean to storms and their feedback to climate are still poorly understood. Using a marine ecosystem model, we examine biogeochemical responses to the storms in the subarctic western North Pacific. The storms in summer through early autumn enhance primary production by wind-induced nutrient injections into the surface waters while the storms in the other seasons reduce primary production by intensifying light limitation on the phytoplankton growth due to vertical dilution of the phytoplankton. The two compensating effects diminish the storm-induced annual change of primary production to only 1%. On the contrary, the storms enhance the annual sea-to-air CO<sub>2</sub> efflux by no less than 34%, resulting from storm-induced strong winds. Our results suggest that previous studies using climatological wind and CO<sub>2</sub> data probably underestimated the sea-to-air CO<sub>2</sub> efflux during storms in the subarctic western North Pacific, and therefore, that continuous observations are required to reduce uncertainties in the global oceanic CO<sub>2</sub> uptake.

## 1 Introduction

Episodic atmospheric disturbances such as storms reduce solar radiation and enhance wind-driven vertical mixing and upwelling in the surface waters (e.g. Price, 1981; Greatbatch, 1985; Cornillon et al., 1987; Shay and Elsberry, 1987; Sakaida et al., 1998; Hong et al., 2003). Such physical changes are expected to impact marine biogeochemistry. For example, decreases in solar radiation and downward transport of phytoplankton out of the euphotic zone are likely to increase light limitation on the phytoplankton growth. On the other hand, wind-induced nutrient injections into the surface waters allow for potentially more growth of phytoplankton.

However, biogeochemical responses of the open ocean to storms are still poorly understood (e.g. Babin et al., 2004). In situ ship-based sampling has greatly enhanced our understanding of the biogeochemistry, but rough weather prevents sampling dur-

**BGD**

5, 65–81, 2008

## Modeling biogeochemical responses to storms

M. Fujii and Y. Yamanaka

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

ing storms. Only a few cruises and mooring buoys have serendipitously encountered episodic strong wind events (e.g. Marra et al., 1990; Bates et al., 1998; Dickey et al., 1998; Nemoto et al., 1999). In most such cases, sudden decrease in the sea surface temperature (SST) and abrupt change in the partial pressure of CO<sub>2</sub> in seawater (pCO<sub>2sea</sub>) have been observed. Recently satellite-derived estimates of wind speed, temperature and chlorophyll-a concentration in the surface waters have provided an opportunity to explore the physical and biological responses to episodic strong wind events (e.g. Babin et al., 2004). However, they cannot clarify whether the observed changes in chlorophyll-a concentration are the result of wind-induced nutrient injections or the result of entrainment of waters with high chlorophyll-a concentration. In addition, wind-induced change in carbon biogeochemistry, which is considered to greatly contribute to ecosystem dynamics and climate, cannot be estimated from observations by satellites at the present stage.

On the other hand, ecosystem modeling allows us to estimate the episodic storm-induced biogeochemical responses at any spatial and temporal scale. Therefore, the application of ecosystem models could significantly advance our understanding of the biogeochemical response to storm passage in the open ocean (e.g. Babin et al., 2004). Biogeochemical responses to storms have hardly been explored in the subarctic western North Pacific, which is considered one of the key regions impacting marine resources and uptake of anthropogenic CO<sub>2</sub> (e.g. Tsurushima et al., 2002). We therefore undertake this study to investigate the storm-induced biogeochemical responses in this region using a marine ecosystem model. We apply the model to time-series Station KNOT (Kyodo North Pacific Ocean Time-series; 44° N, 155° E). More than ten storms capable of deepening the mixed layer depth (MLD) and affecting the biogeochemistry appear at Station KNOT on average each year (Table 1). We investigate the biogeochemical responses to the storms, especially focusing on the impacts to the phytoplankton dynamics and the air-sea CO<sub>2</sub> exchange.

---

**Modeling  
biogeochemical  
responses to storms**M. Fujii and Y. Yamanaka

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 2 Model description and experimental design

We use a 16-compartment marine ecosystem model (Fig. 1; Fujii et al., 2002, 2005, 2007; Yamanaka et al., 2004). In this model, phytoplankton are categorized into two groups: diatoms (PL) and non-diatom small phytoplankton (PS) including coccolithophorids. Phytoplankton components utilize nitrate ( $\text{NO}_3$ ) and ammonium ( $\text{NH}_4$ ) in the process of photosynthesis and produce soft tissue in the form of particulate organic nitrogen (PON). Along with photosynthesis, diatoms utilize silicate ( $\text{Si}(\text{OH})_4$ ) to produce frustules of biogenic silica. Zooplankton are categorized into three groups: microzooplankton (ZS) including foraminifera, mesozooplankton (ZL) and predatory zooplankton (ZP). Seasonal vertical migration of mesozooplankton is taken into account: the mesozooplankton vertically migrate out of the model domain below 330 m-depth at the end of August, and 20% of the mesozooplankton return to the euphotic zone at the beginning of April every year, which is similar to the procedure used by previous studies (Kishi et al., 2001). Coccolithophorids and foraminifera produce hard shells of calcium carbonate ( $\text{CaCO}_3$ ). Total alkalinity (TALK) is calculated by balances of  $\text{CaCO}_3$ ,  $\text{NO}_3$  and  $\text{NH}_4$ . The  $\text{TCO}_2$  is calculated by balances of TALK,  $\text{NO}_3$  and  $\text{NH}_4$  (with a carbon: nitrogen ratio of 6.625; Redfield et al., 1963). Therefore, the  $\text{pCO}_{2\text{sea}}$  is calculated and the air-sea  $\text{CO}_2$  flux can be estimated by this model. Values of biogeochemical parameters in the model are the same as those of Fujii et al. (2007).

The air-sea flux of  $\text{CO}_2$  is calculated in the model using the transfer velocity-wind speed relationships of Wanninkhof (1992) as follows:

$$\text{Air-sea } \text{CO}_2 \text{ flux} = 0.31 U^2 \sqrt{660 / Sc} L \{ \text{pCO}_{2\text{sea}} - \text{pCO}_{2\text{air}} \}, \quad (1)$$

where  $U$  is the wind speed at the sea surface ( $\text{m s}^{-1}$ ),  $Sc$  is the Schmidt number for  $\text{CO}_2$ , expressed as:

$$Sc = 2073.1 - 125.62 \text{SST} + 3.6276 \text{SST}^2 - 0.043219 \text{SST}^3, \quad (2)$$

$L$  is the solubility of  $\text{CO}_2$  calculated from temperature and salinity (Weiss, 1974). The

**BGD**

5, 65–81, 2008

---

### Modeling biogeochemical responses to storms

M. Fujii and Y. Yamanaka

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

$p\text{CO}_{2\text{air}}$  is the partial pressure of  $\text{CO}_2$  in the atmosphere and the monthly in situ observational data collected at Mauna Loa Observatory are used (Keeling et al., 1982).

The model is driven by the wind and solar radiation at the sea surface, and the temperature and salinity at the surface and at the bottom of the model domain (330 m).

We use the National Centers for Environmental Prediction (NCEP) objectively analyzed data (Kalney et al., 1996) for the daily (every 6 h) wind and solar radiation at the sea surface, the Reynolds weekly data for the SST (Reynolds and Smith, 1995), and the KNOT time-series observations for the temperature at the bottom and the salinity at the surface and bottom (Fujii et al., 2002; Tsurushima et al., 2002).

We define storm events as those in which the wind speed is more than  $2\sigma$  value from a 30-d running mean. To examine effects of storms on biogeochemistry, we carried out two experiments, namely Exp-1 (with storms) and Exp-2 (without storms). In Exp-1, the model is driven by the winds including more than  $2\sigma$  values. In Exp-2, the model is driven by the winds in which more than  $2\sigma$  values are filtered out. The simulation is calculated from 1982 to 2000, and the 19-y monthly-mean model results are presented below.

### 3 Results and discussions

The model well reproduces the observed seasonal changes of both physical environments and biogeochemistry at Station KNOT (Fig. 2; Imai et al., 2002; Tsurushima et al., 2002). The strong wind in winter causes the MLD, defined as the depth at which the vertical diffusive coefficient is  $1.0 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$ , to deepen dramatically to more than 100 m in late winter (Fig. 2a, b). The strong wind and  $p\text{CO}_{2\text{sea}}$  elevated by storm-induced normalized  $\text{TCO}_2$  ( $\text{NTCO}_2$ ; normalized to a constant salinity of 35 pss) injections into the surface waters cause tremendous sea-to-air  $\text{CO}_2$  efflux in late winter (Fig. 2a, d, e, f). The primary production is low in winter because of strong light limitation on the phytoplankton growth due to vertical dilution of the phytoplankton, along with low irradiance in this season (Fig. 2c). The primary production is relatively high in

**BGD**

5, 65–81, 2008

## Modeling biogeochemical responses to storms

M. Fujii and Y. Yamanaka

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

spring through early autumn when the light limitation on the phytoplankton growth is alleviated by high irradiance and stratification of the surface waters in these periods. The oceanic region functions as a sink of the atmospheric CO<sub>2</sub> in spring through autumn, supported by large biological uptake of CO<sub>2</sub>, as reported by previous observations (e.g. Tsurushima et al., 2002).

By comparing the model results between Exp-1 (with storms) and Exp-2 (without storms), we find that pCO<sub>2sea</sub> is the most sensitive biogeochemical parameter to the storms, and therefore, that the storms contribute significantly to air-sea exchange of CO<sub>2</sub> (Fig. 2e, f). The sea-to-air CO<sub>2</sub> efflux is enhanced by the storms in late spring, early summer and autumn. For example, the pCO<sub>2sea</sub> abruptly increases by 42 μatm during a storm passage in mid-June 1994 in which the wind speed is more than 3σ value from the 30-day running mean (Fig. 3b). The pCO<sub>2sea</sub> reaches one of its annual peaks during the storm, although the annual maximum generally appears in late winter in the subarctic western North Pacific (Fig. 2e; Tsurushima et al., 2002). The sea-to-air CO<sub>2</sub> efflux abruptly increases during the storm passage (Fig. 3c), accounting for no less than 10% of the annual sea-to-air efflux in 1994. The sudden increase in pCO<sub>2sea</sub> results from the increase in NTCO<sub>2</sub> by 18 mmolC m<sup>-3</sup> (Fig. 3a) because of the storm-induced vertical mixing, which dominates the counteracting effect of the storm-induced sea surface cooling of 0.8°C. This is consistent with a result based on mooring buoy data deployed in the East China Sea during the passage of three typhoons in 1995 that for changes in pCO<sub>2sea</sub> the effect of TCO<sub>2</sub> increase dominated that of SST decrease (Nemoto et al., 1999).

On the contrary, the sea-to-air efflux is reduced slightly by the storms in late summer, because the primary production is enhanced by the storm-induced nutrient injections into the surface waters (Fig. 2c, f). The model result of the decrease in the pCO<sub>2sea</sub> in late summer is consistent with a ship-based result of Bates et al. (1998), who observed a sudden decrease in pCO<sub>2sea</sub> in the Sargasso Sea in the subtropical gyre during a hurricane passage in August 1995. However, the cause of the storm-induced decrease in the sea-to-air CO<sub>2</sub> efflux is different from this study: they observed only a slight in-

---

**Modeling  
biogeochemical  
responses to storms**M. Fujii and Y. Yamanaka

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

crease in  $\text{TCO}_2$ , and the decrease in  $\text{pCO}_{2\text{sea}}$  was followed by a decrease in SST. The difference in responses of  $\text{pCO}_{2\text{sea}}$  to storms between this study and Bates et al. (1998) results from differences in the upper ocean structure and hydrographic conditions between the subarctic and subtropical gyres. In the subarctic gyre, the pycnocline is relatively shallow, and storms occasionally mix the surface waters with  $\text{TCO}_2$ -rich deep waters below the pycnocline. In the subtropical gyre, on the other hand, the pycnocline is usually deep and storm-induced vertical mixing does not reach to the deep waters. The magnitude of the sea surface cooling also depends on the upper ocean structure and hydrographic conditions, as noted in previous studies (e.g. Cornillon et al., 1987; Sakaida et al., 1998). Storm-induced sea surface cooling is more efficient in the subtropical ocean than in the subarctic ocean because of the permanently warm surface waters in the subtropical ocean.

The model result shows that the annual sea-to-air  $\text{CO}_2$  efflux is higher in Exp-1 ( $244.55 [\text{mmolC m}^{-2} \text{y}^{-1}]$ ) than in Exp-2 ( $160.60 [\text{mmolC m}^{-2} \text{y}^{-1}]$ ) (Table 2), and therefore, that the storms enhance the annual sea-to-air  $\text{CO}_2$  efflux by 34%. Previous studies, in which the flux was calculated using the climatological wind and  $\text{pCO}_{2\text{sea}}$  data, probably underestimated the effect of storm events on air-sea  $\text{CO}_2$  exchange and overestimated the role of the entire subarctic western North Pacific in taking up oceanic  $\text{CO}_2$ .

The storms enhance the primary production in summer through early autumn (Fig. 2c), because the nutrient limitation on the phytoplankton growth is alleviated by the storm-induced nutrient injections into the surface waters. This is consistent with previous observation-based results of the oligotrophic subtropical ocean (e.g. Babin et al., 2004), but not as much as in the subtropical ocean, because of higher pre-storm surface nutrient concentrations, and therefore, less nutrient limitation on the phytoplankton growth, in the subarctic ocean than in the subtropical ocean. On the contrary, the storms reduce the primary production in the other seasons, because the storm-induced vertical mixing increases the light limitation on the phytoplankton growth. This result shows that the effect of storms on the phytoplankton dynamics changes with

**BGD**

5, 65–81, 2008

---

**Modeling  
biogeochemical  
responses to storms**

M. Fujii and Y. Yamanaka

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

pre-storm conditions such as the irradiance and nutrient concentrations in the surface waters. The two compensating effects diminish the storm-induced annual change of primary production at Station KNOT to only 1% (Table 2).

#### 4 Conclusion and remarks

Using a marine ecosystem model, we examine the biogeochemical responses to storms at Station KNOT, located in the subarctic western North Pacific. In the simulation, the storms significantly affect both ecosystem dynamics and air-sea  $\text{CO}_2$  exchange, and the effects change with the irradiance, upper ocean structure, hydrographic conditions and nutrient concentrations.

The storms enhance the primary production in summer through early autumn because of the storm-induced nutrient injections into the surface waters. On the contrary, the primary production in the other seasons is reduced by the storm-driven vertical mixing and subsequent limitation of the light limitation on the phytoplankton growth. The two compensating effects diminish the storm-induced annual change of primary production to only 1%.

The storms enhance the sea-to-air  $\text{CO}_2$  efflux in late spring, early summer and autumn by the strong wind, whereas the storms reduce the sea-to-air  $\text{CO}_2$  efflux in late summer because of large biological uptake of  $\text{CO}_2$  stimulated by the storm-induced nutrient injections into the surface waters. The storms are estimated to enhance the annual sea-to-air  $\text{CO}_2$  flux by 34% in this region.

During storms,  $\text{pCO}_{2\text{sea}}$  tends to increase more in the subarctic ocean because of the storm-induced increase in surface  $\text{NTCO}_2$ , whereas  $\text{pCO}_{2\text{sea}}$  tends to decrease in the subtropical ocean because of the storm-induced decrease in the SST. Therefore, it is necessary to perform direct measurements during storm passage in various oceanic regions to elucidate which effects dominate the storm-induced  $\text{pCO}_{2\text{sea}}$  change in each region. Previous studies that calculated the air-sea  $\text{CO}_2$  flux using climatological wind and  $\text{pCO}_{2\text{sea}}$  data probably underestimated the significant contribution of storms to the

**BGD**

5, 65–81, 2008

### Modeling biogeochemical responses to storms

M. Fujii and Y. Yamanaka

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**



air-sea CO<sub>2</sub> exchange. Therefore, to reduce uncertainties in the global oceanic CO<sub>2</sub> uptake, changes in the pCO<sub>2sea</sub> caused by episodic atmospheric disturbances should be measured continuously.

While climate changes associated with global warming may influence the frequency and intensity of storms (e.g. Emanuel, 1987, 2005; Saunders and Harris, 1997; Sugi et al., 2002; Yoshimura and Sugi, 2005; Webster et al., 2005; Yoshimura et al., 2006), storm-induced biogeochemical activity may also contribute to climate. Therefore, to predict future climate change, it is essential that we elucidate the biogeochemical responses to storms. Although it is difficult to conduct direct observations during the rough weather, we would very much like to have high-frequency, in-situ biogeochemical observations with which we could conduct more accurate simulations.

*Acknowledgements.* The authors wish to thank M. N. Aita for providing data necessary for this study. Thanks are extended to S. L. Smith for his proofreading of the English manuscript. M. Fujii was supported by MEXT through Special Coordination Funds for Promoting Science and Technology. A figure was produced by the GFD-DENNOU Library.

## References

- Babin, S. M., Carton, J. A., Dickey, T. D., and Wiggert, J. D.: Satellite evidence of hurricane-induced phytoplankton blooms in an ocean desert, *J. Geophys. Res.*, 109, C03043, doi:10.1029/2003JC001938, 2004.
- Bates, N. R., Takahashi, T., Chipman, D. W., and Knap, A. H.: Variability of pCO<sub>2</sub> on diel to seasonal timescales in the Sargasso Sea near Bermuda, *J. Geophys. Res.*, 103, 15567–15585, 1998.
- Cornillon, P., Stramma, L., and Price, J. F.: Satellite measurements of sea surface cooling during hurricane Gloria, *Nature*, 326, 373–375, 1987.
- Dickey, T. D., Frye, D., Mcneil, J., Manov, D., Nelson, N., Sigurdson, D., Jannasch, H., Siegel, D., Michaels, A. F., and Johnson, R. J.: Upper-ocean temperature response to Hurricane Felix as measured by the Bermuda Testbed Mooring, *Mon. Weather Rev.*, 126, 1195–1201, 1988.

**BGD**

5, 65–81, 2008

---

## Modeling biogeochemical responses to storms

M. Fujii and Y. Yamanaka

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

- Emanuel, K. A.: The dependence of hurricane intensity on climate, *Nature*, 326, 483–485, 1987.
- Emanuel, K.: Increasing destructiveness of tropical cyclones over the past 30 years, *Nature*, 436, 686–688, doi:10.1038/nature03906, 2005.
- 5 Fujii, M., Nojiri, Y., Yamanaka, Y., and Kishi, M. J.: A one-dimensional ecosystem model applied to time series station KNOT, *Deep-Sea Research Part II*, 49, 5441–5461, 2002.
- Fujii, M., Yamanaka, Y., Nojiri, Y., and Kishi, M. J.: Comparison of seasonal characteristics in biogeochemistry among the subarctic North Pacific stations described with a NEMURO-based marine ecosystem model, *Ecol. Model.*, 202, 52–67, 2007.
- 10 Fujii, M., Yoshie, N., Yamanaka, Y., and Chai, F.: Simulated biogeochemical responses to iron enrichments in three high nutrient, low chlorophyll (HNLC) regions, *Prog. Oceanogr.*, 64, 307–324, doi:10.1016/j.pocean.2005.02.017, 2005.
- Geng, Q.-Z. and Sugi, M.: Possible change of extratropical cyclone activity due to enhanced greenhouse gases and sulfate aerosols – Study with a high-resolution AGCM, *J. Climate*, 16, 2262–2274, 2003.
- 15 Greatbatch, R. J.: On the role played by upwelling of water in lowering sea surface temperatures during the passage of a storm, *J. Geophys. Res.*, 90(C6), 11 751–11 755, 1985.
- Hong, C. H. and Yoon, J. H.: A three-dimensional numerical simulation of Typhoon Holly in the northwestern Pacific Ocean, *J. Geophys. Res.*, 108, 3282, doi:10.1029/2002JC001563, 2003.
- 20 Imai, K., Nojiri, Y., Tsurushima, N., and Saino, T.: Time series of seasonal variation of primary productivity at station KNOT (44° N, 155° E) in the sub-arctic western North Pacific, *Deep-Sea Res.*, Part II, 49, 5395–5408, 2002.
- Kalney, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *Bull. Am. Meteorol. Soc.*, 77, 437–471, 1996.
- 25 Keeling, C. D., Bacastow, R. B., Whorf, T. P.: Measurements of the concentration of carbon dioxide at Mauna Loa Observatory, Hawaii, in: *Carbon Dioxide Review*, edited by: Clark, W.C., Oxford University Press, New York, 1982.
- 30 Kishi, M. J., Motono, H., Kashiwai, M., and Tsuda, A.: An ecological-physical coupled model with ontogenetic vertical migration of zooplankton in the northwestern Pacific, *J. Oceanogr.*, 57, 499–507, 2001.

**BGD**

5, 65–81, 2008

---

**Modeling  
biogeochemical  
responses to storms**

M. Fujii and Y. Yamanaka

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

- Marra, J., Bidigare, R. R., and Dickey, T. D.: Nutrients and mixing, chlorophyll and phytoplankton growth, *Deep-Sea Res.*, 37, 127–143, 1990.
- Nemoto, K., Takatani, S., Ogawa, K., Umeda, T., Oyama, J., Midorikawa, T., Kimoto, T., and Inoue, H. Y.: Continuous observations of atmospheric and oceanic CO<sub>2</sub> using the moored buoy in the East China Sea, *Proceedings of the 2nd International Symposium CO<sub>2</sub> in the Oceans*, CGER/NIES, Tsukuba, 545–553, 1999.
- Price, J. F.: Upper ocean response to a hurricane, *J. Phys. Oceanogr.*, 11, 153–175, 1981.
- Redfield, A. C., Ketchum, B. A., and Richards, F. A.: The influence of organisms on the composition of seawater, in: *The Sea*, edited by: Hill, M. N., Wiley, New York, 26–77, 1963.
- Reynolds, R. W. and Smith, T. M.: A high resolution global sea surface temperature climatology, *J. Climate*, 8, 1571–1583, 1995.
- Sakaida, F., Kawamura, H., and Toba, Y.: Sea surface cooling caused by typhoons in the Tohoku Area in August 1989, *J. Geophys. Res.*, 103, 1053–1065, 1998.
- Saunders, M. A. and Harris, A. R.: Statistical evidence links exceptional 1995 Atlantic hurricane season to record sea warming, *Geophys. Res. Lett.*, 24, 1255–1258, 1997.
- Shay, L. K. and Elsberry, R. L.: Near-inertial ocean current response to Hurricane Frederick, *J. Phys. Oceanogr.*, 17, 1249–1269, 1987.
- Sugi, M., Noda, A. and Sato, N.: Influence of the global warming on tropical cyclone climatology: an experiment with the JMA global model, *J. Meteorol. Soc. Jpn.*, 80, 249–272, 2002.
- Takahashi, T., Sutherland, S. C., Sweeney, C., Poisson, A., Metz, N., Tilbrook, B., Bates, N., Wanninkhof, R., Feely, R. A., Sabine, C., Olafsson, J., and Nojiri, Y.: Global sea-air CO<sub>2</sub> flux based on climatological surface ocean pCO<sub>2</sub>, and seasonal biological and temperature effects, *Deep-Sea Res. Part II*, 49, 1601–1622, 2002.
- Tsurushima, N., Nojiri, Y., Imai, K., and Watanabe, S.: Seasonal variations of carbon dioxide system and nutrients in the surface mixed layer at station KNOT (44° N, 155° E) in the subarctic western North Pacific, *Deep-Sea Research Part II*, 49, 5377–5394, 2002.
- Wanninkhof, R.: Relationship between wind speed and gas exchange, *J. Geophys. Res.*, 97(C5), 7373–7382, 1992.
- Webster, P. J., Holland, G. J., Curry, J. A., and Chang, H.-R.: Changes in tropical cyclone number, duration, and intensity in a warming environment, *Science*, 309, 1844–1846, 2005.
- Weiss, R. F.: Carbon dioxide in water and seawater: The solubility of a non-ideal gas, *Mar. Chem.*, 2, 203–215, 1974.
- Yamanaka, Y., Yoshie, N., Fujii, M., Aita, M. N., and Kishi, M. J.: An ecosystem model coupled

**BGD**

5, 65–81, 2008

---

**Modeling  
biogeochemical  
responses to storms**

M. Fujii and Y. Yamanaka

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

with Nitrogen-Silicon-Carbon cycles applied to Station A7 in the Northwestern Pacific, J. Oceanogr., 60, 227–241, 2004.

Yoshimura, J., and Sugi, M.: Tropical cyclone climatology in a high-resolution AGCM – Impact of SST warming and CO<sub>2</sub> increase – , SOLA, 1, 133–136, 2005.

- 5 Yoshimura, J., Sugi, M., and Noda, A.: Influence of global warming on tropical cyclone frequency, J. Meteorol. Soc. Jpn., 84, 405–428, 2006.

**BGD**

5, 65–81, 2008

---

**Modeling  
biogeochemical  
responses to storms**

M. Fujii and Y. Yamanaka

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**EGU**

**Modeling  
biogeochemical  
responses to storms**

M. Fujii and Y. Yamanaka

**Table 1.** Number of days on which the wind speed was more than  $2\sigma$  or  $3\sigma$  values from the 30-d running mean in each year from 1982 to 2000 at Station KNOT.

Year	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	Total
$>2\sigma$	8	10	8	9	10	15	11	12	9	14	8	14	12	6	12	12	10	10	13	203
$>3\sigma$	0	2	0	0	0	0	1	2	0	1	0	0	1	0	0	0	0	0	0	7

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modeling  
biogeochemical  
responses to storms

M. Fujii and Y. Yamanaka

**Table 2.** Annual-mean values of the wind speed, MLD, primary production,  $p\text{CO}_{2\text{sea}}$  and sea-to-air  $\text{CO}_2$  efflux (positive upward) for Exp-1 (with storms), Exp-2 (without storms), differences between the experiments (Exp-1 minus Exp-2), and ratios of the differences to the values for Exp-1, at Station KNOT.

	Exp-1	Exp-2	Difference ((Exp-1)–(Exp-2))	Ratio ((Exp-1)–(Exp-2))/(Exp-1) [%]
Wind speed [ $\text{m s}^{-1}$ ]	7.65	7.50	0.15	1.92
MLD [m]	45.46	40.66	4.80	10.55
Primary production [ $\text{mgC m}^{-2} \text{d}^{-1}$ ]	286.35	283.16	3.19	1.12
$p\text{CO}_{2\text{sea}}$ [ $\mu\text{atm}$ ]	357.45	357.33	0.12	0.03
Sea-to-air $\text{CO}_2$ efflux [ $\text{mmolC m}^{-2} \text{y}^{-1}$ ]	244.55	160.60	83.95	33.63

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Modeling  
biogeochemical  
responses to storms**

M. Fujii and Y. Yamanaka



**Fig. 1.** Schematic view of a 16-compartment marine ecosystem model (from Fujii et al., 2002, 2005, 2007 and Yamanaka et al., 2004).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Modeling biogeochemical responses to storms

M. Fujii and Y. Yamanaka



**Fig. 2.** Seasonal changes of **(a)** wind speed [ $\text{m s}^{-1}$ ], **(b)** MLD [m], **(c)** primary production [ $\text{mgC m}^{-2} \text{d}^{-1}$ ], **(d)**  $\text{NTCO}_2$  [ $\text{mmolC m}^{-3}$ ], **(e)**  $\text{pCO}_{2\text{sea}}$  [ $\mu\text{atm}$ ], and **(f)** sea-to-air  $\text{CO}_2$  efflux [ $\text{mmolC m}^{-2} \text{d}^{-1}$ ] (positive upward) for Exp-1 (with storms; in solid lines) and Exp-2 (without storms; in dotted lines) at Station KNOT. Error bars represent  $1\sigma$  values in Exp-1. Dots: observational data from 1998 to 2000 (Imai et al., 2002; Tsurushima et al., 2002).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 3.** Modeled **(a)** surface NTICO<sub>2</sub> [mmolC m<sup>-3</sup>], **(b)** pCO<sub>2</sub>sea and pCO<sub>2</sub>air [µatm], and **(c)** sea-to-air CO<sub>2</sub> efflux [mmolC m<sup>-2</sup> d<sup>-1</sup>] (positive upward) for Exp-1 (with storms; in red solid lines) and Exp-2 (without storms; in blue dotted lines) in 1994 at Station KNOT. The storm with the wind speed more than 3σ value from the 30-d running mean passed during the hatched period (mid-June).