

***Interactive comment on* “Effects of storms on primary productivity and air-sea CO₂ exchange in the subarctic western North Pacific: a modeling study” by M. Fujii and Y. Yamanaka**

M. Fujii and Y. Yamanaka

Received and published: 27 March 2008

First of all, the authors greatly appreciate the constructive review on our manuscript. We have revised our manuscript, basically according to the reviewers comments in such ways as described below.

Responses to major comments

Because the model used in this study is vertically one-dimensional, it does not reproduce effects of horizontal advection. The authors have excluded the data from high-salinity (>33.2 psu) which are considered to be influenced by subtropical water in our discussion for model-data comparison. This is the same procedure introduced in Tsurushima et al. (2002). Using this procedure, the model results have demon-

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



strated better performance than before, in simulating the mixed layer, net community production, NDIC and pCO₂sea.

There are very limited data for column-integrated net community production in the western North Pacific. Although the modeled net community production is relatively higher than the net community production by Imai et al. (2002), it is not relatively high when comparing other observational data by Shiimoto et al. (1998) and Shiimoto (2000) in the adjacent oceanic region. In the revised manuscript, the authors have added the data by Shiimoto et al. (1998) and Shiimoto (2000) to the figure for reference, which shows the net community production by total phytoplankton in summer is as high as in spring.

The authors have estimated in situ pCO₂air at Station KNOT from Tsurushima et al. (2002). The data has been shown in Figure 2e in the revised manuscript for reference. The data are very similar to those at Mauna Loa (Keeling et al., 1982; Conway et al., 1994), and the authors have kept using the pCO₂air data for calculation in this study. Please see below and an attached figure for the similarity in the pCO₂air in detail.

Responses to moderate comments

The authors have replaced Tsurushima (2002)s raw data in the previous manuscript with the data in Tsurushima et al. (2002). As a result, the number of data has been consistent for NDIC and pCO₂sea. The authors have also removed the data which are considered to be influenced by the subtropical water, as described above.

The authors have mixed up the meaning of [compartment] with that of [state variable] in the previous manuscript. To avoid confusions, the authors have clarified this by describing [16-compartment (16-state variable) marine ecosystem model] in Section 2 and Figure 1 caption in the revised manuscript.

The works of Bates et al. (1998a) and Wanninkhof et al. (2007) have been discussed in Section 3. Interestingly, the conclusion from the two works is quite opposite with

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



regards to the role of storms in interannual CO₂ flux, and our conclusion is closer to that of Bates et al. (1998a). The authors have also mentioned this as follows: [Our model results also suggest that storm events could potentially have a large effect on interannual air-sea CO₂ flux variability globally, which is consistent with Bates et al. (1998a). However, Wanninkhof et al. (2007) induced an opposite conclusion of minimal influence of storms on annual CO₂ flux. Therefore, long-term monitoring of the air-sea CO₂ flux is required in various oceanic regions to assess quantitatively the role of storm events in the interannual air-sea CO₂ flux.]

The authors have added ranges of the wind speeds and highest wind speeds during the storm events to the beginning of Section 3, as follows. [Most of storm events last for no more than one day, and there are no storm events that last for more than five days (Table 2). All the storms have the wind speed of more than 10 (m s⁻¹) with the highest wind speed of 26.3 (m s⁻¹) for the 19 years. However, the storm events which has the wind speed of more than 25 (m s⁻¹) are very limited (Table 3), which is different from situations in the subtropical regions in which typhoons and hurricanes have higher wind speeds more than 30 (m s⁻¹) (e.g. Bates et al., 1998 a, b; DAsaro and McNeil, 2007; Wanninkhof et al., 2007).] To provide supplemental or more detailed information, Tables 2 and 3 have also been added to the revised manuscript. Following the reviewers useful comments, the authors have reviewed the suggested three works of Ho et al. (2006), McNeil and DAsaro (2007), and Fongohr and Woolf (2007). Considering the range of wind speed in our study, Ho et al. (2006) can be used for the wind with <20m s⁻¹ and McNeil and DAsaro (2007) can be applied for the wind with >20m s⁻¹. However, temporal resolution and the wind used for simulation in our study is an hour, it is eligible for the modeling study to use continuous wind speed/gas exchange parameterization in calculating the air-sea CO₂ flux with the wind speed of 26.3 m s⁻¹ at the highest. Considering that the gas transfer velocity is similar between Wanninkhof et al. (1992) and Ho et al. (2007) for the range of wind speed in our study (different by not more than 5-10 per cent), the authors have decided to keep using Wanninkhof et al (1992)s parameterization in this study. Modeling studies with various wind speed/gas

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

exchange parameterizations, along with using various wind data (such as QuickSCAT winds), are our ongoing works (Fujii et al., in prep.).

Following the reviewers comment, the authors have compared $p\text{CO}_2^{\text{air}}$ at Mauna Loa, Hawaii ($19.54^\circ\text{N}, 155.58^\circ\text{W}$), Shemya Island, Alaska ($52.72^\circ\text{N}, 174.10^\circ\text{E}$), Sand Island, Midway ($28.21^\circ\text{N}, 177.38^\circ\text{W}$), and POCN30 in the Pacific Ocean ($30.00^\circ\text{N}, 135.00^\circ\text{W}$). The $p\text{CO}_2^{\text{air}}$ for the four stations have also been compared to the $p\text{CO}_2^{\text{air}}$ by Tsurushima et al. (2002). The comparison result shows that the $p\text{CO}_2^{\text{air}}$ is similar among the stations (see an attached figure). Considering the similarity, it seems okay to use the $p\text{CO}_2^{\text{air}}$ at Mauna Loa for this study, and therefore, the authors have continued to use that. Actually, we need $p\text{CO}_2^{\text{air}}$ data after 1970s to calculate the air-sea CO_2 flux, and Mauna Loa is the only station so far which can provide the long-term data since 1970s.

In the previous manuscript, the authors did not clarify to distinguish terms among [net primary production], [gross primary production] and [net community production]. In the revised manuscript, the authors have unified to the term [net community production], following Fujii et al. (2007).

As mentioned in the manuscript, the Reynolds weekly SST data were used to drive the model (in Section 2), and the modeled SST decreased by 0.8°C in the mid-June, 1994 (in Section 3). Therefore, the Reynolds SST data could allow to pick up the storm signal on temperature although the temporal resolution is not sufficient. Unfortunately, there do not exist in situ or remote-sensing data that validate the decrease in temperature at Station KNOT before mid-1990s.

The authors thank the reviewer for introducing a modeling study of Soetaert et al. (2001). Along with this paper, the authors found several modeling studies which have studied the effect of intraseasonal events on ambient nutrients and chlorophyll, and have added them as references to the revised manuscript (e.g. Kawamiya and Oschlies (2004), McCreary et al. (1996 and 2001), Large and Crawford (1995) and Ridderinkhof

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

(1992)).

The authors have mentioned about the discrepancy between this study and others (Takahashi et al., 2002 and Tsurushima et al., 2002) with regards to extent of the role of the subarctic western North Pacific as a source of CO₂ to the atmosphere, in Sections 3 and 4.

One of the authors used to engage in the VOS project with Dr. Nojiri, and actually used the data as references in this study. The authors have added words to the revised manuscript mentioning that the data were used as references in this study, as well as the URL. Thank you for the comment.

Responses to minor comments

The term [TCO₂] has been replaced with [DIC] in the entire manuscript.

Page 66 Line 13 in the previous manuscript: [continuous observations] → [continuous observations of the wind and CO₂]

Page 66 Line 13 in the previous manuscript: [solar radiation] → [incoming solar radiation]

Page 67 Line 3 in the previous manuscript: [In most such cases] → [In most cases]

Page 67 Line 10 in the previous manuscript: [entrainment] → [horizontal advection]

Page 67 Line 20 in the previous manuscript: Sabine et al. (2004) have been referred in the revised manuscript instead of Tsurushima et al. (2002).

Page 68 Line 14 in the previous manuscript: [shells] → [liths]. Thank you for the useful comment.

Page 68 Line 23 in the previous manuscript: [at the sea surface] → [at 10m height]

Page 69 Line 1 in the previous manuscript: The reviewer is correct in mentioning that pCO₂air is different from xCO₂air. The atmospheric CO₂ data are provided as a form

BGD

5, S256–S267, 2008

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



of xCO_{2air} , and therefore, the xCO_{2air} should be converted to pCO_{2air} by the following formula:

$$pCO_{2air} = (P - pH_2O) \times xCO_{2air} \quad (1)$$

where P is the ambient sea level pressure and pH_2O is the water vapor pressure at 100 per cent humidity calculated from SST and salinity according to Weiss and Price (1980). The authors have checked possible ranges of P and pH_2O at Station KNOT and have also referred to the VOS data which are mentioned above. The P can vary from ca. 0.94 to ca. 1.00, and the pH_2O can change from ca. 0.006 to 0.014. Therefore, the pCO_{2air} can be lower than xCO_{2air} by up to 6 per cent. In the previous manuscript, the pCO_{2air} was not distinguished from the xCO_{2air} . In the revised manuscript, the authors have decided to continue to use the xCO_{2air} for calculating air-sea CO_2 flux. The reason is as follows: Although the sea level pressure with higher frequency than monthly can be obtained from the NCEP objectively analyzed data, the corresponding continuous atmospheric CO_2 data with higher frequency than monthly cannot be obtained. Generally, the sea level pressure is low in late winter when the pCO_{2sea} is higher than the pCO_{2air} in the subarctic western North Pacific. If the pCO_{2air} is used instead of xCO_{2air} , ΔpCO_2 (pCO_{2sea} minus pCO_{2air}) gets larger, and therefore, the sea-to-air CO_2 efflux should be higher as well. This is important and presumably serves as a spur to our claim that previous studies overestimated role of the subarctic western North Pacific as a sink of CO_2 . This argument has been added to Sections 3 and 4 in the revised manuscript. Thank you for the important comment.

Page 69 Line 18 in the previous manuscript: [well reproduces] → [reproduces well]

Page 69 Line 23 in the previous manuscript: [pss] → [psu]

Page 70 Lines 6-9 in the previous manuscript: As the reviewer points out, the phrase [pCO₂sea is the most sensitive biogeochemical parameter to the storms] cannot be justified in this study, and the authors have removed the phrase [that pCO₂sea is the

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



most sensitive biogeochemical parameter to the storms, and therefore,] in the revised manuscript.

Page 70 Line 23 in the previous manuscript: [slihjtly] → [slightly]

Page 71 Line 14 in the previous manuscript: flux values have been rounded (244.55 → 245, 160.60 → 161)

Page 71 Line 15 in the previous manuscript: In the revised manuscript, the authors have referred to two previous studies (Takahashi et al., 2002; Tsurushima et al., 2002).

Page 71 Line 18 in the previous manuscript: [taking up oceanic CO₂] → [taking up atmospheric CO₂]

Page 71 Lines 20 and 27 in the previous manuscript: The reviewers comments [The enhancement of primary production in summer is less than 10 per cent, this is negligible] and [The reduction of net community production in winter and spring is very low, this is even more negligible.] are right, but the model result is for 19-year mean (1982-2000). Therefore, in some years the enhancement is much higher than 10 per cent. The authors have emphasized that the model results shown in Figure 2 is for the 19-year mean in Table 2 and Figure 2 caption and in the main text in the revised manuscript.

Page 72 Line 7 in the previous manuscript: Even considering the authors response above, this seems an over-statement, so the authors have removed a word [significantly] from the sentence in the revised manuscript.

Page 72 Lines 8-9 in the previous manuscript: The reviewer is right in mentioning [the irradiance was not modeled] and [the results of nutrient simulations were not shown] in this study. The authors have removed the terms [the irradiance] and [nutrient concentrations] from the revised manuscript. These modifications do not affect purport of this study.

Figure 1 caption: [schematic view] → [conceptual diagram]

S262

BGD

5, S256–S267, 2008

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Figure 1: [reminerzalization] → [dissolution] for the arrow between [CaCO₃] and [Ca²⁺]

Figure 1: [Ca] → [Ca²⁺]

Figure 1: [shell formation] → [calcification] for the arrow between [PS] and [Ca²⁺] (and also that between [ZS] and [Ca²⁺])

Figure 1: [shell formation] → [frustule formation] for the arrow between [Si(OH)₄] and [PL]

Figure 1: Both flows from PS and PL to NO₃ were set to zero in this study, following Fujii et al. (2007). Modeling diatom sinking is an ongoing study (e.g. Yoshie et al., in prep.). The process seems important especially at the end of the diatom bloom, and this is important because the process accelerates termination of the diatom bloom. The process is partly represented by parameterizing the diatom mortality proportional to the square of diatom abundance.

Figure 2: Following the reviewers comment, Figure 2f has been modified to have two segments.

Figure 3 plot b): The atmospheric CO₂ (pCO₂air) has the seasonality, as the reviewer points out. The pCO₂air in the previous manuscript is not correct and has been revised in the revised manuscript. The seasonality was taken into account in calculating the air-sea CO₂ flux in the previous manuscript, and therefore, the model results do not change.

Figure 3 plot c): The term [efflux] has been replaced with [flux] in Figure 3(c) and the caption, and the main text which explains this figure. The term [positive upward] has been removed from Table 2 caption because it is obvious as long as one uses the term [efflux].

Other revisions

Table 1 in the previous manuscript included some errors in number, and therefore, the

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

numbers have been revised in the revised manuscript. The corresponding statements in the main text have also been revised as well.

Tables 2 and 3 have been added to the revised manuscript. Most of storm events last for no more than one day, and there are no storm events that last for more than five days (Table 2). All the storms have the wind speed of more than 10 (m s⁻¹). However, the storm events which has the wind speed of more than 25 (m s⁻¹) are very limited (Table 3), which is different from situations in the subtropical regions in which typhoons and hurricanes have higher wind speeds more than 30 (m s⁻¹).

Along with references introduced by the reviewers, the authors have referred the following previous studies to the revised manuscript.

Bates, N. R.: Interannual variability in the global uptake of CO₂, *Geophys. Res. Lett.*, 29(5), 10.1029/2001GL013571, 2002.

Chen, C. T., Liu, C., Chuang, W. et al.: Enhanced buoyancy and hence upwelling of subsurface Kuroshio waters after a typhoon in the southern East China Sea, *J. Marine Syst.*, 43, 65-79, 2003.

Cione, J., Black, P., and Houston, S.: Surface observations in the hurricane environment, *Mon. Weather Rev.*, 128, 1550-1561, 2000.

Conway, T. J., Masarie, K. A., and Zhang, Ni: Evidence for interannual variability of the carbon cycle from the National Oceanic and Atmospheric Administration/Climate Monitoring and Diagnostics Laboratory Global Air Sampling Network, *J Geophys. Res.*, 99(D11), 22831-22855, 1994.

DAsaro, E. D., and McNeil, C.: Air-sea gas exchange at extreme wind speeds measured by autonomous oceanographic floats, *J. Mar. Syst.*, 66, 92-109, 2007.

Davis, A., and Yan, X.-H.: Hurricane forcing on chlorophyll-a concentration off the northeast coast of the U.S., *Geophys. Res. Lett.*, 31, L17304, doi:10.1029/2004GL020668, 2004.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Fogel, M. L., Aguilar, C., Cuhel, R., Hollander, D. J., Willey, J. D., and Paerl, H. W., Biological and isotopic changes in coastal waters induced by Hurricane Gordon, *Limnol. Oceanogr.*, 44(6), 1359-1369, 1999.

Iverson, R. I.: Mesoscale oceanic phytoplankton patchiness caused by hurricane effects on nutrient distribution in the Gulf of Mexico, In Andersen, N. R., and Zahuranec, B. J. (eds.), *Oceanic Sound Scattering Prediction*, Plenum Press, New York, 767-776, 1977.

Kataoka, K.: Tracks of tropical cyclones in the mid-latitudes of the western North Pacific, *Tenki*, 50(9), 705-714, 2003 (in Japanese with English abstract).

Kawamiya, M., and Oschlies, A.: Simulated impact of intraseasonal variations in surface heat and momentum fluxes on the pelagic ecosystem of the Arabian Sea, *J. Geophys. Res. Oceans*, 109, C03016, doi:10.1029/2003JC002107, 2004.

Kishi, M. J., Kashiwai, M., Ware, D. M., Megrey, B. A., Eslinger, D. L., Werner, F. E. et al.: NEMURO: a lower trophic level model for the North Pacific marine ecosystem, *Ecol. Modell.*, 202, 12-25, 2007.

Large, W. G., and Crawford, G. B.: Observation and simulations of upper-ocean response to wind events during the ocean storms experiment, *J. Phys. Oceanogr.*, 25, 2831-3852.

Lee, D., and Niller, P: Ocean responses to typhoon Rusa in the south sea of Korea and in the East China Sea, *J. Korean Soc. Oceanogr.*, 38(2), 60-67, 2003.

Lin, I., Liu, W. T., Wu, C.-C. et al.: New evidence as survival alternatives in an unstable environment, *Oceanol. Acta*, 1, 493-509, 1978.

McCreary, J. P., Kohler, K. E., Hood, and Olson, D. B.: A four-compartment ecosystem model of biological activity in the Arabian Sea, *Prog. Oceanogr.*, 37, 193-240, 1996.

McCreary, J. P., Kohler, K. E., Hood, R. R., Smith, S., Kindle, J., Fischer, A. S., and Waller, R. A.: Influences of diurnal and intraseasonal forcing on mixed-layer and bio-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

- logical variability in the central Arabian Sea, *J. Geophys. Res.*, 106, 7139-7155, 2001.
- Platt, T., Bouman, H., Devred, E., Fuentes-Yaco, C., and Sathyendranath, S.: Physical forcing and phytoplankton distributions, *Sci. Mar.*, 69(1), 55-73, 2005.
- Ridderinkhof, H.: On the effects of variability in meteorological forcing on the vertical structure of a stratified water column, *Cont. Shelf Res.*, 12, 25-36, 1992.
- Rodriguez, J., Tintore, J., Allen, J. T. et al.: Mesoscale vertical motion and the size structure of phytoplankton in the ocean, *Nature*, 410, 360-363, 2001.
- Roebber, P. J.: Statistical analysis and updated climatology of explosive cyclones, *Mon. Wea. Rev.*, 112, 1577-1589.
- Sanford, T. B., Black, P. G., Haustein, J. R. et al.: Ocean response to a hurricane, Part I: observations, *J. Phys. Oceanogr.*, 17, 2065-2083, 1987.
- Senjyu, T., and Wanatabe, T.: A sudden temperature decrease along the Sanin coast induced by a typhoon, *UmitoSora*, 75, 1-8, 1999 (in Japanese with English abstract).
- Shiomoto, A.: Chlorophyll-a and net community production during spring in the oceanic region of the Oyashio Water, the north-western Pacific, *J. Mar. Biol. Ass. U.K.*, 80, 343-354, 2000.
- Shiomoto, A., Ishida, Y., Tamaki, M., and Yamanaka Y.: Net community production and chlorophyll a in the western Pacific Ocean in summer, *J. Geophys. Res.*, 103(C11), 24651-24661.
- Soetaert, K., Herman, P. M. J., Middelburg, J. J., Heip, C., Smith, C. L., Tett, P., and Wild-Allen, K.: Numerical modeling of the shelf break ecosystem: reproducing benthic and pelagic measurements, *Deep-Sea Res. Part II*, 48, 3141-3177, 2001.
- Son, S., Platt, T., Bouman, H., Lee, D., and Sathyendranath, S.: Satellite observation of chlorophyll and nutrients increase induced by Typhoon Megi in the Japan/East Sea, *Geophys. Res. Lett.*, 33, L05607, doi:10.1029/2005GL025065, 2006.

BGD

5, S256–S267, 2008

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Son, S., Platt, T., Fuentes-Yaco, C., Bouman, H., Devred, E., Wu, Y., and Sathyendranath, S.: Possible biogeochemical response to the passage of Hurricane Fabian observed by satellites, *J. Plankt. Res.*, 29(8), 687-697, 2007.

Stramma, L., Cornillon, P., and Price, J. F.: Satellite observations of sea surface cooling by Hurricanes, *J. Geophys. Res.*, 91, 5031-5035, 1986.

Subrahmanyam, B., Rao, K. H., Rao, N. S. et al.: Influence of a tropical cyclone on chlorophyll-a concentration in the Arabian Sea, *Geophys. Res. Lett.*, 29, 2065, doi:10.1029/2002GL015892, 2002.

Wanninkhof, R., Olsen, A., and Trinanes, J.: Air-sea CO₂ fluxes in the Caribbean Sea from 2002-2004, *J. Mar. Sys.*, 66, 272-284, 2007.

Wu, Y., Platt, T., Tang, C. et al.: Short-term changes in chlorophyll distribution in response to a moving storm: a modeling study, *Mar. Ecol. Prog. Ser.*, 335, 57-68, 2007.

Yoshida, A., and Asuma, Y.: Structures and environment of explosively developing extratropical cyclones in the northwestern Pacific region, *Mon. Wea. Rev.*, 132, 1121-1142, 2004.

[Interactive comment on Biogeosciences Discuss.](#), 5, 65, 2008.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)