

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

Aerosol fluxes and particle growth above managed grassland

**E. Nemitz¹, J. R. Dorsey², M. J. Flynn², M. W. Gallagher², A. Hensen³,
J.-W. Erisman³, S. M. Owen^{1,4}, U. Dämmgen⁵, and M. A. Sutton¹**

¹Centre for Ecology and Hydrology (Edinburgh), Bush Estate, Penicuik, Midlothian, EH26 0QB, UK

²School for Earth, Atmospheric and Environmental Sciences, University of Manchester, P.O. Box 88, Manchester, M60 1QD, UK

³Netherlands Energy Foundation (ECN), 1755-ZG Petten, The Netherlands

⁴Institute for Environmental and Natural Sciences, University of Lancaster, UK

⁵Institute for Agroecology, Federal Agricultural Research Centre, Braunschweig, Germany

Received: 13 October 2008 – Accepted: 3 November 2008 – Published: 8 January 2009

Correspondence to: E. Nemitz (en@ceh.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Particle deposition velocities (11–3000 nm diameter) measured above grassland by eddy covariance during the EU GRAMINAE experiment in June 2000 averaged 0.24 and 0.03 mm s⁻¹ to long (0.75 m) and short (0.07 m) grass, respectively. After fertilisation with 108 kg N ha⁻¹ as calcium ammonium nitrate, sustained apparent upward fluxes of particles were observed. Analysis of concentrations and fluxes of potential precursor gases, including NH₃, HNO₃, HCl and selected VOCs, shows that condensation of HNO₃ and NH₃ on the surface of existing particles is responsible for this effect. A novel approach is developed to derive particle growth rates at the field scale, from a combination of measurements of vertical fluxes and particle size-distributions. For the first 9 days after fertilization, growth rates of 11 nm particles of 3.5 nm hr⁻¹ and 0.89 nm hr⁻¹ were derived for day and night-time conditions, respectively. This implies total NH₄NO₃ production rates of 1.1 and 0.44 μg m⁻³ h⁻¹, respectively. The effect translates into a small error in measured ammonia fluxes (0.06% day, 0.56% night) and a larger error in NH₄⁺ and NO₃⁻ aerosol fluxes of 3.6% and 10%, respectively. By converting rapidly exchanged NH₃ and HNO₃ into slowly depositing NH₄NO₃, the reaction modifies the total N budget, though this effect is small (<1% for the 10 days following fertilization), as NH₃ emission dominates the net flux. It is estimated that 3.8% of the fertilizer N was volatilised as NH₃, of which 0.05% re-condensed to form NH₄NO₃ particles within the lowest 2 m of the surface layer. This surface induced process would at least scale up to a global NH₄NO₃ formation of ca. 0.21 kt N yr⁻¹ from NH₄NO₃ fertilisers and potentially 45 kt N yr⁻¹ from NH₃ emissions in general.

1 Introduction

Aerosols make an important contribution to the Earth's radiation budget (IPCC, 2007). An important constituent of aerosols in this respect are hygroscopic compounds formed from anthropogenic precursors, such as ammonium sulphates ((NH₄)₂SO₄, NH₄HSO₄),

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ammonium nitrate (NH_4NO_3) and ammonium chloride (NH_4Cl), which are formed by the neutralization of atmospheric ammonia (NH_3) by sulphuric acid (H_2SO_4), nitric acid (HNO_3) and hydrochloric acid (HCl), respectively. Adams et al. (2001) estimated present-day forcing potentials of sulphates and nitrates to be of the order of -0.95 and -0.19 W m^{-2} , respectively. In Europe sulphur (S) emissions have been reduced dramatically, by 61% over the period 1990 to 1998 (Vestreng and Støren, 2000), with the UK figure being 57% (NEGAP, 2001) and the German figure 84% (Umweltbundesamt, 2001). Similar trends are expected for other industrialised countries.

However, as with Germany, a decrease in sulphur emissions is not always mirrored by a decline in aerosol sulphate concentrations, which have been limited by NH_3 emissions (Umweltbundesamt, 2001). European emissions of oxidized nitrogen (NO_x) have declined very little and global NO_x emissions are expected to increase. As a net result, for 2100 Adams et al. (2001) projected global sulphate forcing to decline to -0.85 W m^{-2} , while an increasing amount of the NH_3 will be neutralized by NO_x , increasing the nitrate forcing to -1.28 W m^{-2} . This prognosis is further supported by measurements that indicate that nitrates already dominate the light-scattering in the Netherlands, which are characterized by continental nitrate rich air masses and large NH_3 emissions from high livestock densities (ten Brink et al., 1996).

In addition to the climate effect, the incorporation of relatively fast depositing reactive gaseous pollutants into small particles, that are deposited relatively inefficiently, increases the atmospheric lifetime and thus the transport distance of these acidifying and eutrophying pollutants. This obviously has important consequences for both local and national budgets but and trans-boundary air pollutant transport (Sutton et al., 2007).

A third important effect of changes in the gas/aerosol partitioning concerns the applicability of methods to measure surface/atmosphere exchange fluxes. Micrometeorological flux measurement techniques are formulated for chemically inert tracers and have to be modified if a compound experiences sources or sinks within the atmosphere (e.g. Kramm et al., 1991). Under these conditions, the flux of a reactive tracer may

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



change considerably with height and the vertical transport derived for a measurement height deviate from the true exchange at the surface. Unexpected observations have been made in a number of studies. Since NH_3 exchange with vegetation is potentially bi-directional (e.g. Sutton et al., 1995, 2007), its a priori estimation is often uncertain.

5 HNO_3 and HCl , however, are expected to deposit at the maximum deposition velocity permitted by turbulence (V_{max}), while particles deposit slowly to vegetation, at rates of typically <1 up to 5 mm s^{-1} . Departures from these rates have been observed and attributed to gas-to-particle conversion or the evaporation of NH_4^+ aerosols (Huebert et al., 1988; Nemitz et al., 1996, 2002, 2004; van Oss et al., 1998).

10 The potential for formation of NH_4NO_3 and NH_4Cl is most likely to be found near strong sources of either NH_3 or the acids. Ammonia originates particularly from agricultural point sources such as cattle, pig and poultry units. A large fraction also originates from diffuse surface sources, such as the spreading of manure, organic or mineral fertilizers. These may either cause direct evaporation or raise the nutrient status of the
15 crop or grassland resulting in a raised emission potential, or stomatal compensation point (e.g. Farquhar et al., 1980).

In May/June 2000 a major European integrated experiment, with 12 participating institutes, was organized to investigate NH_3 exchange processes with managed agricultural grassland as part of the EU GRAMINAE project. The strategy of GRAMINAE has been described elsewhere, together with the set-up and overall results of this particular experiment (Sutton et al., 2001, 2008a, b). During the campaign, the experimental
20 grassland was cut, fertilised with 108 kg N ha^{-1} as calcium ammonium nitrate and the course of grass regrowth followed.

As part of this experiment, particle number flux measurements were made by eddy-covariance above long grassland (before the cut), above short grassland (between cut and fertilization) and above the fertilized grassland, to study the behaviour of the aerosol and its interaction with the ammonia exchange. In addition to the observation of slow mean aerosol deposition fluxes to the unfertilized grassland, consistent diurnal patterns of apparent emissions were observed during the first 10 days following fertil-
25

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ization. The particle counter used for these measurements had a lower cut-off of 11 nm and an upper cut-off of effectively $3\text{ }\mu\text{m}$. The observation of upward fluxes of particles could either be caused by the formation of new particles, which grow quickly to a size of $>11\text{ nm}$, or by the condensation of vapours on the surface of existing particles, resulting in the growth of some particles across the 11 nm threshold.

In this paper we represent the particle flux measurements and interpret the results in the context of simultaneous measurements of (i) fluxes of NH_3 , HNO_3 , and selected biogenic volatile organic compounds (BVOCs); (ii) concentrations of gaseous SO_2 , HNO_3 and HCl , as well as aerosol NH_4^+ , NO_3^- , Cl^- and SO_4^{2-} ; (iii) particle size spectra and (iv) meteorological parameters. A novel technique is developed to derive aerosol growth rates at the field scale from a combination of the particle number fluxes and aerosol size distributions. The overall objectives of this paper are:

- (a) to quantify particle deposition rates to (unfertilised) grassland;
- (b) to identify the potential and mechanism of the particle production or growth following fertilisation;
- (c) to quantify particle growth rates;
- (d) to quantify the effect on the measurement of fluxes of NH_3 and total nitrogen;
- (e) to indicate the importance of surface ammonia emissions from fertilizer application for particle production globally.

2 Methods and theory

2.1 Site description and overview of measurements

The measurements were made above a 12 ha experimental agricultural grassland of the Agricultural Research Station Völkenrode near Braunschweig, Germany ($52^\circ 18' \text{N}$,

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



10°26' E; 79 m a.s.l.) during the period 21 May to 16 June 2000. The field was cut on 29 May (06:00–10:00 GMT) and fertilized on 5 June (06:00–07:00 GMT) with 108 kg N ha⁻¹ of calcium ammonium nitrate. The grass was mechanically turned in the morning of 30 May and lifted 31 May (06:00–10:00 GMT). Canopy height before the cut was ≈0.75 m and after the cut 0.07 m. The comprehensive set of instrumentation deployed has been summarized elsewhere (Sutton et al., 2008a; Nemitz et al., 2008a). We here focus on the description of those measurements related to the study of the gas/particle interaction.

Particle fluxes were made with an eddy-correlation system including a TSI condensation particle counter (TSI CPC 3760), with a sample flow rate of 1 l min⁻¹ and a size-range of 11 nm up to 3 μm, similar to the setups used by Dorsey et al. (2002) and Nemitz et al. (2002). A sonic anemometer (Model HS, Gill Instruments, UK) was mounted at a height of 2.02 m. A stainless steel inlet (1/8" OD) led from just below the anemometer to the condensation particle counter, housed in a box at the foot of the measurement mast. The pulse output of the CPC was recorded on a PC based logging system at 10 Hz, and fluxes were calculated according to standard eddy-covariance protocols (Dorsey et al., 2002). In addition, size-segregated particle fluxes over the range 0.1 to 0.4 μm were measured at two different heights using optical particle counters, as presented elsewhere (Dorsey, 2002; Gallagher et al., 2002). In addition, ammonia fluxes were measured by a total of four automated and continuous gradient systems (Milford et al., 2008) and four relaxed eddy accumulation systems (Hensen et al., 2008). Concentrations and gradients of gaseous HNO₃, HCl, SO₂ and aerosol NH₄⁺, NO₃⁻, Cl⁻ and SO₄²⁻ were measured above the grassland with a continuous sampling system combining wet annular denuders and steam jet aerosol collectors with online conductivity analysis for NH₄⁺ and anion chromatography. The system (SJAC/WAD) was a forerunner of the GRAEGOR instrument described by Thomas et al. (2008), and differed from the GRAEGOR in that it used peristaltic pumps rather than syringe pumps to control the liquid flows. In brief, the system consists of two sampling boxes (SB, at 0.7 m and 2.0 m) and a common detector box (DB). In each SB the gas phase is stripped from

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a 15 lpm air sample through a continuously flushed wet annual rotating denuder, while the water soluble aerosol components are collected by passing the sample through steam and collecting the solution in a cyclone. The four channels (gas phase and aerosol phase at two heights) are alternately analysed online in the DB for anions by ion chromatography and for NH_4^+ with a selective ion membrane/conductivity detector. Continuous addition of an internal Br^- standard ensures high precision between measurements. Fluxes were calculated according to the aerodynamic gradient technique as described by Milford et al. (2008) and Thomas et al. (2008). The instrument was operated without a size-segregating cut-off inlet.

In addition, a combination of a tandem differential mobility analyzer (DMPS; Williams, 1999) and an Aerodynamic Particle Sizer (APS 3320, TSI, Minnesota, USA) was deployed to measure the particle size distribution over the range 3 nm to $20\text{ }\mu\text{m}$ at a 10-min resolution. Additional information on the size distribution of the major ions was provided by MOUDI runs (Micro orifice uniform deposit impactor; MSP Corp., Minnesota, USA) runs, integrated over typically 3 days. In order to investigate the effect of the chemical interactions and deposition on concentrations of NH_3 and acid gases, the background concentration of NH_3 , HNO_3 and HCl were measured with an automated rotating wet batch denuder (RBD), providing 40-min batch samples in every hour (Keuken et al., 1988). This denuder was placed on the roof of a disused, 47 m tall building of the Physikalisch Technische Bundesanstalt (PTB), about 1 km ENE of the field site (see map in Sutton et al., 2008a). Daily concentrations of gases (NH_3 , HNO_3 , HCl , HNO_2 and SO_2) and aerosol components (NH_4^+ , NO_3^- , Cl^-) were also measured 250 m north of the grassland site, above wheat, using a KAPS (Kananaskis Air Pollutant Sampler) combination of 3-stage annular denuders and filter pack (PTFE, nylon) (e.g. Dämmgen, 2007). Throughflow was adjusted so that the cut-off point of KAPS is approx. $3\text{ }\mu\text{m}$, and 24-h samples were taken between 0 and 24 h CET. Details of the experimental set-up, the sampling properties and quality parameters are described in Zimmerling et al. (1997). These samplers had shown to be reliable even under extreme weather conditions and yielded good results in sampler intercomparisons (e.g. Wiebe

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 1990). Although the sampling height of 1.2 m was insufficient for concentrations to be fully unaffected by the atmosphere/surface exchange, these concentrations were not affected by the fertiliser application at the grassland site. Analysis was by liquid ion chromatography (IC) at FAL, Braunschweig for the KAPS samplers and at CEH Lancaster for the samples of MOUDI and RBD.

2.2 Concentration and flux measurements of selected biogenic volatile organic compounds (BVOCs)

The rationale behind the BVOC flux measurements was twofold: firstly, BVOCs potentially provide an alternative biogenically controlled precursor gas for particle formation and, secondly, high emissions of leaf alcohols from grass cutting have been reported in the literature (Kirstine et al., 1998; Karl et al., 2001). VOCs were collected onto Tenax TA (200 mg) and Carbotrap (100 mg) packed in steel tubes (6.1 mm OD, 90 mm length, Perkin Elmer). In the relaxed eddy accumulation (REA) approach (e.g. Desjardins, 1977), fast switching valves are used to sample air from up- and down-moving eddies conditionally onto two separate collection tubes. The valves were controlled by the switching logic of the CEH/UMIST REA system for atmospheric NH_3 (Nemitz et al., 2001; Hensen et al., 2008), logging an ultrasonic anemometer (Solent R1012; Gill Instruments, Lymington, UK) at a rate of 20.83 Hz. The sample tubes were tied directly onto the cage of the anemometer, operated at a height of 2.1 m, to minimize flux losses, and mass flow controllers ensured a flow rate of 0.5 l min^{-1} , resulting in a total sample volume of approximately 15 l per channel. Field blanks were taken by exposing a third tube at the same height, without flow. After sampling, tubes were sealed with Swagelok[®] end caps and Teflon ferrules and stored at 4°C, under which conditions the analytes are stable for at least one week.

For gas chromatography – flame ionization detection (GC-FID) analysis, tubes were desorbed using a Perkin-Elmer ATD-400 thermal desorption unit connected to a Hewlett Packard (HP) 5860 series II GC-FID. Separation of the compounds was achieved

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



through the use of an Ultra-2 cross linked 5% phenyl methyl silicone column. The detection limit for the monoterpenes is about 0.05 ng for the column, corresponding to 40 ppt in air for a 1 l sample, with a level of analytical precision of better than +5%.

For gas chromatography – mass spectrometry (GC-MS) analysis, a Perkin-Elmer ATD-400 thermal desorption unit is connected via a heated transfer line held at 200°C to a HP 5890 GC and a HP 5970 mass selective detector. Desorption, prefocussing and separation of the analytes is achieved using the same column and conditions as with the FID system. This system allows positive identification of a wide range of VOC on the basis of peak retention times, mass spectra contained in the library of the National Bureau of Standards and generated from authentic standards. Monoterpenes were quantified by preparing liquid standards by dilution of the pure compounds in methanol. Diluted standards (1–5 µl) were directly injected onto sampling tubes and the methanol removed by passage of ~0.3 l He. Chromatographic data were handled with a VG Minichrom or HP system.

2.3 Chemical reactions, equilibrium constants and time scales

Ammonia reacts reversibly with nitric acid and hydrochloric acid to form the associated ammonium salts:



The dissociation constants ($K_{\text{NH}_4\text{NO}_3}$ and $K_{\text{NH}_4\text{Cl}}$) describe the equilibrium vapour concentration products (K_e), which are expected to form if enough time is provided for aerosol and gas phase to equilibrate. Mozurkewich (1993) and Pio and Harrison (1987) provided expressions for K_e as a function of temperature (T) and relative humidity (h) for pure NH_4NO_3 and NH_4Cl , respectively. The vapour concentration in equilibrium with mixed aqueous solutions, including deliquescent aerosols, is lowered compared with pure solutions, especially by the presence of sulphates (e.g. Stelson and Seinfeld,

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1982), while for ultrafine particles partial pressures may be elevated by the Kelvin effect. The effective vapour concentrations in equilibrium with an externally mixed aerosol consisting of different types and sizes is impossible to predict exactly. K_e is a good approximation for humidities below the relative humidity of deliquescence (about 80% for $(\text{NH}_4)_2\text{SO}_4$ and 60% for NH_4NO_3). Whether equilibrium is attained depends on the comparative time-scales of attainment of equilibrium (τ_c) and the processes perturbing equilibrium, including turbulent diffusive transport (τ_t). Conversely, the Damköhler ratio ($D_r = \tau_t / \tau_c$) is also a measure of the degree to which chemical conversions may affect surface/atmosphere exchange fluxes (Foken et al., 1995). Expressions for τ_c and τ_t can be found elsewhere (e.g. Wexler and Seinfeld, 1990, 1991), as summarized by Nemitz et al. (2000).

2.4 A novel method for the calculation of growth rates from particle flux measurements at the field scale

Particle growth implies that some particles will grow across the lower cut-off of the particle counter ($d_p = 11$ nm or $r = 5.5$ nm) as they deposit. Therefore the concentration of particles > 11 nm may increase towards the ground at the same time as the total number of particles decreases due to deposition, resulting in an apparent upward flux of particles > 11 nm.

The equation of mass conservation of a tracer (χ) is given by (e.g. Fowler and Duyzer, 1989):

$$\frac{\partial F_\chi}{\partial z} = -\frac{\partial \chi}{\partial t} + u \frac{\partial \chi}{\partial x} + w \frac{\partial \chi}{\partial z} + Q_\chi, \quad (3)$$

where F_χ is the vertical flux of the tracer, t is time, u and w are the wind components along the stream-wise and vertical co-ordinates (x and z), and Q_c represents in situ production/destruction of the tracer. The right-hand-side (r.h.s.) terms represent storage, horizontal and vertical advection as well as production/destruction, usually attributed to chemical effects of emissions. Assuming stationary and homogeneous

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



conditions, the first three terms vanish. If we consider particle number (N) as the tracer, the production term may be written as $Q_N = dN/dt$. Despite the similarity to the first r.h.s. term of Eq. (3), this represents production rather than changes in concentration due to storage. For particle number fluxes measured within a certain size bin (for the CPC in the radius bin $5.5 \text{ nm} < r < 1.5 \mu\text{m}$), flux divergence is related to effects that change the concentration within this size bin only:

$$\frac{dF_N(5.5 \text{ nm} < r < 1.5 \mu\text{m})}{dz} = \frac{dN(5.5 \text{ nm} < r < 1.5 \mu\text{m})}{dt} \quad (4)$$

These may either be direct emissions of particles in this size range, coagulation or growth/shrinkage across the lower or upper size cuts. We here assume that there are no emissions sources at the point of measurement and that coagulation is unimportant (see discussion below). Thus the change in $N(5.5 \text{ nm} < r < 1.5 \mu\text{m})$ is only affected by the growth into the size range over the lower size-cut and growth out of the size range across the upper size cut. After application of the chain rule for forming derivatives the vertical flux divergence can be expressed as:

$$\frac{dF_N}{dz} = \frac{dN}{dr} \Big|_{r=5.5 \text{ nm}} \times \frac{dr}{dt} \Big|_{r=5.5 \text{ nm}} \left(- \frac{dN}{dr} \Big|_{r=1.5 \mu\text{m}} \times \frac{dr}{dt} \Big|_{r=1.5 \mu\text{m}} \right) \quad (5)$$

Here F_N and N are particle number flux and number concentration in the radius (r) range 5.5 nm to $1.5 \mu\text{m}$, respectively. The (non-logarithmic) particle size distribution function (dN/dr) at $r=5.5 \text{ nm}$ and $r=1.5 \mu\text{m}$ can be measured directly with appropriate instrumentation. Owing to the small number concentration of super-micron particles, the second r.h.s. term, i.e. the growth across the $1.5 \mu\text{m}$ threshold can be neglected. The growth rate (dr/dt) can be expressed as (Kulmala et al., 2001):

$$\frac{dr}{dt} = \frac{m_v \beta_m D \chi_v}{r \rho_p} \quad (6)$$

where m_v and χ_v are the molecular mass and effective concentration of the condensable vapour, respectively, and ρ_p is the particle density, while β_m is the correction factor

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for mass flux in the transition regime (Fuchs and Sutugin, 1971):

$$\beta_m = \frac{K_n + 1}{0.377 K_n + 1 + 4/3 \alpha^{-1} K_n^2 + 4/3 \alpha^{-1} K_n} \quad (7)$$

Here K_n is the Knudsen number

$$K_n = \frac{\lambda}{r} \quad (8)$$

- 5 with λ being the mean molecular free path in air ($0.065 \mu\text{m}$). Research suggests that the sticking coefficient (α) for NH_3/HNO_3 is in the range 0.3 to 0.8 (Dassios and Pandis, 1999; Cruz et al., 2000; Rudolf et al., 2001); here $\alpha=0.5$ is used, which results in a value of $\beta_m=0.031$ for $r=5.5 \text{ nm}$ and $\beta_m=0.918$ for $r=1.5 \mu\text{m}$. D in Eq. (6) is the binary diffusion coefficient for vapour diffusion towards a particle (Reid et al., 1987):

$$10 \quad D = \frac{RT}{6\pi N_A \nu r} \quad (9)$$

- Here, R is the universal gas constant, T is temperature in K, N_A is the Avogadro constant and ν is the gas dynamic viscosity. A first estimate of the flux change with height (dF_N/dz) is derived from the difference between the measured flux and the flux expected in the absence of chemical reactions, parameterised using an average deposition velocity ($V_d = -F_N/\chi_N$; with χ_N the particle number concentration) of 0.14 mm s^{-1} derived from measurements before fertilization (cf. Sect. 3.1 below):

$$\frac{dF_{\text{Ni}}}{dz} = \frac{F_N + V_d \times \chi_{\text{Ni}}}{z_m} \quad (10)$$

- With Eqs. (5) and (10), the growth rate across the size cut-offs of the particle counter can be estimated from the measured fluxes and particle size spectra. By inverting Eq. (6), the effective vapour concentration available for condensation (χ_v) can be derived. The growth rate across the 11 nm threshold may be extrapolated to derive the

growth rate of all particle sizes, again using Eq. (6). The total mass transport from the gas to the particle phase (j_v) is then given by integration over all size classes (i) as:

$$j_v = \sum_{i=1}^n 4\pi r_i^2 N_i g \frac{\rho_p}{m_v} \frac{dr_i}{dt} \quad (11)$$

where the particle density (ρ_p) is set to that of NH_4NO_3 for consistency. Assuming the mass transport to be due to the combined condensation of NH_3 and HNO_3 , 21.25% of j_v (in terms of mass) is due to the condensation of NH_3 .

3 Results

3.1 Particle number flux measurements before and after fertilisation

The measured particle number unfiltered flux is shown in Fig. 1. Spikes are primarily due to emissions associated with the operations of agricultural machinery in the field (e.g. cutting on 29 May, turning of the cut grass 30 May and removal of the cut grass on 1 June). Otherwise, the measurement show small fluxes, dominated by deposition before the fertilisation, and a diurnal pattern of apparent emission fluxes after fertilisation, which slowly subsided.

The statistics of the concentrations and fluxes before and after fertilisation are summarised in Table 1, after management periods had been removed and data filtered for suitable wind directions. Mean air concentrations were similar during the four periods, ranging from 7190 to 8628 cm^{-3} . Both emission and deposition fluxes were measured during all four periods, but were least frequent (31%) before the cut and most frequent (60%) during the first week after fertilization. The average apparent emission flux after fertilization was larger than the deposition fluxes to the short vegetation. The average (median) deposition velocities were 0.24 (0.24) mm s^{-1} before the cut, 0.03 (0.06) mm s^{-1} between cut and fertilization and 0.06 (0.05) mm s^{-1} 7–10 days

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



after fertilization, qualitatively consistent with the known dependence of deposition velocities on surface roughness (e.g. Gallagher et al., 2002).

3.2 Atmospheric concentrations, equilibrium considerations and potential for NH_4NO_3 formation

5 The time-series of the concentration in air of NH_3 , HNO_3 and HCl at 1 m and 47 m are presented in Fig. 2, clearly showing the effect of fertilisation on the concentration of NH_3 . The average NH_3 concentration at the notional canopy height (z_0') can be estimated from the additional knowledge of the NH_3 flux (F_{NH_3} ; cf. Fig. 7 below – Sutton et al., 1993; Milford et al., 2008), under the simplifying assumption that the flux is
10 unaffected by the chemical conversion:

$$\chi_{\text{NH}_3}(z_0') = \chi_{\text{NH}_3}(1 \text{ m}) + F_{\text{NH}_3} [R_a(1 \text{ m}) + R_b]. \quad (12)$$

Here R_a and R_b are the aerodynamic sub-layer resistance describing the transport through the turbulent and the quasi-laminar part of the atmosphere, respectively. This concentration is also shown in Fig. 2 for comparison. All concentrations undergo a
15 clear diurnal cycle, with daytime maxima. Both acids are depleted over the first two days after fertiliser application on the morning of 5 June. However, this change is also reflected in the background gas measurements (Fig. 2; both by RBD and KAPS), indicating that this decrease is not due to depletion by reaction with NH_3 above the field, but reflects changes in air masses or regional chemistry.

20 $\chi_{\text{NH}_3}(z_0')$ exhibits the behaviour expected for a gas concentration that is governed by equilibrium with dry or aqueous NH_4^+ , with the diurnal pattern reflecting the temperature control and the long-term decrease in soil and plant NH_4^+ after fertilization (cf. Milford et al., 2001; Sutton et al., 2008b). On an equivalent basis, NH_4^+ aerosol concentrations exceeded the NO_3^- concentrations by a factor of two (Fig. 3), the rest of the NH_4^+
25 being neutralized by SO_4^{2-} (not shown). The measurements indicated that NO_3^- concentrations appeared to be increased at the grassland site, compared with the control site 400 m to the north. However, this observation was also made prior to fertilization

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and may at least be partially due to a larger cut-off diameter of the SJAC capturing additional coarse NO_3^- compared with the KAPS.

5 The comparison of measured (K_m) with theoretical equilibrium (K_e) vapour concentration products of NH_3 with the acids shows $K_m \ll K_e$ prior to fertilizer application (Fig. 4). Since the NH_3 concentration above the grassland increased on 5 June, measured products $[\text{NH}_3] \times [\text{HCl}]$ increased also, but were still much smaller than K_e . Measured products of $[\text{NH}_3] \times [\text{HNO}_3]$ were close to their theoretical equilibrium values during 5 to 11 June with exceedances during most nights.

3.3 Concentrations and fluxes of VOCs

10 Financial constraints meant that out of the 27 hourly VOC REA runs performed only 14 could be analyzed, and only for a limited suite of oxygenated compounds and terpenoids (benzaldehyde, nonanal, α -pinene, 1- β -pinene and d-3-carene). Grass cutting has been found to increase emissions of some BVOCs by a factor of 180 (Kirstine et al., 1998; Karl et al., 2001), but the leaf alcohols, the primary BVOCs released, are not very
15 efficient particle precursors until they are processed to form less volatile compounds. By contrast, biogenic alkenes have been found to be efficient aerosol precursors (e.g. Hoffmann et al., 1997). The suite of VOCs was selected to provide information on the magnitude of emissions allowing the estimation of their potential for particle formation/particle growth. Averaged concentrations and fluxes after the cut and before
20 fertilization are contrasted with the measurements after fertilization in Table 2. The flux measurements should be interpreted with caution. However, neither concentrations nor fluxes show significant changes before and after fertilization. Irrespective of fertilization, benzaldehyde showed a clear diurnal cycle with daytime maxima of $8 \mu\text{g m}^{-3}$ and night-time minima of $2 \mu\text{g m}^{-3}$ (not shown).

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.4 Calculation of particle growth rates

Growth rates of 11 nm, total gas-to-particle conversion mass transfer fluxes, estimated vapour concentration and the effect of vapour condensation on the NH_3 flux are presented in Fig. 5 and summarized in Table 3 as typical values for 12-h day and night-time periods following fertilizer application. They are compared with measurements of vapour concentrations and estimated total particle surface area.

3.5 Effects on the measurements of ammonia flux

Even if there is potential for the formation or evaporation of aerosol, the time-scale of this process may be too slow for surface exchange fluxes to be altered significantly. The chemical time-scale (τ_c) (Wexler and Seinfeld, 1990) depends strongly on the surface area of the aerosol, while the turbulent time-scale (τ_t) depends on the turbulence as characterised by the friction velocity (u_*). Chemistry is thought to alter fluxes for Damköhler numbers ($D_r = \tau_t / \tau_c$) greater than 0.1 (Foken et al., 1995). Figure 6 demonstrates that this critical ratio is exceeded regularly, but only during night, when turbulent diffusive time scales are long, and the aerosol concentration (and hence surface area) is enhanced.

According to this time-scale analysis, major effects on the NH_3 flux would be unlikely, at least during daytime. However, the observations presented here indicate that particle growth was considerable, also during daytime. From the growth rate of 11 nm particles, the driving force for the condensation, the effective vapour concentration (χ_v) was estimated using Eq. (6). And the total mass transfer from the gas to the aerosol phase ($J_{\text{NH}_4\text{NO}_3}$) was derived by integration over the measured particle spectrum, using Eq. (11). From this mass transfer the error on the NH_3 flux may be estimated. These parameters are also shown in Table 3, where they are also compared with the actual measured concentrations of NH_3 and HNO_3 , while Fig. 7 shows the time-series of the measured uncorrected and corrected NH_3 flux. The error on the NH_3 flux measurement is small, on average $12.2 \text{ ng m}^{-2} \text{ s}^{-1}$ (1.5%) during day and $12.4 \text{ ng m}^{-2} \text{ s}^{-1}$ (6.6%) during night.

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Discussion

4.1 Deposition rates to unfertilised grassland

Dry deposition rates of particles to vegetation have recently been reviewed by Pryor et al. (2008) and Petroff et al. (2008). At Braunschweig, particle number size distributions were dominated by particles in the size range 10 to 200 nm (Fig. 8) and thus the total particle number flux measurements were also dominated by the Aitken mode.

The mean V_d of 0.03 mm s^{-1} to cut ($h_c=0.07 \text{ m}$), 0.06 mm s^{-1} to partly regrown ($h_c=0.12 \text{ m}$) and 0.26 mm s^{-1} to tall ($h_c=0.75 \text{ m}$) grassland measured at the Braunschweig field site appears to be at the low end of measurements reported in the literature, but only few studies have measured total number fluxes over short vegetation: Nemitz et al. (2002) reported a mean deposition velocity of 0.3 mm s^{-1} to Scottish moorland vegetation ($h_c=0.06 \text{ m}$), measured with a similar setup; the difference is only partly explained by higher wind speeds during the UK measurements with an average u_* of 0.25 m s^{-1} compared with 0.18 m s^{-1} during the present study. Lamaud et al. (1994) measured average V_d of 0.7 mm s^{-1} to semi arid savannah, using an instrument that measured total particle flux, but weighted by surface area. Thus, their measurement is biased towards larger particle than the measurements made here.

Other studies above short vegetation have focused on accumulation mode particles ($0.1\text{--}0.5 \mu\text{m}$) that can be sized with optical techniques (Nemitz et al., 2002; Vong et al., 2004) or mass fluxes, also dominated by the accumulation mode (Wesely et al., 1985): Vong et al. (2004) derived particle deposition velocities to grassland ($0.75 \text{ m} < h_c < 1 \text{ m}$) as a function of particle size and stability. Values of V_d for the smallest size ($d_p=0.34 \mu\text{m}$) ranged from 2 during stable to 6 mm s^{-1} during unstable conditions, while values of V_d for larger particles ($d_p=0.84 \mu\text{m}$) increased from 2 to 7 mm s^{-1} . At the Scottish moorland site, V_d of accumulation mode particles ranged from 0.3 mm s^{-1} ($d_p=0.1 \mu\text{m}$), over 2 mm s^{-1} ($0.5 \mu\text{m}$) to $>10 \text{ mm s}^{-1}$ ($3 \mu\text{m}$). Simi-

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



larly, Wesely et al. (1985) derived an average V_d of 2.2 mm s^{-1} for SO_4^{2-} deposition to grassland, with a likely mass median diameter of $0.4 \mu\text{m}$.

Dorsey (2002) and Gallagher et al. (2002) also presented measurements of size-segregated particle fluxes made at the Braunschweig field site, deriving average V_d for particles with $0.1 \mu\text{m} < d_p < 0.2 \mu\text{m}$ of 0.87 mm s^{-1} for the cut and 1.13 mm s^{-1} for the long grass. It is thus likely that the relative low deposition rates of total particle flux is strongly biased by inefficient deposition of particles $< 0.1 \mu\text{m}$. While the deposition rates presented here are therefore representative for primary organic aerosol which represents much of the Aitken mode, they are probably underestimations for the average deposition of secondary aerosol constituents, which occupies the accumulation mode (e.g. Allan et al., 2003).

4.2 Assessment of alternative mechanism causing apparent particle emission fluxes following fertilisation

The analysis presented so far has implied that the particle emission fluxes were due to combined NH_3 and HNO_3 condensation. We here consider alternative mechanisms.

Biogenic VOC emissions of leaf alcohols and breakdown products, but also of effective precursor gases such as alkenes, have been reported to increase dramatically immediately after grass cutting (Kirstine et al., 1998; Karl et al., 2001). Although some particle emissions were observed at the GRAMINAE experiment after grass cutting, these can be related to agricultural activities on the field (e.g. turning), while persistent apparent particle emissions occurred only after fertilization. The limited number of measurements of BVOC concentrations and, though less certain, their fluxes do not indicate that fertilization had an influence on VOCs. Non-zero water vapour fluxes can also cause artifacts for size-dependent particle flux measurements (Kowalski, 2001), but this effect would also have been similar before and after fertilization.

The only major change induced by the fertilization was a greatly increased emission of NH_3 (Fig. 7), resulting in enhanced concentrations both at the canopy height (z_0)

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and the average measurement height of 1 m (Fig. 2), indicating that the change in NH_3 is almost certainly the cause for the observed apparent particle emissions. Having established the role of NH_3 as a precursor, the particle emission fluxes may still be linked to one of three processes: (i) ternary nucleation of the $\text{H}_2\text{O}-\text{H}_2\text{SO}_4-\text{NH}_3$ system, followed by growth across the 11 nm threshold, (ii) formation of new NH_4NO_3 or NH_4Cl particles, followed by growth across the 11 nm threshold and (iii) particle growth due to the combined condensation of NH_3 and HNO_3 or NH_3 and HCl on the surface of existing particles. It is highly improbable, however, that particle nucleation played a role in the observed apparent emission fluxes, for several reasons:

1. Binary nucleation of volatile compounds such as NH_4NO_3 and NH_4Cl is inefficient under ambient conditions (e.g. Korhonen et al., 1997).
2. Ternary nucleation of the $\text{H}_2\text{O}-\text{H}_2\text{SO}_4-\text{NH}_3$ system can be important, where H_2SO_4 concentrations are large and competing condensational sinks are small (due to low aerosol loading). However, only small NH_3 concentrations in the ng m^{-3} range, virtually always present in agricultural areas, are necessary to promote ternary nucleation of the $\text{H}_2\text{O}-\text{H}_2\text{SO}_4-\text{NH}_3$ system significantly, while the nucleation rate is insensitive to further increases in NH_3 concentrations beyond 100 ppt ($\sim 76 \text{ ng m}^{-3}$) (e.g. Napari et al., 2002). Therefore, the enhanced NH_3 emission after fertilization would not have greatly changed the efficiency of this particle production mechanism.
3. Although individual nucleation events were observed at the measurement site (Sutton et al., 2008a), there was no difference in the average size distribution of the DMPS measurements before and after fertilisation, and average size distributions did not show a clear nucleation mode (Fig. 8). This suggests that the observed nucleation events were regional in nature and independent of local NH_3 emissions.
4. To estimate whether growth of newly nucleated particles, at a diameter of typically 2 nm, would have been sufficiently quick to be detected by the CPC, Eq. (6) can be

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



used to simulate the growth of 2 nm particles at different vapour concentrations. The results (Fig. 9) demonstrate that it takes almost an hour for a 2 nm particle to become detectable, even at $\chi_v=50\text{ }\mu\text{g m}^{-3}$. For comparison, at typical wind speeds of 4 m s^{-1} and an extent of the grassland of between 150 and 500 m, depending on wind direction, the residence time above the field is of the order of 1 to 5 min.

In summary, it is highly unlikely that particle nucleation was involved in causing the observed upward fluxes. Instead, the observation of emission fluxes was almost certainly due to condensation particle growth of particles across the 11 nm cut-off size of the CPC.

4.3 Proposed mechanism leading to apparent particle emission fluxes

After fertilization, close agreement was found between measured concentration products of $[\text{NH}_3]\times[\text{HNO}_3]$ at 1 m and those predicted to be in equilibrium with NH_4NO_3 . For NH_4Cl $K_m\ll K_e$ was always observed, eliminating HCl and supporting HNO_3 as a potential associated precursor. According to the process understanding emerging from the Braunschweig experiment work, direct NH_3 evaporation of fertilizer pellets would have dominated the measured emissions (Fig. 7) directly after fertilizer application, until there was a significant amount of precipitation (Sutton et al., 2008b). Precipitation of 2.6 mm was recorded for the evening of 5 June and 1.3 mm during the late morning of the 7th. From 6 or 7 June NH_3 emission was probably dominated by the enhanced NH_4^+ concentration in soil solution and leaf litter, evident in the measurements, and the gradually increasing emission potential of the plants (Riedo et al., 2002; Sutton et al., 2008b). Small night-time emissions suggest that non-stomatal emissions, such as soil emissions, still contributed to the flux even 5 days after fertilization. Leaf-litter on the soil surface represents an additional NH_3 source which would have been re-captured in the tall canopy, but is able to escape the canopy after the cut (David et al., 2008; Sutton et al., 2008b).

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Because of their solubility in water, the acid gases HNO_3 and HCl are conventionally expected to deposit at the maximum rate permitted by turbulence (V_{max}). A reduction in the deposition velocity of acid gases indicates sources or non-zero equilibrium concentrations at or near the ground. Nemitz et al. (2004) showed that reduced HNO_3 deposition at a Dutch heathland was consistent with evaporation of airborne NH_4NO_3 due to small precursor gas concentration products near the ground. Nemitz et al. (2000) reported emissions of HCl and Neftel et al. (1996) reported upward gradients of HNO_3 , both of which they explained by leaf surface chemistry.

Although not without uncertainty, the HNO_3 gradient measurements at Braunschweig showed reduced deposition rates ($V_d < V_{\text{max}}$) during the afternoons of some of the days after fertilisation (e.g. 9 June) (Fig. 10). The reduced deposition rates of HNO_3 suggest that at least some of the HNO_3 is also surface-derived, with fertiliser pellets and high soil NO_3^- concentrations resulting in a non-zero gas-phase concentration at the ground, at least during warm afternoons. HNO_3 concentrations exhibited a clear diurnal cycle with the maximum concentrations during the day (Figs. 2 and 9), which would be consistent with either its production from NO_x , or the cycle in the equilibrium constant of NH_4NO_3 (Fig. 4).

In general, K_m and K_e change with height, in response to gradients of vapour concentrations, T and h : the increase of NH_3 and h towards the ground increase K_m relative to K_e , while the decrease in HNO_3 and increase of T would have the opposite effect. Unexpected particle fluxes have been attributed to the formation or evaporation of NH_4^+ before: van Oss et al. (1998) simulated over-rapid deposition of NO_3^- aerosol to forest to aerosol evaporation. Nemitz et al. (2004) reported fast deposition of NH_4^+ by gradient technique to a Dutch heathland, while eddy-correlation particle fluxes appeared to detect deposition of particles $>0.2 \mu\text{m}$ at the same time as emission of smaller particles. In a modelling study, Nemitz and Sutton (2004) demonstrated that both observations are consistent with the evaporation of NH_4NO_3 during the deposition process, caused by the depletion of NH_3 and HNO_3 close to the ground.

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In summary, the following picture emerges: a combination of NH_4^+ enriched soil, litter and leaf solutions combined with initial losses directly from the fertilizer pellets leads to emission of NH_3 which partly re-condenses on the surface of atmospheric aerosol particles above the ground, where, at least during the day, temperatures are lower.

5 This condensation of NH_3 is likely to be associated with simultaneous condensation of HNO_3 , which may be partly derived from the atmosphere or from fertilizer volatilization. It is likely that atmospheric HNO_3 was the main source during night, when the potential for HNO_3 emission from the ground was lower. This condensation results in the particles growing during the deposition process (Fig. 11), causing a deposition gradient in
10 the total particle number, but an apparent increase of particles with $r > 5.5$ nm, which is detected as an apparent particle emission flux.

4.4 Time-scales of particle growth

Particle nucleation and growth due to condensation are ubiquitous across the world, with the exception of the tropics, with typical growth rates in the region of $1\text{--}20$ nm hr^{-1} (Kulmala et al., 2004; Holmes, 2007) and can be attributed to a range of precursors.
15 For example, Verheggen et al. (2000) derived growth rates due to H_2SO_4 condensation of 3 to 7 nm hr^{-1} from size-distribution measurements. Growth rates of 2 to 11 nm hr^{-1} were observed following nucleation events during the BIOFOR campaigns above Finnish forests, attributed to the condensation of BVOCs, while faster growth rates of 15
20 to 180 nm hr^{-1} have been derived in the maritime environment during PARFORCE, possibly due to the uptake of biogenic methylated iodine species (Dal Maso et al., 2002). Growth rates linked to the emission of agricultural precursors do not appear to have been presented in the literature. A likely reason is that, until now, most growth rates have been derived by analysis of the time-evolution of size-distributions, typically following nucleation events or emission plumes – an approach, which requires
25 these processes to occur at the regional scale. By contrast, agricultural activity such as fertilisation happens at the field scale. Here we have developed and applied a new

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



approach to derive growth rates at the field scale, through a combination of measurements of particle flux and aerosol size distribution.

5 The growth rates derived in this study for combined $\text{NH}_3\text{--HNO}_3$ uptake (Table 3) are in the range of these previous studies, with 11 nm particles growing at about 3.52 nm hr⁻¹ during the day and 0.89 nm hr⁻¹ at night, with maxima of 20 to 30 nm hr⁻¹ (Fig. 5), which coincided with the maxima of the NH_3 flux. It should be noted that other particle dynamic processes, in particular coagulation, are not considered in the approach presented here (Eq. 5). Since no particular nucleation mode was observed during this study, this process is not considered to be important here. This is in contrast to studies where growth rates are derived from the evolution of size-distributions following nucleation events. At the Braunschweig field site, size-distributions were similar before and after fertilization (Fig. 8) and thus coagulation would also have affected fluxes similarly. Furthermore, there is no reason to believe that coagulation would have acted differently above and below the measurement height. Thus, in the interpretation of CPC flux measurements, coagulation is not normally considered.

10 The approach developed here, to derive growth rates at the field scale from the combined measurement of particle flux and size distribution is sufficiently sensitive to detect growth rates of a few nm hr⁻¹. The main uncertainty relates to the estimate of the dry deposition flux at the surface and thus to the variability in the dry deposition rate, which is here assumed to be constant with time (0.06 mm s⁻¹). Under the conditions encountered here, an increase of V_d to 1 mm s⁻¹ translates into an increase in the estimated growth rate by about 3 nm hr⁻¹.

25 The time-scale analysis (Fig. 6) indicates that chemistry should only be relevant during night-time conditions, while apparent emission fluxes were also observed during daytime. This highlights the limitations of the time-scale analysis based on Damköhler ratios. The analysis here is based on the above-canopy turbulent time-scale. Although chemistry may not be efficient well above the canopy, in the region for which the turbulence time-scale τ_t was estimated, obviously a region exists where τ_t becomes comparable to τ_c . Indeed, radon tracer measurements showed that the time-scale for diffusion

Aerosol fluxes and particle growth above managed grasslandE. Nemitz et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

out of the grass canopy is long (Nemitz et al., 2008b). Thus all relevant time-scales need to be considered when assessing the overall potential for chemistry to modify fluxes between the point of emission and measurement.

4.5 Implications for flux measurements and local budgets of nitrogen

5 There are three ways in which gas-particle conversion affects fluxes and local budgets, which are assessed in this section.

1. *Error for flux measurement.* As the vertical gradient is modified by the chemistry, the aerodynamic gradient technique (AGM) may need to be modified to derive the correct flux. Similarly, the flux measured by micrometeorological techniques that
10 operate at a single measurement height (z_m), such as EC and REA, differs from the true surface exchange.

2. *Error for inferential estimates and deposition modelling.* Deposition and emission are often derived from the air concentration in an inferential approach, using resistance models of a range of complexity. This approach does not usually consider
15 chemical conversion within the resistance analogue. In addition, changes in the gas/particle partitioning modify air concentrations compared with the simulation of an atmospheric transport model that ignores chemical reactions. For example, the NH_3 air concentration is lowered by transfer to the particle phase, further stimulating stomatal emission, which is governed by the difference between stomatal compensation point and atmospheric concentration.
20

3. *Modification of the local N budget.* A fraction of the emitted NH_3 is converted into slowly depositing NH_4NO_3 aerosol, *increasing* the potential for deposition. Similarly, fast depositing HNO_3 is converted into slowly depositing NH_4NO_3 aerosol, *decreasing* this deposition. The net effect of gas-to-particle conversion on the
25 local N budget will depend on the relative magnitudes of the different compounds involved.

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The average NH_4NO_3 production rate below the measurement height of the CPC fluxes was estimated to be 1.11 and $0.44 \mu\text{g m}^{-3} \text{h}^{-1}$ for day- and night-time, respectively. If it is assumed (a) that production above this measurement height is negligible as ammonia concentrations decline with increasing height and (b) that the particle production would have lasted for 14 days after fertilization, then this would equate to a production rate of $5.2 \text{ g NH}_4\text{NO}_3 \text{ ha}^{-1}$ or 0.0017% of the fertilizer N applied. By comparison, 3.2% of the fertilizer N were estimated to be emitted as NH_3 , either by direct volatilisation or, later on, by raising the stomatal compensation point within the first 10 days of fertilisation (Milford et al., 2008), i.e. around 0.05% of the emitted ammonia was converted to ammonium aerosol within the first 2 m above the ground.

Table 4 compares three estimates of the fluxes of the different reactive nitrogen species. The first one is based on the measurements above the field itself. The second estimate summarises the best estimate of the true surface exchange, correcting for the chemical interaction, similar to the correction presented in Fig. 7. The final estimate represents the fluxes that would be expected with an inferential modeling approach, if atmospheric concentrations were not influenced by the chemistry above the field. This is based on the concentrations measured at the background field. The NH_3 emission from the field would be expected to be somewhat suppressed in this scenario as the NH_3 concentration above the field would be slightly larger, but the exact NH_3 emission under these modified conditions is difficult to estimate. Because at this site the total N flux is very much dominated by NH_3 , and the relative effect on the NH_3 flux is small, the difference for the total N flux is small above this particular field. However, the average relative effect on the other forms of N is significant (2.4% for HNO_3 , 3.6% for NH_4^+ and 10% for NO_3^-), suggesting that in other circumstances these interactions could have larger relative effects on local nitrogen budgets.

4.6 Implication for global budgets and climate forcing

From fertiliser consumption statistics it is estimated that 4.5 Mt of $\text{NH}_4\text{NO}_3\text{-N}$ are used in Europe every year and 12.3 Mt globally (Matthews, 1994). Extrapolation of our mea-

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



surements to the globe implies global NH_4NO_3 -N formation/emissions following fertilizer application of 0.21 kt. Again this estimate is based on the assumption that NH_4NO_3 formation is confined to the bottom 2 m, above which precursor gas concentrations are estimated to be sufficiently diluted for continued aerosol formation, which is possibly an underestimate. Volatilisation of NH_4NO_3 fertiliser contributes only 0.46% to the global NH_3 emissions (Bouwman et al., 1997), with the main part of the emissions arising from animal wastes and organic fertilizers; and hence the overall NH_4NO_3 -N formation following NH_3 emissions could easily be as large as 45 kt NH_4NO_3 -N. This reaction will depend on background concentrations of HNO_3 , competition with reaction with sulphur and meteorology. It should also be noted that, unlike with ammonium sulphates, any NH_4NO_3 formed has the potential to re-evaporate if the gas phase concentrations of NH_3 and HNO_3 drop below the equilibrium values. For comparison, the global ammonium sulphate formation is estimated to be in the order of 57 Mt S yr^{-1} (IPCC, 2001).

5 Conclusions

Average deposition velocities to (unfertilised) cut, partially regrown and tall grassland were 0.03 , 0.06 and 0.24 mm s^{-1} , respectively. This is at the lower end of values reported for short vegetation, but consistent with the number flux being dominated by Aitken mode particles. The values are therefore applicable, e.g. to derive deposition of combustion derived primary aerosol. Analysis of sustained apparent emission fluxes after fertilisation, in relation to potential pre-cursor gases shows that these are consistent with condensation of NH_3 and HNO_3 and associated growth across the 11 nm diameter cut-off of the particle counter used for the flux measurements.

We have here developed a novel technique to derive particle growth rates from flux measurements using instruments with well-defined cut-off diameters, together with measurements of the size-distribution. Former estimates of growth rates have usually relied on the presence of a clear nucleation mode, whose size evolution could easily be monitored. By contrast our new approach is universally applicable to derive particle

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



growth rates over homogeneous surfaces at the field scale and could, for example, be used to derive aerosol growth rates due to biogenic secondary organic aerosol formation in natural environments, where bi-directional particle number fluxes are observed, even if this growth happens over a limited area.

5 Above the fertilized field growth rates of 11 nm particles of 3.5 nm hr^{-1} during the day and 0.89 nm hr^{-1} at night are similar to those reported in the literature for the uptake of organic vapours and H_2SO_4 , and illustrate the efficiency of the $\text{NH}_3\text{--HNO}_3$ system as an aerosol precursor. While the chemical interactions significantly change the local flux of gaseous HNO_3 and NH_4NO_3 aerosol, the effect on the total nitrogen flux is small,
10 as this is dominated by the large NH_3 emission during this period for which the relative effect is small.

Acknowledgements. Access to the roof of the 47 m tower was granted by Mr. Deventer from the Physikalisch-Technische Bundesanstalt. Ms L. Thomason provided assistance with the chemical analysis of the VOC tubes at Lancaster University, while K. Gilke, R. Schmelzer,
15 U. Tambor and M. Zerbian from the FAL Braunschweig helped with the analysis of the daily dry denuder/filter pack combinations. This work was funded by the EU projects GRAMINAE and NitroEurope.

References

- 20 Adams, P. J., Seinfeld, J. H., Koch, D., Micley, L., and Jacob, D.: general circulation assessment of direct radiative forcing by the sulfate-nitrate-ammonium-water inorganic aerosol system, *J. Geophys. Res.*, 106(D1), 1097–1111, 2001
- Allan, J. D., Coe, H., Bower, K. N., Williams, P. I., Gallagher, M. W., Alfarra, M. R., Jimenez, J. L., Nemitz, E., and McDonald, A. G.: Quantitative sampling using the Aerodyne Aerosol Mass Spectrometer. Part 2: Measurements of fine particulate chemical composition in two
25 UK cities, *J. Geophys. Res.*, 108, 4091, doi:4010.1029/2002JD002359, 2003
- Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., van der Hoek, K. W. and Olivier, J. G. J.: A global high-resolution emission inventory for ammonia, *Global Biogeochem. Cy.*, 11(4), 561 pp., 1997.

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Cruz, C. N., Dassios, K. G., and Pandis, S. N.: The effect of dioctyl phthalate films on the ammonium nitrate aerosol evaporation rate, *Atmos. Environ.*, 34(23), 3897–3905, 2000.
- Dal Maso, M., Kulmala, M., Lehtinen, K. E. J., Mäkelä, J. M., Aalto, P., and O'Dowd, C. D.: Condensation and coagulation sinks and formation of nucleation mode particles in coastal and boreal forest boundary layers, *J. Geophys. Res.*, 107(D19), 2001JD001053, 2002.
- David, M., Loubet, B., Cellier, P., Mattsson, M., Schjoerring, J. K., Nemitz, E., Roche, R., Riedo, M., and Sutton, M. A.: Ammonia sources and sinks in an intensively managed grassland using dynamic chambers, *Biogeosciences Discuss.*, accepted, 2008.
- Desjardins, R. L.: Description and evaluation of a sensible heat flux detector, *Bound.-Lay. Meteorol.*, 11, 147–154, 1977.
- Dämmgen, U.: Atmospheric nitrogen dynamics in Hesse, Germany: Creating the data base. 2. Atmospheric concentrations of ammonia, its reaction partners and products at Lindenberg, *Landbauforsch. Volk.*, 2(57), 157–170, 2007.
- Dassios, K. G. and Pandis, S. N.: The mass accommodation coefficient of ammonium nitrate aerosol, *Atmos. Environ.* 33(18), 2993–3003, 1999.
- Dorsey, J. R.: Aspects of biosphere/atmosphere exchange and chemical processing of aerosol. Ph.D. thesis, University of Manchester Institute of Science and Technology (UMIST), Manchester, UK, 173 pp., 2002.
- Dorsey, J. R., Nemitz, E., Gallagher, M. W., Fowler, D., Williams, P. I., Bower, K. N., and Beswick, K. M.: Direct measurements and parameterisation of aerosol flux, concentration and emission velocity above a city, *Atmos. Environ.*, 36(5), 791–800, 2002.
- Farquhar, G. D., Firth, P. M., Wetselaar, R., and Weir, B.: On the gaseous exchange of ammonia between leaves and the environment: determination of the ammonia compensation point, *Plant Physiol.*, 66, 710–714, 1980.
- Foken, T., Dlugi, R., and Kramm, G.: On the determination of dry deposition and emission of gaseous compounds at the biosphere-atmosphere interface, *Meteorol. Z.*, 4(3), 91, 1995.
- Fowler, D. and Duyzer, J. H.: Micrometeorological techniques for the measurement of trace gas exchange, in: *Exchange of trace gases between terrestrial ecosystems and the atmosphere*, edited by: Andrea, M. O. and Schimel, D. S., John Wiley & Sons Ltd., New York, 189–207, 1989.
- Fuchs, N. A. and Sutugin, A. G.: Highly dispersed aerosol, in: *Topics in Current Aerosol Research*, edited by: Hydy, G. M. and Brock, J. R., Pergamon, New York, 160 pp., 1971.
- Gallagher, M. W., Nemitz, E., Dorsey, J. R., Fowler, D., Sutton, M. A., Flynn, M., and Duyzer,

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- J.: Measurements and parameterizations of small aerosol deposition velocities to grassland, arable crops and forest: influence of surface roughness length on deposition, *J. Geophys. Res.*, 107(D12), 4154–4163, doi:10.1029/2001JD000817, 2002.
- Hensen, A., Nemitz, E., Flynn, M. J., Blatter, A., Jones, S. K., Srensen, L. L., Hensen, B., Pryor, S., Jensen, B., Otjes, R. P., Cobussen, J., Loubet, B., Erisman, J. W., Gallagher, M. W., Neftel, A., and Sutton, M. A.: Inter-comparison of ammonia fluxes obtained using the relaxed eddy accumulation technique, *Biogeosciences Discuss.*, 5, 3965–4000, 2008, <http://www.biogeosciences-discuss.net/5/3965/2008/>.
- Hoffmann, T., Odum, J. R., Bowman, F., Collins, D., Klockow, D., Flagan, R. C., and Seinfeld, J. H.: Formation of organic aerosol from the oxidation of biogenic hydrocarbons, *J. Atmos. Chem.*, 26, 189–222, 1997.
- Holmes, N. S.: A review of particle formation events and growth in the atmosphere in the various environments and discussion of mechanistic implications, *Atmos. Environ.*, 41, 2183–2201, 2007.
- Huebert, B. J., Luke, W. T., Delany, A. C., and Brost, R. A.: Measurements of concentrations and dry surface fluxes of atmospheric nitrates in the presence of ammonia, *J. Geophys. Res.*, 93, 7127–7136, 1988.
- IPCC: Climate Change 2001 – The scientific basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, New York, 881 pp., 2001.
- IPCC: Climate Change 2007 – The Physical Science Basis. Contribution of Working group I to the Fourth Assessment Report of the IPCC, The Intergovernmental Panel on Climate Change, Cambridge University Press, New York, ISBN 978-0521-88009-1, 1009 pp., 2007.
- Karl, T., Guenther, A., Jordan, A., Fall, R., and Lindinger, W.: Eddy covariance measurement of biogenic oxygenated VOC emissions from hay harvesting, *Atmos. Environ.*, 35(3), 491–495, 2001.
- Keuken, M. P., Schoonebeek, C. A. M., van Wensveen-Louter, A., and Slanina, J.: Simultaneous sampling of NH_3 , HNO_3 , HCl , SO_2 and H_2O_2 in ambient air by a wet annular denuder system, *Atmos. Environ.*, 22, 2541–2548, 1988.
- Kirstine, W., Glabally, I., Ye, Y., and Hooper, M.: Emissions of volatile organic compounds (primarily oxygenated species) from pasture, *J. Geophys. Res.*, 103(D9), 10605–10619, 1998.
- Korhonen, P., Kulmala, M., and Viisanen Y.: A theoretical study of binary homogenous nucle-

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ation of water-ammonium chloride particles in the atmosphere, *J. Aerosol Sci.*, 28(6), 901–917, 1997.

Kowalski, A. S.: Deliquescence induces eddy covariance and estimable dry deposition errors, *Atmos. Environ.*, 35, 4843–4851, 2001.

5 Kramm, G., Müller, H., Fowler, D., Hofken, K. D., Meixner, F. X., and Schaller, E.: A Modified Profile Method for Determining the Vertical Fluxes of NO, NO₂, Ozone, and HNO₃ in the Atmospheric Surface-Layer, *J. Atmos. Chem.*, 13, 265–288, 1991.

Kulmala, M., dal Maso, M., Mäkelä, J. M., Pirjola, L., Väkevä, M., Aalto, P., Mikkulainen, P., Hämeri, K., and O'Dowd, C. D.: On the formation, growth and composition of nucleation
10 mode particles, *Tellus B*, 53(4), 479–490, 2001.

Kulmala, M., Vehkamäki, H., Petaja, T., Dal Maso, M., Lauri, A., Kerminen, V.-M., Birmili, W., and McMurry, P. H.: Formation and growth rates of ultrafine atmospheric particles: a review of observations, *J. Aerosol Sci.*, 35(2), 143–176, 2004.

Matthews, E.: Nitrogen fertilisers: Global distribution of consumption and associated emissions of nitrous oxide and ammonia, *Global Biochem. Cy.*, 8(4), 411–439, 1994.

15 Milford, C., Theobald, M. R., Nemitz, E., and Sutton, M. A.: Dynamics of ammonia exchange in response to cutting and fertilizing in an intensively-managed grassland, *Water Air Soil Poll.: Focus*, 1, 167–176, 2001.

Milford, C., Theobald, M. R., Nemitz, E., Hargreaves, K. J., Horvath, L., Raso, J., Dämmgen, U., Neftel, A., Jones, S. K., Hensen, A., Loubet, B., Cellier, P., and Sutton, M. A.: Ammonia
20 fluxes in relation to cutting and fertilization of an intensively managed grassland derived from an inter-comparison of gradient measurements, *Biogeosciences Discuss.*, 5, 4699–4744, 2008, <http://www.biogeosciences-discuss.net/5/4699/2008/>.

Mozurkewich, M.: The dissociation constant of ammonium nitrate and its dependence on temperature, relative humidity and particle size, *Atmos. Environ.*, 27, 261–270, 1993.

25 Napari, I., Noppel, M., Vehkamäki, H., and Kulmala, M.: Parameterization of ternary nucleation rates for H₂SO₄–NH₃–H₂O vapors, *J. Geophys. Res.*, 107(D19), 4381, doi:10.1029/2002JD002132, 2002.

Neftel, A., Blatter, A., Hesterberg, R., and Staffelbach, T.: Measurements of concentration gradients of HNO₂ and HNO₃ over a semi-natural ecosystem, *Atmos. Environ.*, 30, 3017–3025, 1996.

30 NEG-TAP: Transboundary Air Pollution: Acidification, Eutrophication and Group-Level Ozone in the UK, Report of the National Expert Group on Transboundary Air Pollution (NEG-TAP)

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for the UK Department for Environment, Food and Rural Affairs, Scottish Executive, The National Assembly for Wales/Cynulliad Cenedlaethol Cymru and the Department of the Environment for Northern Ireland, CEH Edinburgh, 2001.

- 5 Nemitz, E. and Sutton, M. A.: Gas-particle interactions above a Dutch heathland: III. Modelling the influence of the $\text{NH}_3\text{--HNO}_3\text{--NH}_4\text{NO}_3$ equilibrium on size-segregated particle fluxes, *Atmos. Chem. Phys.*, 4, 1025–1045, 2004, <http://www.atmos-chem-phys.net/4/1025/2004/>.
- Nemitz, E., Flynn, M., Williams, P. I., Milford, C., Theobald, M. R., Blatter, A., Gallagher, M. W., and Sutton, M. A.: A relaxed eddy accumulation system for the automated measurement of atmospheric ammonia fluxes, *Water Air Soil Poll.: Focus*, 1, 189–202, 2001.
- 10 Nemitz, E., Gallagher, M. W., Duyzer, J. H., and Fowler, D.: Micrometeorological measurements of particle deposition velocities to moorland vegetation, *Q. J. Roy. Meteor. Soc.*, 128, 2281–2300, 2002.
- Nemitz, E., Hargreaves, K. J., Neftel, A., Loubet, B., Cellier, P., Dorsey, J. R., Flynn, M., Hensen, A., Weidinger, T., Meszaros, R., Horvath, L., Frühauf, C., Sutton, M. A. and Gallagher, M. W.: Intercomparison and assessment of turbulent and physiological exchange parameters of grassland, *Biogeosciences Discuss.*, accepted, 2008a.
- 15 Nemitz, E., Loubet, B., Cellier, P., Lehmann, B., Neftel, A., Hargreaves, K. J., Herrmann, B., Jones, S., Ihly, B., Tarakanov, S. V., and Sutton, M. A.: Turbulence characteristics in grassland canopies and implications for tracer transport, *Biogeosciences Discuss.*, accepted, 2008b.
- 20 Nemitz, E., Sutton, M. A., Fowler, D., and Choularton, T.: Application of a NH_3 gas-to-particle conversion model to measurement data, in: Poster proceedings of the international conference on atmospheric ammonia, edited by: Sutton, M. A., Lee, D. S., Dollard, G. J., and Fowler, D., Oxford 2–4 October 1995, Institute of Terrestrial Ecology Edinburgh, UK, 98–103, 1996.
- 25 Nemitz, E., Sutton, M. A., Wyers, G. P., Otjes, R. P., Schjoerring, J. K., Gallagher, M. W., Parrington, J., Fowler, D., and Choularton, T. W.: Surface/atmosphere exchange and chemical interaction of gases and aerosols over oilseed rape, *Agr. Forest Meteorol.*, 105, 427–445, 2000.
- 30 Nemitz, E., Sutton, M. A., Wyers, G. P., Otjes, R. P., Mennen, M. G., van Putten, E. M., and Gallagher, M. W.: Gas-particle interactions above a Dutch heathland: II. Concentrations and surface exchange fluxes of atmospheric particles, *Atmos. Chem. Phys.*, 4, 1007–1024, 2004, <http://www.atmos-chem-phys.net/4/1007/2004/>.

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Petroff, A., Mailliat, A., Amielh, M., and Anselmet, F.: Aerosol dry deposition on vegetation canopies. Part I: Review of present knowledge, *Atmos. Environ.*, 42(16), 3625–3653, 2008.
- Pio, C. A. and Harrison, R. M.: Vapour pressure of ammonium chloride aerosol: effect of temperature and humidity, *Atmos. Environ.*, 21, 2711–2715, 1987.
- 5 Pryor, S. C., Gallagher, M. W., Sievering, H., Larsen, S. E., Barthelmie, R. J., Birsan, F., Nemitz, E., Rinne, J., Kulmala, M., Groenholm, T., Taipale, R., and Vesala, T.: A review of measurement and modelling results of particle atmosphere-surface exchange, *Tellus B*, 60, 42–75, 2008.
- Reid, R. C., Prausnitz, J. M., and Poling, B. E.: *The Properties of gases and Liquids*, McGraw-Hill, New York, 4th edn., 741 pp., 1987.
- 10 Riedo, M., Milford, C., Schmid, M., and Sutton, M. A.: Coupling soil-plant-atmosphere exchange of ammonia with ecosystem functioning in grasslands, *Ecol. Model.*, 158, 83–110, 2002.
- Rudolf, R., Vrtala, A., Kulmala, M., Vesala, T., Viisanen, Y., and Wagner, P. E.: Experimental study of sticking probabilities for condensation of nitric acid – water vapor mixtures, *J. Aerosol Sci.*, 32, 913–932, 2001.
- 15 Stelson, A. W. and Seinfeld, J. H.: Thermodynamic prediction of the water activity, NH_4NO_3 dissociation constant, density and refractive index for the NH_4NO_3 – $(\text{NH}_4)_2\text{SO}_4$ – H_2O system at 25°C, *Atmos. Environ.*, 16, 2507–2514, 1982.
- Sutton M. A., Schjorring J. K., and Wyers, G. P.: Plant – atmosphere exchange of ammonia, *Philos. T. R. Soc. A*, 351, 261–278, 1995.
- 20 Sutton, M. A., Milford, C., Nemitz, E., Theobald, M. R., Hill, P. W., Fowler, D., Schjoerring, J. K., Mattson, M. E., Nielsen, K. H., Husted, S., Erisman, J. W., Otjes, R., Hensen, A., Cellier, P., Loubet, B., David, M., Genermont, S., Neftel, A., Blatter, A., Hermann, B., Jones, S. K., Horvath, L., Führer, E., Mantzanas, K., Koukoura, Z., Gallagher, M. W., Williams, P. I., and Riedo, M.: Biosphere-atmosphere interactions of ammonia with grasslands: experimental strategy and results from a new European initiative, *Plant Soil*, 228(1), 131–145, 2001.
- 25 Sutton, M. A., Fowler, D., Moncrieff, J. B., and Storeton-West, R. L.: The exchange of atmospheric ammonia with vegetated surfaces. II: Fertilized vegetation, *Q. J. Roy. Meteor. Soc.*, 119, 1047–1070, 1993.
- 30 Sutton, M. A., Nemitz, E., Erisman, J. W., Beier, C., Butterbach Bahl, K., Cellier, P. de Vries, W., Cotrufo, F., Skiba, U., Di Marco, C., Jones, S. K., Laville P., Soussana, J. F., Loubet, B., Twigg, M., Famulari, D., Whitehead, J., Gallagher, M. W., Neftel, A., Flechard, C. R., Herrmann, B., Calanca, P. L., Schjoerring, J.K., Daemmgen, U., Horvath, L., Tang, Y. S., Emmett, B. A.,

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Tietema, A., Peñuelas, J., Kesik, M., Brueggemann, N., Pilegaard, K., Vesala, T., Campbell, C. L., Olesen, J. E., Dragosits, U., Theobald, M. R., Levy, P., Mobbs, D. C., Milne, R., Viovy, N., Vuichard, N., Smith, J. U., Smith, P. E., Bergamaschi, P., Fowler, D., and Reis, S.: Challenges in quantifying biosphere-atmosphere exchange of nitrogen species, *Environ. Pollut.*, 150, 125–139, 2007.

Sutton, M. A., Nemitz, E., Milford, C., Erisman, J. W., Hensen, A., Cellier, P., David, M., Loubet, B., Personne, E., Schjoerring, J. K., Mattsson, M. E., Dorsey, J., Gallagher, M., Horvath, L., Weidinger, T., Dämmgen, U., Neftel, A., Herrmann, B., Lehman, B., and Burkhardt, J.: Dynamics of ammonia exchange with cut grassland: Synthesis of results and conclusions, *Biogeosciences Discuss.*, accepted, 2008.

Sutton, M. A., Nemitz, E., Theobald, M. R., Milford, C., Dorsey, J. R., Gallagher, M. W., Hensen, A., Jongejan, P. A. C., Erisman, J. W., Mattsson, M. E., Schjoerring, J. K., Cellier, P., Loubet, B., Roche, R., Neftel, A., Hermann, B., Jones, S., Lehman, B. E., Horvath, L., Weidinger, T., Rajkai, K., Burkhardt, J., Lpmeier, F. J., and Daemmgen, U.: Dynamics of ammonia exchange with cut grassland: strategy and implementation of the GRAMINAE Integrated Experiment, *Biogeosciences Discuss.*, 5, 3347–3407, 2008, <http://www.biogeosciences-discuss.net/5/3347/2008/>.

ten Brink, H. M., Veefkind, J. P., Waijers-Ijpelaan, A., and van der Hage, J. C.: Aerosol light-scattering in the Netherlands, *Atmos. Environ.*, 30, 4251–4261, 1996.

Thomas, R. M., Trebs, I., Otjes, R., Jongejan, P. A. C., ten Brink, H., Phillips, G., Kortner, M., Meixner, F. X., and Nemitz, E.: An automated analyzer to measure surface-atmosphere exchange fluxes of water soluble inorganic aerosol compounds and reactive trace gases, *Environ. Sci. Technol.*, accepted, 2008.

Umweltbundesamt: Jahresbericht 2000 des Messnetzes, UBA Texte 77-01, Umweltbundesamt, Berlin, 174 pp., 2001.

van Oss, R., Duyzer, J., and Wyers, P.: The influence of gas-to-particle conversion on measurements of ammonia exchange over forest, *Atmos. Environ.*, 32(3), 465–471, 1998.

Verheggen, B. and Mozurkewich, M.: Determination of nucleation and growth rates from measurements of atmospheric aerosol size distributions, *J. Aerosol Sci.*, 31(Suppl. 1), S446–S447, 2000.

Vestreng, V. and Støren, E.: Analysis of UN ECE/EMEP emission data, MSC-W Status Report 2000, Oslo, Norwegian Meteorological Institute (EMEP MSC-W Note 1/00) (DNMI Research Note 37), 2000.

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Vong, R. J., Vickers, D., and Covert, D. S.: Eddy correlation measurements of aerosol deposition to grass, *Tellus B*, 56, 105–117, 2004.
- Wesely, M. L., Cook, D. R., Hart, R. L., and Speer, R. E.: Measurements and parameterization of particulate sulfur dry deposition over grass, *J. Geophys. Res.-Atmos.*, 90, 2131–2143, 1985.
- Wexler, A. S. and Seinfeld, J. H.: The distribution of ammonium salts among size and composition dispersed aerosol, *Atmos. Environ.*, 24, 1231–1246, 1990.
- Wiebe, H. A., Anlauf, K. G., Tuazon, E. C., Winer, A. M., Biermann, H. W., Appel, B. R., Solomon, P. A., Cass, G. R., Ellestad, T. G., Knapp, K. T., Peake, E., Spicer, C. W., and Lawson, D. R.: A comparison of measurements of atmospheric ammonia by filter packs, transition flow reactors, simple and annular denuders and Fourier transform infrared spectroscopy, *Atmos. Environ.*, 24, 1019–1028, 1990.
- Williams, P. I.: Construction and validation of a DMPS for aerosol characterization, Ph.D. thesis, University of Manchester Institute of Science and Technology, Manchester, UK, 1999.
- Zimmerling, R., Dämmgen, U., Haenel, H. D., Kusters, A., Max, W., and Jäger, H. J.: The classifying ratio-metric method for the continuous determination of atmospheric flux densities of reactive N- and S-species with denuder filter systems, *J. Appl. Bot.-Angew. Bot.*, 71(1–2), 38–49, 1997.

BGD

6, 341–389, 2009

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Table 1. Summary of the statistics of the concentrations and fluxes measured with the CPC system before the cut and after fertilisation (based on 30-min flux values).

	Period (1): before cut			Period (2): after cut, before fertilisation			Period (3): directly after fertilisation			Period (4): 7–10 days after fertilisation		
	χ_N [# cm ⁻³]	F_N [# cm ⁻² s ⁻¹]	V_d [mm s ⁻¹]	χ_N [# cm ⁻³]	F_N [# cm ⁻² s ⁻¹]	V_d [mm s ⁻¹]	χ_N [# cm ⁻³]	F_N [# cm ⁻² s ⁻¹]	V_d [mm s ⁻¹]	χ_N	F_N	V_d
Min	2903	−4080	−5.39	3004	−3950	−11.7	1800	−5790	−10.8	3231	−4250	−10.8
Max	23 840	5700	4.34	16 526	5820	6.69	23 230	9020	9.26	21 803	6000	4.80
5%-ile	4303	−1340	−0.98	3831	−1110	−1.70	3776	−1090	−3.17	4108	−1160	−1.28
95%-ile	12 930	718	1.74	13 850	1220	1.56	17 510	2570	1.11	13 724	992	1.53
Mean	7190	−181	0.24	7611	−17	0.03	8628	323	−0.46	8301	−63	0.06
Median	6486	−151	0.24	7057	−45	0.06	7689	85	−0.13	7612	−44	0.05
Standard dev.	2665	710	0.95	2920	857	1.37	4146	1220	1.48	3265	835	1.15
Frequency of emission[%]		31			42			60			43	
N	714	701	701	636	580	580	612	598	598	295	290	290

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Table 2. Summary of the concentrations and fluxes of selected VOCs (averages of 9 1-hr runs between cutting and fertilization and 5 1-hr runs after fertilization).

	Concentration [ng m^{-3}]		Flux [$\text{ng m}^{-2} \text{s}^{-1}$]	
	before	after fertilization	Before	after fertilization
Benzaldehyde	5400 (± 220)	5700 (± 430)	−140 (± 69)	−250 (± 61)
Nonanal	590 (± 22)	650 (± 73)	14 (± 8.0)	7.6 (± 16)
α -Pinene	84 (± 9.8)	47 (± 4.6)	1.9 (± 0.6)	1.4 (± 1.1)
1- β -Pinene	75 (± 2.6)	97 (± 26)	−0.33 (± 1.0)	14 (± 16)
d-3-Carene	290 (± 7.4)	300 (± 28)	−4.3 (± 3.6)	8.6 (± 8.4)

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏮](#)
[⏭](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 3. Summary of typical particle growth parameters for various 12-h periods.

Period	Day/ Night	F_N [cm ⁻² s ⁻¹]	A_p [μm ² m ⁻³]	$dr/dt _{dp=}$ =11 nm [nm hr ⁻¹]	$J_{\text{H}_4\text{NO}_3}$ [μg m ⁻³ hr ⁻¹]	ΔF_{NH_3} [ng m ⁻² s ⁻¹]	F_{NH_3} [ng m ⁻² s ⁻¹]
5 June	Day	1789	293	2.18	0.95	0.112	597
5–6 June	Night	178	320	1.42	0.34	0.040	1262
6 June	Day	1102	115	5.73	0.79	0.094	887
6–7 June	Night	54	266	0.50	0.17	0.020	1001
7 June	Day	900	245	2.75	1.14	0.135	443
7–8 June	Night	230	297	1.23	0.47	0.055	1311
8 June	Day	1388	253	17.24	5.33	0.629	549
8–9 June	Night	293	245	0.55	0.26	0.031	193
9 June	Day	285	240	1.09	0.67	0.079	378
9–10 June	Night	149	401	0.71	0.46	0.054	862
10 June	Day	212	298	1.92	0.86	0.102	495
10–11 June	Night	-30	556	-0.02	-0.08	-0.009	160
11 June	Day	99	294	2.13	0.54	0.064	175
11–12 June	Night	114	327	3.13	2.01	0.237	168
12 June	Day	-116	194	-0.77	-0.15	-0.018	202
12–13 June	Night	-6	320	0.59	0.43	0.051	98
13 June	Day	39	240	-0.57	-0.14	-0.016	118
13–14 June	Night	-129	333	-0.12	-0.10	-0.012	99
Average	Day	633	241	3.52	1.11	0.131	427
Average	Night	95	341	0.89	0.44	0.052	573

F_N : measured particle number flux; A_p : aerosol surface area per unit volume of air; $dr/dt|_{dp=11\text{nm}}$: growth rate of 11 nm particles; $J_{\text{NH}_4\text{NO}_3}$: estimated NH_4NO_3 production rate; ΔF_{NH_3} : error on the NH_3 flux measurement due to gas-particle reactions; F_{NH_3} : measured NH_3 flux.

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.

Table 4. Effects of gas-to-particle conversion on flux measurements and budget estimates of reactive nitrogen (excluding NO_x). Fluxes are in kg N ha^{-1} for the period 5 to 15 June 2000; negative values represent deposition fluxes.

Component	Measurements, and parameterisations not corrected for chemistry	Corrected best estimate for surface	No chemistry scenario
NH_3 gas	3.22	3.22	$<3.23^{(a)}$
HNO_3 gas	−0.043	−0.042	−0.033 ^(b)
NH_4^+ aerosol	−0.028	−0.029	−0.027 ^(b)
NO_3^- aerosol	−0.010	−0.011	−0.007 ^(b)
Net N flux	3.14	3.14	<3.20

^(a) Emission would be reduced due to larger air concentration (cf. text).

^(b) Estimated from KAPS filter/denuder measurements over adjacent field.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.




Fig. 1. Time-series of the particle number concentration and fluxes of particles (condensation nuclei, CN) measured with the CPC flux system (11 nm to $\approx 3 \mu\text{m}$).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.




Fig. 2. Evolution of the gas concentrations of NH_3 , HCl and HNO_3 1 m and 47 m above the grassland, compared with the NH_3 concentration extrapolated to the canopy ($\text{NH}_3(z_0')$). See text for details. The fertilizer was applied on 5 June 08:00 to 10:00 GMT. Also indicated are the periods of the MOUDI impactor runs (B to D). The concentrations of NH_3 are based on the best estimate χ_{NH_3} and F_{NH_3} provided by Milford et al. (2008).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.




Fig. 3. Time-series of NH_4^+ and NO_3^- aerosol concentration measured above and upwind of the grassland in both mass concentration and equivalent concentration.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.




Fig. 4. Comparison of measured concentration products at 1 m height of (a) NH_3 with HNO_3 and (b) NH_3 with HCl , compared with concentration products in equilibrium with pure NH_4NO_3 and NH_4Cl aerosol, respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.




Fig. 5. Time-series of the growth rate of 11 nm particles (dr/dt) as derived from CPC flux measurements and DMPS measurements, using Eqs. (5), (6) and (7).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.




Fig. 6. Time-series of the aerosol surface area and the Damköhler ratio ($D_r = \tau_t / \tau_c$). The dashed line represents the critical value of D_r , above which chemistry is expected to alter the vertical gradients.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.




Fig. 7. Time-series of the measured NH_3 flux and the NH_3 flux at the surface, corrected for the fraction of NH_3 converted to NH_4^+ (based on the best flux estimate provided by Milford et al., 2008).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.




Fig. 8. Averaged particle size-distributions during the three management periods.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.




Fig. 9. Time-evolution of the size of particles, with an initial radius (r) of 1 nm, for a range of vapour concentrations (χ_v).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol fluxes and particle growth above managed grassland

E. Nemitz et al.




Fig. 10. Concentration, flux and deposition velocity (V_d) of HNO_3 measured with the SJAC/WAD gradient system, in relation to the maximum deposition velocity (V_{max}).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol fluxes and
particle growth above
managed grassland

E. Nemitz et al.




Fig. 11. Schematic of the proposed processes of emission, gas-particle conversion and deposition at the fertilised grassland site.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)