

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose¹, Y. Feng², G. R. DiTullio³, R. B. Dunbar⁴, C. E. Hare⁵, P. A. Lee³,
M. Lohan⁶, M. Long⁴, W. O. Smith Jr.⁷, B. Sohst⁸, S. Tozzi⁷, Y. Zhang⁵, and
D. A. Hutchins²

¹Biology Department, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

²Biology Department, University of Southern California, Los Angeles, CA 90089-0371, USA

³Hollings Marine Laboratory, College of Charleston, Charleston, SC 29412, USA

⁴Department of Environmental Earth Systems Science, Stanford University, Stanford, CA 94305-2115, USA

⁵College of Marine Studies, University of Delaware, Lewes, DE 19958, USA

⁶School of Earth, Ocean and Environmental Sciences, University of Plymouth, Plymouth PL4 8AA, UK

BGD

6, 5849–5889, 2009

**Synergistic effects of
iron and temperature
on Antarctic plankton
assemblages**

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

⁷Virginia Inst. of Marine Science, College of William and Mary, Gloucester Pt, VA 23602, USA
⁸Department of Ocean, Earth and Atmospheric Sciences, Old Dominion University, Norfolk, VA 23529, USA

Received: 30 May 2009 – Accepted: 9 June 2009 – Published: 19 June 2009

Correspondence to: D. A. Hutchins (dahutch@usc.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

6, 5849–5889, 2009

**Synergistic effects of
iron and temperature
on Antarctic plankton
assemblages**

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Iron availability and temperature are important limiting factors for the biota in many areas of the world ocean, and both have been predicted to change in future climate scenarios. However, the impacts of combined changes in these two key factors on microbial trophic dynamics and nutrient cycling are unknown. We examined the relative effects of iron addition (+1 nM) and increased temperature (+4°C) on plankton assemblages of the Ross Sea, Antarctica, a region characterized by annual algal blooms and an active microbial community. Increased iron and temperature individually had consistently significant but relatively minor positive effects on total phytoplankton abundance, phytoplankton and microzooplankton community composition, as well as photosynthetic parameters and nutrient drawdown. Unexpectedly, increased iron had a consistently negative impact on microzooplankton abundance, most likely a secondary response to changes in phytoplankton community composition. When iron and temperature were increased in concert, the resulting interactive effects were greatly magnified. This synergy between iron and temperature increases would not have been predictable by examining the effects of each variable individually. Our results suggest the possibility that if iron availability increases under future climate regimes, the impacts of predicted temperature increases on plankton assemblages in polar regions could be significantly enhanced. Such synergistic and antagonistic interactions between individual climate change variables highlight the importance of multivariate studies for marine global change experiments.

1 Introduction

The Ross Sea, Antarctica is the location of one of the world's largest annual algal blooms (Smith et al., 2000). These algal blooms are characterized by spatial shifts in taxonomic composition across the region (Arrigo et al., 2000). The colonial prymnesiophyte *Phaeocystis antarctica* typically dominates phytoplankton assemblages of

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the southern and central polynya areas of the Ross Sea during mid- to late austral spring, peaking in early summer (DiTullio and Smith, 1996; Smith et al., 2003). Diatom assemblages, usually composed of *Fragilariopsis* and/or *Thalassiosira*, dominate the portions of the Ross Sea near the edge of the polynya, particularly in the west (Arigo et al., 1999; Leventer and Dunbar, 1996). These and other diatom species such as *Pseudo-nitzschia* spp. are, however, also present throughout the polynya (Garrison et al., 2003). Heterotrophic protists have been reported to achieve high biomass in Antarctic marine systems (Caron et al., 2000; Dennett et al., 2001; Garrison, 1991; Garrison and Gowing, 1993). However, the impact of micro- and mesozooplankton grazing on phytoplankton assemblages dominated by colonial *P. antarctica* appears to be relatively minor (Caron et al., 2000; Haberman et al., 2003; Haberman et al., 2003).

In the two decades since the pioneering experiments of Martin (1990) and deBaar (1990), iron limitation has come to be generally recognized as a primary “bottom-up” control on phytoplankton growth in the Southern Ocean. Both shipboard (Boyd et al., 1999; Hare et al., 2005; Hutchins et al., 2001; Scharek et al., 1997; Takeda, 1998) and in situ (Boyd et al., 2000; Coale et al., 2004; Gervais et al., 2002) experiments have repeatedly demonstrated that iron availability strongly constrains the biology and biogeochemistry of most of the open high nutrient low chlorophyll (HNLC) Southern Ocean.

Parts of the coastal Southern Ocean are also strongly iron-limited, at least on a seasonal basis. As in other iron-limited continental shelf waters off California and Peru (Firme et al., 2003; Hutchins and Bruland, 1998; Hutchins et al., 2002), the combination of limited iron inputs and high biological demand leads to annually recurring summertime iron limitation in the southern Ross Sea (Coale et al., 2003; Martin et al., 1990; Maucher and DiTullio, 2003; Olson et al., 2000; Sedwick et al., 2000). In this region, mixed-layer dissolved Fe concentrations can be <0.1 nM during the growing season (Coale et al., 2003; Fitzwater et al., 2000; Sedwick et al., 2000). Since melting sea ice may be a major source of biologically available iron (Sedwick and DiTullio 1997), future changes in ice cover could lead to large shifts in the availability and timing

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of iron inputs in the Ross Sea.

In general, global change effects are expected to be manifested earlier and most severely in the Southern Ocean (Sarmiento et al., 1998). The Antarctic peninsula has experienced perhaps the fastest rate of warming and ice loss on the planet (Vaughan et al., 2003). In view of the expected minimum global temperature rise of 1–3°C over the next century (Alley et al., 2007), the broad future trends in coastal Antarctic ecosystems seem likely to include further widespread warming and ice retreat. If so, these changes in seasonal ice dynamics and accompanying increased stratification effects will lead to major changes in mixing regimes, nutrient supplies, and light environments experienced by Antarctic phytoplankton (Boyd and Doney, 2002; Montes-Hugo et al., 2009). Experiments in other high latitude regimes such as the Bering Sea and North Atlantic have already demonstrated the potential for significant effects of increased temperature on plankton assemblages, including changes in community composition, photosynthetic rates, nutrient drawdown and microbial food web dynamics (Feng et al., 2009; Hare et al., 2007; Rose et al., 2009).

Less is known for certain about future shifts in iron supplies to the iron-limited phytoplankton communities of the Ross Sea. Since the largest current sources of iron to surface waters here are vertical mixing and seasonal sea ice melting (Coale et al., 2003; Fitzwater et al., 2000; Martin et al., 1990; Sedwick and DiTullio, 1997; Sedwick et al., 2000), iron availability is highly vulnerable to anticipated climate change impacts. Reduction of these sources of iron to the Ross Sea under future global change scenarios would result in intensification of current iron limitation of phytoplankton growth. However, other potential sources of bioavailable iron could increase at the same time, such as melting glacial ice (Overpeck et al., 2006), or even significant local aeolian input as snow cover retreats and exposes new soil on the adjacent continent. The future trends in iron inputs to the Ross Sea cannot be reliably predicted at present, but it seems inevitable that both iron supply and sea surface temperature will differ substantially from today's norms in the Ross Sea, with unknown consequences for regional food webs. In this study, we attempt to address this issue by examining the biological

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and biogeochemical interactions between iron availability and increased temperature in an experiment using a Ross Sea phytoplankton and microzooplankton community.

2 Methods

2.1 Experimental design

5 Experiments were conducted during the CORSACS (Controls On Ross Sea Algal Community Structure) expedition in January 2006 to the Ross Sea, Antarctica, onboard the RVIB Nathaniel B. Palmer (cruise NBP-0601). Water was collected at 75.00° S, 177.36° E using a trace metal clean towed-intake surface water Teflon diaphragm pumping system (Bruland et al., 2005). Water was prescreened through acid-washed
10 200 µm Nitex mesh to eliminate large zooplankton and collected into a 50-L mixing carboy. Collected water was gently mixed and dispensed into 12 4.5-L acid washed trace metal clean clear polycarbonate bottles for incubation. Half of the bottles were spiked with 1.0 nM FeCl₃ (final concentration) at the beginning of the experiment. Bottles were incubated in two temperature controlled deck-board incubators (Feng et al.,
15 2009; Hare et al., 2007). Incubators were screened to 18% of I_0 using two layers of neutral density filter. One incubator was kept at ambient temperature (0°C), while the temperature in the other was gradually increased to 4°C over the course of 24 h. Bottles were incubated for seven days and sampled daily under a laminar flow hood using trace metal-clean techniques. Daily sampling included total chlorophyll, nanophyto-
20 plankton abundance, F_v/F_m and dissolved nutrients. Additional samples were taken at time zero, on day 4 and the final day for >20 µm chlorophyll, particulate nutrients, photosynthesis vs. irradiance curves, diatom and microzooplankton abundance and biogenic sulfur. Samples for phytoplankton pigments (determined by HPLC) and total dissolved iron were taken on the initial and final days of the experiment.

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.2 Plankton assemblage measurements

Chlorophyll samples (50–250 mL) were filtered onto either a GF/F or a 20 µm polycarbonate filter using low vacuum pressure. Samples were extracted for at least 24 h in the dark at –20°C in 90% acetone and read on a Turner Designs fluorometer (Welschmeyer, 1994). Nanophytoplankton abundance was determined using flow cytometry. Two milliliters of sample were preserved in 1% seawater-buffered, 0.2 µm-filtered formalin (final concentration) and frozen at –80°C until analysis. Samples were run on a FACSCalibur flow cytometer for 5 min on the high flow rate setting (Campbell, 2001). Nanophytoplankton were identified on two-dimensional cytograms based on forward scatter (FSC) and red fluorescence (FL3).

A small-volume photosynthetron was used to establish photosynthesis versus irradiance (*P-E*) curves according to van Hilst and Smith (2002). Samples (65 mL) collected from each experimental treatment were mixed with ca. 150 µCi NaH¹⁴CO₃ (pH 9.6). Two mL subsamples of this mix were transferred into each of 32 glass scintillation vials, which were then incubated under a range of irradiances from 0–600 µmol photons m^{–2} s^{–1}. The two-hour incubation was performed using a high-volume recirculating water bath to maintain appropriate experimental temperatures for all subsamples. At the end of the incubation, 0.5 mL of 1.0 N HCl was added to each vial to terminate photosynthesis and samples were vented in a hood to remove inorganic ¹⁴C. After 24 h of venting, 5 mL Ecolume scintillation fluor was added to each sample, and samples were stored in the dark at room temperature for an additional 24 h to reduce chemiluminescence. Samples were then counted on a Packard liquid scintillation counter. Total inorganic ¹⁴C-bicarbonate availability (TA) was measured on a 0.1 mL aliquot (to which 0.05 mL β-phenethylamine, a CO₂ trap, was added) directly in Ecolume.

The *P-E* curves were fit to the Webb et al. (1974) empirical model:

$$P^B = P_m^B \left[1 - e^{-\alpha E / P_m^B} \right] \quad (1)$$

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

P^B = the rate of photosynthesis normalized to chlorophyll-*a* [$\text{mg C (mg chl-}a)^{-1} \text{ h}^{-1}$],
 P_m^B = the maximum rate of photosynthesis in the absence of photoinhibition, α = the
 initial, linear, photosynthetic rate [$\text{mg C (mg chl-}a)^{-1} \text{ h}^{-1}$ ($\mu\text{mol photons m}^{-2} \text{ s}^{-1}$)],
 E = irradiance ($\mu\text{mol photons m}^{-2} \text{ s}^{-1}$). Photoinhibition was insignificant in most of the
 experimental treatments, and the photoinhibition term (Platt et al., 1980) was set to
 zero for all.

Photochemical efficiency of PSII was measured using a MBARI 4th generation
 bench-top fast repetition rate fluorometer (FRRF) (Kolber et al., 1994). Samples were
 collected each day from experimental bottles, immediately placed on ice and kept in
 low light conditions ($5\text{--}10 \text{ mol photons m}^{-2} \text{ s}^{-1}$) for 30–40 min prior to analysis. The
 light and cuvette chamber were constantly flushed with dry nitrogen gas to avoid con-
 densation on the exterior of the cuvette due to the temperature difference between
 the cold seawater and the laboratory air. Minimal (F_0) and maximal (F_m) fluorescence
 and the effective absorption cross section (σ_{PSII}) were calculated from each single
 turnover (ST) saturation curve. The maximum quantum yield efficiency for PSII (F_v/F_m)
 was calculated (Genty et al., 1989) by normalizing F_m by the difference between the
 fluorescence at saturation (F_m) and the minimum fluorescence (F_0):

$$\Phi_{\text{PSII}}^{\text{max}} = \frac{F_m - F_0}{F_m} = \frac{F_v}{F_m} \quad (2)$$

Samples for taxon-specific pigments (600–1000 mL) were filtered under low vacuum
 onto GF/F filters and frozen in liquid nitrogen until analysis using high performance liq-
 uid chromatography (HPLC). An automated Hewlett Packard 1100 HPLC system was
 used to separate pigments with a reverse-phase Waters Symmetry C-8 column and
 a solvent gradient containing methanol, aqueous pyridine, acetone, and acetonitrile
 (Zapata et al., 2000; DiTullio and Geesey, 2002). A diode array detector was used to
 record pigment spectra between the wavelengths 350 and 600 nm, as well as continu-
 ous chromatograms at 410, 440, and 455 nm every 5 s, and chl-*a* and *c* were quantified
 with an HP 1046A fluorescence detector (excitation 421 nm, emission 666 nm). Unial-

gal laboratory cultures with the appropriate pigments were used to generate purified pigment standards for system calibration (DiTullio and Geesey, 2002).

5 Total diatom abundance and community composition was determined for samples preserved with 1% glutaraldehyde (final concentration) and stored in the dark at 4°C until analysis. Twenty-five mL samples were settled in Utermöhl chambers for at least 18 h and enumerated using light microscopy at 400× magnification with a Zeiss Axiovert S100 inverted microscope (Utermöhl, 1958). Diatoms were identified to the genus level. Total microzooplankton abundance and community composition was determined for samples preserved with 10% acid Lugol's solution (final concentration) and stored in the dark at room temperature until analysis (Thronsen, 1978). One hundred mL samples were settled in Utermöhl chambers for at least 18 h and enumerated using light microscopy at 200× magnification. Microzooplankton were identified to the genus level. The use of acid Lugol's solution obscured chl-*a* fluorescence and made it impossible to distinguish phototrophic from heterotrophic dinoflagellates based on autofluorescence. However, certain heterotrophic dinoflagellates such as *Protoperdinium* and *Gyrodinium* could be identified based on morphology, and were thus included in the counts.

2.3 Dissolved and particulate nutrients and dissolved iron

20 Samples for dissolved nutrients were 0.2 µm filtered and immediately analyzed using a Lachat QuikChem 8000 Flow Injection Analysis System. Total dissolved Fe concentrations were filtered through 0.4 µm track-etched polycarbonate filters (Nuclepore Whatman) and acidified to pH 1.7 with 4 mL⁻¹ sub-boiled distilled HCl. Total dissolved Fe was measured using adsorptive cathodic stripping voltammetry (ACSV) (Buck et al., 2007). The voltammetric system consisted of Princeton Applied Research (PAR) 303A interfaced with a computer-controlled AutolabII potentiostat/galvanostat (Eco Chemie). The working electrode was a "large" mercury drop (2.8 mm²), the reference electrode was Ag:saturated AgCl, saturated KCl, and the counterelectrode was a platinum wire.

25 Acidified samples were microwaved 2×15 s at 1100 W to release dissolved Fe from

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ambient organic ligands (Bruland et al., 2005), neutralized once with cool 1 mol L⁻¹ QNH₄OH, and buffered to pH 8.2 with the borate buffer. Once buffered, Fe and SA additions were made and following ACSV analysis Fe concentrations were determined from a linear regression of the standard addition curve. The detection limit for the ACSV method is 0.02 nmol L⁻¹, calculated from three times the standard deviation of a 0.05 nmol L⁻¹ Fe addition, as no peak is observed in either Milli-Q or ultraviolet (UV)-oxidized seawater (from which trace metals and metal-chelating organic ligands are removed from seawater; (Donat and Bruland, 1988) at deposition times of up to 600 s. Deposition times for sample analyses here were between 60 and 400 s, depending on ambient Fe concentrations.

Total particulate carbon and nitrogen samples (150–250 mL) were filtered at low vacuum onto GF/F filters that had been precombusted at 450°C for 2 h. Filters were then dried at 60°C. Samples were analyzed with a Finnigan Delta Plus mass spectrometer and a Carlo Erba NA1500 elemental analyzer/Conflo II. Elemental compositions were measured using the mass 44 beam intensity (V) on the Delta Plus and calibrated against the mass 44 beam intensity of at least five standards also analyzed during each run of 40 samples. Total particulate phosphorus samples (100–200 mL) were gently filtered onto precombusted GF/F filters and rinsed with 2 ml 0.12 mol L⁻¹ Na₂SO₄. Filters were placed overnight in precombusted (450°C, overnight) 20 mL borosilicate glass scintillation vials with 2 mL 0.017 mol L⁻¹ MnSO₄. Vials were covered with aluminum foil, dried at 95°C and stored dessicated until analysis. Total particulate phosphorus content was determined as follows: vials and filters were combusted at 450°C for two h, cooled, and 5 mL 0.2 mol L⁻¹ HCl was added to each vial. Vials were capped and heated to 80°C for 30 min to digest particulate organic phosphorus into inorganic phosphate, and digested samples were analyzed using the standard molybdate colorimetric method (Solorzano and Sharp, 1980). Biogenic silica samples (100–250 mL) were gently filtered onto 0.6 µm polycarbonate filters, dried at 60°C and stored at room temperature until analysis. Samples were analyzed according to Brzezinski and Nelson (1995).

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.4 Biogenic sulfur

Samples for dimethylsulphide (DMS) and dimethylsulphoniopropionate (DMSP) were collected using the methodology of Kiene and Slezak (2006). A small volume of each sample (≤ 15 mL) was gravity filtered through a 25 mm Whatman GF/F filter and the filtrate collected for the immediate determination of DMS. A second gravity-filtered sample (≤ 20 mL) was collected and the filtrate preserved with 50% sulfuric acid for the determination of dissolved DMSP (DMSP_d). Aliquots of the unfiltered samples (≤ 20 mL) were preserved with 50% sulfuric acid (100 μ L per 10 mL of sample) for the determination of total DMSP (DMSP_t). All DMSP samples were stored at 0°C until they could be analyzed, which was within three days of their collection. Upon analysis, the DMSP samples were base-hydrolyzed (2 mol L⁻¹ sodium hydroxide) and measured as DMS using a cryogenic purge and trap systems coupled to either a Hewlett-Packard 5890 Series II gas chromatograph and an Agilent 6890 gas chromatograph that were fitted with flame photometric detectors (DiTullio and Smith, 1995). Both systems were calibrated using constant-temperature DMS permeation devices (Vici Metronics) and DMSP standards (Research Plus Inc). Particulate DMSP (DMSP_p) was calculated as the difference between DMSP_t and DMSP_d.

2.5 Statistical analyses

Main effects of iron and temperature individually and interactions between these two variables were compared across treatments using a modified two-way ANOVA analysis based on a percentile bootstrap method (Wilcox, 2003). This method was chosen over the classic two-way ANOVA because it does not have the assumptions of normality or homoscedasticity and generally has higher power. All tests were done at the $\alpha=0.05$ level. The statistical software program R version 2.5 was used to perform all tests (<http://www.r-project.org>). Diatom and microzooplankton community-level analyses were completed using multivariate statistics. The ecological statistical software program PRIMER v. 6 was used to perform all multivariate analyses (Clarke and Gor-

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ley, 2006; Clarke and Warwick, 2001). A square root transformation was applied to data to slightly increase the contribution of rarer species to the measure of similarity among samples. The Bray-Curtis coefficient was used to generate the similarity matrix. Non-metric multidimensional scaling (MDS) was used to visualize high-level sample relationships in two dimensions. The relationships between individual samples based on whole communities within each were compared using this technique. Each symbol on an MDS plot represents an entire community (in this case, either diatoms or microzooplankton) from a single sample. Two samples with relatively similar assemblages (e.g., replicates from a single treatment at a single time) have symbols located very close to each other on a two-dimensional MDS plot. Two samples with relatively different communities (e.g., samples from different treatments or samples from different times) have symbols located relatively far away from each other on the same two-dimensional MDS plot. The MDS plots illustrate relative differences between samples within a group. Since the distances plotted are relative and not absolute, the axes of an MDS plot are without absolute scale. The significance of treatment effects on community composition were determined using the two-way crossed ANOSIM test (Clarke and Green, 1988). This procedure uses ranked similarities and a permutation test to compare the overall similarity of samples assigned to groups based on individual variables (iron and temperature) to the overall similarity of samples between variable groups to determine the significance level of community composition differences between variables.

3 Results

3.1 Plankton assemblages

Total chlorophyll and the >20 µm chlorophyll size fraction both increased in all treatments over the course of the experiment (Fig. 1). Final day total chlorophyll was lowest in the control treatment, higher in the high temperature treatment, and higher still in the high iron treatment. However, total chlorophyll-*a* was by far highest in the high iron,

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

high temperature (combined treatment) (Fig. 1a). There were significant main effects for both variables and significant interaction between both variables on the final day of the experiment (all $p < 0.001$). On the final day of the experiment, the $>20\ \mu\text{m}$ size fraction constituted the majority of total chlorophyll, from 57% in the control treatment to 63 and 64% in the temperature and iron treatments, respectively, and 76% in the combined treatment (Fig. 1b).

Nanophytoplankton abundance was also significantly greater in all three experimental treatments over controls by the end of the experiment (all $p < 0.001$; Fig. 2a). Both temperature and iron treatments had significant but relatively minor main effects on nanophytoplankton abundance, and there were also significant interactions between these variables, with final abundances in the combined treatment more than double any other treatments. We also observed significant main effects of both variables and interactions between temperature and iron on total diatom abundance at the end of the experiment (Fig. 2b; $p < 0.001$). Final diatom abundance was lowest in controls, double the control abundance in the final high temperature treatment, and more than four times the control abundance in the final high iron treatment. Final diatom abundance in the combined treatment was more than triple that of any other treatments and was more than an order of magnitude greater than the abundance observed in the final control treatment.

The phytoplankton pigment data indicated that the final assemblage in all treatments was dominated by diatoms (as indicated by fucoxanthin; Fuco) rather than haptophytes such as *Phaeocystis antarctica* (as indicated by 19'-hexanoyloxyfucoxanthin; 19-hex; Fig. 3). Final ratios of 19-hex/chl-*a* declined from initial values in all treatments, and only were a fraction of final Fuco concentrations (both normalized to HPLC-based chlorophyll-*a*). Since Fuco pigments are found in both diatoms and haptophytes (but predominantly diatoms in the Ross Sea; DiTullio et al., 2003), and 19-hex pigments are primarily indicative of haptophytes, this result suggests that the major contributors to Fuco pigments in the phytoplankton community were diatoms. Elevated Fuco:Hex ratios were measured in the two iron treatments and were consistent with the decrease

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in the hex:chl-*a* observed in those treatments (Fig. 3c). These data suggest the preferential stimulation of diatoms following Fe addition. The significant increase in the Fuco:Hex treatment in the combined treatment is corroborated by microscope-based counts of total diatom abundance (Fig. 2). The diatom community itself changed significantly over the course of the experiment ($p=0.01$, Fig. 2c, Table 1). Samples significantly grouped according to either iron concentration or temperature based on whole community measurements of similarity ($p=0.01$).

Microscopic diatom cell counts showed a community shift from initial dominance by *Chaetoceros* sp. and *Nitzschia* sp., to a final assemblage largely dominated by *Cylindrotheca* sp. and *Thalassiosira* sp. in all treatments (Table 1). The large chain-forming, heavily-silicified diatoms *Chaetoceros* sp. and *Fragilariopsis* sp. were also relatively important contributors to total diatom numbers in the final samples. For all of these dominant genera, a general pattern was observed in which higher temperature alone increased their abundance significantly above the control values, and iron addition alone increased their numbers equally or even more. However, by far the highest final cell abundances for all four dominant species were seen in the combined high iron, high temperature treatment. For example, final *Cylindrotheca* sp. cell numbers in the combined treatment were 8, 16 and 70 times higher than in the high iron, high temperature, and control treatments, respectively (Table 1).

Photosynthetic efficiency (F_v/F_m) was measured on each day of the experiment and was consistently highest in the two high-iron treatments (Fig. 4). F_v/F_m increased steadily from days 0–3 from 0.44 to ~ 0.6 in both high iron treatments. F_v/F_m remained at this level for the remainder of the experiment in the high iron treatment. In the high iron, high temperature treatment, F_v/F_m peaked at day 3 and steadily declined to initial levels by the end of the experiment. The F_v/F_m in both treatments without added iron slowly and consistently declined over the course of the experiment to 0.33 on the final day. The effects of temperature and iron individually as well as the interaction between these two variables were significant on the final day of the experiment (all $p < 0.001$).

Photosynthesis versus irradiance (PE) curves were generated initially, on day 4, and

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

on the final day of the experiment (Table 2). Maximum biomass-normalized photosynthetic rates (P_{\max}^B) increased in the three experimental treatments between the initial community and mid-point of the experiment. Increases were larger in the two iron treatments, and by far the highest P_{\max}^B was observed in the combined treatment. P_{\max}^B declined between the midpoint and final day of the experiment in the three experimental treatments, but was still highest in the combined treatment on the final day. The initial slope of the P - E curves (α) increased in all four treatments by the middle of the experiment, suggesting photoacclimation to irradiance levels in the deckboard incubators higher than those experienced in situ. The highest α observed at the midpoint was in the control treatment. The α values also declined between the middle and end of the experiment. The highest α observed on the final day of the experiment was in the combined treatment, followed by the high temperature treatment, high iron treatment and then the control treatment.

Microzooplankton abundance followed different trends than those observed for the phytoplankton community (Fig. 5a). Final microzooplankton abundance was higher in the two low-iron treatments, with the highest abundance observed in the high temperature treatment. Microzooplankton abundances declined between the beginning of the experiment and day 4 in the two high-iron treatments, and ended significantly lower than in the two low-iron treatments ($p < 0.001$). All main effects and interactions for microzooplankton abundance were significant on the final day of the experiment ($p < 0.001$). Microzooplankton community composition was also significantly different across the four treatments (Fig. 5b). Microzooplankton communities grouped significantly according to both temperature and iron (both $p = 0.01$), indicating that communities at different temperatures and communities exposed to different iron concentrations were significantly different in their composition.

3.2 Dissolved and particulate nutrients

Dissolved nutrient concentrations declined with increased phytoplankton biomass in all treatments over the course of the experiment (Fig. 6). The greatest decreases in NO_3^- ,

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



PO_4^{3-} and SiOH_4 all occurred in the combined treatment by the final day of the experiment. There were significant main effects and interactions for temperature and iron on all three final day dissolved nutrient concentrations ($p < 0.001$). Total dissolved iron concentrations were considerably lower in all treatments on the final day of the experiment than on the initial day, including the two treatments enriched with 1 nM iron at the beginning of the experiment (Fig. 6h). Final dissolved iron concentrations were lower in the control and high temperature treatments than the high iron and combined treatments, but since replicates were combined to obtain sufficient volume for the measurements, we were unable to determine if these differences were significant. The largest increases in particulate nutrient concentrations consistently occurred in the combined treatment, although effects of single factors were also significant ($p < 0.001$). Interactive effects of temperature and iron were significant for all four particulate nutrients (carbon, nitrogen, phosphorus and biogenic silica).

Average production ratios were calculated for each treatment based on initial and final nutrient concentrations (Table 3). There were significant main effects observed for iron and significant interactions between variables observed for all production ratios calculated ($p < 0.0001$). The C:P and N:P ratios were significantly higher in the two treatments with added iron, especially the combined treatment ($p < 0.001$). BSi:C and BSi:N ratios on the other hand were higher in the two treatments without iron, especially the high temperature treatment ($p < 0.001$). C:N and BSi:P ratios were similar in all the treatments but slightly lower in the high iron bottles. Significant interactive effects were observed for all ratios calculated (all $p < 0.001$).

3.3 Dimethylsulfide and dimethylsulfoniopropionate

DMS concentrations showed only relatively small changes between the start of the experiment and day 4. By the final day, DMS levels in the high temperature and combined treatments were approximately 3- and 4-fold higher than the control (Fig. 7a). Individually, both variables had a significant effect on DMS levels on the final day and a sig-

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

nificant interactive effect between the two variables was also observed (all $p < 0.001$). DMSP_d levels in all treatments were slightly lower at the end of the experiment than the initial concentration, with the combined treatment showing the greatest decline followed by the high temperature treatment (Fig. 7b). However, none of the changes in DMSP_d concentrations were significant.

DMSP_p concentrations increased throughout the course of the incubation with largest increases observed between days 4 and 7 (Fig. 7c). DMSP_p levels were highest in the combined treatment followed by high iron and high temperature treatments. In all cases, the changes in the variables had significant effects ($p < 0.001$) on DMSP_p levels both individually and synergistically. With respect to DMSP_p:chl-*a* ratios, while no change was observed in the control or high temperature treatments relative to starting conditions (Fig. 7d), there was a significant decrease in the high iron treatment and a synergistic decrease in the combined treatment on the final day (both $p < 0.001$). In the case of DMSP_p:PON ratios, values in the high temperature treatment were higher than control on both sampling days although temperature had no significant effect (Fig. 7e). Conversely, the addition of iron caused a significant decline ($p < 0.001$) in DMSP_p:PON ratios, with a significantly higher decrease observed in the high temperature treatment when normalized to PON.

A recent study (Kiene et al., 2007) has noted that samples preserved using the Kiene and Slezak (2006) method can underestimate DMSP_t when *Phaeocystis* sp. is present. Although it is not known quantitatively by how much the DMSP_p levels will be underestimated (del Valle et al., 2009; Rellinger et al., 2009), this finding could impact the DMSP_p results obtained during the experiment. While the phytoplankton community was dominated by diatoms and addition of iron caused the dominance of diatoms to increase (Fig. 3), the differences in DMSP_p concentrations cannot be considered significant as a result of presence of even a minor amount of *Phaeocystis* sp. This underestimation will also result in an underestimation of the decreases in DMSP_p:chl-*a* and DMSP_p:PON ratios. Moreover, the observed trends in DMS and DMSP_d also remain robust since these two parameters are unaffected by the preservation problem.

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4 Discussion

Previous studies of interactive effects of iron limitation with other factors influencing phytoplankton assemblages have focused largely on co-limitation effects with macro- and micronutrients (Bertrand et al., 2007; Leblanc et al., 2005) or light intensity (Boyd et al., 2001; Sunda and Huntsman, 1997). Temperature has also been recognized to play an important role in setting upper limits on physiological rate processes in plankton (Eppley, 1972; Goldman and Carpenter, 1974), and the interactions of temperature with other environmental variables such as rising $p\text{CO}_2$, altered irradiance and nutrient utilization have been examined in a few field and lab studies (Feng et al., 2009; Goldman, 1979; Hare et al., 2007; Lomas and Glibert, 1999; Rose et al., 2009). Temperature has also been hypothesized to constrain growth of heterotrophic protists relative to phototrophic protists at the extreme low temperatures characteristic of polar regions (Rose and Caron, 2007).

Warmer temperatures have been shown to cause large dominance shifts in experimentally manipulated plankton communities from other high-latitude regimes such as the Bering Sea and the North Atlantic (Feng et al., 2009; Hare et al., 2007; Noiri et al., 2005; Rose et al., 2009). The combination of increased iron and temperature was observed to result in shifts in phytoplankton community composition during the SEEDS study in the western subarctic Pacific, but only at the highest temperature examined (18°C ; Noiri et al., 2005). Noiri et al. (2005) did not report interactive effects of temperature and iron on other physiological and biochemical factors, and did not examine potential changes in food web dynamics during their experiments.

We consistently observed nonlinear, synergistic effects of iron and temperature on phytoplankton abundance, phytoplankton physiology and nutrient drawdown in our experiments. These enhancement effects of both variables together were much greater than simply additive, and so could not have been predicted based on the effects of temperature and iron measured individually. The increase in nanophytoplankton abundance in the high iron, high temperature treatment was more than double the increase

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in either the high iron or high temperature treatments, and greater than the additive effects of those single-variable treatments combined. Similar results for total diatom abundance were observed, with the final day abundance in the high temperature, high iron treatment again far exceeding the additive effects of the two variables examined individually.

Increased temperature at high iron concentration appeared to also bring on nutrient limitation earlier in the experiment. F_v/F_m values were nearly identical for both high-iron treatments during the initial days of the experiment and rapidly increased to ~ 0.6 which is expected upon alleviation of iron limitation (Green et al., 1994). However, the high iron, high temperature treatment peaked on day 3 and consistently decreased for the remainder of the experiment, most likely due to the more rapid utilization of added iron and the earlier onset of iron limitation. This trend was not observed when temperatures were increased in the absence of iron addition; decreases in F_v/F_m in the high temperature treatment occurred at a nearly identical rate as those in the control treatment.

Determination of the exact physiological mechanisms responsible for these non-linear impacts of combined iron increases and warming will require further investigation. In general though, physiological iron/temperature synergisms should perhaps not be surprising. Iron plays a central role in many cellular catalytic reactions involved in pathways like photosynthesis and nitrogen assimilation (Hutchins and Fu, 2008), and many of these enzymatic reactions also have a recognized sensitivity to temperature (Geider and Osborne 1992). Since both iron availability and temperature are critical controls on many of the same biochemical pathways, iron/temperature “co-limitation” may be a common physiological feature of marine phytoplankton.

Our observations of iron stimulation of phytoplankton abundance and photosynthetic rates, as well as growth of large diatoms relative to other phytoplankton taxa, are consistent with previous results for bottle incubations and iron enrichment in polar waters with high silicic acid concentrations. Several mesoscale iron enrichment experiments have been conducted in Southern Ocean waters to date (Boyd et al., 2000; Coale

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

et al., 2004; Gervais et al., 2002). An immediate stimulation of F_v/F_m followed by a time-lagged increase in chlorophyll-*a* was observed during all three Antarctic iron enrichment experiments (SOIREE, EisenEx and the SoFex high-Si south patch). A taxonomic shift to large (microplankton-sized) diatoms was observed during SOIREE.

5 During EisenEx and the SoFex south patch, the initial community dominance by diatoms was unchanged by the addition of iron, although authors noted that high iron concentrations in EisenEx favored growth of microphytoplankton relative to nano- and picophytoplankton assemblages.

Our study also points to the possibility of interactive effects of iron and temperature on plankton community elemental stoichiometry (Fig. 6). These suggest that patterns and ratios of nutrient drawdown in the future Ross Sea could be altered by changes in both iron supply and temperature, in addition to other recognized factors such as shifts in dominant taxa (Arrigo et al., 1999). Additionally, both iron and temperature had individual effects on community stoichiometry. For instance, the decrease in Si:N uptake ratios observed in the south patch during SoFex (Coale et al., 2004) was similar to results for the high iron treatments obtained in our study, and have also been observed in other ecosystems (Hutchins and Bruland, 1998). Increasing temperature alone appeared to result in more heavily silicified diatom cells, with significant increases observed in BSi:C, BSi:N and BSi:P. Additionally, greater Si drawdown and BSi production were observed in the high temperature treatment relative to the high iron treatment, although both total chlorophyll and total diatom abundance were lower. This may have been a direct response to temperature but could have also been an indirect response to increased microzooplankton grazing, since microzooplankton abundances were highest in the high temperature treatment by the end of the experiment. Increased diatom cell wall silicification in response to grazing pressure has been documented in the laboratory (Pondaven et al., 2007), but not to our knowledge in natural plankton assemblages.

25 Similarly, the results of this study point to changes in the cycling of DMSP and DMS as result of increases in either iron or temperature, and more critically, a synergistic

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

effect due to both iron and temperature. The observed increases in DMSP_p most likely resulted from increases in biomass since the phytoplankton community structure showed shifts towards increased diatom dominance (or a decrease in DMSP-producing species). While the declines in DMSP_p:chl-*a* and DMSP_p:PON ratios also reflect this shift, it must be remembered that addition of iron would affect all three parameters individually. It would decrease DMSP_p due to a reduction in oxidative stress, alleviate any chlorosis that would result from iron limitation, and increase nitrogen uptake (Allen et al., 2008; Sunda et al., 2002).

Although increasing iron and temperature caused a slight decline in DMSP_d levels, those changes were not quantitatively equivalent to the observed increases in DMS. This observation suggests that the degradation of DMSP and, more importantly, the relative importance of the DMS versus non-DMS producing pathways will be affected by changing iron and temperature regimes in a complex non-linear manner. The changes in the both phytoplankton and microzooplankton communities seen during the experiment will influence changes in predator-prey interactions and the DMSP cycle (Strom et al., 2003; Wolfe and Steinke, 1996; Wolfe et al., 1997) as will variations in the presence of the DMSP-lyase (either microbial or in *Phaeocystis* sp.). Finally, although uncertainties exist as to whether increasing *p*CO₂ will influence DMS (Vogt et al., 2008; Wingenter et al., 2007), the results of this experiment clearly show that increases in iron and temperature will have an important impact on any climate feedback mechanism involving DMS.

Photosynthetic parameters responded to both iron and temperature, but the greatest response was seen in the experiment where both were increased (Table 2). Observed values were well within the range observed in other areas of the Antarctic, and similar to those found for *P. antarctica* and diatom-dominated assemblages in the Ross Sea (Shields and Smith, 2009). Maximum biomass-specific photosynthetic rates are generally considered to be influenced by both temperature and nutrients (e.g., Behrenfeld and Falkowski, 1997), and our results are consistent with the release of iron limitation and increased temperatures that result in increased photosynthetic efficiency. While

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the changes between initial and final values of P_{\max}^B are not exceptionally great, they do represent an increase of nearly 40%, which could give rise to substantial increases in growth rates relative to the unamended controls.

In contrast, the large changes in the microzooplankton community in our experiments were somewhat unexpected. Boyd et al. (2000) reported that ciliate abundances increased after iron enrichment during SOIREE, but that microzooplankton herbivory was primarily focused on the <20 μm size fraction. These authors suggested that microzooplankton were unable to effectively graze the large *Fragilariopsis kerguelensis* that was favored at high iron concentrations. Rates of microzooplankton herbivory have not been reported for EisenEx or SoFex.

It is unlikely that the initial decreases in microzooplankton abundance in the high iron treatments were due to direct inhibition of microzooplankton growth by elevated iron concentrations. The amount of iron added to these experiments was modest, and far below any conceivable toxicity threshold. Iron concentrations greater than 1 nM can occasionally occur naturally in the upper water column of the Ross Sea during ice melt in early spring (Sedwick and DiTullio, 1997). Instead, it seems more likely that the large changes in the diatom community composition observed over the course of the experiment resulted in an algal assemblage that was less able to be grazed by the initial microzooplankton community. The diatom communities in both low iron treatments remained diverse relative to the high iron treatments (Table 1). The communities in the high iron treatments shifted from dominance by *Nitzschia* sp. and *Chaetoceros* sp. to *Thalassiosira* sp. in the high iron treatment and *Cylindrotheca* sp. in the high iron, high temperature treatment. In particular, the very long and narrow morphology of the *Cylindrotheca* may have been unpalatable to microzooplankton.

The microzooplankton assemblage at the beginning of the experiment was relatively diverse, but dominated by small heterotrophic dinoflagellates such as *Oxyphysis* sp. and small species (~20 μm) within the genera *Protoperidinium* and *Gyrodinium*. Heterotrophic dinoflagellates such as *Protoperidinium* are capable of feeding on phytoplankton much larger than themselves using structures such as the pallium (Jacob-

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

son and Anderson, 1986), and these species may have been actively feeding on the *Nitzschia* sp. and *Chaetoceros* sp. that dominated the phytoplankton assemblage initially.

The effects of increased $p\text{CO}_2$ and temperature on polar ecosystems have been a major focus of research to date, but it is important to remember that other physico-chemical factors are also predicted to be affected by a changing climate. The likelihood of changes in iron concentrations in combination with increased temperature highlights the importance of multivariate experiments to identify potential interactive effects of multiple changing factors on future plankton assemblages. The experiments detailed here are an important first step in examining the potential interactive effects of temperature and iron concentration on phytoplankton assemblages in the Ross Sea, Antarctica. It is clear from our results that without concomitant increases in iron inputs, increased temperature alone may have a smaller effect on phytoplankton assemblages. Finally, the potential interactive effects of temperature and iron together indicate that relatively small inputs of iron could disproportionately benefit plankton assemblages growing in a warmer future coastal Antarctic ecosystem.

Acknowledgement. This project was supported by U.S. NSF grants ANT 0528715 to JMR, ANT 0741411, ANT 0741428 and OCE 0825319 to DAH, ANT 0338157 to WOS, ANT 0338097 to GRD, and ANT 0338350 to RBD. The authors would like to thank Dr. Zbigniew Kolber for providing the bench-top MBARI Fast Repetition Rate Fluorometer.

References

- Allen, A. E., LaRoche, J., Maheswari, U., Lommer, M., Schauer, N., Lopez, P. J., Finazzi, G., Fernie, A. R., and Bowler, C.: Whole-cell response of the pennate diatom *Phaeodactylum tricornutum* to iron starvation, Proc. Natl. Acad. Sci. USA, 105, 10438–10443, 2008.
- Alley, R., Berntsen, T., Bindoff, N. L., Chen, Z., Chidthaisong, A., Friedlingstein, P., Gregory, J., Hegerl, G., Heimann, M., Hewitson, B., Hoskins, B., Joos, F., Jouzel, J., Kattsov, V., Lohmann, U., Manning, M., Matsuno, T., Molina, M., Nicholls, N., Overpeck, J., Qin, D., Raga, G., Ramaswamy, V., Ren, J., Rusticucci, M., Solomon, S., Somerville, R., Stocker, T. F.,

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Stott, P., Stouffer, R. J., Whetton, P., Wood, R. A., and Wratt, D.: Climate change 2007: The physical science basis. Summary for policymakers, Intergovernmental Panel on Climate Change, Geneva, 2007.

Arrigo, K. R., Robinson, D. H., Worthen, D. L., Dunbar, R. B., DiTullio, G. R., VanWoert, M., and Lizotte, M. P.: Phytoplankton community structure and the drawdown of nutrients and CO₂ in the Southern Ocean, *Science*, 283, 365–367, 1999.

Arrigo, K. R., DiTullio, G. R., Dunbar, R. B., Robinson, D. H., VanWoert, M., Worthen, D. L., and Lizotte, M. P.: Phytoplankton taxonomic variability in nutrient utilization and primary production in the Ross Sea, *J. Geophys. Res.*, 105, 8827–8846, 2000.

Behrenfeld, M. J. and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll concentration, *Limnol. Oceanogr.*, 42, 1–20, 1997.

Bertrand, E. M., Saito, M. A., Rose, J. M., Riesselman, C. R., Lohan, M. C., Noble, A. E., Lee, P. A., and DiTullio, G. R.: Vitamin B₁₂ and iron colimitation of phytoplankton growth in the Ross Sea, *Limnol. Oceanogr.*, 52, 1079–1093, 2007.

Boyd, P. W., LaRoche, J., Gall, M., Frew, R. D., and McKay, R.: Role of iron, light and silicate in controlling algal biomass in subantarctic waters SE of New Zealand, *J. Geophys. Res.*, 104, 13391–13404, 1999.

Boyd, P. W., Watson, A. J., Law, C. S., Abraham, E. R., Trull, T., Murdoch, R., Bakker, D. C. E., Bowie, A. R., Buesseler, K. O., Chang, H., Charette, M., Croot, P., Downing, K., Frew, R., Gall, M., Hadfield, M., Hall, J., Harvey, M., Jameson, G., LaRoche, J., Liddicoat, M., Ling, R., Maldonado, M. T., McDay, R. M., Nodder, S., Pickmere, S., Pridmore, R., Rintoul, S., Safi, K., Sutton, P., Strzepek, R., Tanneberger, K., Turner, S., Waite, A., and Zeldis, J.: A mesoscale phytoplankton bloom in the polar Southern Ocean stimulated by iron fertilization, *Nature*, 407, 695–702, 2000.

Boyd, P. W., Crossley, C., DiTullio, G. R., Griffiths, F. B., Hutchins, D. A., Queguiner, B., Sedwick, P. N., and Trull, T. W.: Control of phytoplankton growth by iron supply and irradiance in the subantarctic Southern Ocean: experimental results from the SAZ project, *J. Geophys. Res.*, 106, 31573–31583, 2001.

Boyd, P. W. and Doney, S. C.: Modelling regional responses by marine pelagic ecosystems to global climate change, *Geophys. Res. Lett.*, 29, 1806, doi:10.1029/2001GL014130, 2002.

Bruland, K. W., Rue, E. L., Smith, G. J., and DiTullio, G. R.: Iron, macronutrients and diatom blooms in the Peru upwelling regime: brown and blue waters of Peru, *Mar. Chem.*, 93, 81–103, 2005.

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Brzezinski, M. A. and Nelson, D. M.: The annual silica cycle in the Sargasso Sea near Bermuda, *Deep-Sea Res. I*, 42, 1215–1237, 1995.
- Buck, K. N., Lohan, M. C., Berger, C. J. M., and Bruland, K. W.: Dissolved iron speciation in two distinct river plumes and an estuary: Implications for riverine iron supply, *Limnol. Oceanogr.*, 52, 843–855, 2007.
- Campbell, L.: Flow cytometric analysis of autotrophic picoplankton, in: *Methods in Microbiology*, Academic Press, San Diego, CA, 317–341, 2001.
- Caron, D. A., Dennett, M. R., Lonsdale, D. J., Moran, D. M., and Shalapyonok, L.: Microzooplankton herbivory in the Ross Sea, Antarctica, *Deep-Sea Res. II*, 47, 3249–3272, 2000.
- Clarke, K. R. and Green, R. H.: Statistical design and analysis for a “biological effects” study, *Mar. Ecol. Prog. Ser.*, 46, 213–226, 1988.
- Clarke, K. R. and Warwick, R. M.: Change in marine communities: an approach to statistical analysis and interpretation, 2nd edition, Plymouth Marine Laboratory, Plymouth, UK, 2001.
- Clarke, K. R. and Gorley, R. N.: Primer v6: user manual/tutorial, Primer-E Ltd, Plymouth, 190 pp., 2006.
- Coale, K. H., Wang, X., Tanner, S. J., and Johnson, K. S.: Phytoplankton growth and biological response to iron and zinc addition in the Ross Sea and Antarctic Circumpolar Current along 170° W, *Deep-Sea Res. II*, 50, 635–653, 2003.
- Coale, K. H., Johnson, K. S., Chavez, F. P., Buesseler, K. O., Barber, R. T., Brzezinski, M. A., Cochlan, W. P., Millero, F. J., Falkowski, P. G., Bauer, J. E., Wanninkhof, R. H., Kudela, R. M., Altabet, M. A., Hales, B. E., Takahashi, T., Landry, M. R., Bidigare, R. R., Wang, X., Chase, Z., Strutton, P. G., Friederich, G. E., Gorbunov, M. Y., Lance, V. P., Hilting, A. K., Hiscock, M. R., Demarest, M., Hiscock, W. T., Sullivan, K. F., Tanner, S. J., Gordon, R. M., Hunter, C. N., Elrod, V. A., Fitzwater, S. E., Jones, J. L., Tozzi, S., Koblizek, M., Roberts, A. E., Herndon, J., Brewster, J., Ladizinsky, N., Smith, G., Cooper, D., Timothy, D., Brown, S. L., Selph, K. E., Sheridan, C. C., Twining, B. S., and Johnson, Z. I.: Southern Ocean iron enrichment experiment: carbon cycling in high- and low-Si waters, *Science*, 304, 408–414, 2004.
- De Baar, H. J. W., Buma, A. G. J., Nolting, R. F., Cadée, G. C., Jacques, G., and Treguer, P. J.: On iron limitation of the Southern Ocean: experimental observations in the Weddell and Scotia Seas, *Mar. Ecol. Prog. Ser.*, 65, 105–122, 1990.
- del Valle, D. A., Kieber, D. J., Toole, D. A., Bisgrove, J., and Kiene, R. P.: Dissolved DMSO production via biological and photochemical oxidation of dissolved DMS in the Ross Sea, Antarctica, *Deep-Sea Res. I*, 56, 166–177, 2009.

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Dennett, M. R., Mathot, S., Caron, D. A., Smith Jr., W. O., and Lonsdale, D. J.: Abundance and distribution of phototrophic and heterotrophic nano- and microplankton in the southern Ross Sea, *Deep-Sea Res. II*, 48, 4019–4037, 2001.

DiTullio, G. R. and Smith Jr., W. O.: Relationship between dimethylsulfide and phytoplankton pigment concentrations in the Ross Sea, Antarctica, *Deep-Sea Res. I*, 42, 873–892, 1995.

DiTullio, G. R. and Smith Jr., W. O.: Spatial patterns in phytoplankton biomass and pigment distributions in the Ross Sea, *J. Geophys. Res.*, 101, 18467–18478, 1996.

Donat, J. R. and Bruland, K. W.: Direct determination of dissolved cobalt and nickel in seawater by differential pulse cathodic stripping voltammetry preceded by adsorptive collection of cyclohexane-1,2-dione dioxime complexes, *Anal. Chem.*, 60, 240–244, 1988.

Eppley, R. W.: Temperature and phytoplankton growth in the sea, *Fish Bull.*, 70, 1063–1085, 1972.

Feng, Y., Hare, C. E., Leblanc, K., DiTullio, G. R., Lee, P. A., Wilhelm, S. W., Sun, J., Rose, J. M., Nemcek, N., Benner, I., and Hutchins, D. A.: The effects of increased $p\text{CO}_2$ and temperature on the North Atlantic Spring Bloom: I. The phytoplankton community and biogeochemical response, in press, 2009.

Firme, G. F., Rue, E. L., Weeks, D. A., Bruland, K. W., and Hutchins, D. A.: Spatial and temporal variability in phytoplankton iron limitation along the California coast and consequences for Si, N, and C biogeochemistry, *Global Biogeochem. Cy.*, 17, 1016, doi:0.1029/2001GB001824, 2003.

Fitzwater, S. E., Johnson, K. S., Gordon, R. M., Coale, K. M., and Smith Jr., W. O.: Trace metal concentrations in the Ross Sea and their relationship with nutrients and phytoplankton growth, *Deep-Sea Res. II*, 47, 2000.

Garrison, D. L.: An overview of the abundance and role of protozooplankton in Antarctic waters, *J. Mar. Syst.*, 2, 317–331, 1991.

Garrison, D. L. and Gowing, M. M.: Protozooplankton, in: *Antarctic Microbiology*, Friedmann, E. I. (Ed.), Wiley-Liss, New York, 123–166, 1993.

Garrison, D. L., Gibson, A., Kunze, H., Gowing, M. M., Vickers, C. L., Mathot, S., and Bayre, R. C.: The Ross Sea Polynya Project: diatom- and *Phaeocystis*-dominated phytoplankton assemblages in the Ross Sea, Antarctica, 1994 and 1995, in: *Biogeochemistry of the Ross Sea*, edited by: DiTullio, G. R. and Dunbar, R. B., American Geophysical Union, Washington, DC, 53–76, 2003.

Genty, B., Briantais, J. M., and Baker, N. R.: The relationship between the quantum yield

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of photosynthetic electron transport and quenching of chlorophyll fluorescence, *Biochim. Biophys. Acta*, 990, 87–92, 1989.

Gervais, F., Riebesell, U., and Gorbunov, M. Y.: Changes in primary productivity and chlorophyll *a* in response to iron fertilization in the Southern Polar Frontal Zone, *Limnol. Oceanogr.*, 47, 1324–1335, 2002.

Goldman, J. C. and Carpenter, E. J.: A kinetic approach to the effect of temperature on algal growth, *Limnol. Oceanogr.*, 19, 756–766, 1974.

Goldman, J. C.: Temperature effects on steady-state growth, phosphorus uptake, and the chemical composition of a marine phytoplankton, *Microb. Ecol.*, 5, 153–166, 1979.

Green, R. M., Kolber, Z. S., Swift, D. G., Tindale, N. W., and Falkowski, P. G.: Physiological limitation of phytoplankton photosynthesis in the Eastern Equatorial Pacific determined from variability in the quantum yield of fluorescence, *Limnol. Oceanogr.*, 39, 1061–1074, 1994.

Haberman, K. L., Quetin, L. B., and Ross, R. M.: Diet of the Antarctic krill (*Euphausia superba* Dana): I. Comparisons of grazing on *Phaeocystis antarctica* (Karsten) and *Thalassiosira antarctica* (Comber), *J. Exp. Mar. Biol. Ecol.*, 283, 79–95, 2003.

Haberman, K. L., Ross, R. M., and Quetin, L. B.: Diet of the Antarctic krill (*Euphausia superba* Dana): II. Selective grazing in mixed phytoplankton assemblages, *J. Exp. Mar. Biol. Ecol.*, 283, 97–113, 2003.

Hare, C. E., DiTullio, G. R., Trick, C. G., Wilhelm, S. W., Bruland, K. W., Rue, E. L., and Hutchins, D. A.: Phytoplankton community structure changes following simulated upwelled iron inputs in the Peru upwelling region, *Aquat. Microb. Ecol.*, 38, 269–282, 2005.

Hare, C. E., Leblanc, K., DiTullio, G. R., Kudela, R. M., Zhang, Y., Lee, P. A., Riseman, S. F., and Hutchins, D. A.: Consequences of increased temperature and CO₂ for phytoplankton community structure in the Bering Sea, *Mar. Ecol. Prog. Ser.*, 352, 9–16, doi:10.3354/meps07182, 2007.

Hutchins, D. A. and Bruland, K. W.: Iron-limited diatom growth and Si:N uptake ratios in a coastal upwelling regime, *Nature*, 393, 561–564, 1998.

Hutchins, D. A., Campbell, B. J., Cottrell, M. T., Takeda, S., and Cary, S. C.: Response of marine bacterial community composition to iron additions in three iron-limited regimes, *Limnol. Oceanogr.*, 46, 1535–1545, 2001.

Hutchins, D. A., Hare, C. E., Weaver, R. S., Zhang, Y., Firme, G. F., DiTullio, G. R., Alm, M. B., Riseman, S. F., Maucher, J. M., Geesey, M. E., Trick, C. G., Smith, G. J., Rue, E. L., Conn, J., and Bruland, K. W.: Phytoplankton iron limitation in the Humboldt Current and Peru Up-

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- welling, *Limnol. Oceanogr.*, 47, 997–1011, 2002.
- Jacobson, D. M. and Anderson, D. M.: Thecate heterotrophic dinoflagellates: feeding behaviors and mechanisms, *J. Phycol.*, 22, 249–258, 1986.
- Kiene, R. P. and Slezak, D.: Low dissolved DMSP concentrations in seawater revealed by small-volume gravity filtration and dialysis sampling, *Limnol. Oceanogr. Methods*, 4, 80–95, 2006.
- Kiene, R. P., Kieber, D. J., Slezak, D., Toole, D. A., del Valle, D. A., Bisgrove, J., Brinkley, J., and Rellinger, A.: Distribution of cycling of dimethylsulfide, dimethylsulfoniopropionate, and dimethylsulfoxide during spring and early summer in the Southern Ocean south of New Zealand, *Aquat. Sci.*, 69, 305–319, 2007.
- Kolber, Z. S., Barber, R. T., Coale, K. H., Fitzwater, S. E., Greene, R. M., Johnson, K. S., Lindley, S., and Falkowski, G. P.: Iron limitation of the phytoplankton photosynthesis in the equatorial Pacific Ocean, *Nature*, 371, 145–149, 1994.
- Leblanc, K., Hare, C. E., Boyd, P. W., Bruland, K. W., Sohst, B., Pickmere, S., Lohan, M. C., Buck, K., Ellwood, M., and Hutchins, D. A.: Fe and Zn effects on the Si cycle and diatom community structure in two contrasting high and low-silicate HNLC areas, *Deep-Sea Res. I*, 52, 1842–1864, 2005.
- Leventer, A. R. and Dunbar, R. B.: Factors affecting the distribution of diatoms and other algae in the Ross Sea, *J. Geophys. Res.*, 101, 18489–18500, 1996.
- Lomas, M. W., and Glibert, P. M.: Interactions between NH_4^+ and NO_3^- uptake and assimilation: comparison of diatoms and dinoflagellates at several growth temperatures, *Mar. Biol.*, 133, 541–551, 1999.
- Martin, J. H., Gordon, R. M., and Fitzwater, S. E.: Iron in Antarctic waters, *Nature*, 345, 156–158, 1990.
- Maucher, J. M. and DiTullio, G. R.: Chronic iron limitation in the Ross Sea: Flavodoxin:Ferredoxin ratios in the Ross Sea, in: *Biogeochemistry of the Ross Sea*, edited by: DiTullio, G. R. and Dunbar, R. B., Antarctic Research Series, AGU Antarctic Research Series, Washington DC, 209–220, 2003.
- Montes-Hugo, M. A., Doney, S. C., Ducklow, H. W., Fraser, W., Martinson, D., Stammerjohn, S. E., and Schofield, O.: Recent changes in phytoplankton communities associated with rapid regional climate change along the Western Antarctic Peninsula, *Science*, 323, 1470–1473, 2009.
- Noiri, Y., Kudo, I., Kiyosawa, H., Nishioka, J., and Tsuda, A.: Influence of iron and tempera-

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ture on growth, nutrient utilization ratios and phytoplankton species composition in the western subarctic Pacific Ocean during the SEEDS experiment, *Prog. Oceanogr.*, 64, 149–166, 2005.

- Olson, R. J., Sosik, H. M., Chekalyuk, A. M., and Shalapyonok, A.: Effects of iron enrichment on phytoplankton in the Southern Ocean during late summer: active fluorescence and flow cytometric analyses, *Deep-Sea Res. II*, 47, 3181–3200, 2000.
- Overpeck, J. T., Otto-Bliesner, B. L., and Miller, G. H.: Paleoclimatic evidence for future ice-sheet instability and rapid sea-level rise, *Science*, 311, 1747–1750, 2006.
- Pondaven, P., Gallinari, M., Chollet, S., Bucciarelli, E., Sarthou, G., Schultes, S., and Jean, F.: Grazing-induced changes in cell wall silicification in a marine diatom, *Protist*, 158, 21–28, 2007.
- Rellinger, A. N., Kiene, R. P., del Valle, D. A., Kieber, D. J., Slezak, D., Harada, H., Bisgrove, J., and Brinkley, J.: Occurrence and turnover of DMSP and DMS in deep waters of the Ross Sea, Antarctica, *Deep-Sea Res. I*, 56, 686–702, 2009.
- Rose, J. M. and Caron, D. A.: Does low temperature constrain the growth rates of heterotrophic protists? Evidence and implications for algal blooms in cold waters, *Limnol. Oceanogr.*, 52, 886–895, 2007.
- Rose, J. M., Feng, Y., Gobler, C. J., Gutierrez, R., Hare, C. E., Leblanc, K., and Hutchins, D. A.: The effects of increased $p\text{CO}_2$ and temperature on the North Atlantic Spring Bloom. II. Microzooplankton abundance and grazing, in press, 2009.
- Sarmiento, J. L., Hughes, T. M. C., Stouffer, R. J., and Manabe, S.: Simulated response of the ocean carbon cycle to anthropogenic climate warming, *Nature*, 393, 245–249, 1998.
- Scharek, R., van Leeuwe, M. A., and de Baar, H. J. W.: Responses of Southern Ocean phytoplankton to the addition of trace metals, *Deep-Sea Res. II*, 44, 209–227, 1997.
- Sedwick, P. N. and Ditullio, G. R.: Regulation of algal blooms in Antarctic shelf waters by the release of iron from melting sea ice, *Geophys. Res. Lett.*, 24, 2515–2518, 1997.
- Sedwick, P. N., DiTullio, G. R., and Mackey, D. J.: Iron and manganese in the Ross Sea, Antarctica: seasonal iron limitation in Antarctic shelf waters, *J. Geophys. Res.*, 105, 11321–11332, 2000.
- Shields, A. R. and Smith, W. O.: Size-fractionated photosynthesis/irradiance relationships during *Phaeocystis antarctica*-dominated blooms in the Ross Sea, Antarctica, *J. Plankton Res.*, 31, 701–712, 2009.
- Smith Jr., W. O., Marra, J., Hiscock, M. R., and Barber, R. T.: The seasonal cycle of phyto-

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

plankton biomass and primary productivity in the Ross Sea, Antarctica, Deep-Sea Res. II, 47, 3119–3140, 2000.

Smith Jr., W. O., Dennett, M. R., Mathot, S., and Caron, D. A.: The temporal dynamics of the flagellated and colonial stages of *Phaeocystis antarctica* in the Ross Sea, Deep-Sea Res. II, 50, 605–617, 2003.

Solorzano, L. and Sharp, J. H.: Determination of total dissolved phosphorus and particulate phosphorus in natural waters, Limnol. Oceanogr., 25, 756–760, 1980.

Strom, S., Wolfe, G., Holmes, J., Stecher, H., Shimeneck, C., Lambert, S., and Moreno, E.: Chemical defense in the microplankton, I. Feeding and growth rates of heterotrophic protists on the DMS-producing phytoplankter *Emiliana huxleyi*, Limnol. Oceanogr., 48, 217–229, 2003.

Sunda, W. G. and Huntsman, S. A.: Interrelated influence of iron, light, and cell size on growth of marine phytoplankton, Nature, 390, 389–392, 1997.

Sunda, W. G., Kieber, D. J., Kiene, R. P., and Huntsman, S. A.: An antioxidant function for DMSP and DMS in marine algae, Nature, 418, 317–320, 2002.

Takeda, S.: Influence of iron availability on nutrient consumption ratio of diatoms in oceanic waters, Nature, 393, 774–777, 1998.

Thronsen, J.: Preservation and storage, in: Phytoplankton Manual, edited by: Sournia, A., UNESCO, Paris, 69–74, 1978.

Utermöhl, H.: Zur Vervollkommnung der quantitativen Phytoplankton-Methodik, Mitt. Int. Ver. Limnol., 9, 1–38, 1958.

van Hilst, C. M. and Smith, W. O.: Photo synthesis/irradiance relationships in the Ross Sea, Antarctica, and their control by phytoplankton assemblage composition and environmental factors, Mar. Ecol. Prog. Ser., 226, 1–12, 2002.

Vaughan, D. G., Marshall, G. J., Connolley, W. M., Parkinson, C., Mulvaney, R., Hodgson, D. A., King, J. C., Pudsey, C. J., and Turner, J.: Recent rapid regional climate warming on the Antarctic Peninsula, Clim. Change, 60, 243–274, 2003.

Vogt, M., Steinke, M., Turner, S., Paulino, A., Meyerhöfer, M., Riebesell, U., LeQuéré, C., and Liss, P.: Dynamics of dimethylsulphoniopropionate and dimethylsulphide under different CO₂ concentrations during a mesocosm experiment, Biogeosciences, 5, 407–419, 2008, <http://www.biogeosciences.net/5/407/2008/>.

Welschmeyer, N. A.: Fluorometric analysis of chlorophyll *a* in the presence of chlorophyll *b* and phaeopigments, Limnol. Oceanogr., 39, 1985–1992, 1994.

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Wilcox, R. R.: Applying Contemporary Statistical Techniques, Academic Press, New York, 608 pp., 2003.

Wingenter, O. W., Haase, K. B., Zeigler, M., Blake, D. R., Rowland, F. S., Sive, B. C., Paulino, A., Thyrhaug, R., Larsen, A., Schulz, K., Meyerhofer, M., and Riebesell, U.: Unexpected consequences of increasing CO₂ and ocean acidity on marine production of DMS and CH₂Cl.
1. Potential climate impacts, Geophys Res. Lett., 34, L05710, doi:10.1029/2006GL028139, 2007.

Wolfe, G. V. and Steinke, M.: Grazing-activated production of dimethylsulfide (DMS) by two clones of *Emiliana huxleyi*, Limnol. Oceanogr., 41, 1151–1160, 1996.

10 Wolfe, G. V., Steinke, M., and Kirst, G. O.: Grazing-activated chemical defence in a unicellular marine alga, Nature, 387, 894–897, 1997.

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Phytoplankton community composition based on microscopic counts. All values are in cells mL⁻¹. Numbers in parentheses indicate one standard deviation.

	T0	Control	High iron	High temperature	High iron, high temp.
<i>Chaetoceros</i> sp.	5.8 (0.5)	100.0 (26.3)	101.0 (17.9)	99.4 (21.4)	309.7 (102.1)
<i>Pseudo-nitzschia</i> sp.	0.3 (0.3)	12.7 (12.1)	29.7 (14.0)	40.9 (16.5)	48.3 (37.7)
<i>Corethron</i> sp.	2.8 (1.3)	21.4 (3.2)	22.3 (4.9)	21.7 (10.4)	39.6 (17.6)
<i>Fragilariopsis</i> sp.	3.2 (1.7)	66.6 (22.4)	157.3 (15.1)	87.9 (23.5)	365.4 (86.4)
<i>Coscinodiscus</i> sp.	0.3 (0.4)	1.5 (1.1)	1.9 (3.2)	3.1 (1.4)	1.2 (2.1)
<i>Nitzschia</i> sp.	7.8 (3.3)	21.4 (10.7)	22.3 (6.7)	18.6 (4.8)	19.8 (11.4)
<i>Rhizosolenia</i> sp.	0.3 (0.2)	2.2 (1.1)	5.6 (3.2)	2.5 (0.5)	7.4 (3.7)
<i>Cylindrotheca</i> sp.	0.4 (0.1)	61.0 (33.3)	502.3 (117.5)	269.7 (81.8)	4261.1 (262.2)
<i>Silicoflagellate</i> sp.	0.2 (0.1)	0.0	0.6 (1.1)	2.2 (0.5)	1.2 (2.1)
<i>Thalassiosira</i> sp.	0.0	0.9 (1.6)	782.2 (35.9)	49.9 (12.3)	1056.6 (113.6)
<i>Asteromphalus</i> sp.	0.0	0.3 (0.5)	0.6 (1.1)	1.5 (2.7)	1.2 (2.1)

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Photosynthetic parameters as influenced by temperature and iron concentrations.
 P_{\max}^B : mg C (mg chl-*a*)⁻¹ h⁻¹; Alpha: mg C (mg chl-*a*)⁻¹ h⁻¹ (μmol quanta m⁻² s⁻¹)⁻¹.

Control		
Time	P_{\max}^B	Alpha
0	2.04	0.037
4	1.57	0.26
7	2.13	0.026
High iron		
Time	P_{\max}^B	Alpha
0	2.04	0.037
4	8.67	0.14
7	2.41	0.031
High temperature		
Time	P_{\max}^B	Alpha
0	2.04	0.037
4	3.13	0.14
7	2.80	0.047
High iron, high temperature		
Time	P_{\max}^B	Alpha
0	2.04	0.037
4	9.12	0.19
7	3.10	0.065

BGD

6, 5849–5889, 2009

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Table 3. Average production ratios calculated based on initial and final particulate nutrient concentrations. Numbers in parentheses indicate one standard deviation.

	Control	High iron	High temperature	High iron, high temp.
Δ POC/PON	7.58 (0.22)	6.72 (0.12)	7.69 (0.23)	7.73 (0.05)
Δ POC/POP	128 (14.8)	141 (6.6)	117 (9.3)	186 (5.6)
Δ PON/POP	17.0 (2.34)	20.9 (1.12)	15.2 (0.76)	24.1 (0.77)
Δ BSi/POC	0.17 (0.01)	0.12 (0.00)	0.21 (0.01)	0.13 (0.01)
Δ BSi/PON	1.25 (0.06)	0.80 (0.04)	1.62 (0.02)	1.00 (0.06)
Δ BSi/POP	21.1 (1.86)	16.7 (0.40)	24.6 (0.39)	24.1 (1.11)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.




Fig. 1. Size-fractionated chlorophyll-*a* concentrations. **(A)** $0.2\ \mu\text{m}$, total phytoplankton assemblage; **(B)** $>20\ \mu\text{m}$ size fraction. Error bars indicate one standard deviation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.




Fig. 2. Phytoplankton assemblages. **(A)** Nanophytoplankton abundance estimated by flow cytometry; **(B)** total diatom abundance estimated by light microscopy; **(C)** map of relative similarities among initial and final day diatom assemblages plotted by day and treatment using non-metric multidimensional scaling (MDS). The maps illustrate relative distances between samples and thus the axes are without absolute scale. (■) Initial phytoplankton assemblages, (●) control, (▼) high temperature, (○) high iron and (△) high temperature, high iron combined treatment. Error bars indicate one standard deviation.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




Fig. 3. Average **(A)** fuco and **(B)** 19'-hex pigment concentrations normalized to HPLC-derived chlorophyll-*a* concentration, **(C)** ratio of fuco:19'-hex. Error bars indicate one standard deviation.

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.




Fig. 4. Photosystem II efficiency (F_v/F_m) measured over the course of the experiment in all four treatments.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.




Fig. 5. Microzooplankton abundance and community composition. **(A)** Total abundance of microzooplankton over the course of the experiment. Error bars indicate one standard deviation. **(B)** MDS plot of relative similarities among initial and final day microzooplankton assemblages. (■) Initial microzooplankton assemblages, (●) control, (▼) high temperature, (○) high iron and (△) high temperature, high iron combined treatment.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.




Fig. 6. Dissolved and particulate nutrient concentrations over the course of the experiment. **(A)** NO_3^- , **(B)** particulate nitrogen, **(C)** PO_4^{3-} , **(D)** particulate phosphorus, **(E)** SiOH_4 , **(F)** biogenic silica, **(G)** particulate carbon, **(H)** total dissolved iron. Error bars indicate one standard deviation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Synergistic effects of iron and temperature on Antarctic plankton assemblages

J. M. Rose et al.




Fig. 7. Biogenic sulfur parameters over the course of the experiment. Average values for **(A)** DMS, **(B)** DMSP_d, **(C)** DMSP_p, and **(D)** DMSP_p:PON. Error bars indicate one standard deviation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion