

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

**The Arctic Ocean
marine carbon cycle**

N. R. Bates and
J. T. Mathis

The Arctic Ocean marine carbon cycle: evaluation of air-sea CO₂ exchanges, ocean acidification impacts and potential feedbacks

N. R. Bates¹ and J. T. Mathis²

¹Bermuda Institute of Ocean Sciences, Ferry Reach, Bermuda

²School of Fisheries and Ocean Sciences, University of Alaska, Fairbanks, Alaska, USA

Received: 18 June 2009 – Accepted: 21 June 2009 – Published: 9 July 2009

Correspondence to: N. R. Bates (nick.bates@bios.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

At present, although seasonal sea-ice cover mitigates atmosphere-ocean gas exchange, the Arctic Ocean takes up carbon dioxide (CO₂) on the order of -65 to -175 Tg C year⁻¹, contributing 5–14% to the global balance of CO₂ sinks and sources.

Because of this, the Arctic Ocean is an important influence on the global carbon cycle, with the marine carbon cycle and atmosphere-ocean CO₂ exchanges sensitive to Arctic Ocean and global climate change feedbacks. In the near-term, further sea-ice loss and increases in phytoplankton growth rates are expected to increase the uptake of CO₂ by Arctic surface waters, although mitigated somewhat by surface warming in the Arctic. Thus, the capacity of the Arctic Ocean to uptake CO₂ is expected to alter in response to environmental changes driven largely by climate. These changes are likely to continue to modify the physics, biogeochemistry, and ecology of the Arctic Ocean in ways that are not yet fully understood. In surface waters, sea-ice melt, river runoff, cooling and uptake of CO₂ through air-sea gas exchange combine to decrease the calcium carbonate (CaCO₃) mineral saturation states (Ω) of seawater that is counteracted by seasonal phytoplankton primary production (PP). Biological processes drive divergent trajectories for Ω in surface and subsurface waters of Arctic shelves with subsurface water experiencing undersaturation with respect to aragonite and calcite. Thus, in response to increased sea-ice loss, warming and enhanced phytoplankton PP, the benthic ecosystem of the Arctic shelves are expected to be negatively impacted by the biological amplification of ocean acidification. This in turn reduces the ability of many species to produce CaCO₃ shells or tests with profound implications for Arctic marine ecosystems.

1 Introduction

The Arctic Ocean is currently experiencing rapid environmental change due to natural and anthropogenic factors that include warming (e.g., Serreze et al., 2000; Polyakov et

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



al., 2002; Serreze and Francis, 2006; Overland et al., 2008), sea-ice loss (Cavaliere et al., 2003; Maslanik et al., 2007; Giles et al., 2008; Comiso et al., 2008), and other physical changes (Wu et al., 2005; McGuire et al., 2006) as well as biology and ecosystem structure changes (Arrigo et al., 2008; Pabi et al., 2008). These changes and feedbacks could have profound impacts on the Arctic Ocean marine carbon cycle and the importance of the Arctic for the global carbon cycle and the balance of carbon dioxide (CO₂) sinks and sources. In a recent review, Macdonald et al. (2009) compiled inorganic and organic carbon budgets of the Arctic Ocean. Here, we review the present understanding of the marine inorganic carbon cycle, air-sea CO₂ disequilibrium and rates of air-sea CO₂ gas exchange, the physical and biological processes that influence Arctic Ocean CO₂ sinks and sources, the impact of ocean acidification and the potential future drivers of change such as sea-ice loss, phytoplankton primary production (PP) and freshwater inputs. The geographic scope of this review encompasses the Arctic Ocean shelves (i.e., Barents, Kara, Laptev, East Siberian, Chukchi, Beaufort and Canadian Archipelago Seas) and central basin (i.e., Canada and Eurasian Basin), but does not extend to the gateways of the Arctic Ocean including the Greenland, Icelandic, Norwegian, and Bering Seas, and Hudson and Baffin Bays.

As context for this paper, a brief background on the physical and biogeochemical setting is presented in Sect. 2. Relevant information about historical sampling in the Arctic Ocean, methods for estimating air-sea CO₂ exchange rates, considerations about carbonate chemistry and relationships between shelf metabolism (net autotrophy versus net heterotrophy) and CO₂ sink/source issues are summarized in Sect. 3. With a synthesis of published results and data, the present status and magnitude of the Arctic Ocean CO₂ sink on the Arctic shelves and central basin is estimated, with predictions for near-term CO₂ flux rates in an era of rapid change (Sect. 4). The impacts of rapid change in the Arctic and the potential feedbacks on the marine inorganic carbon cycle and air-sea CO₂ exchange rate are also discussed in Sect. 5. Vulnerabilities of the Arctic marine carbon cycle relate to such factors as sea-ice loss and warming, biological and ecosystem changes, and hydrological/freshwater input changes. We also discuss

**The Arctic Ocean
marine carbon cycle**N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the impact of ocean acidification, and biology and physical processes on the marine carbon cycle and calcium carbonate (CaCO_3) mineral saturation states in the Arctic (Sect. 6).

2 The physical and biological setting of the Arctic Ocean

5 There are multiple influences on the marine carbon cycle and air-sea exchange of CO_2 in the Arctic Ocean. The inorganic carbon cycle is dominated by inter-ocean exchanges between the Pacific and Atlantic Oceans (Macdonald et al., 2009) and subsequent biogeochemical modifications and transformations of water while resident in the Arctic during transit between these oceans. In addition, river inputs of freshwater and materials, sea-ice production and melting, and atmosphere-ocean interaction and exchanges also have profound influence.

10 The relatively small Arctic Ocean ($\sim 10.7 \times 10^6 \text{ km}^2$) is almost completely landlocked except for the Bering Strait, Canadian Archipelago, Fram Strait and Norwegian Sea gateways that allow exchanges with the Pacific and Atlantic Oceans (Fig. 1). The relatively broad, generally shallow ($< 200 \text{ m}$ deep) and almost annular continental shelves surrounding the central basin comprise about 53% of the area of the Arctic Ocean, making it unique compared to other ocean basins. All of the Arctic continental shelves are highly varied, unique and difficult to characterize generically.

15 As a framework for this review, the Chukchi and Barents Seas can be characterized as “inflow” shelves (Carmack and Wassmann, 2006) where the import of nutrient-rich water from the Pacific and Atlantic Oceans, respectively, sustains high rates of seasonal phytoplankton PP or net community production (NCP). Other shelves such as the Kara, Laptev, East Siberian and Beaufort Seas constitute “interior shelves”, which are highly influenced by exchanges with other shelves, while the Canadian Archipelago represents an “outflow shelf” where Arctic water is exported to the Atlantic Ocean via Hudson and Baffin Bays. In the central basin of the Arctic Ocean, waters of the Canada Basin or Beaufort Gyre are separated from the Eurasian Basin by the surface trans-

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



polar drift, with inputs and outputs of Atlantic Ocean water through the Fram Strait and shelf-basin exchanges of water across the entire Arctic. Central basin waters are highly stratified vertically with distinct different physical and chemical properties of surface waters and subsurface halocline, Atlantic layer and deep-water layers, with deep waters having limited exchanges with the Atlantic Ocean.

The Arctic Ocean has an important role in the global freshwater cycle (e.g., Aagaard and Carmack, 1989; Wijffels et al., 1992; Woodgate and Aagaard, 2005) and Atlantic Ocean overturning circulation (e.g., Walsh and Chapman, 1990; Mysak et al., 1990; Hakkinen, 1993; Aagaard and Carmack, 1994; Wadley and Bigg, 2002). Surrounding the Arctic Ocean is an extensive land margin and watershed that contributes freshwater and material inputs such as inorganic nutrients and dissolved organic carbon (DOC) to the Arctic Ocean (Opsahl et al., 1999; Dittmar and Kattner, 2003; Benner et al., 2003; Guo et al., 2004). Siberian (Ob, Yenisey, Lena and Kolyma) and North American (Mackenzie, Yukon) rivers contribute significantly to land-ocean inputs (Peterson et al., 2002; McGuire et al., 2006; Serreze et al., 2006; Cooper et al., 2008). Finally, air-sea interaction controls the seasonal sea-ice advance and retreat, and interannual changes of sea-ice distributions and thickness, and sea-ice export from the Arctic Ocean (Carmack and Chapman, 2003). In wintertime, the Arctic is completely covered by sea-ice (except for polynyas and leads) with thick (3–7 m) multi-year ice occupying the interior of the central basin and thinner seasonal sea-ice (1–2 m) across the Arctic shelves and periphery (e.g., significant parts of the Bering Sea shelf; Hudson and Baffin Bays; Greenland, Iceland and Norwegian Sea) that melts and retreats towards the Pole during the summertime. While the seasonal production and melting of sea-ice contributes significantly to the physical structure and mixing of the Arctic, sea-ice melt and brine rejection during sea-ice formation also modifies the biogeochemical properties of surface and halocline layers (Steele and Boyd, 1998; Weingartner et al., 1998; Carmack and Chapman, 2003). The Arctic Ocean sea-ice cover is also particularly sensitive to interannual changes and variability of atmosphere-ocean-sea-ice forcing and feedbacks associated with climate variability such as the Arctic Oscillation (AO) and Arctic

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sea-Ice Oscillation (ASIO) (e.g., Rigor et al., 2002; Overland and Wang, 2005; Wang et al., 2005).

3 Considerations about sampling, carbonate chemistry, air-sea CO₂ exchange calculations and continental shelf CO₂ sink-sources issues

3.1 Marine inorganic carbon cycle sampling in the Arctic Ocean

Compared to many other open-ocean and coastal environments, relatively few studies of the marine inorganic carbon cycle and air-sea CO₂ exchange rates have been conducted in the Arctic. The harsh polar climate and difficult logistical support have limited most studies to opportunistic icebreaker surveys conducted on the Arctic shelves during the summertime sea-ice retreat with a couple of transpolar surveys across the deep basin (e.g., Arctic Ocean Section; Tucker and Cate, 1996; Anderson et al., 1998a; Jutterstrom and Anderson, 2005; Jones et al., 2008; Tanhua et al., 2009; Carbon in the North Atlantic (CARINA) datasets at <http://cdiac.ornl.gov/>). The absence of long-term ocean time-series (e.g., Winn et al., 1994; Bates, 2007; Santana-Casiano et al., 2007), repeat hydrographic surveys (e.g., Sabine et al., 2004; Feely et al., 2004) and underway shipboard surface observations (Takahashi et al., 2002, 2009) that have benefited understanding of seasonal and interannual change in other ocean basins and coastal ocean environments has considerably constrained our knowledge of the marine carbon cycle, and CO₂ sink and source terms in the Arctic Ocean.

3.2 Carbonate chemistry considerations

Studies of the marine inorganic carbon cycle in the Arctic have focused on measurements of one or more of the four directly observable carbonate system parameters, that is either dissolved inorganic carbon (DIC), total alkalinity (TA), partial pressure of CO₂ ($p\text{CO}_2$), or $p\text{H}$. Early historical data of TA and $p\text{H}$ were collected in the Arctic

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Yamamoto-Kawai et al., 2005) but have relatively limited use today due to inherent uncertainties associated with the analytical techniques used. More recently, improvement of analytical techniques, the advent of standardized methods (Dickson and Goyet, 1994; Dickson et al., 2007) and use of seawater standards and calibration have allowed precise and accurate data to be collected in the Arctic Ocean, and subsequently used for assessments of stocks, distributions and transformations of the marine inorganic carbon cycle.

Seawater carbonate chemistry is governed by a series of reactions, such that:



The four directly measureable carbonate system parameters are DIC, TA, $p\text{CO}_2$, and $p\text{H}$. For example, DIC is defined as (Dickson and Goyet, 1994; Dickson et al., 2007):

$$\text{DIC} = [\text{CO}_2^*] + [\text{HCO}_3^-] + [\text{CO}_3^{2-}] \quad (\text{R2})$$

where $[\text{CO}_2^*]$ represents the concentration of all unionized carbon dioxide, present either as H_2CO_3 or as CO_2 . The TA of seawater is defined as:

$$\text{TA} = [\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + [\text{B}(\text{OH})_4^-] + [\text{OH}^-] + [\text{HPO}_4^{2-}] + 2[\text{PO}_4^{3-}] + [\text{SiO}(\text{OH})_3^-] + [\text{HS}^-] + [\text{NH}_3] + [\text{minor species}] - [\text{H}^+] - [\text{HSO}_4^-] - [\text{HF}] - [\text{H}_3\text{PO}_4] - [\text{minor species}] \quad (\text{R3})$$

where $[\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + \text{B}(\text{OH})_4^-$ are the primary components of seawater TA. Other species have minor contribution to TA. DIC and TA are expressed as $\mu\text{moles kg}^{-1}$, $p\text{CO}_2$ is expressed as μatm , while $p\text{H}$ is negative log of the $[\text{H}^+]$ activity generally expressed on the total seawater $p\text{H}$ scale.

The complete seawater carbonic acid system (i.e., $[\text{CO}_2^*]$, $[\text{HCO}_3^-]$, $[\text{CO}_3^{2-}]$, $[\text{H}^+]$ and CaCO_3 mineral saturation states, Ω) can be calculated from a combination of two measured carbonate system parameters (i.e., DIC, TA, $p\text{CO}_2$ and $p\text{H}$). In this paper, we used the thermodynamic model of Zeebe and Wolf-Gladrow (2001), temperature and salinity data, the CO_2 solubility equations of Weiss (1974), and dissociation constants for carbonic acid (i.e., pK_1 and pK_2) of Mehrbach et al. (1973; as refit by Dickson

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Millero, 1997), borate (Dickson, 1990) and phosphate (Dickson and Goyet, 1994). It should be noted that the experimental determinations of carbonic acid dissociation constants (i.e., Mehrbach et al., 1973; Goyet and Poisson, 1989; Roy et al., 1993) have lower limits of -1°C or 0°C , and that the equations governing pK_1 and pK_2 have to be linearly extrapolated for seawater with temperatures less than -0°C . The difference in calculated seawater $p\text{CO}_2$ for different pK_1 and pK_2 determinations is relatively small ($<5\ \mu\text{atm}$) at temperatures less than 0°C (Bates, 2006).

CaCO_3 mineral production and dissolution is governed by the following chemical reaction:



The CaCO_3 mineral saturation state (Ω) is defined as the ion product of carbonate and calcium concentrations, thus:

$$\Omega_{\text{aragonite}} = [\text{Ca}^{2+}][\text{CO}_3^{2-}]/K_{sp}^* \quad (\text{R5})$$

The experimental solubility product (i.e., K_{sp}^*) of Mucci (1983) was used here. In general, CaCO_3 production and dissolution rates vary as a function of saturation state for aragonite ($\Omega_{\text{aragonite}}$) or calcite (Ω_{calcite}). For example, dissolution is generally favored when $\Omega_{\text{aragonite}}$ values are lower than 1, while aragonite formation generally occurs at $\Omega_{\text{aragonite}}$ values greater than 1.

3.3 Air-sea CO_2 exchange rate considerations

Different approaches have also been used to estimate rates of air-sea CO_2 exchange for the Arctic shelves or entire Arctic Ocean. These approaches include: (1) differences in atmospheric and seawater $p\text{CO}_2$ (either by direct measurements of $p\text{CO}_2$ or calculated from other carbonate parameters) that gives $\Delta p\text{CO}_2$ or air-sea CO_2 disequilibrium and, direction of air-sea CO_2 gas exchange (Olsen et al., 2003; Omar et al., 2007; Bates et al., 2006); (2) carbon mass balance assessments (e.g., Anderson et al., 1998; Fransson et al., 2001; Kaltin and Anderson, 2005; Macdonald et al., 2009); and;

(3) direct measurements of air-sea CO₂ exchange through eddy correlation techniques (Semiletov et al., 2006).

In the first approach, $\Delta p\text{CO}_2$ data combined with knowledge of gas transfer velocities commonly parameterized as a function of contemporaneous windspeed data (e.g., Wanninkhof, 1992; Wanninkhof and McGillis, 1999) provide estimates of air-sea CO₂ exchange rates. If surface water $p\text{CO}_2$ values are undersaturated with respect to the atmosphere, $\Delta p\text{CO}_2$ values are negative and the ocean has the potential to uptake CO₂ (i.e., sink of CO₂) with the magnitude of exchange set by gas transfer velocities. If surface water $p\text{CO}_2$ values are oversaturated with respect to the atmosphere, $\Delta p\text{CO}_2$ values are positive and the ocean has the potential to release CO₂ to the atmosphere (i.e., source of CO₂).

In the second approach, air-sea CO₂ exchange rates can be calculated by mass balance assessment of carbon stocks and inputs/outputs of carbon (both inorganic and organic), or temporal/depth changes of DIC that account for temperature and salinity variability and such processes as phytoplankton PP and respiration, vertical entrainment/detrainment and vertical diffusion. Both approaches have considerable uncertainties with caveats and assumptions used in the approaches. The third approach, using eddy correlation techniques, has only been used once in the Arctic Ocean (Semiletov et al., 2007). The results of these different approaches are discussed in more detail in Sect. 4 for individual shelves and the central basin of the Arctic Ocean.

3.4 Continental shelf net metabolism and CO₂ sink-sources issues

The CO₂ source or sink status and net metabolism of the coastal ocean has been highly debated over the last few decades. In early papers, it was first suggested that continental shelves sequestered significant amounts of CO₂ and that the global coastal ocean was net autotrophic (i.e., net consumption of CO₂) and a significant sink for atmospheric CO₂ (Walsh et al., 1985; Walsh, 1991). Smith and Hollibaugh (1993) subsequently argued that the nearshore coastal ocean was net heterotrophic (i.e., net production of CO₂) and offshore shelf coastal ocean neutral, and thus continen-

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tal shelves were potential sources of CO₂ to the atmosphere. Other studies have since suggested that the CO₂ source or sink status (Borges, 2005; Borges et al., 2005), and the metabolic status of continental shelves (i.e., net autotrophy versus net heterotrophy) is highly variable in time and space (e.g., Wollast, 1998; Gattuso et al., 1998).

5 The net metabolism (either net autotrophy or heterotrophy) of a continental shelf is influenced by a balance of transports, transformation and exchanges, such as: the input of organic matter (OM) from terrestrial sources, OM production and respiration in the coastal ocean, retention or export of OM from the shelf and, the net air-sea exchange of CO₂. The balance of these processes dictates the metabolic status (i.e., net
10 autotrophic or heterotrophic), but can also influence the potential CO₂ sink or source status of the continental shelf in question. The spatial and temporal variability of seawater *p*CO₂ on continental shelves is acted upon by a complex interplay of connected and quasi-independent physical and biological factors. For example, the physicochemical process of air-sea CO₂ exchange can proceed independently of the biological processes (e.g., production versus consumption of OM) that dictate the metabolism of the
15 coastal ocean. The variability of *p*CO₂ or Δ*p*CO₂ and separation of biological and physical processes can lead, for example, to a net heterotrophic continental shelf still being a sink for atmospheric CO₂ (Ducklow and McAllister, 2005). Ducklow and McAllister (2005) suggested that the global coastal ocean was net autotrophic, and thus a
20 potential sink for atmospheric CO₂.

Recent syntheses of coastal carbon cycle studies by Chen and Borges (2009) suggest that in general high-latitude and temperate continental shelf seas tend to be net annual sinks of atmospheric CO₂, while upwelling dominated, and subtropical to tropical coastal environments tend to release CO₂ to the atmosphere. In addition, localized shallow nearshore and river influenced areas of the continental shelves tend to be sources of CO₂ to the atmosphere, while the deeper, more expansive areas of the continental shelves tend to be oceanic sinks for atmospheric CO₂ (Chen and
25 Borges, 2009). In summary, Chen and Borges (2009) suggest that the polar continental shelves, for example, are likely to be potential sinks for atmospheric CO₂. However,

**The Arctic Ocean
marine carbon cycle**N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



this potential is also influenced by seasonal sea-ice cover, which act as a barrier to air-sea CO₂ gas exchange compared to open water conditions (Bates, 2006). Whether sea-ice facilitates the uptake or release of CO₂ is an area of active research with no definitive answer (Gosink et al., 1978; Semiletov et al., 2004; Delille et al., 2007). An additional complication is the recent finding that anaerobic microbial production of, and subsequent release of alkalinity can act to decrease seawater pCO₂, which in turn may enhance the uptake of atmospheric CO₂ onto continental shelves where sediments significantly interact with surface waters (Thomas et al., 2009).

4 Present state of knowledge about the marine inorganic carbon cycle and air-sea CO₂ exchange rates of the Arctic Ocean

4.1 Arctic ocean shelves: inflow shelves

4.1.1 Barents sea shelf

The Barents Sea comprises a broad and shallow continental shelf that is dominated by inflowing Atlantic water with minimal freshwater inputs. In the Barents Sea, warm and salty surface Atlantic Ocean water inflows from the Norwegian Sea into the Barents Sea via the Norwegian Atlantic Current (Skelvan et al., 1999; Omar et al., 2007), with ~2 Sv exchange between the Atlantic and Barents Sea (Gammelsrod et al., 2008). Cooling of surface water and brine rejection during sea-ice formation is subsequently thought to transform this surface water into dense subsurface water that is subsequently exported to the Kara Sea and central basin of the Arctic Ocean (Anderson et al., 1994; Fransson et al., 2001; Kaltin et al., 2002; Omar et al., 2007) thereby contributing to the physical and chemical properties of the Eurasian Basin halocline and deeper waters (Jones and Anderson, 1986).

In the earliest study of the inorganic carbon cycle in the Arctic Ocean, Kelley (1970) directly observed undersaturated seawater pCO₂ across the Barents Sea with ΔpCO₂

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



values ranging from ~ -50 to $-150 \mu\text{atm}$. More recently, other studies have shown seawater $p\text{CO}_2$ values of ~ 220 – $350 \mu\text{atm}$ with a seasonal minima occurring during the summertime (May to October) (Omar et al., 2003, 2007; Nakaoka et al., 2007). Given that the $\Delta p\text{CO}_2$ values ranged from ~ -20 to $-110 \mu\text{atm}$, the Barents Sea has a strong potential for significant uptake of CO_2 from the atmosphere.

Nakaoka et al. (2007) and Omar et al. (2007) estimated an annual ocean CO_2 uptake of $46 \pm 18 \text{ g C m}^{-2} \text{ yr}^{-1}$ and $51 \pm 8 \text{ g C m}^{-2} \text{ yr}^{-1}$, respectively, in the Barents Sea (Table 1), accounting for sea-ice cover during the wintertime. Using mass balance approaches, and accounting for the influence of phytoplankton primary production, freshwater inputs and mixing on vertical profiles of DIC and inorganic nutrients, Fransson et al. (2001) computed an annual ocean CO_2 uptake of $44 \pm 10 \text{ g C m}^{-2} \text{ yr}^{-1}$. Kaltin et al. (2002), using similar mass balance approaches applied to inorganic carbon data collected from a section across the Barents Sea estimated an ocean CO_2 uptake of $29 \pm 11 \text{ g C m}^{-2}$ for 3 months of the year (i.e., May to July). For comparison, Takahashi et al. (2009) estimated a climatological mean uptake of ~ 5 – $30 \text{ g C m}^{-2} \text{ yr}^{-1}$ from a very limited dataset of surface underway $p\text{CO}_2$ data, with the oceanic CO_2 sink decreasing in a general trend across the Barents Sea from the Atlantic Ocean inflow to the Kara Sea. Similar spatial trends were also observed by Kelley (1970), with $\Delta p\text{CO}_2$ values generally increasing from $\sim -50 \mu\text{atm}$ close to the coastline to approximately $-150 \mu\text{atm}$ at the outer extents of the Barents Sea shelf.

The seasonal changes in surface $p\text{CO}_2$ and air-sea CO_2 exchange rates for the Barents Sea have been largely attributed to seasonal changes in phytoplankton primary production, warming and cooling, and air-sea CO_2 fluxes. In a general sense, cooling of surface waters during transit from the Atlantic Ocean to the Barents Sea throughout the year acts to decrease seawater $p\text{CO}_2$ at a thermodynamic rate of $\sim 4.1\%$ per $^\circ\text{C}$ (Takahashi et al., 1993; Millero et al., 1995). However, during the sea-ice free summertime, surface seawater $p\text{CO}_2$ decreases by $\sim 100 \mu\text{atm}$ despite significant seasonal warming of the surface layer (~ 5 – 10°C) and ocean uptake of CO_2 through enhanced rates of air-sea CO_2 exchange. As Omar et al. (2007) suggest, these processes are more than

**The Arctic Ocean
marine carbon cycle**N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



compensated for by summertime photosynthetic uptake of CO_2 by phytoplankton PP in the euphotic zone that acts to decrease seawater DIC and $p\text{CO}_2$. Indeed, the annual rate of net PP for the Barents Sea ranges from $\sim <9\text{--}100 \text{ g C m}^{-2} \text{ yr}^{-1}$ (Sakshaug, 2004; Macdonald et al., 2009), with rates of phytoplankton PP greater than other Arctic shelves with the exception of the Chukchi Sea.

The seasonal fall and winter rebound of seawater $p\text{CO}_2$ to values just below equilibrium with the atmosphere observed by Nakaoka et al. (2007) likely results from a continued uptake of CO_2 through gas exchange during sea-ice formation and brine rejection (Anderson et al., 2004; Omar et al., 2005) and vertical entrainment by mixing with CO_2 rich subsurface waters. Despite the absence of supporting data, it is likely that subsurface $p\text{CO}_2$ increases seasonally from fall and winter from the seasonal remineralization of organic matter, produced during earlier summertime enhanced rates of phytoplankton primary production, as observed in other Arctic Ocean marine environments (e.g., Yager et al., 1995; Bates et al., 2009).

4.1.2 Chukchi Sea shelf

The wide and shallow Chukchi Sea shelf, with the shelf break extending hundreds of kilometers northward from the coast, represents a particularly extensive portion of the Arctic shelf system. The Bering Strait acts as the gateway for Pacific Ocean waters to enter the Arctic Ocean via the Chukchi Sea (Coachman et al., 1975; Bjork, 1989). The mean annual inflow of Pacific water at Bering Strait is $\sim 0.8 \text{ Sv}$, with higher transport during the open-water sea-ice free season (Roach et al., 1975; Woodgate et al., 2005a, b; Woodgate and Aagaard, 2005), with minor inputs to the Chukchi Sea from the East Siberian Sea shelf via the intermittent Siberian Coastal Current through Long Strait (Weingartner et al., 1999). In general, there is a northward transport of water across the shallow Chukchi Sea shelf, with four major outflows ($\sim 0.1\text{--}0.3 \text{ Sv}$ each; Woodgate et al., 2005a) from the Chukchi Sea into the Canada Basin and Beaufort Sea shelf through Long Strait, Central Channel, Herald Valley and Barrow Canyon (Paquette and Bourke, 1974; Weingartner et al., 1998, 2005; Woodgate et al., 2005b). As

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**The Arctic Ocean
marine carbon cycle**N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with other Arctic Ocean shelves, physical transformations and seasonal sea-ice cover play a major role in shaping the Chukchi Sea shelf water-masses and ecosystem. During the sea-ice covered season, watermasses on the Chukchi Sea shelf are confined to a small range of temperature-salinity space due to vertical homogenization by ventilation, brine rejection and convective mixing (Woodgate et al., 2005a). During the sea-ice free season, local sea-ice melt contributes to Pacific Ocean water to produce relatively warm, fresher surface mixed layer water (upper 0–~30 m, salinity typically <31; temperature >–1.5) with denser resident winter water remaining in the subsurface. Inflow of inorganic nutrient rich Pacific water supports high rates of phytoplankton PP or NCP on the Chukchi Sea shelf compared to other Arctic shelves (>300 g C m² y⁻¹ or 0.3–2.8 g C m² d⁻¹; Hameedi, 1978; Cota et al., 1996; Sambrotto et al., 1993; Hansell et al., 1993; Wheeler et al., 1996; Gosselin et al., 1997; Hill and Cota, 2005; Bates et al., 2005a; Mathis et al., 2009; Macdonald et al., 2009).

In early studies of the Chukchi Sea, Semiletov et al. (1999) observed undersaturated seawater $p\text{CO}_2$ (~200–350 μatm) during the sea-ice free period across the shelf with lowest values (~150–200 μatm) observed close to the ice edge at the northern shelf break. More recently, other studies have observed similar undersaturated seawater $p\text{CO}_2$ conditions from May to September (~150–350 μatm ; Pipko et al., 2002; Murata and Takizawa, 2003; Bates et al., 2005a; 2006; Bates, 2006; Chen and Gao, 2007). Extremely low $p\text{CO}_2$ conditions (<100 μatm) were observed in the surface layer in the vicinity of Barrow Canyon at the northern edge of the Chukchi Sea shelf in a location with very high rates of phytoplankton primary production (Bates et al., 2005a; Hill and Cota, 2005). Early season observations under near complete sea-ice cover also indicate that Chukchi Sea shelf “winter” surface waters were not as undersaturated with respect to the atmosphere ($\Delta p\text{CO}_2$ values of ~–20–60 μatm) compared to the summertime sea-ice free period (Bates, 2006). In contrast to surface waters, subsurface waters seawater $p\text{CO}_2$ values were highly oversaturated with respect to the atmosphere (Bates, 2006; Bates et al., 2009).

Estimates of the rates of air-sea CO_2 exchange during the sea-ice free period in

**The Arctic Ocean
marine carbon cycle**

N. R. Bates and
J. T. Mathis

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the summertime have ranged from $\sim 20\text{--}90$ mmol CO₂ m⁻² d⁻¹ (Wang et al., 2003; Murata and Takizawa, 2003; Bates, 2006). These studies indicate that the ocean uptake of CO₂ from the atmosphere is greater for the Chukchi Sea shelf than other Arctic shelves. However, in regions of the Chukchi Sea shelf where sea-ice cover remained high (>80%) during the summertime, air-sea CO₂ exchange rates were estimated to be generally low (<1 mmol CO₂ m⁻² d⁻¹; Bates, 2006). Similarly, wintertime air-sea CO₂ exchange rates during complete sea-ice cover were estimated to be minor (<1 mmol CO₂ m⁻² d⁻¹; Bates, 2006). The annual ocean CO₂ uptake for the Chukchi Sea shelf has been estimated at 24–108 g C m⁻² yr⁻¹ (Table 1; Bates, 2006). Similarly, using mass balance approaches, Kaltin and Anderson (2005) estimated that the annual ocean CO₂ uptake for the Chukchi Sea to be 86 ± 22 g C m⁻² yr⁻¹.

The seasonal changes in surface $p\text{CO}_2$ and strong sink for atmospheric CO₂ on the Chukchi Sea shelf have been largely attributed to cooling of surface waters during the northward transit of waters across the Chukchi Sea shelf (Murata and Takizawa, 2003) and high rates of summertime phytoplankton PP that acts to decrease seawater DIC and $p\text{CO}_2$ (Bates, 2006). As in the Barents Sea, the fall and winter rebound of seawater $p\text{CO}_2$ to values just below equilibrium with the atmosphere likely results from a continued uptake of CO₂ through gas exchange during sea-ice formation and brine rejection (Anderson et al., 2004; Omar et al., 2005) and vertical entrainment by mixing with CO₂ rich subsurface waters.

4.2 Arctic Ocean shelves: interior shelves

4.2.1 Siberian Sea shelves (i.e., Kara, Laptev and East Siberian Seas)

The Kara Sea shelf is controlled by exchanges with the Barents Sea and adjoining Eurasian Basin. Freshwater runoff from the Ob and Yenisey Rivers mix with more saline water of Atlantic origin, before entering the Laptev Sea or exported into the central basin of the Arctic Ocean (e.g., Anderson et al., 1994; Jones et al., 1998; Fransson

**The Arctic Ocean
marine carbon cycle**N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2001). The Laptev and East Siberian Sea shelves are subsequently controlled by exchanges with the adjoining Kara and Chukchi Seas, and major freshwater inputs to the Laptev Sea through the Lena River (e.g., Gordeev et al., 1996; Olsson and Anderson, 1997). On all of these shelves, seasonal sea-ice melt and formation, and brine rejection in coastal polynyas are important processes (e.g., Dmitrenko et al., 2008) with sea-ice production in this region occurring at the highest rate compared to other Arctic shelves. There is a general flow from the Laptev Sea across the East Siberian Sea and into the Chukchi Sea through the East Siberian Sea current (e.g., Weingartner, 1999; Nitishinsky et al., 2008), but circulation patterns can reverse with considerable transport from the East Siberian Sea shelf across the northeastern Laptev Sea shelf (e.g., Dmitrenko et al., 2008).

There have been few studies of inorganic carbon on the Siberian Sea shelves, with surface seawater $p\text{CO}_2$ and $\Delta p\text{CO}_2$ values highly variable over time and space. During the sea-ice free period, surface waters of the Kara Sea appear to be undersaturated with respect to the atmosphere. Semiletov et al. (1999) report a range of 273–405 μatm for surface seawater $p\text{CO}_2$ while Kelley (1970) reported $\Delta p\text{CO}_2$ values ranging from –30 to –100 μatm across much of the Kara Sea in 1967. In contrast, supersaturated surface seawater $p\text{CO}_2$ were observed nearshore and close to the outflow of the Ob and Yenisey Rivers (Kelley, 1970; Makkaveev, 1994; Semiletov et al., 1999). In the other Siberian Sea shelves, generally undersaturated surface seawater $p\text{CO}_2$ (216–~400 μatm) for the Laptev Sea and highly variable seawater $p\text{CO}_2$ conditions in the East Siberian Sea (~300–844 μatm) have been reported (Semiletov et al., 1999, 2007). On both shelves, supersaturated surface seawater $p\text{CO}_2$ values were observed for the Lena River that drains into the Laptev Sea and Kolyma Rivers that outflows to the East Siberian Sea shelf (Semiletov et al., 1999, 2007).

Fransson et al. (2001), using carbon mass balance approaches, computed a small annual ocean CO_2 uptake of 1 $\text{g C m}^{-2} \text{yr}^{-1}$ for the Kara Sea (Table 1). For the Laptev Sea, mass balance assessments integrated across the shelf indicate that this sea is a sink for atmospheric CO_2 during the sea-ice free summertime (~2.1 $\text{mmoles C m}^{-2} \text{d}^{-1}$;

July–September; Nitishinsky et al., 2008) despite supersaturated seawater $p\text{CO}_2$ conditions (i.e., sources of CO_2 to the atmosphere) in nearshore bays (Olsson and Anderson, 1997) and Lena River outflow (Semiletov et al., 1999, 2007). Direct eddy correlation measurements of air-sea CO_2 exchange in late summer (September) close to the multiyear sea-ice at the shelf break of the Laptev Sea yielded more contradictory results with air-sea CO_2 exchange rates varying from invasion (i.e., sink of CO_2 ; $\sim 1.7 \text{ mmol C m}^{-2} \text{ d}^{-1}$) to evasion (i.e., source of CO_2 ; $\sim 1.2 \text{ mmol C m}^{-2} \text{ d}^{-1}$; Semiletov et al., 2007). If we extrapolate the results of Nitishinsky et al. (2008) and assume a 100 d sea-ice free period, and negligible air-sea gas exchange during wintertime sea-ice cover, an annual ocean CO_2 uptake of $\sim 2.4 \text{ g C m}^{-2} \text{ yr}^{-1}$ is computed (Table 1). For the East Siberian Sea shelf, the sink or source terms are uncertain. Nitishinsky et al. (2008) estimated that the East Siberian Sea shelf is a small source of CO_2 to the atmosphere ($\sim 0.3 \text{ mmol C d}^{-1}$). However, Semiletov et al. (2007) suggest that the western area of the shelf is a source of CO_2 to the atmosphere ($1 \pm 1.6 \text{ mmol C d}^{-1}$, 2003; and $10.9 \pm 12.6 \text{ mmol C d}^{-1}$, 2004) while the eastern area, influenced by mixing with Pacific Ocean water from the Chukchi Sea shelf, is a sink for atmospheric CO_2 .

The above studies suggest that the Siberian Sea shelves range from minor sinks to minor sources of CO_2 to the atmosphere, with the causes uncertain at present. Nitishinsky et al. (2008) suggest that the drawdown of $p\text{CO}_2$ from summertime photosynthetic uptake more than compensates for seasonal warming, and mixing with river freshwater and high- CO_2 subsurface waters, all of which act to increase seawater $p\text{CO}_2$. Semiletov et al. (2007) showed highly supersaturated values for subsurface waters on all of the Siberian Sea shelves, presumably resulting from remineralization of organic matter introduced from the Siberian Rivers (e.g., Anderson et al., 1990; Cauwet and Sidorov, 1996; Kattner et al., 1999) and from vertical export of organic carbon despite relatively low rates of summertime phytoplankton primary production (compared to other Arctic shelves) in the euphotic zone ($\sim 6\text{--}12 \text{ g C m}^{-2} \text{ yr}^{-1}$; Macdonald et al., 2009). The East Siberian Sea shelf may be more highly influenced by these processes

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and hence more likely to be net heterotrophic and a source of CO₂ to the atmosphere. The Laptev Sea also has a permanent flaw lead polynya (Nitishinsky et al., 2008) that may contribute to ocean CO₂ uptake from the atmosphere during wintertime brine rejection and sea-ice formation as shown elsewhere (Anderson et al., 2004; Omar et al., 2005).

4.2.2 Beaufort Sea shelf

The Beaufort Sea shelf has a relatively narrow (~50 km) continental shelf in contrast to the adjacent Chukchi Sea shelf. The eastern Beaufort Sea shelf is heavily influenced by outflow from the Mackenzie River, and this input of freshwater and terrigenous input dominates the physical and chemical properties of the Beaufort Sea shelf (Carmack and Macdonald, 2002; Macdonald et al., 2009). Compared to the adjacent Chukchi Sea shelf, annual rates of phytoplankton PP for the Beaufort Sea shelf were relatively low (~7–17 g C m⁻² yr⁻¹; Macdonald et al., 2009), similar to other “interior” type shelves (Carmack and Wassman, 2006) such as the Siberian Sea shelves. In the adjoining Canada Basin, wind-driven, clockwise rotation of the Beaufort Gyre controls regional scale sea-ice and upper ocean circulation (Carmack and Chapman, 2003). Along the northern edge of the Chukchi and Beaufort Seas, a narrow eastward boundary current is present (Nikolopoulos et al., 2009) which is constrained to follow the isobaths along the shelf break due to strong dynamic controls (Winsor and Chapman, 2004; Spall, 2007). The generalized regional water circulation along the Beaufort Sea shelf is directed eastward with shelf waters principally renewed principally by Pacific Ocean waters that enter through Bering Strait and exit the Chukchi Sea off the Alaskan coast. Shelf-basin exchange facilitated by the generation of mesoscale eddies (Manley and Hunkins, 1985; Muench et al., 2000; Pickart et al., 2005; Mathis et al., 2007), and winter brine rejection during sea-ice formation are also important for these shelves.

As with the Siberian Sea shelves, few studies have been conducted on the Beaufort Sea shelf. Seawater *p*CO₂ conditions appear to be highly variable (~150–350 μatm) in the western part of the Beaufort Sea shelf (Murata and Takizawa, 2003; Bates, 2006).

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**The Arctic Ocean
marine carbon cycle**N. R. Bates and
J. T. Mathis

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Extremely low seawater $p\text{CO}_2$ conditions ($<100 \mu\text{atm}$) have also been observed in the surface layer of the Beaufort Sea shelf east of Point Barrow in a region of characterized by high proportions of freshwater (10–20%) and sea-ice melt (10–25%) (Cooper et al., 2005; Bates, 2006). Murata and Takizawa (2003) have estimated that rates of air-sea CO_2 exchange were relatively low ($-12 \text{ mmol CO}_2 \text{ m}^{-2} \text{ d}^{-1}$) during the summertime compared to the adjacent Chukchi Sea. In areas of the western Beaufort Sea shelf where sea-ice cover remained high ($>80\%$), air-to-sea CO_2 fluxes were generally low ($<1 \text{ mol CO}_2 \text{ m}^{-2} \text{ d}^{-1}$; Bates, 2006). The annual ocean CO_2 uptake for the Beaufort Sea shelf has been estimated at $1.2 \text{ g C m}^{-2} \text{ yr}^{-1}$, low compared to other Arctic shelves (Table 1; Murata and Takizawa, 2003).

4.3 Arctic Ocean shelves: outflow shelves

4.3.1 Canadian Archipelago shelf

The shelf of the Canadian Archipelago has been poorly sampled for marine inorganic carbon cycle properties. There is little information to estimate rates of air-sea CO_2 gas exchange and the processes that control the inorganic carbon cycle. In the adjacent Hudson Bay, Else et al. (2008) observed mean rates of air-sea CO_2 exchange of $-0.7 \text{ mmol CO}_2 \text{ m}^{-2} \text{ d}^{-1}$ during the sea-ice free period. On annual timescale, a small ocean CO_2 uptake for the Hudson Bay shelf is estimated at $0.07 \text{ g C m}^{-2} \text{ yr}^{-1}$ (Else et al., 2008).

In the CARINA dataset (Fig. 2), computation of $p\text{CO}_2$ from DIC and alkalinity datasets indicate that surface $p\text{CO}_2$ values ranged from $\sim 300\text{--}400 \mu\text{atm}$ on the Canadian Archipelago shelf and generally undersaturated with respect to the atmosphere. During the CASES project in the early 2000's, surface $p\text{CO}_2$ values ranged from $\sim 300\text{--}430 \mu\text{atm}$ with significant spital variability (Mucci et al., 2008). If we assume that air-sea CO_2 exchange rates are similar for the Canadian Archipelago as the “upstream” Beaufort Sea shelf, the annual uptake for the Canadian Archipelago shelf is computed at

-16 to -24 g C m⁻² yr⁻¹ (Table 1).

4.4 Arctic central basin: Canada and Eurasian basins

4.4.1 Canadian Archipelago shelf

The central basin of the Arctic Ocean is strongly stratified (Kinney et al., 1970; Aagaard et al., 1985; Jones and Anderson, 1986; Anderson et al., 1989; Aagaard and Carmack, 1994; Anderson et al., 1994a, b; Schlosser et al., 1995; Codispoti et al., 2005). In the Eurasian and Canada Basin, surface waters of the central basin are derived from a mixture of local sea-ice melt, fresh water inputs, exchanges with shelf waters of Pacific Ocean and Atlantic Ocean origin (Jones et al., 2008). In the Canada Basin, a strong vertical density gradient separates nutrient poor surface waters or polar mixed layer (PML) from underlying nutrient rich halocline waters that can be subdivided into Upper Halocline Layer (UHL) and Lower Halocline Layer (LHL). The shelf surface waters and underlying upper halocline waters have a predominantly Pacific Ocean origin based on characteristic temperature, salinity (Aagaard et al., 1981), inorganic nutrient, dissolved oxygen, barium and alkalinity distributions (e.g., Wallace et al., 1987; Wilson and Wallace, 1990; Jones et al., 1991; Salmon and McRoy, 1994; Falkner et al., 1994; Anderson, 1995; Rudels et al., 1996; Guay and Falkner, 1997; Anderson et al., 1999; Jones et al., 2003). In surface waters of the Canada Basin, the river fraction can be up to 20% (Macdonald et al., 2002; Cooper et al., 2005), while localized to the upper few metres, the sea-ice melt fraction can also exceed 25% (Cooper et al., 2005) during the summertime retreat of sea-ice. In the Eurasian Basin, surface waters and halocline waters are predominantly of Atlantic Ocean origin that have been modified on the Barents Sea and Siberian Seas shelves adjacent to the Eurasian Basin (Anderson et al., 1994). The upper halocline layer is absent from most of the Eurasian Basin (Anderson et al., 1994). At greater depths of the central basin is the Atlantic Water Layer (AWL; ~200–800 m deep) and the Arctic Ocean Deep Water (AODW), both of which have Atlantic Ocean origins (e.g., Jones et al., 1991, 1995).

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**The Arctic Ocean
marine carbon cycle**N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The central basin of the Arctic has been poorly sampled for marine inorganic carbon cycle properties. The Arctic Ocean Section (AOS) expedition in 1994 sampled across the Eurasian Basin and Canada Basin, west of the Chukchi Cap (Anderson et al., 1998a). Calculation of $p\text{CO}_2$ from AOS DIC and TA data (obtained at <http://cdiac.ornl.gov/>) indicate that the surface waters had seawater $p\text{CO}_2$ values of $\sim 300\text{--}330\ \mu\text{atm}$ and $\Delta p\text{CO}_2$ values of $\sim -40\text{--}70\ \mu\text{atm}$, and hence undersaturated with respect to the atmosphere. In the Canada Basin, adjacent to the Chukchi Sea shelf, lower seawater $p\text{CO}_2$ values of $\sim 240\text{--}280\ \mu\text{atm}$ and $\Delta p\text{CO}_2$ values of $\sim -150\pm 25\ \mu\text{atm}$ have also been observed (Bates, 2006; Bates et al., 2006). These observations suggest that the central basin of the Arctic Ocean has the potential to absorb significant amounts of CO_2 (Kaltin and Anderson, 2001; Bates et al., 2006). The capacity for sea-ice to allow direct gas exchange is debated (e.g., Gosink et al., 1976; Semiletov et al., 2004; Delille et al., 2007; Nagurnyi, 2008) and we assume that sea-ice is an effective barrier to gas exchange. In regions of the Canada Basin with 100% sea-ice cover, Bates et al. (2006) estimated an air-sea CO_2 exchange of $< 3\ \text{mmoles CO}_2\ \text{m}^{-2}\ \text{d}^{-1}$ through leads, fractures and brine channels, with much higher rates of exchange in areas without complete sea-ice cover.

4.5 Arctic Ocean (shelves and central basin): annual air-sea CO_2 exchange rate estimates

Early carbon mass balance estimates of the rate of air-sea CO_2 exchange suggested that the entire Arctic Ocean was a sink for CO_2 in the range of -70 to $-129\ \text{Tg C year}^{-1}$ (Table 1: Anderson et al., 1990, 1994; Lundberg and Haugen, 1996). Subsequently, Anderson et al. (1998b) and Kaltin and Anderson (2005) used similar carbon mass balance approaches but revised their estimates of the Arctic Ocean CO_2 sink downward to -24 and $-31\ \text{Tg C year}^{-1}$. More recently, direct $\Delta p\text{CO}_2$ observations and air-sea CO_2 exchange rate estimates revised the Chukchi Sea shelf CO_2 sink upward and the Arctic Ocean CO_2 sink was estimated at $66\ \text{Tg C year}^{-1}$ (Bates, 2006; Bates et al., 2006). A

review of published data suggests that the Arctic Ocean CO₂ sink ranges from approximately -65 to -175 Tg C year⁻¹ depending on lowest to highest flux rates estimates from available studies (Table 1). In the absence of data, if the area of the Canadian Archipelago is scaled to air-sea CO₂ exchange rates observed in the adjacent and “up-stream” Beaufort Sea shelf, estimates of the Arctic Ocean CO₂ sink increase and range from approximately -81 to -199 Tg C year⁻¹ (Table 1). In comparison to the mean annual global ocean CO₂ uptake of approximately -1400 Tg C year⁻¹ (Takahashi et al., 2009), the Arctic Ocean CO₂ sink potentially contributes ~5–14% to the global balance of CO₂ sinks and sources.

5 Feedbacks and vulnerabilities

The Arctic Ocean is currently in the forefront of climate change caused by both natural and anthropogenic factors that influence the cryosphere, hydrological and biogeochemical cycles of the Arctic. However, due to undersampling of the Arctic Ocean, many uncertainties remain about the marine carbon cycle and, as shown earlier in this review, the uptake of CO₂ and the physical and biological processes that control the inorganic carbon cycle on the shelves and in the central basin. It is thus difficult to quantitatively predict changes in processes and feedbacks in the Arctic Ocean. However, here we discuss the potential vulnerabilities of the inorganic carbon cycle due to natural and anthropogenic factors, including: (1) sea-ice loss; (2) warming, circulation and other physical changes; (3) changes in biology and ecosystem structure of the surface and halocline water, and; (4) changes in the hydrological cycle and freshwater inputs to the Arctic Ocean.

5.1 Sea-ice loss

Mean air temperatures over the Arctic have increased by ~2 – 3°C in summertime and by ~4°C in winter since the 1950's (Chapman and Walsh, 2003) with projections for

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

a further 4–5°C increase by the end of the 21st century (ACIA, 2005). In conjunction with these higher temperatures, sea ice cover in the Arctic Ocean has contracted over the past three decades (Levi, 2000; Parkinson, 2000; Cavalieri et al., 2003; Wang et al., 2005; Overland and Wang et al., 2005; Rothrock and Zhang, 2005; Stroeve et al., 2005, 2007), with the pace of decline accelerating beyond model predictions over the last several years (Winton, 2006; Holland et al., 2006; Overland and Wang, 2007; Stroeve et al., 2007; Maslanik et al., 2007). The length of the seasonal sea-ice melt period has declined (e.g., Smith 1998; Rigor et al., 2002; Rigor and Wallace, 2004; Arrigo et al., 2008; Comiso et al., 2008) and sea-ice thickness observed over the central Arctic Ocean has decreased as thick multi-year sea-ice is replaced by thinner first-year sea-ice (e.g., Rothrock et al., 1999; Giles et al., 2008). The decline in Arctic sea-ice cover has been attributed to a combination of factors, including: (1) increased Arctic temperatures (Rothrock and Zhang, 2005; Lindsay and Zhang, 2005); (2) increased advection of warmer water into the Arctic Ocean from the Atlantic Ocean through the Fram Strait and Barents Sea gateways (Steele and Boyd, 1998; Dickson et al., 2000) and from the Pacific Ocean through Bering Strait (Maslowski et al., 2001; Shimada et al., 2006, 2007), and; (3) increased advection of sea-ice out of the Arctic Ocean through Fram Strait due to recent changes in atmospheric circulation patterns (Rigor and Wallace, 2004; Maslanik et al., 2007; Serreze et al., 2007). In combination, these factors contribute to a positive feedback where sea-ice loss reinforces surface warming due to reduced surface albedo and increased shortwave penetration, which in turn inhibits sea-ice formation in wintertime and allows for acceleration of sea-ice loss during the summertime (Perovich et al., 2007).

The loss of sea-ice is expected to both reduce the % sea-ice cover and expose undersaturated surface waters of the Arctic shelves and central basin (i.e., Canada and Eurasian Basins). Prior to 2007, the rate of sea-ice loss from the central Arctic Ocean was estimated at $\sim 36\,000\text{ km}^2\text{ year}^{-1}$ (Cavalieri et al., 2003). In surface waters of the Canada Basin, a mean ocean CO_2 uptake of $46\pm 8\text{ mmol CO}_2\text{ m}^{-2}\text{ d}^{-1}$ was observed in 2002 and 2004 in sea-ice free areas during summertime. These air-sea

CO₂ exchange rates were scaled to the entire central basin and annual reduction of sea-ice extent to estimate an increase of the ocean CO₂ sink of 2.0±0.3 Tg C year⁻¹ due to sea-ice loss with the effect compounding over time (Bates et al., 2006). In 2007 and 2008, sea-ice extent reached a seasonal minima 25% lower than any previously observed in the satellite record (Maslanik et al., 2007; Comiso et al., 2008) constituting an additional exposure of ~600 000 km² of surface waters to air-sea gas exchange. Assuming a *status quo* of inorganic carbon distributions in surface waters of the Arctic, this recent loss of summertime sea-ice may have increased the ocean uptake of CO₂ in the Arctic by an additional 33±10 Tg C year⁻¹. If sea-ice cover was lost completely over the central basin during summertime, surface waters of the Arctic Ocean might absorb an additional ~280 to 1200 Tg of CO₂ (Anderson and Kaltin, 2001; Bates et al., 2006). The range of estimates depends on differences in surface water carbonate chemistry observed between the Canada Basin and Eurasian Basin. For example, Anderson and Kaltin (2001) computed how much CO₂ was needed to equilibrate the surface mixed layer with the atmosphere using vertical profiles of DIC data from the Eurasian Basin.

The above estimates of CO₂ uptake changes due to sea-ice loss have many uncertainties and associated caveats. These studies are only applicable to the near-term future (<decade) since it was assumed that the driving force of gas exchange (e.g., ΔpCO₂ and DIC distributions of surface and underlying halocline layer waters) would not change significantly over the residence time of surface waters in the Arctic (e.g., 2–30 years; Macdonald et al., 2009). However, in the era of rapid change in the Arctic, inorganic carbon distributions and air-sea CO₂ exchange rates are highly likely to change due to a host of other factors. Since the effect of these other factors on the marine inorganic carbon cycle can be opposing or amplifying in nature, future predictions of the trajectory of the Arctic Ocean CO₂ sink/source terms are difficult to make.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5.2 Warming, circulation and other physical changes in the Arctic

Due to simple thermodynamics, warming should increase seawater $p\text{CO}_2$ at a rate of $\sim 4.1\%$ per $^\circ\text{C}$ (Takahashi et al., 1993; Millero et al., 1995). In the central basin, if we assume present day surface water $p\text{CO}_2$ values of $250\text{--}300\ \mu\text{atm}$, a 1°C warming should increase seawater $p\text{CO}_2$ by $\sim 8\text{--}12\ \mu\text{atm}$ per 1°C and reduce $\Delta p\text{CO}_2$ values and the ocean CO_2 sink by $\sim 10\%$. In the recent past (1998–2006), Pabi et al. (2008) observed positive anomalies in surface temperatures of up to 2°C in the regions of significant sea-ice loss (mainly on the “inflow” Arctic shelves). If surface waters equilibrate with the predicted $4\text{--}5^\circ\text{C}$ warming of the atmosphere over the Arctic by the end of the 21st century (ACIA, 2005), and if present-day inorganic carbon distributions remain unchanged, the Arctic Ocean CO_2 sink might reduce by $\sim 40\text{--}50\%$ due to warming.

The loss of sea-ice in the Arctic should result in greater open water area, and increased air-sea interaction, with a variety of consequences. At the same time, increased latitudinal atmospheric gradients could result in increased windspeed and storm events over the Arctic. Although the direction of air-sea gas exchange is forced by $\Delta p\text{CO}_2$ values, the rate of gas exchange of CO_2 and other gases is primarily driven by windspeed and atmospheric boundary layer-surface water interactions. Given these dictates, even though $\Delta p\text{CO}_2$ values might reduce, air-sea CO_2 gas exchange rates can increase since gas transfer velocities increase exponentially relative to windspeed (Wanninkhof, 1992; Wanninkhof and McGillis, 1999) and CO_2 flux in other regions can be dominated by storms and high-wind events (Bates and Merlivat, 2001).

Wind-driven air-sea interaction and eddy formation may also change, including wind-induced upwelling/downwelling (Melling, 1993; Pickart et al., 2006, 2009) and formation of eddies that contribute to exchanges of water and materials between the shelf and basin (Manley and Hunkins, 1985; Muench et al., 2000; Pickart et al., 2005), particularly in the western Arctic between the Beaufort Sea shelf and Canada Basin (Carmack and Macdonald, 2002; Mathis et al., 2007; Spall et al., 2008). Wintertime sea-ice is thinner and there may be potentially greater air-sea gas exchange through

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**The Arctic Ocean
marine carbon cycle**N. R. Bates and
J. T. Mathis

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

sea-ice (Gosink et al., 1978; Semiletov et al., 2004; Delille et al., 2007) with sea-ice a potentially weaker barrier to gas exchange. In wintertime, wind-driven polynyas open up on the Chukchi and Laptev Sea shelves in particular (Cavaliere and Martin, 1994; Nitishinsky et al., 2007), resulting in the formation of dense water via brine-driven convection (Weingartner et al., 1998). Since brine rejection and formation of dense water appears to facilitate uptake of CO₂ from the atmosphere (Anderson et al., 2004; Omar et al., 2005), enhanced opening of polynyas and leads may increase the sink of CO₂ in the Arctic in the near-future depending on inorganic carbon distributions and future air-sea CO₂ disequilibrium. However, the cumulative effect of the present influence and future changes in these processes remain uncertain.

5.3 Changes in biology and ecosystem structure in surface and halocline waters of the Arctic Ocean

The loss of sea-ice in the Arctic and greater open water area should also enhance upwelling at the shelf-break and potentially increase the input of nutrients from subsurface waters to the Arctic shelves (ACIA, 2005). The growing season for phytoplankton should increase with reduced sea-ice extent and longer open-water conditions. As a consequence of increased phytoplankton PP or NCP, the drawdown of $p\text{CO}_2$ and DIC should increase the air-sea CO₂ disequilibrium (i.e., $\Delta p\text{CO}_2$) and increase the net oceanic uptake of CO₂ (Anderson and Kaltin, 2001; Bates et al., 2006). Recent reductions of sea-ice cover, particularly in the western Arctic Ocean, have resulted in a longer phytoplankton growing season and a ~30–60% increase in primary production over the last decade, with a ~10–40% increase in phytoplankton primary production over the last several years (Arrigo et al., 2008). For sea-ice communities that generally account for a relatively small fraction of primary production, the reduction of sea-ice extent and thickness will reduce the contribution of sea-ice algae to primary production (Subba Rao and Platt, 1984; Legendre et al., 1992; Gosselin et al., 1997) and their influence on surface water inorganic carbon distributions and air-sea CO₂ disequilibrium.

Other factors may also influence the marine inorganic carbon cycle and present-day

CO₂ sink in the Arctic Ocean. For example, reduced sea-ice cover has been proposed to favor a “phytoplankton-zooplankton” dominated ecosystem rather than a “sea-ice algae–benthos” dominated ecosystem over the Arctic shelves (Piepenburg, 2005). At present on the highly productive Chukchi Sea shelf, ~10% of NCP is converted to dissolved organic carbon (DOC) and ~15% of NCP is converted to suspended particulate organic carbon (POC) (Mathis et al., 2007b, 2009) that gets exported from the shelf into the Canada Basin beneath the mixed layer (Bates et al., 2005b). The remaining 75% of NCP is exported from the mixed layer to the sea floor as sinking particles that sustain the rich benthos on the sea floor of the Chukchi Sea shelf (Grebmeier et al., 2007). In the Bering Sea, earlier sea-ice loss has led to ecosystem changes and altered pelagic-benthic coupling (e.g., Stabeno et al., 2001, 2002; Hunt et al., 2002; Overland et al., 2001; Napp et al., 2002; Bond et al., 2003; Merico et al., 2004). If there are ecosystem shifts in the future, for example on the Chukchi and Beaufort Seas shelves, the export of organic carbon and pelagic-benthic coupling might decrease, despite concurrent increases in phytoplankton PP.

There may also be changes in the inorganic carbon distributions of subsurface waters on the Arctic shelves and halocline layer waters of the central basin. In the central basin, density stratification generally acts as a barrier to mixing between nutrient-poor surface waters and nutrient-rich deep waters (Aagaard et al., 1981; Jones and Anderson, 1986). Halocline waters generally have much higher $p\text{CO}_2$ and DIC content than surface waters (Jutterstrom and Anderson, 2004), with low rates of upward vertical diffusion between these waters and the surface mixed layer (Wallace et al., 1987; Wilson and Wallace, 1990; Bates, 2006). The mixed layer typically extends to 10–50 m, with depth heavily dependent on seasonality of winds and sea-ice cover (Carmack and Macdonald, 2002). The reduction of sea-ice extent may facilitate deeper mixing and erosion into inorganic nutrient and CO₂ rich halocline waters. Depending on the stoichiometry of nutrients and carbon entrained into the mixed layer, enhancement of phytoplankton PP in typically highly oligotrophic surface waters of the Arctic (e.g., English, 1961) and subsequent vertical export of organic carbon, this mechanism could

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



either enhance or suppress air-sea CO₂ disequilibrium and CO₂ exchange between atmosphere and ocean. The halocline is believed to ventilate predominantly through lateral processes and the transfer of water from the shelves into the interior formation of dense water on the shelves via brine-driven convection (Weingartner et al., 1998).

5 If the marine inorganic carbon cycle on the Arctic shelves is altered, the transfer of inorganic and organic carbon into the halocline may also change. For example, if there is increased transport of DOC from the shelves into the subsurface halocline of the central basin, remineralization of DOC by the microbial community may enhance ρCO_2 and DIC contents of surface waters, and in turn, the upward supply of CO₂ to the surface layer through vertical diffusion and mixing. The release of alkalinity from the
10 sediments, produced by anaerobic microbial decomposition, may also be important for suppressing seawater ρCO_2 contents that could in turn enhance air-sea CO₂ disequilibrium (Thomas et al., 2009). Thus ecosystem changes in the Arctic shelves and in the central basin, both in the surface and underlying halocline layer, may alter inorganic
15 carbon distributions and the CO₂ sink or source terms of the Arctic as a whole.

5.4 Changes in the hydrological cycle and freshwater inputs to the Arctic Ocean

Significant changes in the hydrological cycle of the Arctic are also predicted by the end of the 21st century (ACIA, 2005). Freshwater inputs from Arctic rivers and the transport of sediment and dissolved materials are expected to increase. At present, Arctic rivers
20 influence the hydrographic properties of the Siberian Sea and Beaufort Sea shelves in particular. The eastern Beaufort Sea shelf is heavily impacted by Mackenzie River inflow of freshwater (e.g., Macdonald et al., 1995a, b; 2002, 2009; Kadko and Swart, 2005) and plumes of elevated sediment loads and chromophoric dissolved organic matter (CDOM) can be followed from the Mackenzie outflow into the Canada Basin
25 using satellite remote sensing. However, the impact of increased river flow on the chemical characteristics of the Arctic is uncertain at present. The freshwater input of DOC will presumably increase, but its contribution to inorganic carbon distributions on the shelves from bacterial remineralization of DOC to CO₂ is highly dependent on the

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lability of DOC and its decomposition relative to residence times in the Arctic. While most riverine DOC is thought to be highly refractory (Hansell et al., 2004), enhanced open-water may increase the photochemical breakdown of refractory DOC to CO₂ in surface waters.

5.5 Summary of near-term impacts on Arctic Ocean air-sea CO₂ fluxes

In the near-term, sea-ice loss is expected to increase the uptake of CO₂ by surface waters because the exposed surface waters have a lower CO₂ content than the atmosphere. The loss of sea-ice in the Arctic also is likely to result in greater open water area, and increased air-sea interaction, with the likelihood of increased rates of air-sea gas exchange. Wintertime sea-ice is now thinner than previous decades and there may be potentially greater atmosphere-ocean gas exchange directly through sea-ice, with sea-ice a potentially weaker barrier to gas exchange. As a consequence of increased phytoplankton growth due to sea-ice loss and expansion of open water in the Arctic, the potential for the Arctic Ocean to uptake CO₂ should also increase.

In mitigation, reduced cooling of water during transit poleward and increased absorption of solar radiation resulting in warming of surface water relative to previous decades should act to increase the CO₂ content of seawater. This process may somewhat counteract the impacts of sea-ice loss and increased phytoplankton growth on the atmosphere-ocean exchange of CO₂.

6 Ocean acidification effects and changes in carbonate mineral saturation states

As a consequence of the ocean uptake of anthropogenic CO₂, surface pCO₂ and DIC concentration have increased while pH has decreased in the upper ocean over the last few decades (e.g., Winn et al., 1994; Bates et al., 1996; Bates, 2007; Bates and Peters, 2007; Takahashi et al., 2009). This gradual process, termed *ocean acidification*,

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



has long been recognized (e.g., Broecker and Takahashi, 1966; Broecker et al., 1971; Bacastow and Keeling, 1973). Estimates based on the Intergovernmental Panel on Climate Change (IPCC) projections of future CO₂ emission and subsequent ocean absorption of anthropogenic CO₂ (Solomons et al., 2007) predict that upper ocean pH will decrease by 0.3–0.5 units over the next century and beyond (Caldeira and Wickett, 2003, 2005; Doney, 2006). The effects of ocean acidification are potentially far-reaching in the global ocean, particularly for calcifying fauna (e.g., Buddemeier et al., 2004; Royal Society, 2005; Fabry et al., 2008) with model studies suggesting that CaCO₃ mineral undersaturation due to ocean acidification should impact the Arctic before other regions (Orr et al., 2005; Steinacher et al., 2009).

The predicted worldwide pH changes using the A2 scenario (IPCC, 2007) are expected to increase hydrogen ion concentration by 185% and decrease pH by 0.45 (Steinacher et al., 2009). Decreased pH reduces the saturation states (Ω) of calcium carbonate (CaCO₃) minerals such as aragonite and calcite, and increases the potential for reduced production of CaCO₃ by calcifying fauna (e.g., Buddemeier et al., 2004; Royal Society, 2005; Fabry et al., 2008; Doney et al., 2009) and dissolution of CaCO₃ in the water-column and sediments.

In the Arctic Ocean, sea-ice melt, river runoff and phytoplankton photosynthesis and respiration appear to be the dominant processes that induce divergent directions for CaCO₃ saturation states for surface and subsurface waters. As such, these processes may suppress or enhance Ω values that in turn amplify or mitigate the impacts of ocean acidification in the Arctic.

6.1 Suppression and enhancement of Ω in surface waters

In surface waters, freshwater contributions from sea-ice melt and river runoff appear to be the primary processes that act to decrease CaCO₃ saturation states. Salisbury et al. (2008) have shown that the Ω values of major Arctic rivers are close to zero for the run-off end-member. In the Chukchi Sea and Canada Basin, surface mixed layer waters were generally saturated with respect to aragonite ($\Omega_{\text{aragonite}}$) but $\Omega_{\text{aragonite}}$ values

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**The Arctic Ocean
marine carbon cycle**

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



decreased as runoff fraction increased (Bates et al., 2009). These findings are consistent with observations that a wide variety of Arctic rivers are net heterotrophic (e.g., Chen and Borges, 2009) with high $p\text{CO}_2$ contents (Kelley, 1970; Makkayeev, 1994; Semiletov, 1999; Semiletov et al., 2007; Nitishinsky et al., 2007) and corresponding low Ω values. River DOC remineralization and respiration in combination with many Arctic rivers having high TA contents (Anderson et al., 1983; Olsson and Anderson, 1997; Cooper et al., 2008; Bates et al., 2009) act to decrease Ω values.

Ω values have also been observed to decrease with increasing sea-ice melt fraction (>15%; Bates et al., 2009), and in a few localized surface areas in the Canada Basin, both aragonite and calcite undersaturation have been observed. Similarly, Chierici and Fransson (2009) have observed localized areas where surface waters were undersaturated with respect to aragonite ($\Omega_{\text{aragonite}}$) on the Canadian Archipelago and Beaufort Sea shelves which they attribute to the influence of freshwater contributions from sea-ice melt and river runoff. Cooling and air-sea uptake of CO_2 (which increases $p\text{CO}_2$ and DIC) will also act to decrease Ω values, but are relatively minor impacts compared to sea-ice melt and river runoff inputs to the Arctic. Low Ω values have also been attributed to upwelling of CO_2 -rich subsurface waters off the Chukchi Sea shelf (Chierici and Fransson, 2009).

Oversaturated Ω values are been observed for surface waters across much of the Chukchi Sea shelf and adjacent Canada Basin (Jutterstrom and Anderson, 2005; Bates et al., 2009; Chierici and Fransson, 2009). The oversaturated conditions for both aragonite and calcite have been attributed to the high rates of phytoplankton PP that decrease $p\text{CO}_2$ and DIC, and increase Ω values on the Chukchi Sea shelf (Bates et al., 2009), as well as low DIC:TA ratios and Revelle Factors present in the Canada Basin (Bates, 2006) that facilitates oversaturated Ω conditions.

6.2 Suppression of Ω in subsurface waters

In the North Pacific Ocean, Feely et al. (1988) observed seasonal enhancement of surface Ω values and suppression of subsurface Ω values due to seasonal phytoplankton

PP, export of organic matter and subsequent remineralization to CO_2 . A similar mechanism occurs in the Chukchi Sea where high rates of summertime phytoplankton PP (Hill and Cota, 2005; Bates et al., 2005; Mathis et al., 2009), vertical export of organic carbon (Lepore et al., 2007) and buildup of CO_2 in subsurface waters (primarily through subsurface remineralization of OM to CO_2 , and benthic respiration) suppresses Ω values below saturation. Subsurface waters were observed to be undersaturated with respect to aragonite and calcite, and thus potentially corrosive to CaCO_3 for the Chukchi Sea shelf sediments and benthic ecosystem.

6.3 Amplification and mitigation of ocean acidification impacts in the Arctic Ocean

In many oceanic basins, the oceanic uptake of anthropogenic CO_2 has resulted in the shoaling of the aragonite saturation horizon (i.e., $\Omega_{\text{aragonite}}=1$) by 40 to 200 m (Feely et al., 2004) over the past century. Given the scenarios for pH changes in the Arctic Ocean (Orr et al., 2005; Steinacher et al., 2009), the Arctic shelves will be increasingly impacted by ocean acidification and presence of CaCO_3 mineral undersaturated waters. Superimposed on the ocean acidification effects induced by ocean uptake of anthropogenic CO_2 , seasonal biological processes are likely to mitigate (enhance Ω) and amplify (suppress Ω) the impacts of ocean acidification on the Arctic shelves.

The seasonal changes in Ω induced by biology primarily have been described as a seasonal “Phytoplankton-Carbonate Saturation State” (PhyCASS) interaction that drives divergent trajectories for carbonate chemistry in surface and subsurface waters of Arctic shelves like the Chukchi Sea (Bates et al., 2009). For inflow shelves (Carmack and Wassmann, 2006) like the Chukchi Sea and Barents Sea, high rates of summertime phytoplankton PP or NCP and vertical export of OM during seasonal sea-ice loss may drive a strong seasonal divergence in surface and subsurface Ω . The seasonal divergence in Ω is probably attenuated in the Barents Sea compared to the Chukchi Sea due to the presence of a deeper shelf. For interior shelves such as the Siberian and Beaufort Sea shelves, much lower rates of summertime phytoplankton PP, and

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



much greater influence of river runoff (with lower Ω) is also likely to result in attenuated divergence in Ω for surface and subsurface waters compared to the Chukchi Sea. On all shelves, a combination of Pacific or Atlantic water inflow and wintertime homogenization of shelf waters through wind-mixing and brine rejection processes are likely to “rectify” the seasonal divergences in Ω between surface and subsurface waters.

In an era of rapid change in the Arctic, sea-ice loss, surface warming and increased phytoplankton PP on the Arctic shelves may act to enhance surface Ω and suppress subsurface Ω amplifying the existing impact of ocean acidification. This may be mitigated somewhat by increased freshwater contributions through sea-ice melt and river runoff that suppress surface water Ω . In the central basin of the Arctic Ocean, potentially corrosive waters to aragonite and calcite are found in the upper halocline layer (Jutterstrom and Anderson, 2005; Bates et al., 2009), with highly saturated Ω values in surface water. Increased air-sea interaction in regions experiencing continued seasonal sea-ice loss will likely exhibit increased vertical mixing of CO_2 -rich subsurface waters, increased shelf-basin exchanges and ocean uptake of CO_2 through gas exchange. Combined, these factors will reduce the vertical gradients of Ω between surface and subsurface with the net result of suppressing surface Ω over the residence time of surface water in the central basin (2–30 years; Macdonald et al., 2009).

7 Conclusions

At present, the Arctic Ocean continental shelves and central basin have lower CO_2 content than the atmosphere, and these areas are potential sinks of CO_2 from the atmosphere. There are however localized areas of sea-ice melt and river inputs where the opposite is observed, and these areas are potential sources of CO_2 to the atmosphere. The CO_2 chemistry of the Arctic Ocean is highly influenced by physical and biological processes such as seasonal phytoplankton PP during summertime sea-ice retreat towards the pole, as well as temperature effects (both cooling and warming), shelf-basin exchanges and formation of dense winter waters, and river inputs of fresh-water and carbon.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Although seasonal sea-ice cover provides a barrier to atmosphere-ocean gas exchange, the Arctic Ocean is a sink for CO₂, on the order of –65 to –175 Tg C year⁻¹, contributing 5–14% to the global balance of CO₂ sinks and sources. The Arctic Ocean has become an important influence on the global carbon cycle, with the marine carbon cycle and atmosphere-ocean CO₂ exchanges sensitive to Arctic Ocean and global climate change feedbacks. In the near-term, further sea-ice loss and increases in phytoplankton PP rates are expected to increase the uptake of CO₂ by Arctic surface waters, although mitigated somewhat by warming in the Arctic. Thus, the capacity of the Arctic Ocean to uptake CO₂ is expected to increase and alter in response to environmental changes driven largely by climate. These changes are likely to continue to modify the physics, biogeochemistry, and ecology of the Arctic Ocean in ways that are not yet fully understood. Finally, in response to increased phytoplankton PP and uptake of human-produced CO₂, the benthic ecosystem of the Arctic shelves are expected to be negatively impacted by ocean acidification which reduces the ability of many species to produce calcium carbonate shells or tests with profound implications for Arctic marine ecosystems.

References

- Aagaard, K., Coachman, L. K., and Carmack, E. C.: On the halocline of the Arctic Ocean, *Deep-Sea Res.*, 28, 529–545, 1981.
- Aagaard, K. and Carmack, E. C.: The role of sea ice and other fresh water in the Arctic circulation, *J. Geophys. Res.*, 94(C10), 14 485–14 498, 1989.
- Aagaard, K. and Carmack, E. C.: The Arctic Ocean and Climate: A perspective, in: *The Polar Oceans and Their Role in Shaping the Global Environment*, edited by: Johannessen, O. M., Muench, R. D., and Overland, J. E., American Geophysical Union, Geophysical Monograph, 85, 5–20, 1994.
- Arctic Climate Impact Assessment (ACIA), Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2005.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**The Arctic Ocean
marine carbon cycle**N. R. Bates and
J. T. Mathis

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Anderson, L. G.: Chemical oceanography of the Arctic and its shelf sea, in: Arctic Oceanography: Marginal Ice Zones and Continental Shelves, edited by: Smith Jr., W. O., and Grebmeier, J. M., Coastal and Estuarine Studies, 49, American Geophysical Union, Washington, DC, 1995.
- 5 Anderson, L. G. and Kaltin, S.: Carbon fluxes in the Arctic Ocean – potential impact by climate change, *Polar Res.*, 20(2), 225–232, 2001.
- Anderson, L. G., Jones, E. P., Lindegren, R., Rudels, B., and Sehlstedt, P. I.: Nutrient regeneration in cold, high salinity bottom water of the Arctic shelves, *Cont. Shelf Res.*, 8, 1345–1355, 1988.
- 10 Anderson, L. G., Dyrssen, D., and Jones, E. P.: An assessment of the transport of atmospheric CO₂ into the Arctic ocean, *J. Geophys. Res.*, 95, 1703–1711, 1990.
- Anderson, L. G., Olsson, K., and Skoog, A.: Distributions of dissolved inorganic and organic carbon in the Eurasian Basin of the Arctic Ocean, in: The Polar Oceans and Their Role in Shaping the Global Environment, Johannessen, O. M., Muench, R. D. and Overland, J. E.,
15 American Geophysical Union, Geophysical Monograph, 85, 252–262, 1994.
- Anderson, L. G., Bjork, G., Holby, O., Jones, E. P., Kattner, G., Koltermann, K. P., Liujeband, B., Lindegren, R., Rudels, R., and Swift, J.: Water masses and circulation in the Eurasian Basin: Results from the Oden 91 Expedition, *J. Geophys. Res.-Oceans*, 99, 3273–3283, 1994.
- Anderson, L. G., Olsson, K., and Chierici, M.: A carbon budget for the Arctic Ocean, *Global Biogeochem. Cy.*, 12(3), 455–465, 1998a.
- 20 Anderson, L. G., Olsson, K., Jones, E. P., Chierici, M., and Fransson, A.: Anthropogenic carbon dioxide in the Arctic Ocean: Inventory and sinks, *J. Geophys. Res.*, 103(C12), 27 707–27 716, 1998b.
- Anderson, L. G., Jones, E. P., and Rudels, B.: Ventilation of the Arctic Ocean estimated by a plume entrainment model constrained by CFCs, *J. Geophys. Res.*, 104(C6), 13 423–13 429,
25 1999.
- Anderson, L. G., Falck, E., Jones, E. P., Jutterstrom, S., and Swift, J. H.: Enhanced uptake of atmospheric CO₂ during freezing of seawater: A field study in Storfjorden, Svalbard, *J. Geophys. Res.*, 109(C6), C06004, doi:10.1029/2003JC002120, 2004.
- 30 Arrigo, K. R., van Dijken, G. L., and Pabi, S.: The impact of a shrinking Arctic ice cover on marine primary production, *Geophys. Res. Lett.*, 35, L19603, doi:10.1029/2008GL035028, 2008.

Bacastow, R. D. and Keeling, C. D.: Atmospheric carbon dioxide and radiocarbon in the natural carbon cycle: II. Changes from A.D. 1700 to 2070 as deduced from a geochemical model, in: Carbon and the biosphere, edited by: Woodwell, G. M. and Pecan, E. V., US Atomic Energy Commission, 86–135, 1973.

5 Bates, N. R.: Air-sea CO₂ fluxes and the continental shelf pump of carbon in the Chukchi Sea adjacent to the Arctic Ocean, *J. Geophys. Res.-Oceans*, 111, C10013, doi:10.129/2005JC003083, 2006.

Bates, N. R.: Interannual variability of the oceanic CO₂ sink in the subtropical gyre of the North Atlantic Ocean over the last two decades, *J. Geophys. Res.-Oceans*, 112, C09013, doi:10.1029/2006JC003759, 2007.

10 Bates, N. R. and Merlivat, L.: The influence of short-term wind variability on air-sea CO₂ exchange, *Geophys. Res. Lett.*, 28(17), 3281–3284, 2001.

Bates, N. R. and Peters, A. J.: The contribution of atmospheric acid deposition to ocean acidification in the subtropical North Atlantic Ocean, *Mar. Chem.*, 107, 547–558, 2007.

15 Bates, N. R., Michaels, A. F., and Knap, A. H.: Seasonal and interannual variability of the oceanic carbon dioxide system at the U.S. JGOFS Bermuda Atlantic Time-series Site, *Deep-Sea Res. Pt. II*, 43(2–3), 347–383, 1996.

Bates, N. R., Best, M. H. P., and Hansell, D. A.: Spatio-temporal distribution of dissolved inorganic carbon and net community production in the Chukchi and Beaufort Seas, *Deep-Sea Res. Pt. II*, 52, 3303–3323, 2005a.

20 Bates, N. R., Hansell, D. A., Moran, S. B., and Codispoti, L. A.: Seasonal and spatial distributions of particulate organic matter (POM) in the Chukchi Sea, *Deep-Sea Res. Pt. II*, 52, 3324–3343, 2005b.

25 Bates, N. R., Moran, S. B., Hansell, D. A., and Mathis, J. T.: An increasing CO₂ sink in the Arctic Ocean due to sea-ice loss?, *Geophys. Res. Lett.*, 33, L23609, doi:10.1029/2006GL027028, 2006.

Bates, N. R., Mathis, J. T., and Cooper, L.: The effect of ocean acidification on biologically induced seasonality of carbonate mineral saturation states in the Western Arctic Ocean, *J. Geophys. Res.-Oceans*, in press, 2009.

30 Björk, G.: A one-dimensional time-dependent model for the vertical stratification of the upper Arctic Ocean, *J. Phys. Oceanogr.*, 19, 52–67, 1989.

Bond, N. A., Overland, J. E., Spillane, M., and Stabeno, P.: Recent shifts in the state of the North Pacific, *Geophys. Res. Lett.*, 30(23), 2183, doi:10.1029/2003GL018597, 2003.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Broecker, W. S. and Takahashi, T.: Calcium carbonate precipitation on the Bahama Banks, *J. Geophys. Res.*, 71, 1575–1602, 1966.
- Broecker, W. S., Li, Y.-H., and Peng, T.-H.: Carbon dioxide – man’s unseen artifact, in: Impingement of man on the oceans, edited by: Hood, D. W., John Wiley and Sons, Inc, 287–324, 1971.
- Buddemeier, R. W., Kleypas, J. A., and Aronson, R. B.: Coral Reefs and Global Climate Change: Potential Contributions of Climate Change to Stresses on Coral Reef Ecosystems, Pew Center on Climate Change, available at: http://www.pewclimate.org/docUploads/Coral_Reefs.pdf, 2004.
- Caldeira, K. and Wickett, M. E.: Anthropogenic carbon and ocean pH, *Nature*, 425(6956), 365–365, 2003.
- Caldeira, K. and Wickett, M. E.: Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean, *J. Geophys. Res.-Oceans*, 110(C9), C09S04, doi:10.1029/2004JC002671, 2005.
- Carmack, E. C. and Chapman, D.: Wind-driven shelf/basin exchange on an Arctic shelf: The joint roles of ice cover extent and shelf-break bathymetry, *Geophys. Res. Lett.*, 30, 1778, doi:10.1029/2003GL017526, 2003.
- Carmack, E. C. and Macdonald, R. W.: Oceanography of the Canadian shelf of the Beaufort Sea: A setting for marine life, *Arctic*, 55, Supplement 1, 29–45, 2002.
- Carmack, E. and Wassman, P.: Food webs and physical-biological coupling on pan-Arctic shelves: Unifying concepts and comprehensive perspectives, *Progress in Oceanography*, 71, 446–477, 2006.
- Cavaliere, D. J. and Martin, S.: The contribution of Alaskan, Siberian, and Canadian coastal polynyas to the cold halocline of the Arctic Ocean, *J. Geophys. Res.*, 99, 18343–18362, 1994.
- Cauwet, G. and Sidorov, I.: The biogeochemistry of Lena River: organic carbon and nutrients distribution, *Mar. Chem.*, 53, 211–227, 1996.
- Chapman, W. L. and Walsh, J. E.: Observed climate change in the Arctic: updated from Chapman and Walsh, 1993: Recent variations of sea ice and air temperatures in high latitudes, *B. Am. Meteorol. Soc.*, 74(1), 33–47, 2003.
- Chen, C. T. A. and Borges, A. V.: Reconciling opposing views on carbon cycling in the coastal ocean: Continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO₂, *Deep-Sea Res. Pt. II*, in press, 2009.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Chen, L. and Gao, Z.: Spatial variability in the partial pressures of CO₂ in the northern Bering and Chukchi seas, *Deep-Sea Res. Pt. II*, 54, 2619–2629, 2007.
- Chierici, M. and Fransson, A.: Calcium carbonate saturation in the surface water of the Arctic Ocean: undersaturation in freshwater influenced shelves, *Biogeosciences Discuss.*, 6, 4963–4991, 2009,
5 <http://www.biogeosciences-discuss.net/6/4963/2009/>.
- Coachman, L. K., Aagaard, K., and Tripp, R. B.: *Bering Strait: The Regional Physical Oceanography*, University of Washington Press, Seattle, 1975.
- Comiso, J. C., Parkinson, C. L., Gersten, R., and Stock, L.: Accelerated decline in the Arctic sea ice cover, *Geophys. Res. Lett.*, 35, L01703, doi:10.1029/2007GL031972, 2008.
- 10 Cooper, L. W., Benner, R., McClelland, J. R., Peterson, B. J., Holmes, R. M., Raymond, P. A., Hansell, D. A., Grebmeier, J. M., and Codispoti, L. A.: The linkage between runoff, dissolved organic carbon, and the stable oxygen isotope composition of seawater and other water mass indicators in the Arctic Ocean, *J. Geophys. Res.*, 110, G02013, doi:10.1029/2005JG000031, 2005.
- 15 Cooper, L. W., McClelland, J. W., Holmes, R. M., Raymond, P. A., Gibson, J. J., Guay, C. K., and Peterson, B. J.: Flow-weighted tracer content (¹⁸O, DOC, Ba, alkalinity) of the six largest Arctic rivers, *Geophys. Res. Lett.*, 35, L18606, doi:10.1029/2008GL035007, 2008.
- Cota, G. F., Pomeroy, L. R., Harrison, W. G., Jones, E. P., Peters, F., Sheldon W. M., and Weingartner T. R.: Nutrients, primary production and microbial heterotrophy in the south-eastern Chukchi Sea: Arctic summer nutrient depletion and heterotrophy, *Mar. Ecol.-Prog. Ser.*, 135(1–3), 247–258, 1996.
- 20 Delille, B., Jourdain, B., Borges, A. V., Tison, J.-L., and Delille, D.: Biogas (CO₂, O₂, dimethylsulfide) dynamics in spring Antarctic fast ice, *Limnol. Oceanogr.*, 52, 1367–1379, 2007.
- 25 Dickson, A. G.: Thermodynamics of the dissolution of boric acid in synthetic seawater from 273.15°K to 318.15°K, *Deep-sea Res.*, 37A(5), 755–766, 1990.
- Dickson, A. G. and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media, *Deep-Sea Res.*, 34, 1733–1743, 1987.
- Dickson, A. G. and Goyet, C.: *Handbook of Methods for the Analysis of the Various Parameters of the Carbon Dioxide System in Seawater*; version 2.0, US Department of Energy CO₂ Science Team Report, 1994.
- 30 Dickson, A. G., Sabine, C. L., and Christian, J. R.: *Guide to best practices for ocean CO₂ measurements*, Sidney, British Columbia, North Pacific Marine Science Organization, PICES

BGD

6, 6695–6747, 2009

**The Arctic Ocean
marine carbon cycle**N. R. Bates and
J. T. Mathis

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Special Publication 3, 2007.

Dmitrenko, I. A., Kirilov, S. A., and Trembley, B. L.: The long-term and interannual variability of summer fresh water storage over the eastern Siberian shelf: Implications for climate change, *J. Geophys. Res.-Oceans*, 113, C03007, doi:10.1029/2007JC004304, 2008.

5 Doney, S. C.: The dangers of ocean acidification, *Scientific American*, 58–65, 2006.

Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean acidification: The other CO₂ problem, *Annual Review of Marine Science*, 1, 169–192, 2009.

Ducklow, H. W. and McAllister, S. L.: Biogeochemistry of carbon dioxide in the coastal oceans, in: *The Sea, Volume 13, The Global Coastal Ocean-Multiscale Interdisciplinary Processes*, edited by: Robinson, A. R. and Brink, K., J. Wiley and Sons, NY, 2005.

10 Else, B.G.T., Papakyriakou, T.N., Granskog, M.A., and Yackel, J.J., 2008. Observations of sea surface *f*CO₂ distributions and estimated air-sea CO₂ fluxes in the Hudson Bay region (Canada) during the open water season, *J. Geophys. Res.-Oceans*, 113, C08026, doi:10.1029/2007JC004389, 2008.

15 Fabry, V. J., Seibel, B. A., Feely, R. A., and Orr, J. C.: Impacts of ocean acidification on marine fauna and ecosystem processes, *ICES J. Mar. Sci.*, 65, 414–432, 2008.

Falkner, K.K., MacDonald, R.W., Carmack, E.C., and Weingartner, T., 1994. The potential of barium as a tracer of Arctic water masses, in: *Polar Oceans and Their Role in Shaping the Global Environment*, edited by: Johannessen, O. M., Muench, R. D., and Overland, J. E., American Geophysical Unions, Geophysical Monograph, 85, 63–76, 1994.

20 Feely, R. A., Byrne, R. H., Acker, J. G., Betzer, P. R., Chen, C.-T. A., Gendron, J. F., and Lamb, M. F.: Winter summer variations of calcite and aragonite saturation in the northeast Pacific, *Mar. Chem.*, 25(3), 227–241, 1988.

25 Feely, R. A., Sabine, C. L., Lee, K., Berelson, W., Kleypas, J., Fabry, V. J., and Millero, F. J.: Impact of anthropogenic CO₂ on the CaCO₃ system in the oceans, *Science*, 305(5682), 362–366, 2004.

Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D., and Hales, B.: Evidence for upwelling of corrosive “acidified” water onto the continental shelf, *Science*, 320, 1490–1492, 2008.

30 Fransson, A., Chierici, M., Anderson, L. C., Bussmann, I., Kattner, G., Jones, E. P., and Swift, J. H.: The importance of shelf processes for the modification of chemical constituents in the waters of the Eurasian Arctic Ocean: implications for carbon fluxes, *Cont. Shelf Res.*, 21(3), 225–242, 2001.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Gammelsrod, T., Leikvin, O., Lienb, V., Budgeit, W. P., Loeng, H., and Maslowski, W.: Mass and heat transports in the NE Barents Sea: Observations and models, *J. Marine Syst.*, 75(1–2), 56–69, 2008.
- 5 Gattuso, J. P., Frankignoulle, M., and Wollast, R.: Carbon and carbonate metabolism in coastal aquatic ecosystems, *Annu. Rev. Ecol. Syst.*, 29, 405–434, 1998.
- Giles, K. A., Laxon, S. W., and Ridout, A. L.: Circumpolar thinning of Arctic sea ice following the 2007 record ice extent minimum, *Geophys. Res. Lett.*, 35(22), L22502, doi:10.1029/2008GL035710, 2008.
- 10 Gordeev, V. V., Martin, J. M., Sidorov, I. S., and Sidorova, M. V.: A reassessment of the Eurasian River input of water, sediment, major elements, and nutrients to the Arctic Ocean, *Am. J. Sci.*, 296, 664–691, 1996.
- Gosink, T. A., Pearson, J. G., and Kelley, J. J.: Gas movement through sea ice, *Nature*, 263, 41–42, 1976.
- Goyet, C. and Poisson, A. P.: New determination of carbonic acid dissociation constants in seawater as a function of temperature and salinity, *Deep-Sea Res.*, 36, 1635–1654, 1989.
- 15 Gosselin, M., Levasseur, M., Wheeler, P. A., Horner, R. A., and Booth, B. C.: New measurements of phytoplankton and ice algal production in the Arctic Ocean, *Deep-Sea Res. Pt. II*, 44, 1623–1644, 1997.
- Grebmeier, J. M., Bates, N. R., and Devol, A.: Continental Margins of the Arctic Ocean and Bering Sea, in: *North American Continental Margins: A Synthesis and Planning Workshop*, edited by: Hales, B., Cai, W.-J., Mitchell, B. G., Sabine, C. L., and Schofield, O., US Carbon Cycle Science Program, Washington DC, 120 pp., 61–72, 2008.
- 20 Guay, C. K. and Falkner, K. K.: Barium as a tracer of Arctic halocline and river water, *Deep-Sea Res. Pt. II*, 44, 1543–1570 1997.
- 25 Häkkinen, S.: An Arctic source for the Great Salinity Anomaly: A simulation of the Arctic ice-ocean system for 1955–1975, *J. Geophys. Res.*, 98, 16 397–16 410, 1993.
- Hameedi, M. J.: Aspects of water column primary productivity in Chukchi Sea during summer, *Mar. Biol.*, 48(1), 37–46, 1978.
- Hansell, D. A., Whitledge, T. E., and Goering, J. J.: Patterns of nitrate utilization and new production over the Bering-Chukchi shelf, *Cont. Shelf Res.*, 13, 601–628, 1993.
- 30 Hill, V. J. and Cota, G. F.: Spatial patterns of primary production in the Chukchi Sea in the spring and summer of 2002, *Deep-Sea Res. Pt. II*, 52, 3344–3354, 2005.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hunt, G. L., Stabeno, P., Walters, G., Sinclair, E., Brodeur, R. D., Napp, J. M., and Bond, N. A.: Climate change and control of the southeastern Bering Sea pelagic ecosystem, *Deep-Sea Res. Pt. II*, 49, 5821–5853, 2002.

IPCC: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Avery, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 996 pp., 2007.

Jones, E. P. and Anderson, L. G.: On the origin of chemical properties of the Arctic Ocean halocline, *J. Geophys. Res.*, 91, 10 759-10 767, 1986.

Jones, E. P., Anderson, L.G., and Wallace, D. W. R.: Tracers of near-surface, halocline and deep waters in the Arctic Ocean: Implications for circulation, *J. Marine Syst.*, 2, 241–255, 1991.

Jones, E. P., Rudels, B., and Anderson, L. G.: Deep waters of the Arctic ocean – origin and circulation, *Deep-Sea Res. Pt. I*, 42, 737–760, 1995.

Jones, E. P., Swift, J. H., Anderson, L. G., Lipzer, M., Civatarese, G., Falkner, K. K., Kattner, G., and McLaughlin, F.: Tracing Pacific water in the North Atlantic Ocean, *J. Geophys. Res.*, 108(C4), 3116, doi:10.1029/2001JC001141, 2003.

Jones, E. P., Anderson, L. G., Jutterstrom, S., Mintrop, L., and Swift, J. H.: Pacific freshwater, river water and sea-ice meltwater across the Arctic Ocean basin: Results from the 2005 Beringia Expedition, *J. Geophys. Res.*, 113, C08012, doi:10.1029/2007JC004124, 2008.

Jutterström, S. and Anderson, L. G.: The saturation of calcite and aragonite in the Arctic Ocean, *Mar. Chem.*, 94, 101–110, 2005.

Kadko, D. and Swart, P.: The source of the high heat and freshwater content of the upper ocean at the SHEBA site in the Beaufort Sea in 1997, *J. Geophys. Res.*, 109, C01022, doi:10.1029/2002JC001734, 2004.

Kaltin, S. and Anderson, L. G.: Uptake of atmospheric carbon dioxide in Arctic shelf seas: Evaluation of the relative importance of processes that influence $p\text{CO}_2$ in water transported over the Bering-Chukchi Sea shelf, *Mar. Chem.*, 94, 67–79, 2005.

Kaltin, S., Anderson, L. G., Olsson, K., Fransson, A., and Chierici M.: Uptake of atmospheric carbon dioxide in the Barents Sea, *J. Marine Syst.*, 38(1–2), 31–45, 2002.

Kattner, G., Lobbes, J. M., Fitznar, H. P., Engbrodt, R., Nothig, E.-M., and Lara, R. J.: Tracing dissolved organic substances and nutrients from the Lena River through Laptev Sea (Arctic),

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Mar. Chem., 65, 25–39, 1999.
- Kelley, J. J.: Carbon dioxide in the surface waters of the North Atlantic and the Barents and Kara Sea, *Limnol. Oceanogr.*, 15, 80–87, 1970.
- Kinney, P., Arhelger, M. E., and Burrell, D. C.: Chemical characteristics of water masses in the Amerasian Basin of the Arctic Ocean, *J. Geophys. Res.*, 75, 4097–4104, 1970.
- Legendre, L., Ackley, S. F., Dieckmann, G. S., Gullicksen, B., Horner, R., Hoshiai, T., Melnikov, I. A., Reeburgh, W. S., Spindler, M., and Sullivan, C. W.: Ecology of sea ice biota: 2. Global significance, *Polar Biol.*, 12, 429–444, 1992.
- Lindsay, R. W. and Zhang, J.: The thinning of Arctic sea ice, 1988–2003: Have we passed a tipping point?, *J. Climate*, 18, 4879–4894, 2005.
- Lundberg, L. and Haugen, P. M.: A Nordic Seas – Arctic Ocean carbon budget from volume flows and inorganic carbon data, *Global Biogeochem. Cy.*, 10, 493–510, 1996.
- Macdonald, R. W., Paton, D. W., Carmack, E. C., and Omstedt, A.: The fresh-water budget and under-ice spreading of Mackenzie River water in the Canadian Beaufort Sea based on salinity and O^{18}/O^{16} measurements in water and ice, *J. Geophys. Res.*, 100(C1), 895–919, 1995.
- Macdonald, R. W., Carmack, E. C., and Paton, D. W. Using the delta O^{18} composition in landfast ice as a record of Arctic estuarine processes, *Mar. Chem.*, 65(1–2), 3–24, 1995.
- Macdonald, R. W., McLaughlin, F. A., and Carmack, E. C.: Fresh water and its sources during the SHEBA drift in the Canada Basin of the Arctic Ocean, *Deep Sea Res. Pt. I*, 49, 1769–1785, 2002.
- Macdonald, R. W., Anderson, L. G., Christensen, J. P., Miller, L. A., Semiletov, I. P., and Stein, R.: The Arctic Ocean, in: *Carbon and Nutrient Fluxes in Continental Margins: A Global Synthesis*, edited by: Liu, K. K., Atkinson, L., Quinones, R., and Talue-McManus, L., *Global Change – The IGBP Series*, Springer, New York, 291–303, 2009.
- Macklin, S. A., Hunt, G. L., and Overland, J. E.: Variability in the Bering Sea ecosystem, *Prog. Oceanogr.*, 55, 1–4, 2002.
- Makkaveev, P. N.: The dissolved inorganic carbon in the Kara Sea and in the mouths of the Ob and Yenisei Rivers, *Okeanologiya+*, 34, 668–672, 1994.
- Manley, T. O. and Hunkins, K.: Mesoscale eddies of the Arctic Ocean, *J. Geophys. Res.*, 90, 4911–4930, 1985.
- Maslanik, J., Drobot, S., Fowler, C., Emery, W., and Barry, R.: On the Arctic climate paradox and the continuing role of atmospheric circulation in affecting sea ice conditions, *Geophys.*

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Res. Lett., 34, L03711, doi:10.1029/2006GL028269, 2007.

Maslowski, W., Marble, D. C., Walczowski, W., and Semtner, A. J.: On large-scale shifts in the Arctic Ocean and sea-ice conditions during 1979–98, *Ann. Glaciol.*, 33, 545–550, 2001.

Mathis, J. T., Pickart, R. S., Hansell, D. A., Kadko, D., and Bates, N. R.: Eddy transport of organic carbon and nutrients from the Chukchi Shelf: Impact on the upper halocline of the western Arctic Ocean, *J. Geophys. Res.*, 112, C05011, doi:10.1029/2006JC003899, 2007a.

Mathis, J. T., Hansell, D. A., Kadko, D., Bates, N. R., and Cooper, L. W.: Determining net DOC production in the hydrographically complex western Arctic Ocean, *Limnol. Oceanogr.*, 52(5), 1789–1799, 2007b.

Mathis, J. T., Hansell, D. A., and Bates, N. R.: Interannual variability of dissolved inorganic carbon distribution and net community production during the Western Arctic Shelf-Basin Interactions Project, *Deep-Sea Res. Pt. II*, 56, 1213–1222, 2009.

Mehrbach, C., Culberson, C. H., Hawley, J. E., and Pytkowicz, R. M.: Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure, *Limnol. Oceanogr.*, 18, 897–907, 1973.

Merico, A., Tyrrell, T., Lessard, E. J., Oguz, T., Stabeno, P. J., Zeeman, S. I., and Whittedge, T. E.: Modelling phytoplankton succession on the Bering Sea shelf: role of climate influences and trophic interactions in generating *Emiliania huxleyi* blooms 1997–2000, *Deep-Sea Res. Pt. I*, 51(12), 1803–1826, 2004.

Millero, F. J.: Thermodynamics of the carbon dioxide system in the oceans, *Geochim. Cosmochim. Ac.*, 59, 661–677, 1995.

Mucci, A.: The solubility of calcite and aragonite in seawater at various salinities, temperatures, and one atmosphere total pressure, *Am. J. Sci.*, 283, 780–799, 1983.

Mucci, A., Forest, A., Fortier, L., Fukuchi, M., Grant, J., Hattori, H., Hill, P., Lintern, G., Makbe, R., Magen, C., Miller, L., Sampei, M., Sasaki, H., Sundby, B., Walker, T., and Wassman, P.: Chapter 7: Organic and inorganic fluxes, in: *On Thin Ice: A Synthesis of the Canadian Arctic Shelf Exchange Study (CASES)*, edited by: Fortier, L., Barber, D., and Michaud, J., University of Manitoba Publication, 113–141, 2008.

Muench, R. D., Gunn, J. T., Whittedge, T. E., Schlosser, P., and Smethie, W.: An Arctic Ocean cold core eddy, *J. Geophys. Res.*, 105(C10), 23 997–24 006, 2000.

Murata, A. and Takizawa, T.: Summertime CO₂ sinks in shelf and slope waters of the western Arctic Ocean, *Cont. Shelf Res.*, 23(8), 753–776, 2003.

Mysak, L. A., Manak, D. K., and Marsden, R. F.: Sea ice anomalies in the Greenland and

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Labrador Seas during 1901-1984 and their relation to an interdecadal Arctic climate cycle, *Clim. Dynam.*, 5, 111–133, 1990.

Nagurnyi, A. P.: On the role of Arctic sea-ice in seasonal variability of carbon dioxide concentration in Northern Latitudes, *Russ. Meteorol. Hydrol.*, 33(1), 43–47, 2008.

5 Nakaoka, S., Aiki, S., Nakazawa, T., Hashida, G., Morimoto, S., Yamanouchi, T., and Yoshikawa-Inoue, H.: Temporal and spatial variations of oceanic $p\text{CO}_2$ and air-sea CO_2 flux in the Greenland Sea and the Barents Sea, *Tellus*, 58(2), 148–161, 2006.

Napp, J. M., Baier, C. T., Brodeur, R. D., Coyle, K. O., Shiga, N., and Mier, K.: Interannual and decadal variability in zooplankton communities of the southeast Bering Sea shelf, *Deep-Sea Res. Pt. II*, 49(26), 5991–6008, 2002.

10 Nikolopoulos, A., Pickart, R. S., Fratantoni, P. S., Shimada, K., Torres, D. J., and Jones, E. P., 2009. The western Arctic boundary current at 152oW: Structure, variability, and transport, *Deep-Sea Res. Pt. II*, in press, 2009.

Nitishinsky, M., Anderson, L. G., and Holemann, J. A.: Inorganic carbon and nutrient fluxes on the Arctic Shelf, *Cont. Shelf Res.*, 27, 1584–1599, 2007.

15 Olsen, A., Bellerby, R. G. J., Johannessen, T., Omar, A. M., and Skjelvan, I.: Interannual variability in the wintertime air-sea flux of carbon dioxide in the northern North Atlantic, 1981–2001, *Deep-Sea Res. Pt. I*, 50(10–11), 1323–1338, 2003.

Omar, A. M., Johannessen, T., Kaltin, S., and Olsen, A.: Anthropogenic increase of oceanic $p\text{CO}_2$ in the Barents Sea surface water, *J. Geophys. Res.*, 108(C12), 3388, 2003.

20 Omar, A., Johannessen, T., Bellerby, R. G. J., Olsen, A., Anderson, L. G., and Kivimäe, C.: Sea ice and brine formation in Storfjorden: Implications for the Arctic wintertime air-sea CO_2 flux, in: *The Nordic Sea: An Integrated Perspective*, Geophysical Monograph, 158, 177–187, 2005.

25 Omar, A. M., Johannessen, T., Olsen, A., Kaltin, S., and Rey, F.: Seasonal and interannual variability of the air-sea CO_2 flux in the Atlantic sector of the Barents Sea, *Mar. Chem.*, 104, 203–213, 2007.

Overland, J. E. and Wang, M.: The Arctic climate paradox: The recent decrease of the Arctic Oscillation, *Geophys. Res. Lett.*, 32(6), L06701, doi:10.1029/2004GL021752, 2005.

30 Pabi, S., van Dijken, G. L., and Arrigo, K. R.: Primary production in the Arctic Ocean, 1998–2006, *J. Geophys. Res.*, 113, C08005, doi:10.1029/2007JC004578, 2008.

Paquette, R. G. and Bourke, R. H.: Observations on the coastal current of arctic Alaska, *J. Mar. Res.*, 32, 195–207, 1974.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Parkinson, C. L.: Variability of Arctic sea ice: The view from space, and 18-year record, *Arctic*, 53, 341–358, 2000.
- Perovich, D. K. Light, B., Eicken, H., Jones, K. F., Runciman, K., and Nghiem, S. V.: Increasing solar heating of the Arctic Ocean and adjacent seas, 1979–2005: Attribution and role in the ice-albedo feedback, *Geophys. Res. Lett.*, 34, L19505, doi:10.1029/2007GL031480, 2007.
- Pickart, R. S., Weingartner, T. J., Zimmermann, S., Torres, D. J., and Pratt, L. J.: Flow of winter-transformed Pacific water into the western Arctic, *Deep-Sea Res. Pt. II*, 52, 3175–3198, 2005.
- Pickart, R. S., Fratantoni, P. S., Goldsmith, R. A., Moore, G. W. K., Agnew, T., and Vandeweghe, J.: Upwelling in the western Arctic boundary current north of Alaska, *EOS T. Am. Geophys. Un.*, 87(36), Ocean Sci. Meet. Suppl., Abstract OS350-07, 2006.
- Pickart, R. S., Moore, G. W. K., and Yang, J.: On the nature of upwelling storms in the Western Arctic, *J. Geophys. Res.*, submitted, 2009.
- Piepenburg, D.: Recent research on Arctic benthos: common notions need to be revised, *Polar Biol.*, 28(10), 733–755, 2005.
- Pipko, I. I., Semiletov, I. P., Tishchenko, P. Y., Pugach, S. P., and Christensen, J. P.: Carbonate chemistry dynamics in Bering Strait and the Chukchi Sea, *Prog. Oceanogr.*, 55(1–2), 77–94, 2002.
- Rigor, I. G., Wallace, J. M., and Colony, R. L.: Response of sea ice to the Arctic Oscillation, *J. Climate*, 15(18), 2648, 2002.
- Rigor, I. G. and Wallace, J. M.: Variations in the age of Arctic sea ice extent, *Geophys. Res. Lett.*, 31(9), L09401, doi:10.1029/2004GL019492, 2004.
- Roach, A. T., Aagaard, K., Pease, C. H., Salo, S. A., Weingartner, T. J., Pavlov, V., and Kulakov, M.: Direct measurements of transport and water properties through Bering Strait, *J. Geophys. Res.*, 100, 18 443–18 457, 1995.
- Rothrock, D. A. and Zhang, J.: Arctic Ocean sea ice volume: What explains its recent depletion?, *J. Geophys. Res.*, 110, C01002, doi:10.1029/2004JC002282, 2005.
- Rothrock, D. A., Zhang, J., and Yu, Y.: The arctic ice thickness anomaly of the 1990s: A consistent view from observations and models, *J. Geophys. Res.*, 108, 3083, doi:10.1029/2001JC001208, 2003
- Roy, R. N., Roy, L. N., Vogel, R., Moore, C. P., Pearson, T., Good, C. E., Millero, F. J., and Campbell, D.: Determination of the ionization constants of carbonic acid in seawater, *Mar. Chem.*, 44, 249–268, 1993.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Royal Society: Ocean acidification due to increasing atmospheric carbon dioxide, The Clyvedon Press, Ltd, Cardiff, UK, 2005.
- Rudels, B., Anderson, L. G., and Jones, E. P.: Formation and evolution of the surface mixed layer and halocline of the Arctic Ocean, *J. Geophys. Res.-Oceans*, 101(C4), 8807–8821, 1996.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F.J., Peng, T.-H., Kozyr, A., Ono, T., and Rios, A. F.: The oceanic sink for anthropogenic CO₂, *Science*, 305, 367–371, 2004.
- Sakshaug, E.: Primary and secondary production in the Arctic seas, in: *The Organic Carbon Cycle in the Arctic Ocean*, edited by: Stein, R. and Macdonald, R. W., Springer Publishing Company, Heidelberg, 57–81, 2004.
- Salisbury, J. E., Green, M., Hunt, C., and Campbell, J.: Coastal acidification by rivers: A threat to shellfish?, *EOS T. Am. Geophys. Un.*, 89(50), 513–528, 2008.
- Salmon, D. K. and McRoy, C. P.: Nutrient-based tracers in the Western Arctic: A new lower halocline water defined, in: *The Polar Oceans and Their Role in Shaping the Global Environment*, edited by: Johannessen, O. M., Muench, R.D., and Overland, J.E., American Geophysical Union, Geophysical Monograph, 85, 47–61, 1994.
- Sambrotto, R. N., Goering, J. J., and McRoy, C. P.: Large yearly production of phytoplankton in the western Bering Sea, *Science*, 225, 1147–1155, 1984.
- Santana-Casiano, J. M., Gonzalez-Davila, M., Rueda, M.-J., Llinas, O., and Gonzalez-Davila, E.-F.: The interannual variability of oceanic CO₂ parameters in the western Atlantic subtropical gyre at the ESTOC site, *Global Biogeochem. Cy.*, 21, GB1015, doi:10.1029/2006GB002788, 2007.
- Schlosser, P., Swift, J. H., Lewis, D., and Pfirman, S.: The role of the large-scale Arctic Ocean circulation in the transport of contaminant, *Deep-Sea Res. Pt. II*, 42, 1341–1368, 1995.
- Semiletov, I. P.: Aquatic sources of CO₂ and CH₄ in the Polar regions, *J. Atmos. Sci.*, 56(2), 286–306, 1999.
- Semiletov, I., Makshatas, A.I., Akasofu, S.-I., and Andreas, E. L.: Atmospheric CO₂ balance: The role of arctic sea ice, *Geophys. Res. Lett.*, 31, L05121, doi:10.1029/2003GL017996, 2004.
- Semiletov, I. P., Pipko, I. I., Repina, I., and Shakhova, N. E.: Carbonate chemistry dynamics and carbon dioxide fluxes across the atmosphere-ice-water interface in the Arctic Ocean, *J. Marine Syst.*, 66, 204–226, 2007.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Serreze, M. C., Holland, M. M., and Stroeve, J.: Perspectives on the Arctic's shrinking sea-ice cover, *Science*, 315(5818), 1533–1536, 2007.
- Shimada, K., Kamoshida, T., Itoh, M., Nishino, S., Carmack, E., McLaughlin, F., Zimmermann, S., and Proshutinsky, A.: Pacific Ocean inflow: influence on catastrophic reduction of sea ice cover in the Arctic Ocean, *Geophys. Res. Lett.*, 33(8), L08605, doi:10.1029/2005GL025624, 2006.
- Skjelvan, I., Olsen, A., Anderson, L. G., Bellerby, R. G. J., Falck, E., Kasajima, Y., Kivimäem, C., Omar, A., Rey, F., Olsson, K. A., Johannessen, T., Heinze, C.: A review of the biogeochemistry of the Nordic Seas and Barents Sea – with focus on the inorganic carbon cycle, in: *The Nordic Seas: An Integrated Perspective Oceanography, Climatology, and Modelling*, edited by: Drange, H., Dokken, T. M., Furevik, T., et al., American Geophysical Union, Washington DC, 157–175, 2005.
- Skjelvan, I. M., Johannessen, T., and Miller, L.: Interannual variability of $f\text{CO}_2$ in the Greenland and Norwegian Seas, *Tellus B*, 51, 477–489, 1999b.
- Smith, D. M.: Recent increase in the length of the melt season of perennial Arctic sea ice, *Geophys. Res. Lett.*, 25(5), 655–8, 1998.
- Smith, S. V. and Hollibaugh, J. T.: Coastal metabolism and the oceanic organic carbon balance, *Rev. Geophys.*, 31, 75–89, 1985.
- Spall, M. A., Pickart, R. S., Fratantoni, P. S., and Plueddemann, A. J.: Western Arctic shelfbreak eddies: Formation and transport, *J. Phys. Oceanogr.*, 38, 1644–1668, 2008.
- Solomons, S., Qin, D., Manning, M., Marquis, M., Averyt, K., Tignor, M. M. B., Miller, H. L., and Chen, Z.: Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge Univ. Press, Cambridge and New York, 2007.
- Stabeno, P. J., Bond, N. A., Kachel, N. B., Salo, S. A., and Schumacher, J. D.: On the temporal variability of the physical environment over the southeastern Bering Sea, *Fish. Oceanogr.*, 10, 81–98, 2001.
- Stabeno, P. J., Kachel, N. B., Sullivan, M., and Whittedge, T. E.: Variability of physical and chemical characteristics along the 70-m isobath of the southeastern Bering Sea, *Deep-Sea Res. Pt. II*, 49, 5931–5943, 2002.
- Steele, M. and Boyd, T.: Retreat of the cold halocline layer in the Arctic Ocean, *J. Geophys. Res.*, 103(C5), 10 419–10 435, 1998.
- Steinacher, M., Joos, F., Frölicher, T. L., Plattner, G.-K., and Doney, S. C.: Imminent ocean acid-

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ification in the Arctic projected with the NCAR global coupled carbon cycle-climate model, *Biogeosciences*, 6, 515–533, 2009, <http://www.biogeosciences.net/6/515/2009/>.

5 Stroeve, J., Holland, M. M., Meier, W., Scambos, T., and Serreze, M.: Arctic sea ice decline: Faster than forecast, *Geophys. Res. Lett.*, 34(9), L09501, doi:10.1029/2007GL029703, 2007.

Subba Rao, D. V. and Platt, T.: Primary productivity of Arctic waters, *Polar Biol.*, 3, 191–201, 1984.

10 Takahashi, T., Olafsson, J., Goddard, J. G., Chipman, D. W., and Sutherland, S. G.: Seasonal variation of CO₂ and nutrients in the high-latitude surface oceans: a comparative study, *Global Biogeochem. Cy.*, 7(4), 843–878, 1993.

Takahashi, T., Sutherland, S. G., Sweeney, C., Poisson, A. P., Metz, N., Tilbrook, B., Bates, N. R., Wanninkhof, R. H., Feely, R. A., Sabine, C. L., and Olafsson, J.: Biological and temperature effects on seasonal changes of pCO₂ in global ocean surface waters, *Deep-Sea Res. Pt. II*, 49, 1601–1622, 2002.

15 Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G. E., Chavez, F. P., Watson, A. J., Bakker, D. C. E., Schuster, U., Metz, N., Yoshikawa-Inoue, H., Olafsson, J., Arnarson, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R. J., Nojiri, Y., Wong, C. S., Delille, B., and Bates, N. R., and de Baar, H. J. W.: Climatological mean and decadal change in surface ocean pCO₂, and net sea-air CO₂ flux over the global oceans, *Deep-Sea Res. Pt. II*, 56, 554–577, 2009.

20 Thomas, H., Schiettecatte, L.-S., Suykens, K., Koné, Y. J. M., Shadwick, E. H., Prowe, A. E. F., Bozec, Y., de Baar, H. J. W., and Borges, A. V.: Enhanced ocean carbon storage from anaerobic alkalinity generation in coastal sediments, *Biogeosciences*, 6, 267–274, 2009, <http://www.biogeosciences.net/6/267/2009/>.

25 Tucker, W. and Cate, D.: The 1994 Arctic ocean Section: The first major scientific crossing of the Arctic Ocean, US Army Cold Regions Research and Engineering Laboratory, Special Report, 96–23, 117 pp, 1996.

30 Wadley, M. R. and Bigg, G. R.: Impact of flow through the Canadian Archipelago and Bering Strait on the North Atlantic and Arctic circulation: An ocean modelling study, *Q. J. Roy. Meteor. Soc.*, 128, 2187–2203, 2002.

Walsh, J. E. and Chapman, W. L.: Arctic contribution to upper ocean variability in the North Atlantic, *J. Climate*, 3, 1462–1473, 1990.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Walsh, J. J.: Importance of continental margins in the marine biogeochemical cycling of carbon and nitrogen, *Nature*, 350, 53–55, 1991.
- Walsh, J. J., Premuzic, E. T., Gaffney, J. S., Rowe, G. T., Harbottle, G., Stoenner, R. W., Balsam, W. L., Betzer, P. R., and Macko, S. A.: Organic storage of CO₂ on the continental slope off the Mid-Atlantic bight, the southeastern Bering Sea, and the Peru coast, *Deep-Sea Res.*, 32, 853–883, 1985.
- Wallace, D. W. R., Moore, R. M., and Jones, E. P.: Ventilation of the Arctic Ocean cold halocline: rates of diapycnal and isopycnal transport, oxygen utilization and primary production inferred using chlorofluoromethane distributions, *Deep-Sea Res.*, 34, 1957–1979, 1987.
- Wang, J., Ikeda, M., Zhang, S., and Gerdes, R.: Linking the northern hemisphere sea-ice reduction trend and the quasi-decadal arctic sea-ice oscillation, *Clim. Dynam.*, 24(2–3), 115–130, 2005.
- Wang, W. Q., Yang, X. T., Huang, H. B., and Chen, L. Q.: Investigation on distribution and fluxes of sea-air CO₂ of the expedition areas in the Arctic Ocean. *Sci. China Ser. D*, 46, 569–580, 2003.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.-Oceans*, 97, 7373–7382, 1992.
- Wanninkhof, R. and McGillis, W. R.: A cubic relationship between air-sea CO₂ exchange and windspeed, *Geophys. Res. Lett.*, 26(13), 1889–1892.
- Weingartner, T. J., Cavalieri, D. J., Aagaard, K., and Sasaki, Y.: Circulation, dense water formation and outflow on the northeast Chukchi Sea shelf, *J. Geophys. Res.*, 103, 7647–7662, 1998.
- Weingartner, T. J., Danielson, S., Sasaki, Y., Pavlov, V., and Kulakov, M.: The Siberian Coastal Current: A wind and buoyancy-forced arctic coastal current, *J. Geophys. Res.*, 104, 29 697–29 713, 1999.
- Weingartner, T., Aagaard, K., Woodgate, R., Danielson, S., Sasaki, Y., and Cavalieri, D.: Circulation on the north central Chukchi Sea shelf, *Deep-Sea Res. Pt. II*, 52, 3150–3174, 2005.
- Wheeler, P. A., Gosselin, M., Sherr, E., Thibault, D., Kirchman, D. L., Benner, R., and Whitledge, T. E.: Active cycling of organic carbon in the central Arctic Ocean, *Nature*, 380(6576), 697–699, 1996.
- Wijffels, S. E., Schmitt, R. W., Bryden, H. L., and Stigebrandt, A.: Transport of freshwater by the oceans, *J. Geophys. Res.*, 22, 155–162, 1992.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Wilson, C. and Wallace, D. W. R.: Using the nutrient ratio NO/PO as a tracer of continental shelf waters in the central Arctic Ocean. *J. Geophys. Res.*, 95, 22 193–22 208, 1990.
- Winn, C. D., Mackenzie, F. T., Carillo, C. J., Sabine, C. L., and Karl, D. M.: Air-sea carbon dioxide exchange in the North Pacific subtropical gyre: Implications for the global carbon budget, *Global Biogeochem. Cy.*, 6, 157–163, 1994.
- Winsor, P. and Chapman, D. C.: Pathways of Pacific water across the Chukchi Sea: A numerical model study, *J. Geophys. Res.*, 109, CO3002, doi:10.1029/2003JC001962, 2004.
- Wollast, R.: Evaluation and comparison of the global carbon cycle in the coastal zone and in the open ocean, in: *The Sea 10. The Global Coastal Ocean*, edited by: Brink, K. H. and Robinson, A. R., Wiley, New York, 213–252, 1998.
- Woodgate, R. A. and Aagaard, K.: Revising the Bering Strait freshwater flux into the Arctic Ocean, *Geophys. Res. Lett.*, 32, L02602, doi:10.1029/2004GL021747, 2005.
- Woodgate, R. A., Aagaard, K., and Weingartner, T. J.: A year in the physical oceanography of the Chukchi Sea: Moored measurements from autumn 1990–1991, *Deep-Sea Res. Pt. II.*, 52, 3116–3149, 2005.
- Woodgate, R. A., Aagaard, K., and Weingartner, T. J.: Monthly temperature, salinity and transport variability of the Bering Strait Throughflow, *Geophys. Res. Lett.*, 32(4), L04601, doi:10.1029/2004GL021880, 2005.
- Yager, P. L., Wallace, D. W. R., Johnson, K. M., Smith, W. O., Minnett, P. J., and Deming, J. W.: The Northeast Water Polynya as an atmospheric CO₂ sink: A seasonal rectification hypothesis, *J. Geophys. Res.*, 100, 4389–4398, 1995.
- Yamamoto-Kawai, M., Tanaka, N., and Pivovarov, S.: Freshwater and brine behaviors in the Arctic Ocean deduced from historical data of delta O-18 and alkalinity (1929–2002 AD). *J. Geophys. Res.*, 110, C10003, doi:10.1029/2004JC002793, 2005.

BGD

6, 6695–6747, 2009

The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Areas, depths, residence times, air-sea CO₂ exchange rates expressed in mmol C m⁻² d⁻¹ and g C m⁻² d⁻¹, and annual air-sea CO₂ exchange rate expressed in Tg C (10¹² g C). Negative air-sea CO₂ exchange rates indicates ocean uptake of CO₂ (i.e., CO₂ sink).

	Area ^a (km ²)	Depth ^a (m)	Residence ^a (years)	Air-Sea CO ₂ flux	Air-Sea CO ₂ flux	Annual ^f CO ₂ flux	Reference
				Cm ⁻² d ⁻¹ (mmol (g C m ⁻² yr ⁻¹)) (Tg C yr ⁻¹)			
Barents Sea	1512000	200	1 ^b	n/a	-46±18	-70±27	Nakaoka et al. (2007)
				n/a	-51±8	-77±12	Omar et al. 2007
				n/a	-44±10	-67±15	Fransson et al. (2001) ^g
				n/a	-29±11	-44±16	Kaltin et al. (2002) ^g
Kara Sea	926000	131	2.5	n/a	-1	-1.0	Fransson et al. (2001) ^g
				n/a	n/a	-5.7	Anderson et al. (1998a, b) ^g
Laptev Sea	498000	48	3.5±2	-2.1	-2.4	-1.2	Nitishinsky et al. (2007) ^g
				n/a	n/a	-4.3	Anderson et al. (1998a, b) ^g
East Siberian Sea	987000	58	3.5±2	-1 to -10.9	n/a	-1.2/-13	Semiletov et al. (2007)
				+0.3	n/a	+0.3	Nitishinsky et al. (2007) ^g
				n/a	n/a	-5.9	Anderson et al. (1998a, b) ^g
Chukchi Sea	620000	80	0.2-1.2	-12	-17 ^d	-11	Murata and Takizawa (2003)
				-40±22	-48±10 ^d	-36±6	Bates et al. (2006)
				n/a	-61±10 ^g	-46±6	Bates et al. (2006)
				n/a	-86±22	-53±14	Kaltin and Anderson (2001) ^g
Beaufort Sea	178000	124	0.5-1.0	n/a	n/a	-2.9	Anderson et al. (1998a, b) ^g
				-12	-17 ^d	-2	Murata and Takizawa (2003)
Canadian Archipelago	1490000	290	0.5-12	?	?	-16/-24	scaled to Beaufort Sea shelf
Central basin	4489000	2748	2-30 ^c	<-1 to -3	-4 to -12	-6 to -19	Bates et al. (2006)
Arctic Ocean	10700000					-65	low estimate from above
						-175	high estimate from above
						-81	low estimate from above ^k
						-199	high estimate from above ^k
						-129±65	Anderson et al. (1990) ^g
						-70±65	Anderson et al. (1994) ^g
Arctic Ocean (other studies)						-110	Lundberg and Haugen (1996) ^{g,h}
						-24±17	Anderson et al. (1998b) ^g
						-41±18	Anderson et al. (1998b) ^{g,i}
						-31	Kaltin and Anderson (2005) ^g
						-66	Bates (2006)

Notes: ^a Macdonald et al. (2009) estimates of shelf areas, mean depth (m) and residence times; ^b Kaltin et al. (2002) estimate of Barents Sea shelf water residence time; ^c residence time of surface and halocline waters (upper 0-300 m only) from Macdonald et al. (2009); ^d summertime open water only (~100 d); ^e including wintertime air-sea CO₂ exchange; ^f annual flux estimated from annual uptake (g C yr⁻¹) multiplied by area of Arctic shelf from Macdonald et al. (2009); ^g mass balance assessment; ^h includes Norwegian Sea (Lundberg and Haugen, 1996); ⁱ includes river contribution (Anderson et al., 1998b), and; ^k includes air-sea CO₂ exchange rates for the Canadian Archipelago scaled to estimates of the Beaufort Sea shelf.

The Arctic Ocean marine carbon cycle

N. R. Bates and J. T. Mathis

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**The Arctic Ocean
marine carbon cycle**

N. R. Bates and
J. T. Mathis




Fig. 1. Schematic of the Arctic Ocean, central basin (Canada and Eurasian Basins) and Arctic continental shelves (with approximate boundaries of each Arctic shelf sea), and major rivers draining into the region. Inset: characterization of Arctic Ocean continental shelves according to Carmack and Wassman (2006).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The Arctic Ocean marine carbon cycle

N. R. Bates and
J. T. Mathis




Fig. 2. Surface seawater $p\text{CO}_2$ values computed from DIC and TA data collected in the Arctic Ocean and part of the CARINA dataset (data obtained at <http://cdiac.ornl.gov/>).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

