

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

An iron budget during the natural iron fertilisation experiment KEOPS (Kerguelen Islands, Southern Ocean)

**An iron budget
during the natural
iron fertilisation
experiment KEOPS**

F. Chever et al.

F. Chever^{1,2}, **G. Sarthou**^{1,2}, **E. Bucciarelli**^{1,2}, **S. Blain**^{3,4}, and **A. R. Bowie**⁵

¹Université Européenne de Bretagne, France

² Université de Brest, CNRS, IRD, UMR 6539 LEMAR, IUEM, Technopôle Brest Iroise, Place Nicolas Copernic, 29280 Plouzané, France

³Laboratoire Océanographie Biologique de Banyuls sur Mer, Université Pierre et Marie Curie, Paris 06, quai Fontaulé, 66650 Banyuls sur Mer, France

⁴Laboratoire Océanographie Biologique de Banyuls sur Mer, CNRS, quai Fontaulé, 66650 Banyuls sur Mer, France

⁵Antarctic Climate and Ecosystems Cooperative Research Centre (ACE CRC), University of Tasmania, Hobart, Australia

Received: 29 May 2009 – Accepted: 22 June 2009 – Published: 10 July 2009

Correspondence to: G. Sarthou (geraldine.sarthou@univ-brest.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Total dissolvable iron (TDFe) was measured in the water column above and in the surrounding of the Kerguelen Plateau (Indian sector of the Southern Ocean) during the Kerguelen Ocean Plateau compared Study (KEOPS) cruise. TDFe concentrations ranged from 0.90 to 65.6 nmol L⁻¹ above the plateau and from 0.34 to 2.23 nmol L⁻¹ offshore of the plateau. Station C1 located south of the plateau, near Heard Island, exhibited very high values (329–770 nmol L⁻¹). Apparent particulate iron (Fe_{app}), calculated as the difference between the TDFe and the dissolved iron measured on board (DFe) represented 95±5% of the TDFe above the plateau, suggesting that particles and refractory colloids largely dominated the iron pool. This paper presents a budget of DFe and Fe_{app} above the plateau. Lateral advection of water that had been in contact with the continental shelf of Heard Island seems to be the predominant source of Fe_{app} and DFe above the plateau, with a supply of 9.7±2.3×10⁶ and 8.3±6.7×10³ mol d⁻¹, respectively. The residence times of 1.7 and 48 days estimated for Fe_{app} and DFe, respectively, indicate a rapid turnover in the surface water. A comparison between Fe_{app} and total particulate iron (TPFe) suggests that the total dissolved fraction is mainly constituted of small refractory colloids. This fraction does not seem to be a potential source of iron to the phytoplankton in our study. Finally, when taking into account the lateral supply of dissolved iron, the seasonal carbon sequestration efficiency was estimated at 154 000 mol C (mol Fe)⁻¹, which is 4-fold lower than the previously estimated value in this area but still 18-fold higher than the one estimated during the other study of a natural iron fertilisation experiment, CROZEX.

1 Introduction

Iron (Fe) is essential for the growth of marine phytoplankton (Sunda, 1989), and plays an important role in biochemical reactions such as photosynthesis and nitrate reduction (Rueter and Ades, 1987; Kutska et al., 2002). Artificial and natural Fe fertilisations in

BGD

6, 6803–6837, 2009

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

High Nutrient Low Chlorophyll (HNLC) regions of the World's oceans have shown that iron inputs enhance phytoplankton growth and partly control the major biogeochemical cycles of elements important for Earth's climate (Boyd et al., 2000; Coale et al., 2004; Blain et al., 2007; Pollard et al., 2009). However, many uncertainties exist following such fertilisation experiments, for example the carbon sequestration efficiency, i.e. the amount of carbon exported to the deep ocean per unit iron added.

Moreover, the biogeochemical cycle of iron is still not well understood. This is partly due to its complex physico-chemical speciation. Iron exists in different forms and the availability of these forms to phytoplankton is not well established (Bruland and Rue, 2001). The physical speciation is operationally defined as the partitioning between dissolved iron (DFe, $< 0.2 \mu\text{m}$) and particulate iron (PFe, $> 0.2 \mu\text{m}$). Although particulate fraction may be the dominant pool of total iron in the water column (de Baar and de Jong, 2001), the majority of the studies had focused on the dissolved phase which is thought to be a proxy for the bioavailable form (Frew et al., 2006; Morel et al., 2008). Exchanges between the different physical pools of iron take place in seawater, and an understanding of the distribution of iron in those pools, as well as of the fluxes between them is needed to better constrain the iron cycle in the ocean (Frew et al., 2006). Only a few studies have shown that part of the particulate iron may be used by phytoplankton (Johnson et al., 2001, Maldonado et al., 2001). Small-particles below the defined colloidal size range ($0.02\text{--}0.2 \mu\text{m}$; Wu et al., 2001), and included in the dissolved phase, have also been considered to be a possible bioavailable form of iron to phytoplankton although less available than the truly soluble iron (Wu et al., 2001).

Acidified unfiltered samples allow the determination of total dissolvable iron (TDFe). That fraction represents the sum of the dissolved iron (i.e. the labile dissolved iron which is the fraction that is released after a few days of acidification) plus the refractory colloidal iron (i.e., that fraction that dissolved after more than a few days of acidification) plus the fraction of particulate material that dissolves during extended (>6 months) acid storage (iron adsorbed on lithogenic or biogenic particles, contained within biogenic particles and labile lithogenic particulate iron) (de Baar et al., 1999; Löscher et al.,

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1997; Bowie and Sedwick, 2004). In the literature, the set of TDFe observations is much smaller than that of DFe. Studies of TDFe can however give important information about the iron cycle, notably on the sources of iron to the ocean. Indeed, lithogenic material deposited on continental shelf sediments and atmospheric dust inputs, which are the most important Fe sources in the surface waters of the open ocean, have been correlated with elevated concentrations of TDFe (Jickells et al., 2005; Elrod et al., 2004; Sarthou et al., 1997; Sedwick et al., 2008; Croot et al., 2004).

During the KEOPS (Kerguelen Ocean and Plateau compared Study) cruise, a multi-tracer approach was used to identify and quantify natural iron fertilisation over the plateau. DFe ($>0.2 \mu\text{m}$) was analysed on board and it represented the labile dissolved iron (i.e., soluble Fe and colloidal Fe dissolving after a few days of acid storage). The study of this fraction clearly demonstrated the existence of a dissolved iron-rich reservoir above the plateau below the mixed layer (Blain et al., 2007). At the date of the cruise (January–February 2005), DFe enrichment was partly due to inputs from the sediment above the plateau and/or to regeneration of biogenic particles. Predominant mechanisms allowing DFe transport into surface waters were shown to be diapycnal diffusive flux and winter mixing (Blain et al., 2008). However, other geochemical tracers have suggested that the dissolution of lithogenic material transported over the plateau by lateral advection might also be a source of iron to this region (Zhang et al., 2008; van Beek et al., 2008; Jacquet et al., 2008).

The present work focuses on the distribution of TDFe above the Kerguelen plateau and in the surrounding waters, and uses this parameter as an additional tracer to better constrain the Fe cycle in a naturally fertilised area. The apparent particulate iron ($\text{Fe}_{\text{app}} = \text{TDFe} - \text{DFe}$, de Baar et al., 1999). A budget of DFe and Fe_{app} for the region located above the plateau is presented and the fluxes exchanged between these two pools are calculated.

An iron budget during the natural iron fertilisation experiment KEOPSF. Chever et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2 Material and methods

2.1 Study area

During the KEOPS cruise (18 January to 13 February 2005), three transects were studied (A, B and C). Seven stations located above the Kerguelen plateau (A3, B1, B3, B5, B7, C1 and C5; water depth from 150 to 607 m) and two stations located outside the plateau (B11, C11; water depth > 3000 m) were sampled for TDFe (unfiltered samples) (Fig. 1).

The hydrology and the circulation around and above the Kerguelen Plateau have been described by Park et al. (2008a,b). Briefly the Kerguelen Plateau constitutes a barrier to the eastward flowing Antarctic Circumpolar Current (ACC). Most of the ACC is deflected north of the Kerguelen Islands but a remainder passes between the Kerguelen Islands and Antarctica. Above the plateau, the remainder of the ACC comes from the western part of the plateau (see Fig. 1). Currents travel along the western flank of the plateau, passing south and east of Heard Island, before riding up above the plateau. South of the plateau, a branch of the Fawn Trough Current (FTC) flows toward the north along the eastern flank of the plateau (near stations B11 and C11).

2.2 Sampling and analyses

Samples were collected with acid-cleaned 12L Go-Flo bottles mounted on a Kevlar cable. All sampling was carried out in a clean-room container. Unfiltered samples were collected in acid-cleaned 125 mL high density polyethylene (HDPE) bottles and immediately acidified with ultrapure hydrochloric acid (HCl, Merck, 250 μ L, final pH 1.7). Samples were stored at room temperature and analysed 18 months later in the shore-based laboratory (LEMAR, Brest, France), in order to release the most refractory Fe species into the dissolved form (Löscher et al., 1997; Bowie and Sedwick, 2004).

TDFe analyses were performed by flow injection analysis (FIA) with on line preconcentration and chemiluminescence detection (Obata et al., 1993; Sarthou et al., 2003),

BGD

6, 6803–6837, 2009

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

identical to the method used for DFe (Blain et al., 2008). The mean blank was equal to $0.07 \pm 0.07 \text{ nmol L}^{-1}$ ($n = 9$) and the detection limit equal to three times the standard deviation of the blank was $0.02 \pm 0.02 \text{ nmol L}^{-1}$ ($n = 9$). The individual contributions to the total blank from ultrapure® hydrochloric acid (MERCK), suprapure® ammonia (MERCK), and ammonium acetate buffer purified three times through a 8-HQ column were determined by addition of increasing amounts of these reagents to the sample and were lower than our detection limit. The accuracy of the method was assessed by analysing the DFe standards collected during the Sampling and Analysis of Fe (SAFE) cruise (Johnson et al., 2007). The two samples analysed (S1 and D2) gave values of 0.117 ± 0.009 ($n = 3$) and 0.82 ± 0.06 ($n = 3$) nmol L^{-1} , respectively, for the surface and the deep waters standards, which are in agreement with the certified values of 0.099 ± 0.020 and $0.91 \pm 0.07 \text{ nmol L}^{-1}$, respectively (Johnson et al., 2006).

Samples for total particulate iron (TPFe) were collected by over-pressurising the 12 L GoFlo bottles and filtering as much of the whole sample as possible through a $0.2 \mu\text{m}$ pore size and 47 mm diameter polycarbonate membrane. Filters were acid extracted in Teflon-PFA vials (Savillex, Minnetonka USA) in 1 mL of concentrated ultrapure nitric acid (HNO_3 , Seastar Baseline) and heated for 4 h on a Teflon-coated hot plate at 120°C , before analysis using Magnetic Sector Inductively Coupled Plasma-Mass Spectrometry (Finnigan ELEMENT, Bremen, Germany) (following adaptation of the methods reported in Cullen and Sherrell 1999; Townsend et al., 2000, and Lannuzel et al., submitted, 2009). Given that a mixture of strong acids (HF , HCl , HNO_3) was not used, we assume that the TPFe fraction measured here therefore represents the most leachable particulate Fe pool, probably mostly comprised of material of biogenic origin (Lannuzel et al., submitted, 2009).

3 Results

Total Dissolvable Fe concentrations are reported in Table 1. TDFe vertical profiles for the 9 stations are presented in Fig. 2. At station C1, close to Heard Island, very

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



high values were observed throughout the water column (concentrations ranged from 329 ± 10 to 770 ± 8 nmol L⁻¹). This atypical station is discussed separately from the others.

TDFe concentrations at stations located above the plateau (A3, B1, B3, B5, B7 and C5) were higher than offshore stations (B11 and C11). Above the plateau, concentrations varied from 0.90 ± 0.01 (B7, 80 m) to 65.6 ± 1.00 nmol L⁻¹ (C5, 450 m) whereas outside the plateau values ranged from 0.34 ± 0.05 (B11, 40 m) to 2.23 ± 0.07 nmol L⁻¹ (C11, 600 m).

Stations outside the plateau exhibited concentrations typical of the open Southern Ocean. Our values are in the same range as those measured by Sarthou et al. (1997) in the Indian sector and Sedwick et al. (1997) in the Australian sector (see Table 2).

Higher TDFe concentrations were measured at stations located above the plateau, especially at stations located in the southeastern part, East of Heard Island: 66 and 41 nmol L⁻¹ at 450 and 500 m, respectively, at C5 station (bottom depth 561 m); 38 nmol L⁻¹ at 400 m at B3 station (bottom depth 450 m), see Table 1 and Fig. 2. In the literature, such high values ($38\text{--}66$ nmol L⁻¹) were not often reported even in Fe enriched waters (see Table 2). Concentrations reaching 6.2 nmol L⁻¹ suggested lateral advection of iron-rich shelf water (Sarthou et al., 1997). Values up to 12.5 nmol L⁻¹ were measured south of Tasmania, supporting a sedimentary source (Sedwick et al., 2008). Concentrations higher than 10 nmol L⁻¹ and 25 nmol L⁻¹ were reported by Chase et al. (2005) and Johnson et al. (2001), respectively, in the upwelling system of Monterey Bay, California. The higher values determined by Löscher et al. (1997) in the Atlantic sector of the Southern Ocean are explained by the upwelling of iron rich Antarctic waters.

Apparent particulate iron (Fe_{app}) concentrations, calculated by subtracting DFe values measured during the cruise (Blain et al., 2008) from TDFe, are reported in Table 1. Fe_{app} is defined as the iron adsorbed on lithogenic or biogenic particles, the biogenic iron and the labile lithogenic particulate iron (de Baar et al., 1999) plus the refractory colloids included in the dissolved fraction and that were not measured on board after

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

a few days of acid storage. Fe_{app} represents between 82 and 99% of the TDFe (mean value $95\pm 5\%$) above the plateau and between 83 and 93% (mean value $86\pm 3\%$) outside the plateau with no difference between surface and deep water (see Table 1). This suggests that particles or refractory colloids largely dominated the iron pool. Such high percentages were previously observed in the open ocean and in enriched waters but with a larger variability. Löscher et al. (1997), Sedwick et al. (1997) and Croot et al. (2004) observed percentages of Fe_{app} varying between 0 and more than 80% in the Atlantic sector of the Southern Ocean, the Australian sector of the Southern Ocean and the Equatorial sector of the Atlantic Ocean, respectively (Table 2).

4 Discussion

The onboard study of the concentration of DFe above the plateau highlighted the existence of a deep dissolved iron-rich reservoir below the mixed layer (Blain et al., 2007). The main sources of DFe in the surface waters were identified to be diapycnal diffusive mixing and the utilisation of the winter stock (Blain et al., 2008), but a comparison of iron supply versus iron demand of phytoplankton suggested that an additional iron source was present, and that was possibly the dissolution of lithogenic particles (Blain et al. 2007, 2008; Sarthou et al., 2008). Additionally, the study of the rare-earth elements (REE) showed that Heard Island could be a significant source of lithogenic material in the water column above the plateau (Zhang et al., 2008). This source is also evidenced by elevated ^{228}Ra activities above the plateau (van Beek et al., 2008). The lateral advection of water masses that have been in contact with the continental shelf of Heard Island should thus be considered as a possible source of particulate and dissolved iron above the plateau. In the following section, we focus on the Fe_{app} and the DFe fractions. The different sources of particulate and dissolved iron above the plateau are investigated and a budget of iron is presented. The question of the bioavailability of these particles is also discussed.

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.1 Budget of iron above the Kerguelen plateau

4.1.1 Description of the model

A two box-model including pools and fluxes of dissolved and particulate iron was used to determine an iron budget above the Kerguelen plateau. The model is illustrated in Fig. 3a. The area covered by these boxes is the surface plateau (i.e. 45 000 km²). The surface box represents the upper 150 m of the water column. This depth stratum was chosen in accordance with the study of Blain et al. (2008) who observed a higher vertical gradient of DFe below 150 m for the stations located above the plateau. The deep box represents the depth stratum 150 m – bottom depth, with increasing depth from the incoming (325 m depth) to the exit (500 m depth) of the box in accordance with the topography of the plateau (Park et al., 2008a). All the sources and sinks of dissolved and particulate iron in the two boxes are listed below with the corresponding inflowing and outflowing fluxes. The sedimentary source which could play a significant role is taken into account for the dissolved pool but data are lacking for the particulate flux calculation. Finally, the isopycnal mixing is neglected. Indeed, Maraldi et al. (2009) studied the influence of the short term (~ less than a week) lateral mixing on the phytoplankton bloom over the Kerguelen Plateau. They concluded that the spatial pattern of the bloom was delimited by a barrier of high lateral mixing due essentially to tides, with minor contributions from Ekman transport, geostrophy or barotropic atmospheric forced currents. Mongin et al. (2008) also showed that the effect of lateral diffusion has little effect on the overall carbon and iron budgets. We have therefore decided to make the same assumptions for our iron budget model.

The four following equations are used to describe the variation with time of dissolved and particulate iron in surface and deep waters.

$$\frac{d[\text{DFe}]_1}{dt} \times V1 = Ad \times S + (Fw_1 \times [\text{DFe}]_c)_1 + DMd + WSd - Fw_1 \times [\text{DFe}]_{\text{plateau}1} - E1 \quad (1)$$

BGD

6, 6803–6837, 2009

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$$\frac{d[\text{DFe}]_2}{dt} \times V2 = (Fw_1 \times [\text{DFe}]_C)_2 + (Fw_2 \times [\text{DFe}]_C)_2 + Sed_d - DMd - WSd - Fw_2 \times [\text{DFe}]_{\text{plateau2}} - E2 \quad (2)$$

$$\frac{d[\text{FE}_{\text{app}}]_1}{dt} \times V1 = Ap \times S + (Fw_1 \times [\text{FE}_{\text{app}}]_C)_1 + DMp + E1 - Fw_1 \times [\text{FE}_{\text{app}}]_{\text{plateau1}} - F1 \quad (3)$$

$$\frac{d[\text{FE}_{\text{app}}]_2}{dt} \times V2 = (Fw_1 \times [\text{FE}_{\text{app}}]_C)_2 + (Fw_2 \times [\text{FE}_{\text{app}}]_C)_2 + E2 + F1 - DMp - Fw_2 \times [\text{FE}_{\text{app}}]_{\text{plateau2}} - F2 \quad (4)$$

5 Numbers 1 and 2 correspond to the surface and the deep layer, respectively. “V” and “S” represent the box volume and the box surface. Letters “d” and “p” refer to the dissolved and the particulate fluxes. “A” represents the atmospheric inputs. “Fw” represents the water flux. “Plateau” represents the mean concentration above the plateau and “C” represents the mean concentration along transect C. “DM” represents the inflowing flux in the surface layer coming from the diapycnal mixing, “WS” refers to the winter stock. “E” corresponds to the net fluxes exchanged between dissolved and particulate fraction. “F” refers to the particle fluxes at the bottom of each box.

10 To estimate lateral inflowing and outflowing fluxes of iron in our model, we used the mean current velocity estimated by Park et al. (2008b) above the plateau (4.0±0.5 cm s⁻¹). The corresponding hydrological fluxes (Fw) are calculated by multiplying this current by the section of the box (350 km width and 150 m depth for the upper box, 350 km width and 175 m and 350 m depth for the entrance and the exit of the deep box, respectively). These water fluxes are multiplied by the mean dissolved or particulate iron concentrations along the transect C or above the plateau, in the upper or deep boxes. These mean concentrations are reported in Table 3. We assumed that the water flux transporting dissolved and apparent particulate iron from the transit C in the surface box is uniform over the 500 m water column leading to the two different fluxes (Fw₁*[DFe]_C)₁ and (Fw₁*[DFe]_C)₂ for the DFe, and (Fw₁*[FE_{app}]_C)₁ and

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$(Fw_1^*[Fe_{app}]_C)_2$ for the Fe_{app} fraction (Fig. 3a). Figure 3b displays the results of the calculated stocks and fluxes.

4.1.2 Budget calculation

Atmospheric inputs

5 One of the main external sources of iron to the surface waters of the open ocean is aeolian dust deposition (Jickells et al., 2005). During the cruise, aerosols were collected to estimate atmospheric fluxes to the Southern Ocean, and in particular iron fluxes over the study area (Wagener et al., 2008). A total iron flux (dry + wet) of $23 \pm 9 \text{ nmol m}^{-2} \text{ d}^{-1}$ was estimated. Assuming a solubility ranging from 1 to 10% (Bonnet and Guieu, 2004; Baker and Jickells, 2006), the atmospheric dissolved iron flux
10 over the Kerguelen plateau can be estimated to be $0.2\text{--}3.2 \text{ nmol m}^{-2} \text{ d}^{-1}$ (Wagener et al., 2008). A mean value of $1.7 \pm 2.1 \text{ nmol m}^{-2} \text{ d}^{-1}$ is considered hereafter. These fluxes are low compared to those calculated during the Crozex and Fe Cycle cruises, which were $100 \text{ nmol m}^{-2} \text{ d}^{-1}$ (Planquette et al., 2007) and $7.58\text{--}75.8 \text{ nmol m}^{-2} \text{ d}^{-1}$, respectively, assuming a 1–10% dissolution of the total flux given by Boyd et al. (2005).
15 The particulate iron flux (90–99% of the total flux) during KEOPS was estimated to be $12.6\text{--}31.7 \text{ nmol m}^{-2} \text{ d}^{-1}$ with a mean value of $22.2 \pm 13.5 \text{ nmol m}^{-2} \text{ d}^{-1}$. Over an area of $45\,000 \text{ km}^2$, the atmospheric fluxes $Ad \times S$ and $Ap \times S$ equal $76.5 \pm 94.5 \text{ mol d}^{-1}$ and $999 \pm 607 \text{ mol d}^{-1}$ for the dissolved and particulate iron, respectively. Given that the
20 other particulate fluxes calculated in our budget represent apparent particulate fluxes, this atmospheric flux may be overestimation.

Diapycnal diffusive fluxes

The enrichment of DFe and Fe_{app} for the stations located above the plateau (see Fig. 2) clearly indicates an input of iron from the bottom of the water column. Park et al. (2008b) estimate a vertical eddy diffusivity (Kz) of $3.8 \pm 2.4 \cdot 10^{-4} \text{ m}^2 \text{ s}^{-1}$ in the seasonal pycnocline ($80 \text{ m} < z < 180 \text{ m}$) at A3 using a Thorpe scale analysis. By multiplying
25

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

this coefficient by the vertical gradient of Fe_{app} calculated at this station (8.6 nmol m^{-4} between 80 and 200 m), a diapycnal mixing of $283 \pm 178 \text{ nmol m}^{-2} \text{ d}^{-1}$ can be estimated. Above the plateau, K_z was only estimated at A3 station. Assuming that the diapycnal mixing is homogeneous over all the plateau area, we calculate a flux of particles (DMp) of $12735 \pm 8010 \text{ mol d}^{-1}$. The vertical supply calculated for DFe at A3 is $31 \text{ nmol m}^{-2} \text{ d}^{-1}$ (Blain et al., 2008). With the same assumption that this mixing is homogeneous, we calculate a dissolved flux (DMd) of 1.395 mol d^{-1} .

Winter Mixing

The second mechanism of fertilization mentioned by Blain et al. (2007, 2008) for dissolved iron is the utilisation of the winter stock above the plateau. During winter, the strong winds allow a greater mixing of the surface waters. The thickness of the mixed layer can reach 200 m. As deep water contains higher concentration of iron than surface water (Blain et al., 2008), at the beginning of the spring, surface water is enriched in iron. This stock is called winter stock. To calculate the stock of iron potentially available for phytoplankton growth, the thermal structure of the water column has to be known. The temperature minimum is indicative of the winter concentration (C_{winter}). The mean concentration of DFe measured within the mixed layer is called the summer concentration (C_{summer}). The winter stock utilization ($\text{nmol m}^{-2} \text{ d}^{-1}$) is calculated using the following equation $(C_{winter} - C_{summer}) * \text{MLD} / 90$ where MLD represents the mean mixed layer depth (70 m at A3) and 90 days is the duration of the bloom. Winter stock utilization was $52 \text{ nmol m}^{-2} \text{ d}^{-1}$ for the DFe (Blain et al. 2008). Assuming that this flux is homogeneous over all the surface of the plateau, we calculate a winter mixing flux (WSd) of 2340 mol d^{-1} for the dissolved iron.

Lateral advection

Lateral advection is a potential source of dissolved and particulate iron above the plateau. Apart from some sporadic mesoscale intrusions (Zhang et al., 2008), the circulation above the KEOPS study area rules out any transport of waters from the

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Kerguelen Island to the plateau (Park et al., 2008a), whereas a lateral advection from the south can be considered. General mean circulation above the plateau is described in Fig. 1. Waters that have interacted with the continental shelf of Heard Island can be transported onto the plateau. This source is also recognised for other tracers like ^{228}Ra (van Beek et al., 2008), lithogenic Ba (Jacquet et al., 2008) and rare-earth elements (Zhang et al., 2008).

Using the mean current velocity estimated by Park et al. (2008b) above the plateau in the 500 m water column, the hydrological fluxes (F_w) are calculated by multiplying this current by the section of the box, leading to a surface water flux of $1.81 \times 10^{14} \text{ L d}^{-1}$. For the deep box, values of $2.12 \times 10^{14} \text{ L d}^{-1}$ for the inflowing flux and $4.23 \times 10^{14} \text{ L d}^{-1}$ for the outflowing flux are calculated. In the surface box, multiplying the water flux by the DFe concentrations from the transect C , $(F_{w_1} * [\text{DFe}]_C)_1$ and $(F_{w_1} * [\text{DFe}]_C)_2$ give values of $23.9 \pm 1.6 \times 10^3 \text{ mol d}^{-1}$ and $55.9 \pm 3.8 \times 10^3 \text{ mol d}^{-1}$, respectively. Concerning the Fe_{app} fraction, $(F_{w_1} * [\text{Feapp}]_C)_1$ and $(F_{w_1} * [\text{Feapp}]_C)_2$ give values of $10\,138 \pm 2\,122 \times 10^3 \text{ mol d}^{-1}$ and $23\,656 \pm 4\,966 \times 10^3 \text{ mol d}^{-1}$, respectively. In the deep box, $(F_{w_2} * [\text{DFe}]_C)_2$ and $(F_{w_2} * [\text{Feapp}]_C)_2$ give flux values of 85.9 ± 29.6 and $7\,142 \pm 4\,905 \times 10^3 \text{ mol d}^{-1}$, respectively (Fig. 3b).

Sedimentary inputs

Over the plateau, inputs of DFe in the water column from the sediment were measured during the cruise and comprised of a flux of $136 \mu\text{mol m}^{-2} \text{ d}^{-1}$ (Blain et al., 2008), higher than previous fluxes calculated along the Californian coast (Elrod et al., 2004). This equates to a flux Sed_d of $6\,120 \times 10^3 \text{ mol d}^{-1}$, assuming it is homogeneous over the plateau.

4.1.3 Dissolved and particulate budgets

Results are reported in Fig. 3b. Atmospheric deposition is a negligible source of Fe_{app} above the plateau, as already observed for DFe (Blain et al., 2008). Lateral

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

advection is the predominant source of iron above the plateau, for Fe_{app} but also for DFe (although the uncertainty in the calculation of this flux is important). The corresponding fluxes are equal to $9\,697 \pm 2\,311 \times 10^3 \text{ mol d}^{-1}$ ($= 10\,138 - 441 \times 10^3 \text{ mol d}^{-1}$) and $8.3 \pm 6.7 \times 10^3 \text{ mol d}^{-1}$ ($= 23.9 - 15.6 \times 10^3 \text{ mol d}^{-1}$) for Fe_{app} and DFe, respectively.

They represent 99% and 69% of the total net fluxes of Fe_{app} and DFe that enter the surface box, respectively.

The biogenic pool of Fe of $80 \pm 9 \text{ p mol L}^{-1}$ calculated by Sarthou et al. (2008) represents $540 \times 10^3 \text{ mol}$ in our surface box (150 m depth, area of $45\,000 \text{ km}^2$). It is only 3.3% of the apparent particulate iron stock calculated in our surface box ($16\,402 \times 10^3 \text{ mol}$, see Fig. 3b), which suggests that the biogenic fraction is not predominant.

At steady-state, left members of the Eqs. (1)–(4) are equal to 0. Values of E1, E2, F1 and F2 can be calculated by resolving these 4 equations and are reported in Fig. 3b. Values of $12.1 \pm 6.8 \times 10^3$ and $6\,160 \pm 65 \times 10^3 \text{ mol d}^{-1}$ were calculated, respectively for E1 and E2.

The positive values of E1 and E2 suggest a predominant transport of iron from the dissolved to the apparent particulate fraction. Processes involved in such a transport are biological uptake, colloidal aggregation, adsorption on phytoplankton cells and scavenging by non-living particles (through adsorption, precipitation or aggregation), whereas processes responsible for the transfer of apparent particulate iron to the dissolved iron pool are due to the dissolution and regeneration of particulate or colloidal iron (Sarthou and Jeandel, 2001).

Using the radiotracer ^{55}Fe , the phytoplankton net Fe demand in the bloom (uptake minus regeneration) was equal to $208 \pm 77 \text{ nmol m}^{-2} \text{ d}^{-1}$ (Sarthou et al., 2008). Adsorption on phytoplankton cells and scavenging by non-living particles represented between 20 and 37% of the total biogenic iron (Sarthou et al., 2008), leading to a total biogenic iron flux of $260 - 330 \text{ nmol m}^{-2} \text{ d}^{-1}$. Assuming an area of $45\,000 \text{ km}^2$, this gives a flux varying between 11.7×10^3 and $14.9 \times 10^3 \text{ mol d}^{-1}$. These values are close to E1 ($12.1 \pm 6.8 \times 10^3 \text{ mol d}^{-1}$), suggesting that the dissolution of Fe_{app} may not be significant.

Taking into account the maximum value of the flux calculated from the ^{55}Fe

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

experiment ($14.9 \times 10^3 \text{ mol d}^{-1}$) and the minimum value of the E1 flux (i.e. 12.1–6.8 = $5.3 \times 10^3 \text{ mol d}^{-1}$), a maximum dissolution of $9.5 \times 10^3 \text{ mol d}^{-1}$ can however be estimated. As the biogenic pool represents 3.3% of the total pool (see above), the maximum dissolution of lithogenic material is estimated to $9.2 \times 10^3 \text{ mol d}^{-1}$. It represents 0.09% of the total apparent particulate flux that enters in the upper box ($10\,138 \times 10^3 \text{ mol d}^{-1}$). If we assume that all the DFe at C1 comes from the dissolution of Fe_{app} , a dissolution of 0.2% is calculated. These values are in the same range as those calculated by Bonnet and Guieu (2004) and listed by Jickells et al. (2001) for the dissolution of lithogenic Fe from atmospheric dusts.

The exchange flux between the dissolved and apparent particulate fraction in deep waters (E2) also shows a positive value, two orders of magnitude higher than in surface waters. This suggests that scavenging and/or aggregation are predominant in the deep waters. This could be due to the high transport of particulate material in the deep box and to the inputs from the sediments.

Values of $9\,723 \pm 2.326 \times 10^3$ and $43\,868 \pm 15\,710 \times 10^3 \text{ mol d}^{-1}$ were calculated for the particulate sinking fluxes F1 and F2, respectively. This budget suggests that most of the particles that flow above the plateau are removed from the water column by sedimentation.

The calculation of F1 allows us to estimate the residence time (RT) of apparent particulate iron (RT = PFe stock/ downward PFe flux, Boyd et al., 2005). It is equal to 1.7 days in surface waters. This value is lower than that of 100 days estimated during the Fe Cycle experiment (Boyd et al., 2005). Residence times varying between 6 (range 2–12 days) and 62 (range 21–186) days for total iron (dissolved + particulate) were calculated in the Equatorial sector of the Atlantic Ocean (Croot et al., 2004). The shortest residence times were associated with high particulate Fe flux. In our budget, the high supply of Fe_{app} from the weathering of Heard Island, which is rapidly removed by sinking, may explain the very low residence times calculated.

Considering a mean dissolved iron stock in the upper 150 m of the water column above the plateau of $581 \times 10^3 \text{ mol}$, and a supply of DFe of $12.1 \times \text{mol d}^{-1}$

($= (A_d * S + DMd + WS_d + HS_d)$), with HS_d equal to the horizontal supply of iron above the plateau ($= (23.9 - 15.6) \times 10^3 \text{ mol d}^{-1}$), we estimated a DFe residence time equal to 48 days. Such a value is consistent with residence times of ~ 10 days – 1 year observed in the North and Equatorial Atlantic Ocean (Jickells, 1999; de Baar and de Jong, 2001; Sarthou et al., 2003; Sarthou et al., 2007).

4.2 Importance of the colloidal fraction to the bioavailability of Fe

Vertical profiles of total and apparent particulate iron at stations A3 and C11 are plotted in Fig. 4. For both stations, higher values are observed in the Fe_{app} pool compared to the TPF_e pool. Indeed, Fe_{app} concentrations are 1.3 to 3.1 times higher than TPF_e.

Given that some very refractory particulate Fe remains unreactive to the TDFe handling and analysis (Powell et al., 1995), Fe_{app} fraction should exhibit smaller concentrations than TPF_e. Several reasons might explain this difference. First, for the 2 stations TPF_e was sampled 10 days before TDFe and DFe. Given that the residence time of iron particles is 1.7 days, changes in the transport of particles could have occurred during this period. Second, an incomplete digestion of particulate material, leading to an underestimation of the TPF_e values might have occurred, since the digestion protocol did not include a strong attack with HF acid. Finally, another possibility would be that a substantial fraction of colloidal iron was not analysed on board in the DFe pool, but was reactive to the more acidic attack and to the time storage of the TDFe samples.

A significant fraction of dissolved iron is actually thought to be in the colloidal pool (Kuma et al., 1998; Wu et al., 2001; Cullen et al., 2006). Wu et al. (2001) and Cullen et al. (2006) observed that 80–90% of the dissolved iron is present under colloidal fraction in the surface waters and 70% in the deep waters in the Atlantic Ocean.

A few duplicates of DFe samples were stored at room temperature and were analysed at the same time as the TDFe samples at the laboratory (i.e. 18 months later). DFe concentrations increased by 37–90% during these 18 months of storage. This result implies that a substantial fraction of DFe is colloidal and is only released during long acid storage. This observation confirms the work of Zhang et al. (2008) who

BGD

6, 6803–6837, 2009

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



suggest that a large pool of DFe is probably colloidal. However Fe_{app} still dominates the total dissolvable pool after 18 months of storage. Indeed, Fe_{app} represents between 54 and 99% of the TDFe (mean value $85 \pm 11\%$, $n = 27$) when considering the dissolved iron samples analysed at the laboratory.

5 Only two duplicates of DFe were stored at A3 and it was possible to re-calculate the concentrations of Fe_{app} by using the values of DFe measured at the laboratory and to compare them with the TPF_e concentrations. The new estimations of these Fe_{app} concentrations are presented on Fig. 4 and seem to explain the difference between the two pools. This result suggests that a significant portion of refractory colloids is present
10 in the DFe fraction and that only a long acid storage (18 months) allows measuring this fraction.

Assuming that the colloids came from the dissolution of particulate lithogenic Fe at C1 and that the mean current velocity above the plateau is $4.0 \pm 0.5 \text{ cm s}^{-1}$ (Park et al., 2008b), it takes nearly three months for the colloids to reach A3. Chen and
15 Wang (2001) showed that freshly precipitated colloids were available to phytoplankton but aging processes (15 days) markedly reduced their availability.

Moreover, the E1 flux calculated in our budget in the surface box is consistent with the uptake by phytoplankton estimated by Sarthou et al. (2008). These calculations seem to indicate that the refractory colloidal iron present in an apparent particulate
20 fraction is not taken up by phytoplankton.

These results give some information on the measurements of the physical speciation of iron. First, the most labile fraction of DFe measured after few hours of acidification could be the fraction that is bioavailable to the phytoplankton. Second, more than a few days of acidification are needed to measure the total dissolved iron fraction, especially in a region with high input of refractory iron. Our study indicates that 18
25 months of acid storage allow the measurement of the refractory colloids and determine the total particulate Fe from TDFe and DFe analyses. However, it should be noted that only digestion including HF acid (which was not used here) ensures the recovery of the most refractory and crystalline particles determined in the TPF_e fraction. These

An iron budget during the natural iron fertilisation experiment KEOPSF. Chever et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

observations are in agreement with the study of Bowie et al. (2004) who considered it useful to extend the storage of acidified samples to determine the total dissolved iron pool.

4.3 Carbon sequestration efficiency

5 Carbon sequestration efficiency is based on the ratio of the excess of carbon exported to the dissolved iron supplied (de Baar et al., 2005). An iron supply of $5 \times 10^{-3} \text{ nmol m}^{-2}$, corresponding to the vertical supply and the winter stock utilization, has already been calculated by Blain et al. (2007). Taking into account the lateral advection which supplies $8.3 \times \text{mol Fe d}^{-1}$ ($= (23.9 - 15.6) \times 10^3 \text{ mol d}^{-1}$) over an area of
10 $45\,000 \text{ km}^2$ during 90 days, the contribution of the lateral supply can be estimated to be equal to $16.6 \times 10^{-3} \text{ nmol m}^{-2}$ leading to a new estimate of dissolved iron supply of $21.6 \times 10^{-3} \text{ nmol m}^{-2}$. This value is probably overestimated because of a possible impact of this lateral advection on the stations located outside the bloom (C11 station for example), which would lead to a smaller excess of iron. Using the excess of particulate organic carbon (POC) calculated by Blain et al. (2007) of $3\,317 \text{ nmol m}^{-2}$, the
15 new estimated sequestration efficiency is equal to $154\,000 \text{ mol C (mol Fe)}^{-1}$. With our assumptions, consideration of the lateral advection reduces the first estimate of the sequestration efficiency by 4-fold. However, this value is still approximately 18-fold higher than the value calculated during the other study of a natural iron fertilisation experiment
20 CROZEX ($8\,640 \text{ mol C (mol Fe)}^{-1}$, Pollard et al., 2009) and is also higher than the two artificial iron fertilisation experiments SOFeX ($3\,300 \text{ mol C (mol Fe)}^{-1}$, Buesseler et al., 2004) and SERIES ($\sim 500 \text{ mol C (mol Fe)}^{-1}$, Boyd et al., 2004). Blain et al. (2007) explained the higher carbon sequestration efficiency calculated during KEOPS than during SOFeX and SERIES by two reasons. First, it is suspected that the end of the bloom was reached during KEOPS but not during SOFeX, leading to a higher excess of carbon during KEOPS. Secondly, during mesoscale enrichment experiments a large amount of the DFe added to seawater is rapidly loss contrary to the natural iron enrichment where the input of iron is slow and continuous (Blain et al. 2007).
25

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The comparison of the carbon sequestration efficiency between the CROZEX and the KEOPS experiments is difficult because of the different methods used during the two studies to calculate iron supply and carbon export. However, Pollard et al. (2009) observed that during KEOPS, the supply of iron was 8-fold lower and the carbon exported was 10-fold higher than during CROZEX. They suggested that either a vertical mixing or a more intense lateral advection of lithogenic material could have occurred but before the late-summer observation period of KEOPS. Our study allowed us to better constrain the lateral fluxes of Fe over the plateau. Considering these fluxes, the sequestration efficiency previously calculated by Blain et al. (2007) decreases, but still remains largely higher than during CROZEX.

5 Conclusions

Our results suggest that TDFe in our study area predominantly consists of particles. This study also highlights the importance of the refractory colloidal fraction in the dissolved pool and emphasises the importance of extended storage to better constrain the physical speciation of Fe. The most labile fraction of DFe measured after a few days of acidification could be the fraction that is bioavailable to the phytoplankton. The total particulate Fe may be determined from TDFe and DFe analyses with sufficient acid storage (18 months). The study of the TDFe fraction above the Kerguelen Plateau gives information on iron sources, the physical speciation and the bioavailability of iron.

Our Fe budget also shows that the predominant source of Fe_{app} and DFe above the Kerguelen Plateau is the lateral advection of waters that have been in contact with the continental shelf of Heard Island. It represents 99% and 69% of the total apparent particulate and dissolved iron supply, respectively.

By taking into account the lateral advection of dissolved iron, we calculate a revised seasonal carbon sequestration efficiency of $154\,000 \text{ mol C (mol Fe)}^{-1}$, 4 times lower than the one previously calculated by Blain et al. (2007), but still 18-fold higher than the one calculated during the natural iron fertilisation experiment CROZEX (Pollard et al.,

BGD

6, 6803–6837, 2009

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2009). This result is also significantly higher than the carbon sequestration efficiencies obtained during artificial iron fertilisation experiments. Two reasons may explain these differences: first, if Fe is supplied in a continuous way or not, second if the bloom reached its maximum or not.

5 *Acknowledgements.* We are grateful to the project coordinator S. Blain, to B. Quéguiner, the chief scientist, and to the captain and crew of the R. V. M. Dufresne for their support on board. We also acknowledge T. Wagener for his help with sampling clean seawater. This work was supported by the Institut National des Sciences de l'Univers (INSU) and the French Polar Institute (Institut Polaire Emile Victor, IPEV). This research was partly supported by the Australian Government Cooperative Research Centres Program through the Antarctic Climate and Ecosystems CRC (ACE CRC) and the French-Australian Science and Technology (FAST) programme (contract number FR040170). We thank ACE CRC internal reviewers B. Griffiths and D. Lannuzel for their comments.



The publication of this article is financed by CNRS-INSU.

References

- Baker, A. R., Jickells, T. D., Witt, M., and Linge K. L.: Trends in the solubility of iron, aluminium, manganese and phosphorus in aerosol collected over the Atlantic Ocean, *Mar. Chem.*, 98, 43–58, 2006.
- Blain, S., Quéguiner, B., Armand, L., Belviso, S., Bombled, B., Bopp, L., Bowie, A. R., Brunet, C., Brussaard, C., Carlotti, F., Christaki, U., Corbière, A., Durand, I., Ebersbach, F., Fuda, J. L., Garcia, N., Gerringa, L., Griffiths, B., Guigue, C., Guillerm, C., Jacquet, S., Jeandel, C.,

BGD

6, 6803–6837, 2009

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**An iron budget
during the natural
iron fertilisation
experiment KEOPS**

F. Chever et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Laan, P., Lefèvre, D., Lo Monaco, C., Malits, A., Mosseri, J., Obernesterer, I., Park, Y. H., Picheral, M., Pondaven, P., Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scoarnec, L., Souhaut, M., Thuiller, D., Timmermans, K., Trull, T., Uitz, J., van Beek, P., Veldhuis, M., Vincent, D., Viollier, E., Vong, L., and Wagener, T.: Effect of natural iron fertilization on carbon sequestration in the Southern Ocean, *Nature*, 446, 1070–1074, 2007.

Blain, S., Sarthou, G., and Laan, P.: Distribution of dissolved iron during the natural iron-fertilization experiment KEOPS (Kerguelen Plateau, Southern Ocean), *Deep Sea Res. II*, 55, 5–7, 594–605, 2008.

Bonnet, S. and Guieu C.: Dissolution of atmospheric iron in seawater, *Geophys. Res. Lett.*, 31, 3, L03303, doi: 10.1029/2003GL018423, 2004.

Bowie, A. R., Sedwick, P. N., and Worsfold, P. J.: Analytical intercomparison between flow injection–chemiluminescence and flow injection – spectrophotometry for the determination of picomolar concentrations of iron in seawater, *Limnol. Oceanogr.-Meth.*, 2, 42–54, 2004.

Boyd, P. W., Watson, A. J., Law, C. S., Abraham, E. R., Trull, T., Murdoch, R., Bakker, D. C. E., Bowie, A., Buesseler, K. O., Chang, H., Charette, M., Croot, P., Downing, K., Frew, R., Gall, M., Hadfield, M., Hall, J., Harvey, M., Jameson, G., LaRoche, J., Liddicoat, M., Ling, R., Maldonado, M. T., McKay, R. M., Nodder, S., Pickmere, S., Pridmore, R., Rintoul, S., Safi, K., Sutton, P., Strzepek, R., Tanneberger, K., Turner, S., Waite, A., and Zeldis, J.: A mesoscale phytoplankton bloom in the polar Southern Ocean stimulated by iron fertilization, *Nature*, 407, 695–702, 2000.

Boyd, P. W., Law, C. S., Wong, C. S., Nojiri, Y., Tsuda, A., Ievasseur, M., Takeda, S., Rivkin, R., Harrison, P. J., Strzepek, R., Gower, J., McKay, R. M., Abraham, E., Arychuk, M., Barwell, C. J., Crawford, W., Crawford, D., Hale, M., Harada, K., Johnson, K., Kiyosawa, H., Kudo, I., Marchetti, A., Miller, W., Needoba, J., Nishioka, J., Ogawa, H., Page, J., Robert, M., Saito, H., Sastri, A., Sherry, N., Soutar, T., Sutherland, N., Taira, Y., Whitney, F., Wong, S. K. E., and Yoshimura, T.: The decline and fate of an iron-induced subarctic phytoplankton bloom, *Nature*, 428, 549–553, 2004.

Boyd, P. W., Law, C. S., Hutchins, D. A., Abraham, E. R., Croot, P. L., Ellwood, M., Frew, R. D., Hadfield, M., Hall, J., Handy, S., Hare, C. E., Higgins, J., Hill, P., Hunter, K. A., Leblanc, K., Maldonado, M. T., Mckay, R. M., Mioni, C., Oliver, M., Pickmere, S., Pinkerton, M., Safi, K., Sander, S., Sanudo-Wilhelmy, S. A., Smith, M., Strzepek, R., Tavor-Sanchez, A., and Wilhelm S. W.: FeCycle: Attempting an iron biogeochemical budget from a mesoscale SF6 tracer experiment in unperturbed low iron waters, *Global Biogeochem. Cy.*, 19, GB4S20,

doi:10.1029/2005GB002494, 2005.

Bruland, K. W. and Rue, E. L.: Analytical Methods for the Determination of Concentrations and Speciation of Iron, Biogeochemistry of Fe in Seawater, edited by: Turner, D. R. and Hunter, K. A., SCOR/IUPAC series, J. Wiley, UK, Chapter 6, 255–290, 2001.

5 Buesseler, K. O., Andrews, J. E., Pike, S. M., and Charette, M. A.: The effects of iron fertilization on carbon sequestration in the Southern Ocean, *Science*, 304, 5669, 414–417, 2004.

Chase, Z., Johnson, K. S., Elrod, V. A., Plant, J. N., Fitzwater, S. E., Pickella, L., and Sakamoto, C. M.: Manganese and iron distributions off central California influenced by upwelling and shelf width, *Mar. Chem.*, 95, 235–254, 2005.

10 Chen, M. and Wang, W.-X.: Bioavailability of natural colloid-bound iron to marine plankton: Influences of colloidal size and aging, *Limnol. Oceanogr.*, 46(8), 1956–1967, 2001.

Coale, K. H., Johnson, K. S., Chavez, F. P., Buesseler, K. O., Barber, R. T., Brzezinski, M. A., Cochlan, W. P., Millero, F. J., Falkowski, P. G., Bauer, J. E., Wanninkhof, R. H., Kudela, R. M., Altabet, M. A., Hales, B. E., Takahashi, T., Landry, M. R., Bidigare, R. R., Wang, X. J.,
15 Chase, Z., Strutton, P. G., Friederich, G. E., Gorbunov, M. Y., Lance, V. P., Hiltig, A. K., Hiscock, M. R., Demarest, M., Hiscock, W. T., Sullivan, K. F., Tanner, S. J., Gordon, R. M., Hunter, C. N., Elrod, V. A., Fitzwater, S. E., Jones, J. L., Tozzi, S., Koblizek, M., Roberts, A. E., Herndon, J., Brewster, J., Ladizinski, N., Smith, G., Cooper, D., Timothy, D., Brown, S. L., Selph, K. E., Sheridan, C. C., Twining, B. S., and Johnson, Z. I.: Southern ocean
20 iron enrichment experiment: Carbon cycling in high- and low- Si waters, *Science*, 304, 5669, 408–414, 2004.

Croot, P. L., Streu, P., and Baker, A. R.: Short residence time for iron in surface seawater impacted by atmospheric dry deposition from Saharan dust events, *Geophys. Res. Lett.*, 31, L23S08, doi: 10.1029/2004GL020153, 2004.

25 Cullen, J. T. and Sherrell, R. M.: Techniques for determination of trace metals in small samples of size-fractionated particulate matter: phytoplankton metals off central California, *Mar. Chem.*, 67, 3–4, 233–247, 1999.

Cullen, J. T., Bergquist, B. A., and Moffett, J. W.: Thermodynamic characterization of the partitioning of iron between soluble and colloidal species in the Atlantic Ocean, *Mar. Chem.*, 98, 295–303, 2006.

30 de Baar, H. J. W., de Jong, J. T. M., Nolting, R. F., Timmermans, K. R., van Leeuwe, M. A., Bathmann, U., van der Loeff, M. R., and Sildam, J.: Low dissolved Fe and the absence of diatom blooms in remote Pacific waters of the Southern Ocean, *Mar. Chem.*, 66, 1–34, 1999.

BGD

6, 6803–6837, 2009

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**An iron budget
during the natural
iron fertilisation
experiment KEOPS**

F. Chever et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

de Baar, H. J. W. and de Jong, J. T. M.: Distributions, Sources and Sinks of Iron in Seawater, Biogeochemistry of Fe in Seawater, edited by: Turner, D. R. and Hunter, K. A., J Wiley, SCOR-IUPAC series, Baltimore, USA, Chapter 5, 123–253, 2001.

de Baar, H. J. W., Boyd, P. W., Coale, K. H., and Tsuda, A.: Synthesis of eight in situ iron fertilizations in high nutrient low chlorophyll waters confirms the control by wind mixed layer depth of phytoplankton blooms, *J. Geophys. Res.*, 110, 700–724, 2005.

Elrod, V. A., Berelson, W. M., Coale, K. H., and Johnson, K. S.: The flux of iron from continental shelf sediments: A missing source for global budgets, *Geophys. Res. Lett.*, 31, L12307, doi:10.1029/2004GL020216, 2004.

Fitzwater, S. E., Johnson, K. S., Elrod, V. A., Ryana, J. P., Coletta, L. J., Tanner, S. J., Gordon, R. M., and Chavez, F. P.: Iron, nutrient and phytoplankton biomass relationships in upwelled waters of the California coastal system, *Cont. Shelf Res.*, 23, 1523–1544, 2003.

Frew, R. D., Hutchins, D. A., Nodder, S., Sanudo-Wilhelmy, S., Tovar-Sanchez, A., Leblanc, K., Hare, C. E., and Boyd, P. W.: Particulate iron dynamics during FeCycle in sub-antarctic waters southeast of New Zealand, *Global. Biogeochem. Cy.*, 20, GB1S93, doi:10.1029/2005GB002558, 2006.

Jacquet, S. H. M., Dehairs, F., Savoye, N., Obernosterer, I., Christaki, U., Monnin, C., and Cardinal, C.: Mesopelagic organic carbon remineralization in the Kerguelen Plateau region tracked by biogenic particulate Ba, *Deep Sea Res. II*, 55, 5–7, 868–879, 2008.

Jickells, T.: The inputs of dust derived elements to the Sargasso Sea: a synthesis, *Mar. Chem.*, 68, 5–14, 1999.

Jickells, T. and Spokes, L. J.: Atmospheric iron inputs to the oceans. The biogeochemistry of Iron in seawater, D. R. Turner and K. A. Hunter, J Wiley, Baltimore, USA, SCOR-IUPAC series, Chapter 4, 85–121, 2001.

Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., La Roche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global Iron Connections Between Desert Dust, Ocean Biogeochemistry, and Climate, *Science*, 308, 67–71, 2005.

Johnson, K. S., Chavez, F. P., Elrod, V. A., Fitzwater, S. E., Pinnington, J. T., Buck, K. R., and Walz, P. M.: The annual cycle of iron and the biological response in central California coastal waters, *Geophys. Res. Lett.*, 28(7), 1247–1250, 2001.

Johnson, K. S., Boyle, E., Bruland, K., Coale, K., Measures, C., and Moffett, J.: Sampling and

**An iron budget
during the natural
iron fertilisation
experiment KEOPS**

F. Chever et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Analysis of Fe: The SAFE Iron Intercomparison Cruise, Eos Trans. AGU, 87, 36, Ocean Sci. Meet. Suppl., Abstract OS11N-02, 2006.

Johnson, K. S., Boyle, E., Bruland, K., Coale, K., Measures, C., and Moffett, J.: Developing Standards for Dissolved Iron in Seawater, Eos Trans. AGU, 88, 11, 131–132, 2007.

5 Kuma, K., Katsumoto, A., Nishioka, J., and Matsunaga, K.: Size-fractionated iron concentrations and Fe(III) hydroxide solubilities in various coastal waters, Estuarine, Coastal and Shelf Science, 47, 275–283, 1998.

Kutska, A. B., Carpenter, E. J., and Sanudo-Wilhelmy, S. A.: Iron and marine nitrogen fixation: progress and future directions, Res. Microbiol., 153, 255–262, 2002.

10 Lannuzel, D., Remenyi, T., Lam, P., Townsend, A., Ibsanmi, E., Butler, E., Wagener, T., Schoemann, V., and Bowie, A. R.: Distributions of dissolved and particulate iron in the sub-Antarctic and polar frontal Southern Ocean (Australian sector), Deep-Sea Res. Pt. II, in review, 2009.

Löscher, B. M., de Baar, H. J. W., de Jong, J. T. M., Veth, C., and Dehairs, F.: The distribution of Fe in the Antarctic Circumpolar Current. Ecology and biogeochemistry of the Antarctic Circumpolar Current during austral spring: Southern Ocean JGOFS cruise ANT X/6 of R.V. Polarstern, Deep Sea Res. II, 44, 1–2, 143–187, 1997.

15 Maldonado, M. T., and Price, N. M.: Reduction and transport of organically bound iron by *Thalassiosira oceanica* (Bacillariophyceae), J. Phycol., 37, 2, 298–309, 2001.

Maraldi, C., Mongin, M., Coleman, R., and Testut, L.: The influence of lateral mixing on a phytoplankton bloom: Distribution in the Kerguelen Plateau region, Deep Sea Res. I., 56, 6, 963–973, 2009.

Mongin, M., Molina, E., and Trull, T. W.: Seasonality and scale of the Kerguelen plateau phytoplankton bloom: A remote sensing and modeling analysis of the influence of natural iron fertilization in the Southern Ocean, Deep Sea Res. II, 55, 880–892, 2008.

25 Morel, F. M. M., Kustka, A. B., and Shaked, Y.: The role of unchelated Fe in the iron nutrition of phytoplankton, Limnology&Oceanography, 53, 1, 400–404, 2008.

Obata, H., Karatani, H., and Nakayama, E.: Automated determination of iron in seawater by chelating resin concentration and chemiluminescence, Anal. Chem., 65, 1524–1528, 1993.

Park, Y.-H., Roquet, F., Durand, I., and Fuda, J. L.: Large-scale circulation over and around the Northern Kerguelen Plateau, Deep Sea Res. II, 55, 5–7, 566–581, 2008a.

30 Park, Y.-H., Fuda, J.-L., Durand, I., and Naveira Garabato, A. C.: Internal tides and vertical mixing over the Kerguelen Plateau, Deep Sea Res. II, 55, 5–7, 582–593, 2008b.

Planquette, H., Statham, P. J., Fones, G. R., Charette, M. A., Moore, C. M., Salter, I., Nedelec,

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

F. H., Taylor, S. L., French, M., Baker, A. R., Mahowald, N., and Jickells, T. D.: Dissolved iron in the vicinity of the Crozet Islands, Southern Ocean, Deep Sea Res. II, 54, 1999–2019, 2007.

Pollard, R. T., Salter, I., Sander, S., Lucas, M. I., Moore, C. M., Mills, R. A., Statham, P. J., Allen, J. T., Baker, A. R., Bakker, D. C. E., Charette, M. A., Fielding, S., Fones, G. R., French, M., Hickman, A. E., Holland, R. J., Hughes, J. A., Jickells, T. D., Lampitt, R. S., Morris, P. J., Nedelec, F. H., Nielsdottir, M., Planquette, H., Popova, E. E., Poulton, A. J., Read, J. F., Seeyave, S., Smith, T., Stinchcombe, M., Taylor, S. L., Thomalla, S., Venables, H. J., Williamson, R., and Zubkov, M.: Southern Ocean deep-water carbon export enhanced by natural iron fertilization, Nature, 457, 577–58, 2009.

Powell, R. T., King, D. W., and Landing, W. M.: Iron distributions in the surface waters of the south Atlantic, Mar. Chem., 50, 1–4, 13–20, 1995.

Rueter, J. G. and Ades, D. R.: The role of iron nutrition in photosynthesis and nitrogen assimilation in *Scenedesmus quadeicauda* (Chlorophyceae), J. Physicol., 23, 452–457, 1987.

Sarthou, G., Jeandel, C., Brisset, L., Amouroux, D., and Donard, O. F. X.: Fe and H₂O₂ distributions in the upper 250 m of the water column in the Indian sector of the Southern Ocean, Earth Planet Sc. Lett., 147, 83–92, 1997.

Sarthou, G. and Jeandel, C.: Seasonal variations of iron concentrations in the Ligurian Sea and iron budget in the Western Mediterranean Sea, Mar. Chem., 74, 2–3, 115–129, 2001.

Sarthou, G., Baker, A. R., Blain, S., Achterberg, E. P., Boye, M., Bowie, A. R., Croot, P. L., Laan, P., de Baar, H. J. W., Jickells, T. D., and Worsfold, P. J.: Atmospheric iron deposition and sea-surface dissolved iron concentrations in the eastern Atlantic Ocean, Deep Sea Res. I, 50, 10–11, 1339–1352, 2003.

Sarthou, G., Baker, A. R., Kramer, J., Laan, P., Laës, A., Ussher, S., Achterberg, E. P., de Baar, H. J. W., Timmermans, K. R., and Blain, S.: Influence of atmospheric inputs on the iron distribution in the subtropical North-East Atlantic Ocean, Mar. Chem., 104, 186–202, 2007.

Sarthou, G., Vincent, D., Christaki, U., Obernesterer, I., Timmermans, K. R., and Brussaard, C. P. D.: The fate of biogenic iron during a phytoplankton bloom induced by natural fertilization: Impact of copepod grazing, Deep Sea Res. II, 55, 5–7, 734–751, 2008.

Schlitzer, R.: Ocean Data View, available at: <http://odv.awi.de>, 2007.

Sedwick, P. N., Edwards, P. R., Mackey, D. J., Griffiths, F. B., and Parslow, J. S.: Iron and manganese in surface waters of the Australian subantarctic region, Deep-Sea Res. I, 44, 1239–1253, 1997.

Sedwick, P. N., Bowie, A. R., and Trull, T. W.: Dissolved iron in the Australian sector of the Southern ocean (CLIVAR SR3 section): Meridional and seasonal trends, *Deep Sea Res. I*, 55, 911–925, 2008.

Sunda, W. G.: Trace metal interactions with marine phytoplankton, *Biol. Oceanogr.*, 6, 411–442, 1989.

Townsend, A. T., Miller, K. R., McLean, S., and Aldous, S.: The determination of copper, zinc, cadmium and lead in urine by high resolution ICP-MS, *J. Anal. Atom. Spectrom.*, 13, 1213–1219, 1998.

van Beek, P., Bourquin, M., Reyss J.-L., Souhaut, M., Charrette, M. A., and Jeandel, C.: Radium isotopes to investigate the water mass pathways on the Kerguelen Plateau (Southern Ocean), *Deep Sea Res. II*, 55, 5–7, 622–637, 2008.

Wagener, T., Guieu, C., Losno, R., Bonnet, S., and Mahowald, N.: Revisiting atmospheric dust export to the Southern Hemisphere ocean: Biogeochemical implications, *Global Biogeochemical Cycles*, 22, GB2006, doi:10.1029/2007GB002984, 2008.

Wu, J., Boyle, E., Sunda, W., and Wen, L.-S.: Soluble and colloidal iron in the oligotrophic North Atlantic and North Pacific, *Science*, 293, 847–849, 2001.

Zhang, Y., Lacan, F., and Jeandel, C.: Dissolved rare earth elements tracing lithogenic inputs over the Kerguelen Plateau (Southern Ocean), *Deep Sea Res. II*, 55, 5–7, 638–652, 2008.

BGD

6, 6803–6837, 2009

**An iron budget
during the natural
iron fertilisation
experiment KEOPS**

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Total dissolvable iron (TDFe), apparent particulate iron (Fe_{app}) and location of the sampling stations. Fe_{app} is calculated using dissolved iron (DFe) values from Blain et al. (2008). Uncertainties on the TDFe concentrations correspond to standard deviation of triplicate measurements of the sample measured 3 times. Uncertainties on the Fe_{app} correspond to the sum of the TDFe and DFe uncertainties. n.d. represents “not determined” values.

Station	Latitude	Longitude	Depth (m)	TDFe (nmol L ⁻¹)	Fe_{app} (nmol L ⁻¹)	% Fe_{app} /TDFe			
A3	50.65° S	72.07° E	40	1.22 ± 0.01	0.97 ± 0.02	90%			
			80	1.07 ± 0.01	0.82 ± 0.01	87%			
			120	1.73 ± 0.09	n.d.	n.d.			
			150	2.18 ± 0.04	1.96 ± 0.04	95%			
			200	2.16 ± 0.01	1.85 ± 0.01	91%			
			300	3.45 ± 0.03	3.15 ± 0.03	94%			
			350	5.09 ± 0.15	4.75 ± 0.15	95%			
			400	6.48 ± 0.06	n.d.	n.d.			
			450	9.63 ± 0.03	n.d.	n.d.			
B1	51.5° S	73.0° E	40	2.72 ± 0.04	2.64 ± 0.04	97%			
			80	1.75 ± 0.02	1.70 ± 0.02	97%			
			120	1.96 ± 0.06	1.89 ± 0.06	97%			
			150	2.97 ± 0.04	2.90 ± 0.04	98%			
			250	4.35 ± 0.02	4.19 ± 0.03	96%			
			300	5.68 ± 0.06	5.51 ± 0.06	97%			
			350	7.35 ± 0.21	7.15 ± 0.21	97%			
			B3	51.3° S	73.8° E	40	2.22 ± 0.02	2.10 ± 0.02	95%
						80	2.92 ± 0.02	2.81 ± 0.03	96%
120	2.60 ± 0.14	2.52 ± 0.14				97%			
150	3.07 ± 0.02	2.96 ± 0.03				96%			
200	2.83 ± 0.02	2.68 ± 0.02				95%			
300	3.97 ± 0.05	3.72 ± 0.06				94%			
350	13.64 ± 0.44	13.40 ± 0.45				98%			
400	38.10 ± 1.32	37.75 ± 1.32				99%			
B5	51.1° S	74.58° E				43	2.73 ± 0.14	n.d.	n.d.
			80	4.67 ± 0.02	4.62 ± 0.02	99%			
			93	3.79 ± 0.02	n.d.	n.d.			
			120	3.39 ± 0.09	3.34 ± 0.10	98%			
			150	2.39 ± 0.08	2.30 ± 0.08	96%			
			200	3.34 ± 0.05	3.26 ± 0.05	97%			
			300	4.26 ± 0.11	4.09 ± 0.11	96%			
			B7	50.9° S	75.4° E	40	4.30 ± 0.07	4.25 ± 0.08	99%
						80	0.90 ± 0.01	0.76 ± 0.02	85%
150	2.18 ± 0.06	n.d.				n.d.			
200	1.59 ± 0.01	1.33 ± 0.02				84%			
300	2.43 ± 0.02	2.12 ± 0.03				87%			
400	3.90 ± 0.01	3.57 ± 0.03				91%			
450	4.93 ± 0.03	4.63 ± 0.04				94%			
500	3.83 ± 0.02	3.49 ± 0.03				91%			

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Continued.

Station	Latitude	Longitude	Depth (m)	TDFe (nmol L ⁻¹)	Fe _{app} (nmol L ⁻¹)	%Fe _{app} /TDFe
B11	50.5° S	77.0° E	40	0.34 ± 0.05	n.d.	n.d.
			150	0.43 ± 0.02	n.d.	n.d.
			200	0.71 ± 0.01	n.d.	n.d.
			400	0.39 ± 0.01	n.d.	n.d.
			600	0.89 ± 0.02	n.d.	n.d.
			800	0.46 ± 0.01	n.d.	n.d.
			1000	0.62 ± 0.01	n.d.	n.d.
C1	53.18° S	73.85° E	2000	1.01 ± 0.01	n.d.	n.d.
			20	364.44 ± 9.10	363.66 ± 9.11	100%
			40	328.60 ± 9.50	327.79 ± 9.50	100%
			80	366.67 ± 26.30	365.89 ± 26.35	100%
			100	770.22 ± 8.10	n.d.	n.d.
C5	52.42° S	75.6° E	120	420.08 ± 16.10	419.27 ± 16.10	100%
			40	2.19 ± 0.05	2.13 ± 0.05	97%
			80	5.02 ± 0.03	4.95 ± 0.03	99%
			120	2.22 ± 0.03	2.14 ± 0.03	96%
			150	4.10 ± 0.02	4.02 ± 0.02	98%
			200	18.42 ± 3.80	18.16 ± 3.80	99%
			300	17.29 ± 2.40	16.99 ± 2.40	98%
			400	15.97 ± 1.50	15.60 ± 1.51	98%
C11	51.65° S	77.73° E	450	65.59 ± 1.00	65.07 ± 1.02	99%
			500	51.42 ± 8.00	50.84 ± 8.01	99%
			40	0.97 ± 0.03	0.84 ± 0.03	87%
			80	0.98 ± 0.01	0.91 ± 0.02	93%
			120	1.00 ± 0.01	n.d.	n.d.
			200	0.56 ± 0.02	0.49 ± 0.02	87%
			400	0.53 ± 0.01	0.47 ± 0.01	89%
			500	1.74 ± 0.02	n.d.	n.d.
			600	2.23 ± 0.07	1.99 ± 0.07	89%
			800	1.55 ± 0.01	1.28 ± 0.01	82%
1000	2.02 ± 0.01	1.76 ± 0.01	87%			

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Table 2. Range of total dissolvable iron (TDFe), apparent particulate iron (Fe_{app}) and percentage of Fe_{app} in the total dissolvable fraction in the water column for different seasons and sectors of the Southern Ocean ((a) Sarthou et al., 1997, (b) Löscher et al., 1997, (c) Sedwick et al., 1997, (d) Sedwick et al., 2008), for a coastal environment in the Pacific Ocean ((e) Chase et al., 2005, (f) Johnson et al., 2001, (g) Fitzwater et al., 2003) and for the Atlantic Ocean ((h) Croot et al., 2004).

[Fe] (nmol L ⁻¹) on the whole water column	Total dissolvable Location	Apparent particulate Iron (nmol L ⁻¹)	Iron (nmol L ⁻¹)	%(Fe_{app} /TDFe)
Open Ocean	Indian Sector. summer (this study)	0.34–2.23	0.47–1.99	83–93 (mean: 86±3)
	Indian Sector. summer (a)	0.6–4.8		
	Atlantic Sector. spring (b)	0.5–8.9	0–6.86	0–97
Fe rich waters	Australian Sector. summer (c)	0.17–1.3	0.02–1.07	0–81
	Indian Sector. summer (this study)	0.9–65.6	0.8–65.1	82–99 (mean: 95±5)
	Indian Sector. summer (a)	6.2		
	Australian Sector. spring (d)	12.5		
	Monterey Bay. California. summer (e)	0.3–17.4		
	Monterey Bay. California. spring (f)	25		~80
	Monterey Bay. California (g)	0.80–6.57		
	Equatorial Atlantic (h)	0.3–6	0–5	0–80 (mean: 46±12)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.

Table 3. Mean dissolved and apparent particulate iron concentrations calculated above and below 150 m, for all the stations located above the plateau and for the stations located along transect C.

	Upper box		Deep box	
	Fe _{app} (nmol L ⁻¹)	DFe (nmol L ⁻¹)	Fe _{rapp} (nmol L ⁻¹)	DFe (nmol L ⁻¹)
Transect C*	186±39	0.44±0.03	33.74±23.17	0.41±0.14
Above the plateau	2.43±1.04	0.09±0.03	6.66±8.13	0.23±0.08

*Values in the upper box correspond to the mean of the two stations C1 and C5. Values in the deep box correspond to the mean of the single station C5.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.




Fig. 1. (a) The black rectangle denotes the location of the KEOPS study area in the Southern Ocean, (b) Location of the stations. Black arrows describe the general mean circulation around and above the plateau (Park et al., 2008b). Scale represents the bathymetry of the study area (in m). ACC represents the Antarctic Circumpolar Current and FTC represents the Fawn Through Current. Figure prepared using Ocean Data View (Schlitzer, 2007).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.




Fig. 2. Vertical profiles of total dissolvable iron (TDFe, nmol L^{-1}) at C1 station (near Heard Island), above the plateau (A3, B1, B3, B5, B7 and C5) and outside the plateau (B11 and C11). Depicted are mean values $\pm 1\text{SD}$. Depth and concentration scales are not the same for all the stations. For stations located above the plateau, the dashed lines represent the bottom depth.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.




Fig. 3a. Dissolved and particulate iron 2-box budgets above the Kerguelen Plateau. Letters used in the figure are explained in the text. We consider a surface layer of 150 m and a deep layer below 150 m with increasing depth from the transect C to the plateau. Dissolved iron fluxes are represented with numbers in italic. Particulate iron fluxes are represented with numbers in bold. Numbers in bold and italic are calculated from the resolution of the 4 equations describing the variation of dissolved and particulate iron with time.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.




Fig. 3b. Numerical values of dissolved (in italic) and particulate (in bold) iron fluxes (in 10³ mol d⁻¹) and stocks (in 10³ mol). Stocks are obtained by multiplying the mean concentrations calculated above the plateau in Fig. 3a by the volume of the boxes (6.75 × 10¹² m³ for the surface box and 1.81 × 10¹³ m³ for the deep box).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An iron budget during the natural iron fertilisation experiment KEOPS

F. Chever et al.




Fig. 4. Vertical profiles of total particulate (crosses) and apparent particulate iron (circles) iron at A3 (left) and C11 (right) stations. Depicted are mean values \pm 1SD. The two triangles represented at station A3 correspond to the Fe_{app} concentrations calculated with the dissolved samples measured at the shore-based laboratory, 18 months after sampling (Fe_{app2}) (see text for more details).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion