

Interactive
Comment

Interactive comment on “Influence of elevated CO₂ concentrations on cell division and nitrogen fixation rates in the bloom-forming cyanobacterium *Nodularia spumigena*” by J. Czerny et al.

J. Czerny

jczerny@ifm-geomar.de

Received and published: 9 July 2009

Thank you very much for the constructive remarks on our manuscript that helped us to improve it substantially.

1. Given the current controversy about altering CO₂ by acidifying the medium vs. bubbling with CO₂, some justification for choosing acidification, with the change in alkalinity this brings about, could have been included.

See comment Nr. 5 in the answer to comment by D. Hutchins.

C1097

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



2. I wonder whether, given the propensity of *Nodularia* to form large blooms and cell clumps, extension of the study to compare aggregated vs. homogeneous cultures might be warranted and discussed at least.

See comment Nr. 3 in the answer to comment by D. Hutchins.

3. I am intrigued by the suggestion that another cyanobacterium, *Anabaena*, may show a different response - is this closer to that of *Trichodesmium* or a different response again? A bit more information would be useful.

See comment Nr. 11 in the answer to comment by D. Hutchins.

Interactive comment on *Biogeosciences Discuss.*, 6, 4279, 2009.

BGD

6, C1097–C1098, 2009

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C1098

