

Interactive  
Comment

## ***Interactive comment on “Coastal hypoxia responses to remediation” by W. M. Kemp et al.***

**R. Diaz (Referee)**

diaz@vims.edu

Received and published: 28 July 2009

Additional Literature: Baird, D., R.R. Christian, C.H. Peterson, G.A. Johnson. 2004. Consequences of hypoxia on estuarine ecosystem function: Energy diversion from consumers to microbes. *Ecol. Appl.* 14:805-822.

Interactive comment on Biogeosciences Discuss., 6, 6889, 2009.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper






FIG. 3. Calculated summertime hypoxic volumes for Chesapeake Bay during 1950–2001 as reported in Table 3 and corresponding model predictions calculating using observed river flow (Eq. 1). Trend lines indicate model-predicted (e.g., 1) hypoxic volumes assuming low winter–spring average Susquehanna River flow ( $1,300 \text{ m}^3 \text{ s}^{-1}$ ), average flow ( $1,700 \text{ m}^3 \text{ s}^{-1}$ ), and high flow ( $2,100 \text{ m}^3 \text{ s}^{-1}$ ). Note the differences in y-axis scales.

Figure B. From Hagy et al. 2004.

Fig. 1. Figure B for Kemp et al. review

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

