

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Differences in community composition of bacteria in four deep ice sheets in western China

L. An¹, Y. Chen¹, S.-R. Xiang^{2,3}, T.-C. Shang⁴, and L.-De Tian^{2,3}

¹Key Laboratory of Arid and Grassland Agroecology (Ministry of Education) School of Life Sciences, Lanzhou University, Lanzhou, Gansu 730000, China

²Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Science, Beijing 100085, China

³State Key Laboratory of Cryospheric Science, Cold and Arid Regions Environmental Engineering Research Institute, Chinese Academy of Science, Lanzhou, Gansu 730000, China

⁴Department of Plant Pathology, Gansu Agricultural University, Lanzhou, Gansu 730070, China

Received: 29 January 2010 – Accepted: 8 February 2010 – Published: 17 February 2010

Correspondence to: S.-R. Xiang (srxiang@ns.lzb.ac.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Microbial community patterns vary in glaciers world wide, presenting unique responses to global climatic and environmental changes. Four bacterial clone libraries were established by 16S rRNA gene amplification from four ice layers along the 42-m-long ice core MuztB drilled from the Muztag Ata Glacier. A total of 152 bacterial sequences obtained from the ice core MuztB were phylogenetically compared with the 71 previously reported sequences from three ice cores extracted from ice caps Malan, Dundee, and Puruoganri. The six functional clusters *Flavisolibacter*, *Flexibacter* (*Bacteroidetes*), *Acinetobacter*, *Enterobacter* (*Gammaproteobacteria*), *Planococcus/Anoxybacillus* (*Firmicutes*), and *Propionibacter/Luteococcus* (*Actinobacteria*) frequently occurred along the Muztag Ata Glacier profile. Sequence analysis showed that most of the sequences from the ice core clustered with those from cold environments, and the sequences from the same glacier formed a distinct cluster. Moreover, bacterial communities from the same location or similarly aged ice formed a cluster, and were clearly separate from those from other geographically isolated glaciers. In a summary, the findings provide preliminary evidence of zone distribution of microbial community, support our hypothesis of the spatial and temporal biogeography of microorganisms in glacial ice.

1 Introduction

Variation in the communities of microorganisms in the deep ice sheets (ice cores) world wide reflects a response of microorganisms to global climatic and environmental change (Christner et al., 2000; Muller et al., 2004; Yergeau et al., 2007; Miteva et al., 2009). Analysis of a 102-m-long ice core drilled from the Malan ice cap (Xiang et al., 2004), a shallow ice core extracted at an elevation of 6350 m from the Muztag Ata Glacier (Xiang et al., 2005) as well as ice cores from the Arctic, Antarctica and other mountain glaciers (Christner et al., 2000) showed a great difference in the phylogenetic relationship of bacteria between the glacier ice and mild environments. These results

BGD

7, 1167–1199, 2010

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

suggest an important selective effect of the extreme cold glacier environments on microorganisms in glacier ice. However, it is not well known how changes in the patterns of microbial communities in glacier ice are related to climates and the environments.

Recent analysis of glacial snow pits from Kuytun 51 (Xiang et al., 2009b), Guoqu (Liu et al., 2009) and Rongbuk (Liu et al., 2006) revealed apparent differences in community composition in the different glacial snow layers, suggesting the effects of seasonal conditions on the microbial communities. Phylogenetic comparison of bacterial communities in the surface snow and snow pits among the isolated glaciers showed less of a shift between seasonal communities than between those extending over a large spatial scale (Xiang et al., unpublished). A recent investigation of the Greenland ice core GISP2 demonstrated a great difference in the proportion of the main phylogenetic phyla during the distinct geographical periods which occurred from 30–80 Ka, suggesting a strong temporal effect of aeolian activities on community composition of microorganisms in Greenland (Miteva et al., 2009). These results suggest strong spatial and temporal effects on the microbial communities in glacier ice-snow.

To investigate microbial biogeography in glacial ice over extended scales of time and space, the present study extended previous preliminary investigations on the glacier surface (Xiang et al., 2009c) and deep snow (Xiang et al., 2009b), and further investigated the composition of bacterial communities along the four ice core profiles extracted from geographically isolated glaciers. First, to investigate the seasonal community changes in glacier ice, four bacterial clone libraries were established by 16S rRNA gene amplification from the ice core MuztB drilled at an elevation 7000-m Muztag Ata Glacier. Second, the geographical effects on the evolution of microorganisms in glacier ice were preliminarily evaluated by a sequence comparison between the glaciers and the surrounding mild environments, and phylogenetic comparison among the various geographically isolated glaciers. Finally, to investigate the community shift at a large spatial scale, four clone libraries from the ice core MuztB were phylogenetically compared with eight previously recovered bacterial clone libraries from ice cores drilled from three ice caps Dunde (Zhang et al., 2009), Malan (Xiang et al., 2004;

**Differences in
community
composition of
bacteria**L. An et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Zhang et al., 2002) and Puruogangri (Zhang et al., 2009). This study attempted to provide interpretations of biogeography of microorganisms at a wide range of altitudes. The limitation of conclusions on biogeography of microorganisms in ice cores was also discussed in this study.

2 Materials and methods

2.1 Study sites and sample collection

The data used in this study were collected from four ice cores from the Muztag Ata Glacier (38°17' N, 75°04' E, Tian et al., 2006), Dundee ice cap (38°06' N, 96°24' E, Zhang et al., 2009), Puruogangri ice cap (33°44'–34°04', 89°20'–89°50' E, Zhang et al., 2009), and Malan ice cap (35°48.40' N, 90°35.34' E, Xiang et al., 2004), respectively. The Muztag Ata Glacier is located in the most western periphery of the Tibetan Plateau, where precipitation is derived from the air masses from arid and semi-arid regions, including the deserts Sary-Ishykotrau, Muyun Kum, Kyzyl Kum, Kara Kum, Taklimakan and Gurbantunnut (Fig. 1). The Dundee ice cap is located in the north margin of the Qaidam Basin and in the Qilian mountain region on the northeastern Tibetan Plateau (Fig. 1). Precipitation results from the incursion of westerly depressions along the southern slopes of the Hymalayas during the winter (Murakami, 1987; Davis et al., 2005). During the summer, Indian monsoon circulation transports moisture from the Bay of Bengal to the central Hymalayas, and extend to the Qaidam Basin (Davis et al., 2005). Moreover, the numerous large depressions in the Takalamakan Desert and Daidam Basin cause strong winds and snowstorms as well (Dregne, 1968; Chen and Bowler, 1986). The Malan and Puruogangri ice caps are located in the center of the Tibetan Plateau, where precipitation is derived from a westerly direction in winter, and Indian monsoons in summer (Wang, 1989; Wake et al., 1993; Shi and Liu, 2000).

The first ice core MuztB (42-m-long) was extracted at 7010 m a.s.l. (above sea level) from the Muztagata Glacier in the summer of 2003 (Tian et al., 2006), the second ice

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



core Dundee (140-m-long) was extracted at an elevation of 5325 m on the Dundee ice cap in 1987 (Thompson et al., 1990). The third ice core Puruogangri (86-m-long) was extracted at an elevation of 5970 m from the Puruogangri ice cap in October, 2000 (Thompson et al., 2006), and the fourth ice core, Malan (102-m-long), was extracted at an elevation of 5620 m on the Malan ice cap in 1999 (Wang et al., 2003). The visible stratigraphic features were recorded immediately after ice core drilling. All ice cores were returned frozen to the freezer room (air temperature between -18 to -24 °C) at the Key Laboratory of the Ice Core and Cold Regions Environment of the Chinese Academy of Sciences. Each ice column of the obtained ice cores was split lengthwise into four sections, and stored in a refrigerated room at -18 °C to -24 °C. All ice core sub-samples were always handled at temperatures below 20 °C within a sterile and positive-pressure laminar flow hood by following the procedure described previously (Xiang et al., 2005). An annulus (10 mm) was (cut) successively cut three times from the surface of each core sample using three clean, sterilized saw-tooth knives. The remaining inner core was washed, and samples were allowed to melt at 4 °C in covered, autoclaved containers and then used for further analysis. Microbial data of the three ice cores Dundee, Puruogangri, and Malan were collected from published reports (Zhang et al., 2009; Xiang et al., 2004). The ice columns used in this study were subsections of ice cores MuztB, Malan, Puruogangri and Dundee, and dated to AD 1970 to 1988 (Tian et al., 2006), 1600 to 1800 (Wang et al., 2003), 1750 to 1920 (Thompson et al., 2006) and 1780 to 1830 (Thompson et al., 1990), respectively.

2.2 Biomass analysis and clone library establishment of the bacterial 16S rRNA gene amplified from the Muztagata Glacier

The fresh melt-water (10 ml) obtained from the ice core MuztB was used for the determination of the total and live biomass by flow cytometric (FCM) analysis. Approximately 400 ml of melt-water was used for DNA extraction. The FCM analysis, DNA extraction and further clone library establishment procedure were conducted by following the same protocols as previously used in a microbial analysis of the Kuytun 51

**Differences in
community
composition of
bacteria**L. An et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Glacier samples (Xiang et al., 2009c). To avoid possible bias, the pooled PCR products were used to establish clone library from each ice column. A total of 151 clones were sequenced by *Hae*III-based ARDRA (amplified rRNA restriction analysis) out of the 352 clones from the ice core MuztB obtained from the Muztag Ata Glacier. Each sequence was named using the initial of Muztag Ata Glacier (MuztB, B was noted for the ice core drilled at an elevation of 7010 m to distinguish it from the Muztag Ata ice core MuztA at 6350 m a.s.l. in the summer of 2002, Xiang et al., 2005), along with the column tube number (13, 16, 28 and 30) followed by the clone number (1 to 148). For example, clones MuztB13-132, MuztB16-36, MuztB28-125, MuztB30-118 were representative of the ice core columns 13 (1988 spring), 16 (1984 autumn–1985 winter), 28 (1972 spring–summer), and 30 (1970 winter) which were obtained from the ice core MuztB drilled at 7010 m a.s.l. of the Muztag Ata Glacier. The time series of the ice core MuztB was determined based on the visible annual layers, seasonal fluctuation in the oxygen isotope ratios, and the beta activity in the melt water as a reference (Tian et al., 2006). The accession numbers of the cloned sequences obtained from the Muztag Ata ice core B in GenBank are: GU246831–GU246982.

2.3 Statistical analysis of the bacterial communities in the four deep ice sheets in western China

To investigate the biogeography of microorganisms in glacier ice over extensive spatial and temporal scales, all 151 sequences from the Muztag Ata Glacier were compared with the 71 previously reported sequences obtained from the three glaciers Malan (accession number AY322483–AY322493 in Xiang et al., 2004 and AY121823–AY121830 in Zhang et al., 2002), Dunde (accession number AY313918, AY313919 and DQ076445–DQ076456, Zhang et al., 2009), and Puruogangri (accession number AY313907–AY313917, AY313920–AY313922, and DQ076420–DQ076444, Zhang et al., 2009), as well as other known reference species obtained by Blast search (Altschul et al., 1990) and aligned with reference sequences obtained using ClustalX (Thompson et al., 1997). A Neighbor-Joining (NJ, Tamura et al., 2004) phylogeny for

**Differences in
community
composition of
bacteria**L. An et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the aligned sequences was constructed using MEGA 4.0 (Tamura et al., 2007: <http://www.megasoftware.net/>) (with) using the pairwise deletion mode for gaps and (with) the Maximum Composite Likelihood (MCL) method for substitutions. The 16S rDNA sequences from *Methanosaeta harundinacea* strain 8Ac (accession no. AY817738) and *Methanosaeta concilii* strain NW-1 (accession no. DQ150255) were used as out-group references on all trees. The obtained sequences displaying similarities of >97% with known species were identified as the reported species. Most of the obtained clones were related to known cultivated genera or genus clones (e.g., *Acinetobacter* sp., *Cryobacterium* sp., and *Sphingomonas* sp.). A few clones had <97% similarity with reported species, and thus were designated separately.

The diversity (Shannon-Wiener index H') and evenness (E) indices were based upon the distribution of unique sequence OTUs (operational taxonomic units) obtained from the clone libraries using equations: $H' = -\sum\{p_i \cdot \ln(p_i)\}$ and $E = H'/\ln(S)$, respectively (Hill et al., 2003), where p_i = the proportion of the i^{th} clone in the total clones in each individual library, and S is the single unique sequence richness. Bacterial phylotype abundance and coverage estimators were calculated with the software program EstimateS (Kemp and Aller, 2004, the EstimateS program is available online at <http://www.aslo.org/lomethods/free/2004/0114a.html>).

To further investigate the biogeographic effect on the bacterial communities in the ice cores drilled from the geographically different glaciers in western China, the representative sequences were compared among four glaciers Muztag Ata Glacier, Malan (Xiang et al., 2004; Zhang et al., 2002), Dundee (Zhang et al., 2009), and Puruogangri (Zhang et al., 2009). The bacterial community composition was statistically analyzed using the UniFrac software package (Lozupone and Knight, 2005).

3 Results

3.1 Differences in biomass and diversity of the dominant bacteria along the Muztag Ata Glacier depth profile

There was a great variation in diversity of the dominant bacteria along the Muztag Ata Glacier profile, although there was a little change in the total and live biomass at a range from 1.03×10^5 to 1.99×10^5 cells/ml and 4.28×10^3 to 4.98×10^5 cells/ml, respectively, in the four tested ice layers (Table 1). The ice layer in the warm spring-summer season of 1972 contained the most diverse bacteria, as estimated by the Schao1 value, the winter 1970 ice layer contained the highest Shannon index value, and the ice layer from autumn 1984 to winter 1985 contained the lowest OUTs of bacteria in both the Schao1 and Shannon indices (Table 1). This indicated a great heterogeneity of bacteria along the glacier depth profile.

The dominant bacteria in the four ice layers were investigated by 16S rRNA gene amplification, as well as clone library and sequencing techniques. BLAST results of the 16S rDNA sequences obtained from the Muztag Ata ice core showed that the bacterial phyla *Gammaproteobacteria*, *Betaproteobacteria*, *Actinobacteria*, *Bacteroidetes*, and *Firmicutes* frequently occurred throughout the glacier depth profile (Fig. 2). However, there was a great difference in the proportion of the main phylogenetic groups along the glacier depth profile (Fig. 2). Phyla *Alphaproteobacteria* and *Firmicutes* were dominant in the spring ice layer of 1988, *Gammaproteobacteria*, *Betaproteobacteria*, *Actinobacteria*, and *Bacteroidetes* phyla were dominant in the ice layer from 1984 autumn to 1985 winter, *Gammaproteobacteria*, *Betaproteobacteria*, *Actinobacteria*, *Bacteroidetes*, and *Firmicutes* were common in the 1972 spring-summer ice layer, and *Gammaproteobacteria*, *Betaproteobacteria*, and *Actinobacteria* dominated the community in the 1970 winter ice layer. Interestingly, the *firmicutes* phylum was dominant in the spring seasons of both 1988 and 1972, but was rare in the winter seasons of 1970 and 1985.

BGD

7, 1167–1199, 2010

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.2 Similarities and differences in the main phylogenetic bacterial groups among the four isolated glaciers

To investigate the biogeography of microorganisms in glacier ice, the sequences obtained from the Muztag Ata Glacier were compared with those from the geographically isolated glaciers Puruogangri, Dundee and Malan. The sequence data showed that all of the ice core clones were related (with 97% to 100% similarity) to those from the different environments of urban aerosols, soil, from the river to the Antarctic (Fig. 3a, b, c, and d). For example, two clones MuztB30-40 and TD-21 were 97% similar to *Comamonadaceae* sp. clone XJ-L144 (with accession number EU817496, a betaproteobacterial family from the Xiangjiang River, clone MuztB30-61, which was 100% identical to a *Pelomonas saccharophila* strain (AF368755) from the ultrapure water from an industrial system, and the two clones MuztB28-21 and MuztB28-32, which are closely similar (99% similarity) to a *Methylibium fulvum* isolate (AB245356) from ginseng field soil (Fig. 3a). However, most of the glacier bacterial clones were closely clustered with those from the cold environments, such as the glaciers and Antarctica (Fig. 3a, b, c, d, and e). The two clones MuztB28-21 and MuztB28-32 were easily clustered with clone Kuy-SL-42 (EU263707) from the Kuytun 51 Glacier in the *Methylibium* sp. group within the family of *Betaproteobacteria* (Fig. 3a). Clones MuztB28-125, MuztB16-36, P80-5, P80-18, P200-23 and P60-49 were closely clustered with the betaproteobacterial clone KuyT-IWPB-41 (EU263727) from the Kuytun 51 Glacier and a clone (DQ675477) from the Hymilayan glacier, respectively (Fig. 3a). Other clones from the Muztag Ata, Puruogangri, and Dundee glacier clustered with the Antarctic clones H12_ELL02 (EF220189) and H07_ELL02 (EF220180) within the *Flexibacter* sp. group in the family of *Bacteroidetes* (Fig. 3c).

There were certain common species which occurred in the different glaciers. For example, in the *Rhodoferrax* sp. group in the *Betaproteobacteria* phylum, clone MuztB28-95 from the Muztag Ata Glacier was 99% similar to clone Malan B-48 from the Malan Glacier (Fig. 3a). Clone Malan D-10 clustered with the Muztag Ata glacier clone group

BGD

7, 1167–1199, 2010

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(e.g., clones MuztB 28-104 and MuztB13-2) and fell into the *Acinetobacter* sp. group among the members of the *Gammaproteobacteria* phylum (Fig. 3a).

The common bacterial clones not only occurred in the geographically isolated glaciers, but were also present in the different ice layers throughout the ice core profile, which was very evident along the MuztB ice core profile (Fig. 3a, b, c, d, and e). In the *Schlegella* sp. cluster within the *Betaproteobacteria* phylum, two clones MuztB28-39 and MuztB28-81 from the ice column 28 (1972 spring–summer) closely clustered with clone MuztB30-36 from the 1970 winter column 30 (Fig. 3a). Two clones (MuztB30-15 and MuztB30-3) from ice column 30 (1970 winter), MuztB30-15 and MuztB30-3, were 98% similar to two clones from the spring 1988 ice column 13 in the *Xanthomonas* sp. subgroup within the members of *Gammaproteobacteria* subphylum (Fig. 3a). In *Chamaeshiphon* sp. cluster within the *Cyanobacteria* phylum, clone MuztB16-114 from ice column 16 (1984 autumn–1985 winter) was 100% identical to the clone MuztB28-129 from ice column 28 (1972 spring–summer) (Fig. 3c). In the *Thermus* sp. cluster (*Thermus*), clone MuztB30-117 from ice column 30 (1970 winter) was associated with 100% similarity with clones MuztB13-91, MuztB13-59 and MuztB13-58 from ice column 13 (1988 spring) and clone MuztB28-56 from the 1972 ice column 28 (Fig. 3c). Similarly, *Anoxybacillus* sp. (*Firmicutes*) occurred throughout the ice columns of MuztB 13 (e.g., clones MuztB13-102, MuztB13-14, and MuztB13-78 etc.), MuztB 16 (e.g., clone MuztB16-76), MuztB 28 (e.g., clones MuztB28-123, MuztB28-16, MuztB28-5, MuztB28-57) and MuztB 30 (e.g., clones MuztB30-28 and MuztB30-136), in the ice layers spring 1988, autumn 1984–winter 1985, spring–summer 1972 and winter 1970, respectively (Fig. 3d).

However, a comparison of sequences among the four geographically isolated glaciers showed that the bacterial clones from the same location readily formed a group, and were clearly distinct from those recovered from the geographically isolated glaciers (Fig. 3a, b, c, d, and e), which presented a phenomenon of zone distribution of the microorganisms in glacier ice on a large spatial scale. This was apparent in the dominant clusters, as indicated by the gray shaded areas within the members of

**Differences in
community
composition of
bacteria**L. An et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Differences in
community
composition of
bacteria**L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Betaproteobacteria, *Gammaproteobacteria*, *Bacteroidetes* and *Actinobacteria* phylum (Fig. 3a, b, c, d, and e). In the *Rhodoferrax* sp. cluster within the *Betaproteobacteria* phylum, the Muztag Ata Glacier clones MuztB28-118 and MuztB28-125 were clearly distinct from the two clones TD-16 and TD-46 from the Dunder Glacier (Fig. 3a). Four clones, MuztB28-85, MuztB13-132, MuztB30-98, and MuztB16-36 from the Muztag Ata Glacier, formed a cluster distinct from four clones from the Puruogangri Glacier in the *Polaromonas* sp. cluster within the *Betaproteobacteria* phylum (Fig. 3a). The two clones Malan D-11 and Malan C-37 from the Malan Glacier were clearly distinct from the eight Muztag Ata Glacier clones MuztB16-20, MuztB16-66, MuztB13-46, MuztB28-104, MuztB13-2, MuztB30-62, MuztB13-92 and MuztB13-25 in the *Acinetobacter* sp. cluster within the *Gammaproteobacteria* phylum (Fig. 3a). Similarly, the clones from the Muztag Ata Glacier (e.g., clones MuztB16-2 and MuztB16-14), were clearly separated from the clones from the Puruogangri Glacier (e.g., clones P60-11 and P60-25) in the *Flexibacter* sp. cluster within the *Bacteroidetes* phylum (Fig. 3c). The Malan Glacier clone Malan A38 distinct from the Dunde Glacier clone TD-77 and the Muztag Ata Glacier clones (e.g., clones MuztB28-30 and MuztB13-99) in the *Flavisolibacter* sp. cluster within the *Bacteroidetes* phylum (Fig. 3c); while the two clones MuztB28-18 and MuztB28-111 from the Muztag Ata Glacier were distinct from clone P80-33 from the Puruogangri Glacier in the *Cryobacteria/Frigoribacter* cluster within the *Actinobacteria* phylum (Fig. 3d).

The proportion of the main phylogenetic groups varied throughout the depth profile of the Muztag Ata Glacier (Fig. 4). Six clusters, *Planococcus/Anoxybacillus*, *Propionibacter/Luteococcus*, *Flavisolibacter* sp., *Flexibacter* sp., *Acinetobacter* sp., and *Enterobacter* sp. were common throughout the ice core profile, but they appeared at different frequencies along the profile. *Propionibacter/Luteococcus* dominated in the cold winter ice layers in 1970 (Fig. 4d) and 1985 (Fig. 4b), but rarely occurred in the spring ice layers in 1972 (Fig. 4c) and 1988 (Fig. 4a), indicating their prevalence in dry cold seasons. *Flexibacter* sp. dominated in the ice layers in summer 1972 and winter 1985, while *Enterobacter* sp. dominated in the ice layer in winter 1970. Other bacteria

only occurred in certain specific seasons, as indicated by the solid triangles (Fig. 4), likely indicating an opportunist bacterial event in the glacier. A combination of prevailing, opportunistic, and other minor species constituted the community of bacteria in the ice-snow layers of specific seasons along the Muztag Ata Glacier profile.

3.3 Community comparison of bacteria among the four geographically isolated glaciers

UniFrac analysis of the four glaciers Muztag Ata, Dundee, Malan, and Puruoganri revealed a biogeography of bacterial communities (Fig. 5). Generally, the obtained 9 clone libraries of bacteria generated four clusters corresponding to the geographical distance, with the bacterial communities from the same glacier clustering with one another, clearly separated from those of other geographically isolated glaciers. This was particularly evident for the bacterial communities from the ice layers along the Puruoganri and Muztag Ata ice core profiles. All four bacterial clone libraries from the Muztag Ata Glacier fell into the same cluster, and the same trend was found for the two clone libraries from the Puruogangri Glacier.

Interestingly, the bacterial communities appeared to cluster with one another not only on a spatial but also a temporal scale (Fig. 5). This was clearly evident for the bacterial communities across the four geographically glaciers. Five bacterial clone libraries from the ancient ice layers of >200 years were clearly separated from four clone libraries from the modern ice layers of <40 years. Two clone libraries from around AD 1600 and AD 1800 ice layers in the Malan Glacier formed two distant branches, and distinct from other younger ice layers in the Muztag Ata Glacier. The clone library from the ice layer TD obtained from the Dundee Glacier of AD 1780 - 1830 clustered with the two clone libraries from the Puruogangri Glacier of similar ice age, i.e. AD 1750–1800 and 1850–1920, respectively.

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4 Discussion

Previous reports of the deep ice sheets (ice cores) demonstrated a great variability in the proportion of the main phyla of bacteria in glaciers worldwide (Christner et al., 2000; Xiang et al., 2005; Liu et al., 2009), which was also evident in our present analysis of four ice cores extracted from the four geographically isolated glaciers, Muztag Ata, Dunde, Puruogangri and Malan. This suggests a biogeographic effect on microbial communities in glacier ice. In this current study, sequence comparison of the bacterial 16S rRNA genes amplified from the four ice cores indeed showed clear zone distribution of bacterial communities. This supports our hypothesis of the spatial and temporal biogeography of microorganisms in glacier ice.

4.1 Methodological considerations

Ice core records have been used to reconstruct a history of climatic and glacial changes, presenting consistent results with the available climatic data (Naftz et al., 2002; Thompson et al., 1989, 2000; White et al., 1997; Yao, et al., 1999). The quality of the ice core records at the extremely high altitude (>5300 m a.s.l.) is much less influenced by the interruption of snow-melting and snow-layer-overturning and sublimation than at low altitude. Moreover, all the four ice cores used in this study were cautiously dated to 48 to 400 years before today (Wang et al., 2003; Thompson et al., 1990, 2006; Tian et al., 2006), and the MuztB ice dating was confirmed by the peak in β activity (Tian et al., 2006). The evident correlation (coefficient of 0.67) of annual variation in values of oxygen isotope ratios with the annual air temperature changes at the nearby meteorological station Taxkorgen (Tian et al., 2006) indicated a reliability of temperature records in the ice core MuztB. The high quality of ice core records makes validity of microbial data from modern and ancient ice, and sufficient for community comparison of microorganisms in the geographically isolated glaciers.

BGD

7, 1167–1199, 2010

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.2 The ubiquity and biogeography of bacteria in glacier ice

Previous investigations showed that most of the bacterial sequences from glaciers worldwide are closely related to those from certain environments, such as agricultural soil, river water and urban aerosol (Christner et al., 2000; Xiang et al., 2005; Miteva et al., 2004), indicating the ubiquity of microorganisms on the earth. The ubiquity of microorganisms was also apparent in the bacteria found in the four geographically isolated glaciers, with 96% to 100% sequence similarity to those from mild environments (Fig. 3a, b, c, d, and e). Previous studies, however, also reported phylogenetic differences between bacteria from glacier ice and the surrounding mild environments. These glacier clone sequences closely cluster with those from cold environments such as sea ice, Antarctic soil and other snow-ice (Xiang et al., 2005; Zhang et al., 2009). This reveals the transportation of microorganisms from outside environments onto the glacier surface (Gloster et al., 1982; Shuval et al., 1989; Abyzov, 1993; Prospero et al., 2005), and strengthens the concept of adaptation and acclimation of microorganisms to the extremely cold glacier environments (Morgan-Kiss et al., 2006; Vincent, 2000). This was obvious for the biogeographic properties of the microorganisms in glacier ice when all of the sequences from the four geographically isolated glaciers Muztag Ata, Malan, Dundee and Puruogangri, along with the nearest relatives from other environments, were subjected to phylogenetic analysis (Fig. 3a, b, c, d, and e). Moreover, 60% of the total bacterial clones from the same glacier (e.g., the Muztag Ata Glacier) were easily clustered together, clearly separating them from those from other, geographically isolated glaciers, or the surrounding environments (Fig. 3a, c, and d). The biogeography of microorganisms in glacier ice may be attributed to the geographic distance and the consequently isolated effect on the evolution of microorganisms in glacier snow-ice, which will be discussed below.

BGD

7, 1167–1199, 2010

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.3 Biogeography of bacterial community in glacier ice

Several bacterial genera frequently occurred throughout the depth profile from the Muztag Ata Glacier, indicating their prevalence in this specific glacier (Fig. 4). The six functional clusters *Planococcus/Anoxybacillus*, *Propionibacter/Luteococcus*, *Flavisolibacter* sp., *Flexibacter* sp., *Acinetobacter* sp., and *Enterobacter* sp. constituted the main community constituents throughout the depth profile, suggesting they are the prevalent bacteria in the Muztag Ata mountain regions. In particular, *Acinetobacteria* sp. not only was found in all four of the established clone libraries from the ice core MuztB profile extracted at an elevation of 7010 m in the Muztag Ata Glacier (Fig. 4a, b, c, and d), but was also frequently isolated by a culture method from four ice layers from the ice core drilled at an elevation of 6350 m in the same location (Xiang et al., 2005). This confirmed that the culture-independent based results on the dominant bacteria in the ice core was consistent with our previous culture-based data from the Muztag Ata Glacier, and thus it was sufficient for a comparison of dominant bacteria in the different seasons (Fig. 4). The prevalent bacteria in the ice core may represent the biological indicators of climatic and environmental conditions in the air mass over the Muztag Ata Glacier, and also on the subsequent glacier surface during the current deposition period. The phenomenon of prevalent bacteria in the local regions can be found in other reports as well. For example, *Bacillus* sp., and *Microbacter/Arthrobacter* sp. were frequently isolated from the ice layers along the Guliya Glacier profile (Christner et al., 2000), potentially serving as a microbial indicator of the air mass over the glacier and in the surface snow-ice. *Propionibacterium* sp. and *Bacillus* sp. frequently appear in three clear ice layers along the Greenland GISP2 ice core profile (Miteva et al., 2009), constitute the prevalent bacteria at the specific deposition time, and thus may be considered as the biological indicator of the prevailing air mass over Greenland during the current deposition period. Likely, *Flexibacter* sp. and *Polaromonas* sp. frequently occurred at three ice layers along the Puruoganri ice core profile (Fig. 3a and c), constituted the main components of bacterial community in the ice core (Fig. 1 in reference of Zhang et al.,

BGD

7, 1167–1199, 2010

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2009), and thus may be the biological indicator in the prevailing air mass over the Puruogangri Glacier. Three clusters *Arthrobacter* sp. (*Actinobacteria*), *Pseudomonas* sp. (*Gammaproteobacteria*) and *Sphingomonas* sp. (*Alphaproteobacteria*) were the dominant components of microbial community in the Guoqu (Geladaindong) Glacier (Yao et al., 2008).

4.4 Climatic and environmental implications of microbial communities in glacier ice

Microorganisms vary in geographically isolated glaciers, not only at the species-genera level (Fig. 3a, b, c, d, and e), but also at the whole community level (Fig. 4 and 5). The Muztag Ata Glacier community is dominated by the six functional clusters *Planococcus/Anoxybacillus*, *Propionibacter/Luteococcus*, *Flavisolibacter* sp., *Flexibacter* sp., *Acinetobacter* sp., and *Enterobacter* sp. (Fig. 4). In contrast, the Puruogangri Glacier community is dominated by three main functional clusters, *Cryobacterium/Aeromicrobium* sp., *Polaromonas* sp., and *Flexibacter* sp. (Fig. 3a, c, d, and e; Zhang et al., 2009). This distinct functional community composition may be attributed to a combination of microbial deposition through aeolian activities over a glacier as well as post-deposition selection on the community structure of microorganisms in the surface snow-ice (Xiang et al., 2009a). On the one hand, the Muztag Ata Glacier is located in the most western periphery of the Tibetan Plateau, surrounded by the vast arid and semi-arid regions of central Asia (Fig. 1). It receives precipitation mainly derived from the western dry air mass (Wang, 1989; Wake et al., 1993), in which the frequent dust storms (Li et al., 2003) may carry abundant microorganisms to the glacier surface. The Dunde, Malan, and Puruogangri ice caps are located in the northern to middle part of the Tibetan Plateau, where the air masses are derived from the cold westerly mass in winter and mild south Asia monsoon in the warm summer seasons. They are also affected by the powerful winds and snowstorms caused by the numerous large depressions in the Tibetan Plateau, and in the local mountain regions (Fig. 1, Wang, 1989; Wake et al., 1993; Dregne, 1968; Chen and Bowler, 1986). The dramatic changes of

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Differences in
community
composition of
bacteria**L. An et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the air masses may lead to differences in the microbial species pool and thus result in the distinct community composition of microorganisms across the glaciers. On the other hand, differences in the local climatic and environmental conditions, such as temperature, light intensity, melt-water availability and nutrient concentrations in the glacier ice (Takeuchi et al., 1998, 2001; Takeuchi and Li, 2008; Takeuchi and Koshima, 2004) may cause a great variation in the growth rate of tolerant microorganisms, which in turn leads to the subsequent changes in the community composition of microorganisms in glacier ice. Variations in the functional population pool as a result of both aeolian and post-deposition processes lead to the apparent zone distribution of microbial communities, which clearly corresponds to the distances across the four geographically isolated glaciers (Fig. 5). Compared with the ancient ice layers in the Malan, Puruoganri, and Dunde glaciers, the modern ice layers in the Muztag Ata Glacier contain distinct communities at a very fine lineage level, as shown by the UniFrac distance in Fig. 5. However, there is an uncertainty of the differences between the modern ice and ancient ice because lack of a consistent ice core records, which may be a result of spatial effect on the microbial communities across the four glaciers (Fig. 5). The evident spatial patterns of the microbial communities across the four deep ice sheets supports our hypothesis of an ecological selection effect on the functional community composition of microorganisms in the air mass and under the glacier system over extended spatial scales.

The results suggest the spatial and temporal biogeography of dominant bacteria across four geographically isolated glaciers, including three characteristics. First, most bacterial species from the same glacier form a distinct cluster (gray shaded areas in Fig. 3a, c, and d). Second, several functional phylogenetic clusters contributed to the community shift (Fig. 4) across the glaciers on the Tibetan plateau. Finally, variations in the proportion of the main functional phylogenetic clusters at the species/genera level resulted in a seasonal community shift along the glacier depth profile. Although the high quality of ice core records at the extremely high altitude showed seasonal changes of major species, it can not absolutely excluded the uncertainty of minor species trend

**Differences in
community
composition of
bacteria**L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in different seasons since our data were a series of single-ice-core based results. In this current study, the community shift of microorganisms was related to the changes in the air masses over the glaciers on the Tibetan plateau. This has important implications for research on global climatic and environmental changes using microbial indices in glacial ice. However, it is not clear how the prevailing aeolian and post-deposition processes influence the climatic and environmental changes in the glacial regions considered. The preliminary investigations of temporal and spatial patterns of microbial communities across the four glaciers provide only a glimpse of microbial biogeography and its relation to global climatic and environmental changes. This current study was based on the limited group of thirteen clone libraries established from four glaciers. More data on the meteorologic, physico-chemical, and biological characteristics of the surface snow and repeatable ice core data of microbial analysis will be crucial for truly understanding the dynamics of microorganisms in glacier ice. More sequence data based on multiple-loci genes such as *recA* (encoding the multi-functional DNA-binding protein involved in homologous recombination, Story et al., 1992), *gyrB* (encoding the DNA gyrase B protein, Kumar et al., 1993), *wzm* (encoding the inner membrane protein, Reeves et al., 1996), and *rpoB* (encoding the β -subunit of RNA polymerase B, Landick et al., 1990) of the chromosome of microorganisms in glacier ice may provide the necessary detailed information to advance microbial biogeography, since the sequence data in this study were based on only a single 16S rRNA gene analysis.

Acknowledgements. We thank all of the members of the Muztag Ata Glacier expedition for assistance in the field sample collection. We also thank for Pacific Edit for the improvement of English use in this manuscript prior to submission. This work was supported by the NSF project of China (Grant 40471025 and 40871046).

References

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J.: Basic local alignment search tool, *J. Mol. Biol.*, 215, 403–410, 1990.

**Differences in
community
composition of
bacteria**

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Chen, K. and Bowler, J. M.: Late Pleistocene evolution of salt lakes in the Qaidam Basin, Qinghai Province, China, *Paleogeog. Paleoclimat. Paleoeol.*, 54, 87–104, 1986.
- Christner, B. C., Mosley-Thompson, E., Thompson, L. G., Zagorodnov, V., Sandman, K., and Reeve, J. N.: Recovery and identification of viable bacteria immured in glacial ice, *Icarus*, 144, 479–485, 2000.
- Davis, M. E., Thompson, L. G., Yao, T., and Wang, N. L.: Forcing of the Asian monsoon on the Tibetan Plateau: evidence from high-resolution ice core and tropical coral records, *J. Geophys. Res.*, 110, D04101. doi:10.1029/2004JD004933, 2005.
- Dregne, H. E.: Surface materials of desert environments, in: *Deserts of the world*, edited by: McGinnies, W. G., Goldman, B. J., and Paylore, P., University of Arizona Press, Tuscon, 286–377, 1968.
- Gloster, J., Sellers, R. F., and Donaldson, A. I.: Long distance transport of foot-and-mouth disease virus over the sea, *Veterin. Record*, 110, 47–52, 1982.
- Hallett, P., Grimshaw, A. J., Wigley, D. B., and Maxwell, A.: Cloning of the DNA gyrase genes under tac promoter control: overproduction of the gyrase A and B proteins, *Gene.*, 93, 139–142, 1990.
- Hill, T. C. J., Walsh, K. A., Harris, J. A., and Moffett, B. F.: Using ecological diversity measures with bacterial communities, *FEMS Microb. Ecol.*, 43, 1–11, 2003.
- Kemp, P. F. and Aller, J. Y.: Estimating prokaryotic diversity: When are 16S rDNA libraries large enough?, *Limnol. Oceanogr. Methods*, 2, 114–125, 2004.
- Kumar, K. A., Mahalakshmi, S., and Muniyappa, K.: DNA-induced conformational changes in RecA protein, *J. Biol. Chem.*, 268, 26162–26170, 1993.
- Landick, R., Stewart, J., and Lee, D. N.: Amino acid changes in conserved regions of the beta-subunit of *Escherichia coli* RNA polymerase alter transcription pausing and termination, *Genes. Dev.*, 4, 1623–1636, 1990.
- Li, X., Qin, D., Jiang, G., Duan, K., and Zhou, H.: Atmospheric pollution of a remote area of Tianshan Mountain: ice core record, *J. Geoph. Res.*, 108, 4406–4416, 2003.
- Liu, Y., Yao, T.-D., Kang, S., Jiao, N., Zeng, Y., Shi, Y., Luo, T., Jing, Z., and Huang, S.: Seasonal variation of snow microbial community structure in the East Rongbuk glacier, Mt. Everest, *Chin. Sci. Bull.*, 51, 1476–1486, 2006.
- Liu, Y., Yao, T., Jiao, N., Kang, S., Xu, B., Zeng, Y., Huang, S., and Liu, X.: Bacterial diversity in the snow over Tibetan Plateau Glaciers, *Extremoph.*, 13, 411–23, 2009.
- Lozupone, C. and Knight, R.: UniFrac: a new phylogenetic method for comparing microbial

- communities, *Appl. Environ. Microbiol.* 71, 8228–8235, 2005.
- Miteva, V., Sheridan, P. P., and Brenchley, J. E.: Phylogenetic and physiological diversity of microorganisms isolated from a deep Greenland glacier ice core, *Appl. Environ. Microbiol.*, 70, 202–213, 2004.
- 5 Miteva, V., Teacher, C., Sowers, T., and Brenchley, J.: Comparison of the microbial diversity at different depths of the GISP2 Greenland ice core in relationship to deposition climates, *Environ. Microbiol.*, 11, 640–656, 2009.
- Morgan-Kiss, R. M., Priscu, J. C., Pockock, T., Gudynaite-Savitch, L., and Huner, N. P. A.: Adaptation and acclimation of photosynthetic microorganisms to permanently cold environments, *Microb. Molecul. Biol. Rev.*, 70, 222–252, 2006.
- 10 Mueller, D. R. and Pollard, W. H.: Gradient analysis of cryoconite ecosystems from two polar glaciers, *Polar. Biol.*, 27, 66–74, 2004.
- Murakami, T.: Effects of the Tibetan Plateau, in: *Monsoon Meteorol.*, edited by: Chang, C. P. and Krishnamurti, T. N., Oxford University Press, New York, 235–270, 1987.
- 15 Naftz, D. L., Susong, D. D., Schuster, P. F., Cecil, L. D., Dettinger, M. D., Michel, R. L., and Kendall, C.: Ice core evidence of rapid air temperature increases since 1960 in alpine areas of the Wind River Range, Wyoming, United States, *J. Geophys. Res.*, 107(D13), 4171, doi:10.1029/2001JD000621, 2002.
- Prospero, J. M., Blades, E., Mathison, G., and Naidu, R.: Interhemispheric transport of viable fungi and bacteria from Africa to the Caribbean with soil dust, *Aerobiol.*, 12, 1–19, 2005.
- 20 Shi, Y. F. and Liu, S. Y.: Estimation on the response of glaciers in China to the global warming in the 21st century, *Chin. Sci. Bull.*, 45, 668–672, 2000.
- Shuval, H. I., Guttman-Bass, N., Applebaum, J., and Fattal, B.: Aerosolized enteric bacteria and viruses generated by spray irrigation of wastewater, *Water Sci. Technol.*, 21, 131–135, 1989.
- 25 Story, R. M., Weber, I. T., and Steitz, T. A.: The structure of the *E. Coli* RecA protein monomer and polymer, *Nature*, 355, 318–325, 1992.
- Takeuchi, N. and Kohshima, S.: A snow algal community on a Patagonian glacier, Tyndall glacier in the southern Patagonia Icefield, *Arct. Antarct. Alp. Res.*, 36, 91–98, 2004.
- 30 Takeuchi, N. and Li, Z.: Characteristics of surface dust on Ürümqi Glacier No. 1 in the Tien Shan Mountains, China, *Arct. Antarct. Alp. Res.*, 40, 744–750, 2008.
- Takeuchi, N., Kohshima, S., and Fujita, K.: Snow algae community on a Himalayan glacier, Glacier AX010 East Nepal: Relationship with glacier summer mass balance, *Bull. Glacier*

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Res., 16, 43–50, 1998.

Takeuchi, N.: The altitudinal distribution of snow algae on an Alaska glacier (Gulkana Glacier in the Alaska Range), *Hydrol. Process.*, 15, 3447–3459, 2001.

Tamura, K., Dudley, J., Nei, M., and Kumar, S.: MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) Software Version 4.0., *Mol. Biol. Evol.*, 24, 1596–1599, 2007.

Tamura, K., Nei, M., and Kumar, S.: Prospects for inferring very large phylogenies by using the Neighbor-Joining method, *PNAS.*, 101, 11030–11035, 2004.

Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F., and Higgins, D. G.: The CLUSTAL–X Windows interface – flexible strategies for multiple sequence alignment aided by quality analysis tools, *Nucleic Acids Res.*, 24, 4876–4882, 1997.

Thompson, L. G., Mosley-Thompson, E., Davis, M. E., Bolzan, J. F., Yao, T., Gundestrup, N., Wu, X., Klein, L., and Xie, Z.: 100000 year climate record from Qinghai-Tibetan Plateau ice cores, *Science*, 246, 474–477, 1989.

Thompson, L. G., Yao, T., Mosley-Thompson, E., Davis, M. E., Henderson, K. A., and Lin P.-N.: A high-resolution millennial record of the South Asian Monsoon from Himalayan ice cores, *Science*, 289, 1916–1919. 2000.

Thompson, L. G., Mosley-Thompson, E., Davis, M., Bolzan, J., Dai, J., Yao, T., Gundestrup, N., Wu, X., Klein, L., and Xie, Z.: Glacial stage ice-core records from the subtropical Dunde ice cap, China, *Ann. Glaciol.*, 14, 288–297, 1990.

Thompson, L. G., Yao, T., Davis, M. E., Mosley-Thompson, E., Lin, P. N., Mashiotta, T. A., Mikhalenko, V. N., and Zagordonov, V. S.: Holocene climate variability archived in the Puruogangri ice cape from the central Tibetan Plateau, *Ann. Glaciol.*, 43, 61–69, 2006.

Tian, L.-D., Yao, T.-D, Li, Z., MacClune, K., Wu, G.-J., Xu, B.-Q., Li, Y. F., Lu, A.-X., and Shen, Y.-P.: Recent rapid warming trend revealed from the isotopic record in Muztagata ice core, eastern Pamirs, *J. Geophys. Res.*, 111, D13103, doi:10.1029/2005JD006249, 2006.

Vincent, W. F.: Evolutionary origins of Antarctic microbiota: invasion, selection and endemism, *Antarct. Sci.* 12, 374–385, 2000.

Wake, C. P., Mayewski, P. A., Xie, Z., Wang, P., and Li, Z.: Regional distribution of monsoon and desert dust signals recorded in asian glaciers, *Geophys. Res. Lett.*, 20, 1411–1414, 1993.

Wang, N., Thompson, L. G., Davis, M. E., Mosley-Thompson, E., Yao, T., and Pu, J.: Influence of variations in NAO and SO on air temperature over the northern Tibetan Plateau as recorded by $\delta^{18}\text{O}$ in the Malan ice core, *Geophys. Res. Lett.*, 30(22), 2167–2170, doi:10.1029/2003GL018188, 2003.

BGD

7, 1167–1199, 2010

Differences in community composition of bacteria

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Differences in
community
composition of
bacteria**L. An et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Wang, Z.-T.: Recent variations of glaciers and the influence of climate and glacier fluctuations on runoff in China, Snow cover and variations (Proceedings of the Blatimore symposium, Maryland, May), IAHS Publ., 183, 45–52, 1989.
- White, J. W. C., Barlow, L. K., Fisher, D., Grootes, P. M., Jouzel, J., Johnsen, S. J., Stuiver, M., and Clausen, H. B.: The climate signal in the stable isotopes of snow from Summit, Greenland: Results of comparisons with modern climate observations, *J. Geophys. Res.*, 102, 26425–26439, 1997.
- Xiang, S.-R., Shang, T.-C., Chen, Y., and Yao, T.-D.: Deposition and post-deposition mechanisms as possible drivers of microbial population variability in glacier ice, *FEMS Microb. Ecol.*, 70, 9–20, 2009a.
- Xiang, S.-R., Shang, T.-C., Chen, Y.,m Jing, Z.-F., and Yao, T.-D.: Changes in diversity and biomass of bacteria along a shallow snow pit from Kuytun 51 Glacier, Tianshan Mountains, China, *J. Geophys. Res. Biogeosci.*, 114, G04008, doi:10.1029/2008JG000864, 2009b.
- Xiang, S.-R., Shang, T.-C., Chen, Y., Jing, Z.-F., and Yao, T.-D.: Dominant bacteria and biomass in the Kuytun 51 Glacier, *Appl. Environ. Microbiol.*, 75, 7287–7290, 2009c.
- Xiang, S.-R., Yao, T.-D., An, L.-Z., Xu, B.-L., and Wang, J.-X.: 16S rRNA Sequences and differences in bacteria isolated from the Muztag Ata glacier at increasing depths, *Appl. Environ. Microbiol.*, 71, 4619–4627, 2005.
- Xiang, S.-R., Yao, T.-D., An, L.-Z., Xu, B.-Q., Li, Z., Wu, G.-J., Wang, Y.-Q., Ma, S., and Chen, X.-R. Bacterial diversity in Malan ice core from the Tibetan Plateau, *Folia Microbiol.*, 49, 269–276, 2004.
- Yao, T.-D., Liu, Y., Kang, S., Jiao, N., Zeng, Y., Liu, X., and Zhang, Y.: Bacteria variabilities in a Tibetan ice core and their relations with climate change, *Global Biogeochem. Cy.*, 22, GB4017, doi:10.1029/2007GB003140, 2008.
- Yao, T., Masson, V., Jouzel, J., Stievenard, M., Sun, W., and Jiao, K.: Relationship between $\delta^{18}\text{O}$ in precipitation and surface air temperature in the Urumqi River Basin, East Tianshan Mountain, China, *Geophys. Res. Lett.*, 26, 3473–3480, 1999.
- Yergeau, E., Newsham, K. K., Pearce, D. A., and Kowalchuk, G. A.: Patterns of bacterial diversity across a range of Antarctic terrestrial habitats, *Environ. Microbiol.*, 9, 2670–2682, 2007.
- Zhang, X., Ma, X., Wang, N., and Yao, T.: New subgroups of Bacteroidetes and diverse microorganisms in Tibetan plateau glacial ice provide a biological record of environmental conditions, *FEMS Microbiol. Ecol.*, 67, 21–29, 2009.

Zhang, X., Yao, D., Ma, X., and Wang, N.: Microorganisms in a High Altitude Tibet Glacier Ice, Folia Microbiol., 47, 241–245, 2002.

BGD

7, 1167–1199, 2010

**Differences in
community
composition of
bacteria**

L. An et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Differences in
community
composition of
bacteria**

L. An et al.

Table 1. Biomass and diversity of dominant bacteria in the Muztag Ata Glacier as assessed based on 16S rRNA gene sequence analysis.

Clone library (ice age, years)	MuztB 13 (1988 spring)	MuztB 16 (1984 summer–1985 winter)	MuztB 28 (1972 spring–summer)	MuztB 30 (1970 winter–1971 spring)
Total cells (10^5 cells ml^{-1})	1.23	1.95	1.99	1.03
Live cells (10^3 cells ml^{-1})	4.47	4.90	4.98	4.28
No. of OTUs predicted (S_{chao1})	55	36	76	58
Coverage C_{ACE} (%)	84	84	74	84
No. of OTUs observed	45	23	49	31
Shannon index	3.60	2.90	3.05	4.01
Evenness	0.94	0.94	0.78	1.10

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Differences in community composition of bacteria

L. An et al.




Fig. 1. Map showing the locations of four geographically isolated glaciers Muztag Ata, Puruoganri, Malan and Dunde.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Differences in community composition of bacteria

L. An et al.




Fig. 2. Relative abundance of the main bacteria phyla based on the Blast result of 16S rRNA gene sequences in each of the clone libraries in the four different ice layers along the Muztag Ata Glacier profile.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Differences in community composition of bacteria

L. An et al.




Fig. 3a. Phylogenetic analysis of the 16S rRNA genes for *Betaproteobacteria* and *Gammaproteobacteria* clones from the four ice cores and the closest relatives. The tree was generated by the neighbourjoining method after sequence alignment, and rooted with two *Methanosaela harundinacea* strains (accession no. AY817738 and DQ150255). Bootstrap values (100 replications) were specified for each Node. Numbers of the obtained snow-ice clones (had the same ARDRA pattern to the sequenced representatives listed on the tree) and relative sequence affiliations corresponding to GenBank accession number were provided in parentheses. Scale bar indicated 0.05 substitutions per nucleotide. The sequences discussed in this study were noted bold. The Muztag Ata Glacier clones were represented by MuztB13 (1988 spring), MuztB16 (1984 autumn to 1985 winter), MuztB 28 (1972 spring–summer) and MuztB 30 (1970 winter), the Puruogangri Glacier clones were noted by P60 (at 52 m depth, AD 1850–1920), P80 (62 m depth, AD 1750–1800), and P200 (89 m depth, AD 1600–1700), while the Malan Glacier clones were noted by Malan A (35 m depth, AD 1800), Malan B (64 m depth, AD 1600), Malan C (70 m depth, AD 1650), and Malan D (82 m depth, AD 1400). The typical endemic cluster was indicated in the gray shaded areas (same for the following Figs). See a detailed description for the assigned sequence references and numbers in materials and methods.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Differences in community composition of bacteria

L. An et al.




Fig. 3b. Phylogenetic analysis of the 16S rRNA genes for the *Alphaproteobacteria*, *Deltaproteobacteria*, OP10 and OP11 clones from the four geographically isolated glaciers and the closest relatives. The tree was constructed by following the protocol as described in Fig. 3a. Scale bar indicated 0.05 substitutions per nucleotide.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Differences in community composition of bacteria

L. An et al.




Fig. 3c. Phylogenetic analysis of the 16S rRNA genes for the *Bacteroidetes*, TM7, *Cyanobacteria*, *Deinococcus*, *Chloroflexi*, and *Thermus* clones from the four geographically isolated glaciers and the closest relatives. The tree was constructed by following the protocol as described in Fig. 3a. Scale bar indicated 0.05 substitutions per nucleotide.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Differences in community composition of bacteria

L. An et al.




Fig. 3d. Phylogenetic analysis of the 16S rRNA genes for the *Actinobacteria* and *Firmicutes* clones from four geographically isolated glaciers and the closest relatives. The tree was constructed by following the protocol as described in Fig. 3a. Scale bar indicated 0.05 substitutions per nucleotide.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Differences in community composition of bacteria

L. An et al.




Fig. 3e. Phylogenetic analysis of the 16S rRNA genes for the *Bacteroidetes*, *Betaproteobacteria*, *Actinobacteria*, TM7, and *Verrucomicrobia* clones (with later portion of the 16S rRNA gene sequences, corresponding to 800-1452 of the *Escherichia coli* 16S rRNA molecule) from the Puruogangri and Dunde Glaciers and the closest relatives. The tree was constructed by following the protocol as described in Fig. 3a. Scale bar indicated 0.05 substitutions per nucleotide.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Differences in community composition of bacteria

L. An et al.




Fig. 4. Proportion of the main phylogenetic clusters (genera) based on the Blast result of 16S rRNA gene sequences in each of the clone libraries in the different ice layers along the MuztB ice core profile. *Alph*, *Beta*, and *Gamma*=*Alpha*-, *Beta*-, and *Gamma*-proteobacteria.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Differences in community composition of bacteria

L. An et al.




Fig. 5. Hierarchical cluster showing the overall phylogenetic distances amongst the clone libraries from four geographically isolated glaciers Puruogangri (two clone libraries containing the forward portion of 16S rRNA gene sequence corresponding to 1 to ~800 of *Esherichia coli* 16S rRNA molecule, with sequence accession number AY313907–AY313917, DQ076421–DQ076430, and DQ076441–DQ076444, Zhang et al., 2009), Dunde (with sequence accession number AY313918, AY313919 and DQ076445–DQ076456, Zhang et al., 2009), Malan (two clone libraries containing >7 unique clones with sequence accession number AY322483–AY322489 in Xiang et al., 2004 and AY121823–AY121830 in Zhang et al., 2002) and Muztag Ata (this study). Distances were estimated with the weighted Unifrac algorithm (Lozupone and Knight, 2005). A sequence jackknifing technique was applied to each cluster to determine the sensitivity of the relationships to sample size. UniFrac distance indicated difference amongst the bacterial communities.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

