Biogeosciences Discuss., 7, 1675–1706, 2010 www.biogeosciences-discuss.net/7/1675/2010/ © Author(s) 2010. This work is distributed under the Creative Commons Attribution 3.0 License.



This discussion paper is/has been under review for the journal Biogeosciences (BG). Please refer to the corresponding final paper in BG if available.

Quantifying nitrous oxide emissions from Chinese grasslands with a process-based model

F. Zhang 1,2 , J. Qi 2,1 , F. M. Li 1 , C. S. Li 3 , and C. B. Li 2,4

¹Key Laboratory of Arid and Grassland Agro-Ecology (MOE), Lanzhou University, Lanzhou, Gansu, 730000, China

²Center for Global Change and Earth Observations, Michigan State University, East Lansing, MI, 48823, USA

³Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, NH 03824, USA

⁴Key Laboratory of Western China's Environmental Systems (MOE), Lanzhou University, Lanzhou, Gansu, 730000, China

Received: 2 January 2010 - Accepted: 5 February 2010 - Published: 5 March 2010

Correspondence to: F. M. Li (fmli@lzu.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

1675

Abstract

As one of the largest land cover types, grassland can potentially play an important role in the ecosystem services of natural resources in China. Nitrous oxide (N₂O) is a major greenhouse gas emitted from grasslands. Current N₂O inventory at regional or

- ⁵ national level in China relies on the emission factor method, and is based on limited measurements. To improve inventory accuracy and capture the spatial variability of the N₂O emissions under the diverse climate, soil and management conditions across China, we adopted an approach that uses a process-based biogeochemical model, DeNitrification-DeComposition (DNDC) in this study, to map the N₂O emissions from
- China's grasslands. The DNDC was linked to a GIS database of spatially distributed information of climate, soil, vegetation and management at county-level for all grass-lands in China. Daily weather data from 2000–2007 based on the national network of 670 meteorological stations were utilized in the model simulations. The results were validated against observations from several grasslands in China and from other coun-
- tries. The modelled results showed a clear geographic pattern of N₂O emissions from China's grasslands. A high-emission strip was found that stretched from northeast to central China, along the eastern boundary of the temperate grassland region adjacent to the major agricultural regions. The grasslands in the western mountain regions, however, emitted much less N₂O. The regional average of N₂O emission rates was 0.23,
- ²⁰ 0.11 and 0.39 kg N ha⁻¹ y⁻¹ for the temperate, montane and tropical/subtropical grasslands, respectively. The national N₂O emission was 76.5 Gg N from the 337 million ha of grasslands in China. The modelled results were in good agreement with observations ($R^2 = 0.64$ for 11 datasets), suggesting that the process-based model can be used to capture the spatial dynamics of N₂O emissions as an effective alternative to
- ²⁵ statistical method currently used in China.

1 Introduction

According to the Intergovernmental Panel on Climate Change (IPCC), the global average surface temperature has increased by around 0.74 ± 0.18 °C during the 20th century, and global atmospheric concentrations of greenhouse gases (GHGs) have in-

- ⁵ creased markedly as a result of human activities since 1750 (IPCC, 2007). Increased GHG emissions have altered the energy balance of the climate system and are one of the major causes of climate change (IPCC, 2007). On a global basis, carbon dioxide (CO₂), methane (CH₄) and nitrous oxide (N₂O), contribute 60, 15 and 5%, respectively, to the anthropogenic GHGs' effect (Rodhe, 1990). The instantaneous flux rates of the
- ¹⁰ GHGs were important clues to lead an accurate assessment of the terrestrial ecosystem contributions to climate change (Du et al., 2008). N₂O is a major contributor to global warming because of its long atmospheric lifetime and large radiative forcing potential (Cicerone, 1989; Mummey et al., 2000; IPCC, 2007); however, large uncertainty exists about the global N₂O budget (Bouwman et al., 2000; Chapuis-Lardy et al., 2007).
- ¹⁵ Grasslands, where herbaceous plants form the dominant climax community (Coupland, 1992; Mummey et al., 2000), play a significant role in the regional climate and global carbon cycle both in tropical and temperate regions (Scurlock and Hall, 1998). However, compared with forest ecosystems, little attention has been paid to N₂O exchange in grassland ecosystems, despite the fact that grassland covers about one-fifth of the
- world's land surface and about 40% of China's national land cover (Allard et al., 2007; Kang et al., 2007). It is important, therefore, to quantify N₂O emissions from grass-lands for more accurate assessment of the N₂O budget (Mummey et al., 2000) and to gain a better understanding of the potential N₂O production and emission processes in grassland ecosystems for future climate change mitigation (Guo and Zhou, 2007).
- The IPCC method was frequently used to estimate N₂O emissions although less capable of quantifying impacts of climate or management on the gas fluxes. Consequently process-oriented models were proposed to improve N₂O emission estimates (Boeckx and Van Cleemput, 2001). At a national scale, process-based models have

1677

proven useful in reducing uncertainty and helping understand the complex biogeochemical processes involved in trace gas production (Barnsley, 2007). Spatially distributed process-oriented models have also been effective in achieving this objective (Chen et al., 2008), which is the focus of this paper.

- Grassland is widely distributed in north and northwest China (Kang et al., 2007), and is under threat of serious degradation and over grazing. Increasing numbers of scholars have advocated changing grassland management and adopting intensive management (Nan, 2005). Grassland management practices have recently undergone some significant changes over the past decades because of economic development (Chen
- and Chen, 2007). To slow down degradation and maintain sustainability, some grassland laws have been introduced and some management practices were promoted by the Chinese central government (Chen and Chen, 2007; Unkovich and Nan, 2008). These include converting reclaimed land to pasture lands, practicing rotational grazing method using fencing; promoting conservation practices; tending and controlling ro-
- dents and insect pests; promoting rangeland improvements such as shallow-ploughing, fertilization and irrigation; and promoting restorations such as aerial-sowing, cultivation practices, converting cultivated lands to pastures, and establishment of a forage base (Yang, 1992; Chen and Chen, 2007). Among of them, fertilization and improved grazing management practices are the most adopted options (Conant et al., 2001). These
- ²⁰ management options tend to have the potential to change the spatial distribution of N₂O emissions, thus making the national estimation of N₂O difficult. Therefore, the preliminary researches indicated that the magnitudes of N₂O fluxes from Chinese grassland varied greatly in space and time (Du et al., 1997, 2006; Dong et al., 2000; Xu et al., 2003). There is a need to use spatially explicit process-based models to account for the
- ²⁵ dynamic nature of grasslands in China in order to arrive at a more accurate estimate of the N₂O emissions at the national level.

The objective of this study was to estimate N_2O emissions and their spatial distributions across the grasslands in China using a spatially explicit process-based DNDC model, with a goal of improving the national estimate of N_2O emissions. We adopted a

four-phase method that includes (1) database development, (2) scaling up, (3) model validation and (4) sensitivity analysis.

2 Method

2.1 Database development

- To apply DNDC at a national scale, we developed a GIS database to store the spatially distributed information of climate, soil and grassland types. Counties were used as the basic unit for the simulation. The database contained 2368 counties, excluding those with small areas less than 500 square kilometres. The database consisted of: (1) 18 grassland types with growth properties (e.g. maximal production, maximal height, and
- root/shoot ratio); (2) soil properties (e.g. maximum and minimum soil surface organic carbon content, bulk density, clay fraction and pH); (3) daily climate data (e.g. maximum and minimum air temperatures, and precipitation); and (4) areas and geographic locations of grassland types at county level. These data were obtained from the Commission for the Integrated Survey of Natural Resources; the Institute of Soil Science,
- ¹⁵ Chinese Academy of Sciences; and the Chinese Meteorological Administration.

2.1.1 Grassland dataset

There is a large uncertainty concerning the total grassland area in China. Fang et al. (1996) estimated 569.9 million ha based on agricultural atlases and land use maps of the 1980s. Ni (2001) used 405.9 million ha in his report, and Fan et al. (2008) estimated 331 million ha of grassland area. Because of the large discrepancies, in

estimated 331 million ha of grassland area. Because of the large discrepancies, in this study, we digitized the 1:1 000 000 grassland resource maps (Commission for Integrated Survey of Natural Resources, 1995), and classified all grasslands 18 vegetation types, according to Ni (2002). Considering the extensive changes in land use over the last decade in China (Lin and Ho, 2003), we applied the 1:100 000 National Land Cover

1679

Dataset (NLCD) of China acquired in 1999 and 2000 to modify the grassland area by retaining only those areas identified as grassland in the NLCD. The final grassland area of our new database was 336.98 million ha (Fig. 1).

2.1.2 Soil dataset

- ⁵ We used a 1:1 000 000 scale soil database developed by the Institute of Soil Science, Chinese Academy of Sciences, which was compiled based on the second national soil survey conducted in 1979–1994 covering all the counties. The database contains three attributes: locations, attributes, and reference systems. It contains multi-layer soil properties (e.g. organic matter, pH and bulk density), soil texture (sand, silt and clay) and spatial information (Shi et al., 2004; Yu et al., 2007). In this study, we used data in
- the upper 0–10 cm soil profile as the soil surface properties for model simulations.

2.1.3 Climate dataset

We used 10 km-resolution daily national climate data from 2000–2007, which were interpolated from 670 meteorological stations using Ordinary Kriging method. The climate data included daily precipitation, maximum and minimum temperatures (http:

//data.cma.gov.cn/).

2.1.4 Grazing data

Based on the statistical data in 2000 at the national scale from the National Bureau of Statistics of China (NBSC, http://www.stats.gov.cn/english/statisticaldata/), we as-

²⁰ sumed that the grasslands in China experienced a similar grazing practice, and livestock was evenly distributed among all grasslands. The average national grazing stock value was applied (total livestock divided by total area). Specifically, the grazing stocks were 0.32, 0.06 and 0.72 head/ha of cattle, horses and sheep, respectively, and no fertilizer application was assumed.

2.2 Model upscaling

To estimate the national N_2O emissions from grasslands in China we linked DNDC to the database containing climate data (2000–2007), soil properties and grassland types and total grassland area. Because SOC content was one of the most sensitive factors

⁵ affecting N₂O fluxes, we ran the DNDC model for each grid-cell with the maximum and minimum SOC content values to quantify the uncertainty in the upscaling (see details in Li (2007). Then, the eight years of N₂O emission average rates were reported and compared with field and previously published results.

2.3 Model validation

- ¹⁰ DNDC is a process-based model of carbon and nitrogen biogeochemistry for terrestrial ecosystems. The model was originally developed for estimating N₂O emissions from agricultural ecosystems (Li, 1992a, b; Li et al., 2004). Detailed processes of nitrification and denitrification were built in the model to track the kinetic processes of N₂O production and consumption driven by the soil climate and substrate concentrations
- (Li, 2007). DNDC consists of two main components. The first component consists of soil climate, crop growth and decomposition sub-models converts primary drivers such as climate, soil, vegetation and human activities into soil environmental factors (e.g. soil temperature, humidity, pH, Eh and substrate concentration gradients). The second component of DNDC includes nitrification, denitrification and fermentation sub-
- ²⁰ models calculates N₂O and CH₄ production and consumption (Li, 2000; Li et al., 2000). The DNDC model has been successfully tested and applied for N₂O inventory in many countries across climatic zones, soil types and management regimes (Frolking et al., 1998; Li, 2000; Li et al., 2000, 2001, 2005; Saggar et al., 2004; Beheydt et al., 2007; Saggar et al., 2007), and is recognized as one of the most widely used N₂O models
- in the world (Chen et al., 2008). DNDC has been also applied in China for estimating N₂O emissions from grassland at the site-scale (Xu et al., 2003).

1681

We further tested DNDC for its applicability to grasslands in China. Ten grassland sites, including eight in China and two in the United States, where N_2O fluxes were measured and the annual emission rates were reported in publications (Table 1) were used for validation tests. The sites were natural grasslands as defined by Coupland

- $_5$ (1992) without any fertilizer applied. The modelled annual N_2O emissions rates were consistent with reports for the sites (Fig. 2), except for the alpine meadow (Du et al., 2008) for which the modelled N_2O flux was only 1/10 of observation. However, Du's observed values were higher than that reported by other researchers for montane grasslands in China (Pei et al., 2003, 2004) or other countries (Mosier et al., 1993), and was
- ¹⁰ also higher than that from fertilized grassland in Europe (Levy et al., 2007). The high N₂O emissions from the site of Du et al. (2008) could be related to its high soil moisture and high emissions from plants.

2.4 Sensitivity analysis

- The sensitivity analysis was conducted by varying a single model input parameter in a predefined range, which was commonly observed in a unit of study, while keeping all other input parameters constant. Simulated annual N₂O flux sensitivities were evaluated for soil factors (i.e. pH, soil texture and soil organic C content), grazing intensity, and climate (i.e. precipitation and temperature). The baseline simulation was done using the average values of the entire grassland region (Table 2). The sensitivity tests
- were done to assess the model responses to soil attributes, management options, and climate variables

3 Results and Discussion

3.1 Model sensitivities to environmental variables

Table 2 lists the parameters in the sensitivity analysis. It is clear that the model responded to changes in both environmental factors as well as to human management options. This suggests that the model can be used to assess how climate change and human management practices affect N_2O emissions from the vast grasslands in China.

3.2 Sensitivity to soil attributes

The modelled N₂O emission rates were well correlated with the initial soil organic carbon (SOC) contents (Fig. 3c), which is similar to what was reported by Babu et al. (2006). Variation of initial SOC content showed significant effects on production and emission of N₂O. More N₂O gas was produced at pH 7 than from alkaline soils (Fig. 3a). Soil texture was also a sensitive factor due to its effects on soil aeration status; sandy loam soil was more likely to produce N₂O than clay soil (Fig. 3b).

10 3.3 Sensitivity to management options

The modelled N_2O emission was very sensitive to grazing management options, such as grazing intensity and duration. Enhanced grazing intensity increased N_2O emissions (Fig. 3d, e). The result was in agreement with that reported by Flechard et al. (2007). Nitrogen deficit is one of the major factors limiting grass growth in most

¹⁵ Chinese grasslands (Nan, 2005; Unkovich and Nan, 2008). When fertilizer application rates increased from 0 to 45 kg N ha^{-1} , N₂O emissions increased from 0.1 to $0.25 \text{ kg N ha}^{-1} \text{ y}^{-1}$ (Fig. 3f).

3.4 Sensitivity to climate conditions

The N₂O emissions were susceptible to changes in rainfall and temperature, which is in agreement with previous studies that have shown the N₂O emissions were positively correlated with air or soil temperature (Yamulki et al., 1997; Dong et al., 2000). In this study, precipitation was either increased or decreased by 20% of the baseline value (520 mm y⁻¹). Increases in precipitation elevated N₂O emissions (Fig. 3g). Modelled data showed that the high precipitation stimulated denitrification, which is a major pro-

 $_{\rm 25}$ cess producing $\rm N_2O$ and embedded in DNDC. The effect of temperature was examined

1683

by running a sequence of simulations with daily temperatures at 1 or 2°C increment; N₂O emissions did not follow a simple relationship with change in temperature. With an increase of 1°C in daily temperature, the N₂O emissions did not change; but when the increase was 2°C, the total N₂O emission changed significantly (Fig. 3h). When

temperature decreased, N₂O emissions also decreased clearly, since a decrease in temperature decreased the rate of organic matter decomposition and therefore decreased nitrification, which is an important source of N₂O (Li et al., 2000).

3.5 National inventory of N₂O emissions from China's grasslands

Based on the modelled annual N₂O-fluxes for the eight years (Table 3), a multi-year average N₂O emission value was calculated for each grid-cell as well as for all grasslands in China. The mean annual N₂O emission rates from all the grasslands were < $0.5 \text{ kg N ha}^{-1} \text{ y}^{-1}$ with the highest for swamp grassland and the lowest for Alpine desert grassland (0.48 and $0.02 \text{ kg N ha}^{-1} \text{ y}^{-1}$, respectively) and the national average 0.22 kg N ha⁻¹ y⁻¹. National inventory of N₂O emissions from Chinese grassland was calculated based on the area and N₂O emission rates for each type of grassland (Table 3).

The results showed that the 337 million ha of grasslands in China emitted 76.5 Gg N_2O . N y⁻¹ on a multi–year average basis.

The modelled N₂O fluxes were analyzed by combining the 18 types of grassland into three categories based on ecological climate zone. The three categories were temperate grasslands, montane grasslands and tropical/subtropical grasslands (Table 4, Fig. 4). Each of the three categories possessed specific features and was basically consistent with climate patterns. The greatest N₂O flux (0.39 kg N ha⁻¹ y⁻¹) occurred in the tropical/subtropical grassland region, in which both moisture and temperature were more favourable for N₂O production. In the montane grassland region, the av-

erage N₂O flux was the lowest at 0.11 kg N ha⁻¹ y⁻¹; this region is humid but with low temperature. The temperate grassland region had a medium emission flux, averaging 0.23 kg N ha⁻¹ y⁻¹. Even though the temperate grassland's N₂O flux rate was in

the mid-range, it comprised 49% of the 337 million ha of total grassland area in China (Table 4), and thus this category's contribution is large, accounting for 57% of national emissions. Despite the large amount of montane grasslands (39% of the total grassland area in China), due to the lower N₂O emission rate, it only contributed 24% of the total N₂O flux. The tropical/subtropical grasslands occupy only 12% of total grassland

areas, but account for 19% of the total N₂O flux.

3.6 Spatial and temporal distribution of N₂O emissions

The spatial distribution of the grassland N_2O fluxes was basically consistent with the climate patterns. In China, mean annual precipitation and temperature tend to decrease

- from east to west and from south to north. N_2O flux rates also followed this climate change trend, gradually decreasing from eastern to western regions. In the western high-altitude region, the N_2O flux rates were lower than in the eastern and southern region, especially in eastern Inner Mongolia and the sparsely distributed grasslands of the south, where the N_2O flux (Fig. 5) was higher.
- ¹⁵ On the national scale, during 2000–2007 the annual N₂O emission rate fluctuated year-to-year, and had an increasing trend (Fig. 6). In 2001 and 2005, the modelled national N₂O emissions were higher than in other years. In 2005, the total N₂O had an abrupt emission peak compared with other years. We compared the accumulated annual temperature >0 °C and the annual accumulated precipitation between 2005 and
- ²⁰ 2001 (Fig. 7a, b). These comparisons suggested that climate change seems to be the main reason for increased N₂O emissions. In the northern grassland region, the precipitation and the temperature significantly increased (Fig. 7a, b), which is believed to have caused the N₂O emissions increase. Another reason was that temperate grassland had a higher base emission rate than other types of grassland.

1685

3.7 Comparison of N₂O with worldwide grassland

In China, most grasslands are located in semi-arid and arid areas, which are classified as ecologically fragile zones (Liu, 2001). The poor soils and low precipitation could limit N_2O emissions due to the depressed nitrification and denitrification processes. However, some grasslands such as meadow and swamp grasslands have relatively high humidity and the modelled N_2O fluxes were relatively high in these areas. To estimate the reliability of our modelled results, we compared our modelled N_2O fluxes with those reported for other parts of the world, based on the similarity in grassland types.

10 3.7.1 Comparison with North American grasslands

The temperate grassland in Inner Mongolia was similar to the North Great Plains grassland in the US. The two grasslands are located at similar latitudes. Mummey et al. (2000) reported 0.24 kg N ha⁻¹ y⁻¹ emitted from Great Plains grassland, and Mosier et al. (1996) reported 0.1–0.2 kg N ha⁻¹ y⁻¹ based on long-term experiments in Col-

orado. These values were very similar to our $0.23 \text{ kg N ha}^{-1} \text{ y}^{-1}$ for temperate grassland in China. For the subalpine meadow experiment site in the US, values were 0.11 and 0.22 kg N ha⁻¹ y⁻¹ for 1991 and 1992, respectively (Mosier et al., 1993), for China's similar montane grasslands, this value was 0.11 kg N ha⁻¹ y⁻¹. The comparisons indicate that the DNDC estimated values were within reasonable ranges.

20 3.7.2 Comparison with European and New Zealand grasslands

About 40% of the agricultural land in Europe is grassland. Some grasslands are tilled and reseeded to support productive grass species (Pinto et al., 2004). Flechard et al. (2007) reported a mean emission of $0.93 \text{ kg N ha}^{-1} \text{ y}^{-1}$, higher than the 0.23 kg N ha⁻¹ y⁻¹ we modelled for China's temperate grassland. However, Flechard et al. (2007) pointed out that intensively managed systems emitted more N₂O than

the extensive management grasslands in Europe. Allard et al. (2007) also reported the extensive grassland (similar to natural grassland) had a lower N₂O emission rate (average 0.13 kg N ha⁻¹ y⁻¹) which is close to results from China's grasslands.

Fertilizer use is common for European semi-natural grassland as well as grassland in
 New Zealand that leads to high N₂O emissions (Boeckx and Van Cleemput, 2001; Saggar et al., 2004, 2007; Levy et al., 2007). In China, grasslands are sparsely fertilized or irrigated (Huang et al., 2003; Nan, 2005; Unkovich and Nan, 2008).

3.7.3 Comparison with African grassland

Savannahs are the most widespread vegetation in Africa (White, 1983; Rees et al., 2006), and the grass species in savannahs are quite different from those in temperate grasslands (http://www.bcgrasslands.org/library/world.htm). Brummer et al. (2008) found that in an African savannah natural-reserve site (southwest of Burkina Faso), the N₂O emission rate was 0.52 and 0.67 kg N ha⁻¹ y⁻¹ in 2005 and 2006, respectively. For

Zimbabwean savannah, the measured N₂O flux was 0.25–0.5 kg N ha⁻¹ y⁻¹ (Rees et al., 2006). In China, there are no real savannahs as found in Africa, and the most similar vegetation type is Chinese tropical/subtropical grassland. Our estimate for these grasslands was 0.39 kg N ha⁻¹ y⁻¹, within the range of values reported in Africa.

3.7.4 Global comparison

Prentice et al. (1993) estimated that the present total area of global grassland is 4.16 ²⁰ billion ha, and China has 8% of this. If we apply the modelled average N₂O emission rate (0.22 kg N ha⁻¹ y⁻¹) based on this ratio, the global grassland N₂O emission would be 0.92 Tg N y⁻¹. Globally, the total anthropogenic emission of N₂O is estimated at 6.3– 6.7 Tg N y⁻¹ (Khalil and Rasmussen, 1992; IPCC, 2000). Of that, grassland accounts for 14% of annual total anthropogenic N₂O–N emission. This may underestimate the

²⁵ grassland N₂O emission contribution to the atmosphere, as most grasslands in China have relatively low emission rates compared to semi-natural grasslands.

1687

In China, grassland has a long history of anthropogenic use, and is currently experiencing serious degradation. Moderating the degradation and improving the grassland conditions are urgent tasks for sustainable development in China. New policies have been proposed to encourage farmers and herdsmen to maintain high productivity of

- ⁵ grassland by converting farmland to grasslands or adopting grassland–farming rotations system (Nan, 2005). Specifically, intensive management practices have been advocated, which can improve the grassland and increase carbon sequestration (Conant et al., 2001), including fertilizer application and mowing (Committee on Scholarly Communication with the People's Republic of China and National Research Council
- ¹⁰ 1992). However, such practices can also change the N₂O flux. More research is needed to determine optimal management strategies that achieve a balance between N₂O emission and carbon-sequestration. The present study was an attempt to apply process-based models such as DNDC for assessing N₂O inventory and mitigation potentials at a national scale.

15 4 Conclusions

The N₂O emissions from various grassland types in China differed. The emissions from temperate, montane, and tropical/subtropical grasslands were 0.23, 0.11 and 0.39 kg N ha⁻¹ y⁻¹, respectively. Of the 337 million ha of grasslands nationwide in China, annual N₂O emissions were 76.5 Gg N. The N₂O emissions from the entire

20 grassland ecosystem varied year-by-year and have increased constantly from 2000–2007, with climate change playing the main role in this process. Grasslands in China have been intensively developed recently, thus future emission estimates from grasslands may need to include more specific grassland management practices, land use change and spatially distributed grazing rates such as fertilizer application rates.

Acknowledgements. This research was funded by "973" Program (2007CB106804); MAIRS and China projects, which funded by the NASA (NNX08AH50G, NNG05GD49G) and by NSF (BCS-0624018). Additional supports were provided by MAES project funded through Department of Geography at Michigan State University, "111" Program and Innovative Team Program

of Ministry of Education. 5

References

10

25

25

- Allard, V., Soussana, J. F., Falcimagne, R., Berbigier, P., Bonnefond, J. M., Ceschia, E., D'Hour, P., Henault, C., Laville, P., Martin, C., and Pinares-Patino, C.: The role of grazing management for the net biome productivity and greenhouse gas budget (CO₂, N₂O and CH₄) of semi-natural grassland, Agr. Ecosyst. Environ., 121, 47–58, doi:10.1016/j.agee.2006.12.004,
- 2007. Babu, Y., Li, C., Frolking, S., Nayak, D., and Adhya, T.: Field Validation of DNDC Model for Methane and Nitrous Oxide Emissions from Rice-based Production Systems of India, Nutr.
- Cvcl. Agroecosvs. 74, 157-174, 2006. Barnsley, M. J.: Environmental Modeling: A Practical Introduction, Taylor & Francis, Inc, 2007. Beheydt, D., Boeckx, P., Sleutel, S., Li, C., and Van Cleemput, O.: Validation of DNDC for 22 long-term N₂O field emission measurements, Atmos. Environ., 41, 6196-6211, 2007.
- Boeckx, P. and Van Cleemput, O.: Estimates of N₂O and CH₄ fluxes from agricultural lands in various regions in Europe, Nutr. Cycl. Agroecosys., 60, 35-47, 2001.
- Bouwman, A. F., Taylor, J. A., and Kroeze, C.: Testing hypotheses on global emissions of nitrous oxide using atmospheric models, Chemosphere Global Change Sci., 2, 475-492, 2000.
 - Brummer, C., Bruggemann, N., Butterbach-Bahl, K., Falk, U., Szarzynski, J., Vielhauer, K., Wassmann, R., and Papen, H.: Soil-atmosphere exchange of N₂O and NO in near-natural savanna and agricultural land in Burkina Faso (W. Africa), Ecosystems, 11, 582-600, doi:10.1007/s10021-008-9144-1, 2008.
- Chapuis-Lardy, L., Wrage, N., Metay, A., Chotte, J.-L., and Bernoux, M.: Soils, a sink for N₂O? A review, Global. Change. Biol., 13, 1–17, doi:10.1111/j.1365-2486.2006.01280.x, 2007.
 - Chen, B. and Chen, G. Q.: Resource analysis of the Chinese society 1980-2002 based on exergy-Part 4: Fishery and rangeland, Energ. Policy, 35, 2079-2086, 2007.

1689

- Chen, D., Li, Y., Grace, P., and Mosier, A.: N₂O emissions from agricultural lands: a synthesis of simulation approaches, Plant Soil, 309, 169-189, 2008.
- Chen, G. X., Huang, B., Xu, H., Zhang, Y., Huang, G. H., Yu, K. W., Hou, A. X., Du, R., Han, S. J., and VanCleemput, O.: Nitrous oxide emissions from terrestrial ecosystems in China, Chemosphere Global Change Sci., 2, 373-378, 2000.
- Cicerone, R. J.: Analysis of Sources and Sinks of Atmospheric Nitrous Oxide (N₂O), J. Geophys. Res. Atmos., 94, 18265-18271, 1989.
- Commission for Integrated Survey of Natural Resources, C. A. o. S.: Atlas of Grassland Resources of China (1:1 million), Chinese Academy of Science, Beijing, 10-75, 1995.
- Conant, R. T., Paustian, K., and Elliott, E. T.: Grassland management and conversion into grassland: Effects on soil carbon, Ecol. Appl., 11, 343-355, 2001.
- Council, C. o. S. C. w. t. P. s. R. o. C. N. R.: Grasslands and Grassland Sciences in Northern China : A Report of the Committee On Scholarly Communication With the People's Republic of China, National Academies Press, Washington, D.C., 1992.
- Coupland, R. T.: Natural Grasslands: Introduction and Western Hemisphere, Ecosystems of the World, edited by: Coupland, R. T., Elsevier, New York, 1992.
 - Dong, Y. S., Zhang, S., Qi, Y. C., Chen, Z. Z., and Geng, Y. B.: Fluxes of CO₂, N₂O and CH₄ from a typical temperate grassland in Inner Mongolia and its daily variation, Chin. Sci. Bull., 45, 1590-1594, 2000.
- Du, R., Chen, G., Lu, D., and Wang, G.: The Primary Research on in situ Measurements of N₂O and CH₄ Fluxes from the Inner Mongolia Grassland Ecosystem, Climatic and Environmental Ressarch, 2, 264-272, 1997.
 - Du, R., Lu, D., and Wang, G.: Diurnal, seasonal, and inter-annual variations of N₂O fluxes from native semi-arid grassland soils of inner Mongolia, Soil Biol. Biochem., 38, 3474–3482, 2006.
- Du, Y., Cui, Y., Xu, X., Liang, D., Long, R., and Cao, G.: Nitrous oxide emissions from two alpine meadows in the Qinghai-Tibetan Plateau, Plant Soil, 311, 245-254, 2008.
- Fan, J., Zhong, H., Harris, W., Yu, G., Wang, S., Hu, Z., and Yue, Y.: Carbon storage in the grasslands of China based on field measurements of above- and below-ground biomass, Clim. Change, 86, 375-396, 2008.
- Fang, J., Liu, G., and Xu, S.: Carbon pools in terrestrial ecosystems in China, edited by: Wang, R. S., Fang, J. Y., Gao, L., and Feng, Z. W.: Hot spots in modern ecology, China Sciences and Technology, Beijing, 1996.

- Flechard, C. R., Ambus, P., Skiba, U., Rees, R. M., Hensen, A., van Amstel, A., Dasselaar, A. v. d. P.-v., Soussana, J. F., Jones, M., Clifton-Brown, J., Raschi, A., Horvath, L., Neftel, A., Jocher, M., Ammann, C., Leifeld, J., Fuhrer, J., Calanca, P., Thalman, E., Pilegaard, K., Di Marco, C., Campbell, C., Nemitz, E., Hargreaves, K. J., Levy, P. E., Ball, B. C., Jones, S. K.,
- van de Bulk, W. C. M., Groot, T., Blom, M., Domingues, R., Kasper, G., Allard, V., Ceschia, E., Cellier, P., Laville, P., Henault, C., Bizouard, F., Abdalla, M., Williams, M., Baronti, S., Berretti, F., and Grosz, B.: Effects of climate and management intensity on nitrous oxide emissions in grassland systems across Europe, Agric. Ecosyst. Environ., 121, 135–152, 2007.
- Frolking, S. E., Mosier, A. R., Ojima, D. S., Li, C., Parton, W. J., Potter, C. S., Priesack, E., Stenger, R., Haberbosch, C., Dörsch, P., Flessa, H., and Smith, K. A.: Comparison of N₂O
- emissions from soils at three temperate agricultural sites: simulations of year-round measurements by four models, Nutr. Cycl. Agroecosys, 52, 77–105, 1998.
- Guo, J. and Zhou, C.: Greenhouse gas emissions and mitigation measures in Chinese agroecosystems, Agr. Forest. Meteorol., 142, 270–277, 2007.
- Huang, B., Chen, G., Huang, G., and Hauro, T.: Nitrous oxide emission from temperate meadow grassland and emission estimation for temperate grassland of China, Nutr. Cycl. Agroecosys, 67, 31–36, 2003.
 - IPCC: Emissions Scenarios, A Special Report of IPCC Working Group III, Intergovernmental Panel on Climate Change, 3–20, 2000.
- IPCC: Climate change 2007: synthesis report, IPCC Fourth Assessment Report, Intergovernmental Panel on Climate Change, 30–40, 2007.
 - Kang, L., Han, X., Zhang, Z., and Sun, O. J.: Grassland ecosystems in China: review of current knowledge and research advancement, Philosophical Transactions of the Royal Society B: Biological Sciences, 362, 997–1008, 2007.
- ²⁵ Khalil, M. A. K. and Rasmussen, R. A.: The Global Sources of Nitrous-Oxide, J. Geophys. Res. Atmos., 97, 14651–14660, 1992.
 - Levy, P. E., Mobbs, D. C., Jones, S. K., Milne, R., Campbell, C., and Sutton, M. A.: Simulation of fluxes of greenhouse gases from European grasslands using the DNDC model, Agric. Ecosyst. Environ., 121, 186–192, 2007.
- Li, C., Zhuang, Y., Cao, M., Crill, P., Dai, Z., Frolking, S., Moore, B., Salas, W., Song, W., and Wang, X.: Comparing a process-based agro-ecosystem model to the IPCC methodology for developing a national inventory of N₂O emissions from arable lands in China, Nutr. Cycl. Agroecosys, 60, 159–175, 2001.

1691

- Li, C., Cui, J., Sun, G., and Trettin, C.: Modeling Impacts of Management on Carbon Sequestration and Trace Gas Emissions in Forested Wetland Ecosystems, Environ. Manage., 33, S176–S186, 2004.
- Li, C.: Quantifying greenhouse gas emissions from soils : Scientific basis and modeling approach(<Special Section>Greenhouse Gas Emissions from Agroecosystems in Monsoon Asia), Soil Sci. Plant Nutr., 53, 344–352, 2007.
 - Li, C. S.: A model of nitrous-oxide evolution from soil driven by rainfall events, 1. model structure and sensitivity, J. Geophys. Res. Atmos., 97, 9759–9776, 1992a.
- Li, C. S.: A model of nitrous-oxide evolution from soil driven by rainfall events, 2. model applications, J. Geophys. Res. Atmos., 97, 9777–9783, 1992b.
- Li, C. S.: Modeling Trace Gas Emissions from Agricultural Ecosystems, Nutr. Cycl. Agroecosys, 58, 259–276, 2000.
- Li, C. S., Aber, J., Stange, F., Butterbach-Bahl, K., and Papen, H.: A process-oriented model of N₂O and NO emissions from forest soils: 1. Model development, J. Geophys. Res. Atmos., 105, 4369–4384, 2000.
- Li, C. S., Frolking, S., Xiao, X. M., Moore, B., III, Boles, S., Qiu, J., Huang, Y., Salas, W., and Sass, R.: Modeling impacts of farming management alternatives on CO₂, CH₄, and N₂O emissions: a case study for water management of rice agriculture of China, Global. Biogeochem. Cy, 19, GB3010, doi:10.1029/2004gb002341, 2005.

15

- Lin, G. C. S. and Ho, S. P. S.: China's land resources and land-use change: insights from the 1996 land survey, Land Use Policy, 20, 87–107, 2003.
 - Liu, J.: Integrated Ecosystem Assessment of Western China, Millennium Ecosystem Assessment., Institute of Geographical Sciences and Natural Resources Research, CAS; Research Institute of Forest Resource Information Techniques, CAF; Information Cen-
- ter, SEPA;Research Center for Eco-Environmental Sciences, CAS;Xishuangbanna Tropical Botanical Garden, CAS; Institute of Mountain Hazards and Environment, CAS; Xinjiang Institute of Ecology and Geography, CAS; Cold and Arid Regions Environmental and Engineering Research, CAS; National Institute for Environmental Studies of Japan., 52, 2001.
- Mosier, A. R., Klemedtsson, L. K., Sommerfeld, R. A., and Musselman, R. C.: Methane and nitrous-oxide flux in a Wyoming sub-alpine meadow, Global. Biogeochem. Cy, 7, 771–784, 1993.
 - Mosier, A. R., Parton, W. J., Valentine, D. W., Ojima, D. S., Schimel, D. S., and Delgado, J. A.: CH₄ and N₂O Fluxes in the Colorado Shortgrass Steppe: 1. Impact of Landscape and

Nitrogen Addition, Global. Biogeochem. Cy, 10, 387-399, 1996.

5

- Mummey, D. L., Smith, J. L., and Bluhm, G.: Estimation of Nitrous Oxide Emissions from US Grasslands, Environ. Manage., 25, 169–175, 2000.
- Nan, Z.: The grassland farming system and sustainable agricultural development in China, Grassland Science, 51, 15–19, 2005.
- Ni, J.: Carbon Storage in Terrestrial Ecosystems of China: Estimates at Different Spatial Resolutions and Their Responses to Climate Change, Clim. Change, 49, 339–358, 2001.
 Ni, J.: Carbon storage in grasslands of China, J. Arid Environ., 50, 205–218, 2002.
- Pei, Z., Ouyang, H., Zhou, C., and Xu, X.: Fluxes of CO₂, CH₄ and N₂O from alpine grassland in the Tibetan Plateau, J. Geog. Sci., 13, 27-34, 2003.
 - Pei, Z. Y., Ouyang, H., Zhou, C. P., and Xu, X. L.: N₂O exchange within a soil and atmosphere profile in alpine grasslands on the Qinghai-Xizang Plateau, Acta Bot. Sin., 46, 20–28, 2004.
 Pinto, M., Merino, P., del Prado, A., Estavillo, J. M., Yamulki, S., Gebauer, G., Piertzak, S., Lauf,
- J., and Oenema, O.: Increased emissions of nitric oxide and nitrous oxide following tillage of a perennial pasture, Nutr. Cycl. Agroecosys, 70, 13–22, 2004.
- Prentice, I. C., Sykes, M. T., Lautenschlager, M., Harrison, S. P., Denissenko, O., and Bartlein, P. J.: Modelling Global Vegetation Patterns and Terrestrial Carbon Storage at the Last Glacial Maximum, Global. Ecol. Biogeogr, 3, 67–76, 1993.
- Rees, R. M., Wuta, M., Furley, P. A., and Li, C. S.: Nitrous oxide fluxes from savanna (miombo) woodlands in Zimbabwe, J. Biogeogr., 33, 424–437, doi:10.1111/j.1365-2699.2005.01423.x, 2006.

Rodhe, H.: A Comparison of the Contribution of Various Gases to the Greenhouse Effect, Science, 248, 1217–1219, doi:10.1126/science.248.4960.1217, 1990.

Saggar, S., Andrew, R. M., Tate, K. R., Hedley, C. B., Rodda, N. J., and Townsend, J. A.:
 Modelling nitrous oxide emissions from dairy-grazed pastures, Nutr. Cycl. Agroecosys, 68, 243–255, 2004.

Saggar, S., Giltrap, D. L., Li, C., and Tate, K. R.: Modelling nitrous oxide emissions from grazed grasslands in New Zealand, Agric. Ecosyst. Environ., 119, 205–216, 2007.

Scurlock, J. M. O. and Hall, D. O.: The global carbon sink: a grassland perspective, Global. Change. Biol., 4, 229–233, doi:10.1046/j.1365-2486.1998.00151.x, 1998.

Shi, X. Z., Yu, D. S., Warner, E. D., Pan, X. Z., Petersen, G. W., Gong, Z. G., and Weindorf, D. C.: Soil database of 1:1,000,000 digital soil survey and reference system of the Chinese Genetic Soil Classification System, Soil Survey Horizons, 45, 129–136, 2004.

1693

- Unkovich, M. and Nan, Z.: Problems and prospects of grassland agroecosystems in western China, Agric. Ecosyst. Environ., 124, 1–2, 2008.
- Wang, Y., Ma, X., Ji, B., Du, R., Chen, Z., Wang, G., Wang, Y., and Wan, X.: Diurnal and seasonal variation in methane and nitrous oxide fluxes in meadow steppe of Inner Mongolia, Zhiwu Shengtai Xuebao, 27, 792–796, 2003.
- White, F.: The Vegetation of Africa: a descriptive Memoir to accompany the Unesco/AETFAT/UNSO Vegetation Map of Africa, 356, United Nations Educational, Scientific and Cultural Organization (UNESCO), Paris, 356 pp., 1983.
- Xu, R., Wang, M., and Wang, Y.: Using a modified DNDC model to estimate N₂O fluxes from semi-arid grassland in China, Soil Biol. Biochem., 35, 615–620, 2003.

Yamulki, S., Harrison, R. M., Goulding, K. W. T., and Webster, C. P.: N₂O, NO and NO₂ fluxes from a grassland: Effect of soil pH, Soil Biol. Biochem., 29, 1199–1208, 1997.

Yang, D.: Degradation and protection of grassland on the Qinghai-Tibet Plateau, Erosion, debris flow and environment in mountain regions. Proceedings of the Chengdu Symposium, China, 1992.

Yu, D. S., Shi, X. Z., Wang, H. J., Sun, W. X., Chen, J. M., Liu, Q. H., and Zhao, Y. C.: Regional patterns of soil organic carbon stocks in China, J. Environ. Manage., 85, 680–689, 2007.

Table 1. DNDC model results co	ompared with the literature data.
--------------------------------	-----------------------------------

Longitude	Latitude	Reference	N_2O	Modelled results	Location	Grassland Type
$(kg N ha^{-1} y^{-1})$						
123.01-119.15	44.01-43.03	Huang et al. (2003)	0.72	0.70	Songnen Plain, China	Temperate Grassland
116-117	43-44	Du et al. (1997)	0.28	0.45	Inner Mongolia, China	Temperate Grassland
116-117	43.5-44	Dong et al. (2000)	0.56	0.45	Inner Mongolia, China	Temperate Grassland
116.7	43.62	Xu et al. (2003)	0.18	0.37	Inner Mongolia, China	Temperate Grassland
93.83	35.22	Pei et al. (2003, 2004)	0.05	0.06	Qinghai, China	Montane Grassland
115.5-117.2	43.5-44.6	Wang et al. (2003)	0.09	0.40	Inner Mongolia, China	Temperate Grassland
116.66	43.5	Du et al. (2006)	0.73	0.64	Inner Mongolia, China	Temperate Grassland
101.25	37.5	Du et al. (2008)	2.08	0.16	Qinghai, China	Montane Grassland
116.66	43.5	Chen et al. (2000)	0.17	0.40	Inner Mongolia, China	Temperate Grassland
а		Mummey et al. (2002)	0.28	0.22	US	Temperate Grassland
а		Mosier et al. (1993)	0.17	0.16	Wyoming, US	Montane grasslands

^a For China domestic, the value extracted according to the coordinate of literature data, and for the other places of the world, the compared value was use the average value of similar grassland type in China. If the literature study area was smaller than the DNDC simulation unit, then the average literature value was applied.

1695

Table 2. Baseline values for sensitivity tests.

Property	Baseline value
Annual mean temperature (°C)	9
Total annual precipitation (mm)	520
Clay fraction (%)	0.19
Field capacity (%)	0.49
Wilting point (%)	0.22
Porosity (%)	0.451
Initial soil C fraction (kg C/kg soil)	0.0025
Bulk density (g/cm ³)	1.22
Soil pH	7.4
Number of days grazed (per year)	12
Grazing hours per day	12
Cattle grazing intensity (head/ha)	0.3
Sheep grazing intensity (head/ha)	3

Table 3. Modelled annual N_2O emission rates for 18 types of grassland in China (kg N ha	$(^{-1} y^{-1}),$
grassland area and total N_2O emission per year.	

Grassland Type	Emission rates $(kg N ha^{-1} y^{-1})$	Grassland Area (million ha)	Total N_2O emission (Gg N y ⁻¹)
Temperate Meadow Steppe (TMS)	0.26	18	4.7
Temperate Steppe (TS)	0.33	47	15.5
Temperate Desert Steppe (TDS)	0.16	16	2.6
Alpine Meadow Steppe (AMS)	0.10	5	0.5
Alpine Steppe (AS)	0.08	30	2.4
Alpine Desert Steppe (ADS)	0.03	6	0.2
Temperate Steppe Desert (TSD)	0.11	7	0.8
Temperate Desert (TD)	0.09	20	1.8
Alpine Desert (AD)	0.03	5	0.2
Warm Temperate Tussock (WTT)	0.28	11	3.1
Warm Temperate Shrub Tussock (WTST)	0.30	12	3.6
Tropical Tussock (TT)	0.37	19	7
Tropical Shrub Tussock (TST)	0.37	17	6.3
Dry Tropical Shrub Tussock with Savanna (DTSTS)	0.32	1	0.3
Lowland Meadow (LM)	0.32	35	11.2
Mountain Meadow (MM)	0.23	22	5.1
Alpine Meadow (AM)	0.16	64	10.2
Swamp	0.48	2	1
Average rate/Total area/Total emission	0.22	337	76.5

Table 4. Grassland categories, zonal climate and average N_2O flux.

Categories	Grassland Type	Annual Precipitation (mm)	Annual Temperature (°C)	Area (million ha) [%]	N_2O Emission Rate (kg N ha ⁻¹ year ⁻¹)	Total N ₂ O Emission (Gg N y ⁻¹) [%]
Temperate Grassland	TMS TS TDS TSD TD WTT WTST LM	282.11	5.54	165.92 [49]	0.23	43.3 [57]
Montane grasslands	AMS AS ADS AD AM MM	400.77	-1.25	132.56 [39]	0.11	18.6 [24]
Tropical/ subtropical grasslands	TT TST DTSTS Swamp	1109.49	16.22	39.50 [12]	0.39	14.6 [19]



Fig. 1. China's 18 types grassland distributions.





Fig. 2. DNDC results compared with literature reports. Dashed line is 1:1 line (Du et al., 2008 was excluded).



Fig. 3. Effect of changing a single factor of soil, management and climate on sensitive analysis scenario.





Fig. 4. Three grassland ecological categories of Chinese grasslands.



Fig. 5. The spatial distribution of $\rm N_2O$ flux for Chinese grassland (the blank area is non-grassland).





Fig. 6. Chinese grassland $\mathrm{N_2O}$ emission rate yearly changes for 2000–2007.



Fig. 7a. The annual accumulated precipitation difference between 2005 and 2001. Positive values indicate precipitation increased from 2001 to 2005, and negative values indicate a decrease.





Fig. 7b. The accumulated temperature > 0 °C difference between 2005 and 2001. Positive values indicate accumulated temperature increased from 2001 to 2005, and negative values indicate a decrease.