

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

**Abrupt sea surface
pH change at the end
of the Younger Dryas**

E. Douville et al.

Abrupt sea surface pH change at the end of the Younger Dryas in the central sub-equatorial Pacific inferred from boron isotope abundance in corals (*Porites*)

E. Douville¹, M. Paterne¹, G. Cabioch², P. Louvat³, J. Gaillardet³,
A. Juillet-Leclerc¹, and L. Ayliffe⁴

¹Laboratoire des Sciences du Climat et de l'Environnement, Institut Pierre Simon Laplace (LSCE/IPSL), UMR8212 CEA-CNRS-UVSQ, Domaine du CNRS, Av. de la Terrasse, 91198 Gif/Yvette, France

²Institut de Recherche pour le Développement (IRD), UMR IPSL/LOCEAN, UPMC/CNRS/IRD/MNHN, 32 avenue Henri Varagnat, 93143 Bondy Cedex, France

³Institut Physique du Globe de Paris (IPG-Paris), Laboratoire de Géochimie et de Cosmochimie, Place Jussieu, 75252 Paris, France

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



⁴ Research School of Earth Sciences, Australian National University, ACT 0200, Australia

Received: 24 February 2010 – Accepted: 4 March 2010 2007 – Published: 18 March 2010

Correspondence to: E. Douville (eric.douville@lsce.ipsl.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

7, 1959–1993, 2010

**Abrupt sea surface
pH change at the end
of the Younger Dryas**

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The “ $\delta^{11}\text{B}$ -pH” technique was applied to modern and ancient *Porites* from the sub-equatorial Pacific areas (Tahiti and Marquesas) spanning a time interval from 0 to 20720 calendar years to determine the amplitude of pH changes between the Last Glacial Period and the Holocene. Boron isotopes were measured by Multi-Collector-Inductively Coupled Plasma Mass Spectrometry (MC-ICPMS) with an external reproducibility of 0.25‰, allowing a precision of ± 0.025 pH-units. The boron concentration [B] and isotopic composition of modern samples indicate that the temperature strongly controls the partition coefficient K_D for different aragonite species. Modern coral $\delta^{11}\text{B}$ values and the reconstructed sea surface pH values for different Pacific areas match the measured pH expressed on the Sea Water Scale and confirm the calculation parameters that were previously determined by laboratory calibration exercises. Most ancient sea surface pH reconstructions near Marquesas are higher than modern values. These values range between 8.20 and 8.26 for the Holocene and reached 8.31 at the end of the last glacial period (20.7 kyr BP). At the end of the Younger Dryas (11.50 \pm 0.1 kyr BP), the central sub-equatorial Pacific experienced a dramatic drop of up to 0.2 pH-units from the average pH of 8.2 before and after this short event. Using the CO2SYS program, we recalculated the aqueous $p\text{CO}_2$ to be 400 \pm 24 ppmV at around 11.5 kyr BP for corals at Marquesas and \sim 500 ppmV near Tahiti where it was assumed that $p\text{CO}_2$ in the atmosphere was 250 ppmV. Throughout the Holocene, the difference in $p\text{CO}_2$ between the ocean and the atmosphere at Marquesas ($\Delta p\text{CO}_2$) indicates that the surface waters behave as a moderate CO_2 sink (–67 to –11 ppmV) during El Niño-like conditions. In contrast, during the last glacial/interglacial transition, this area was a moderate source of CO_2 (–9 to 56 ppmV) for the atmosphere, highlighting predominant La Niña-like conditions. Such conditions were particularly pronounced at the end of the Younger Dryas with a large amount of CO_2 released with $\Delta p\text{CO}_2$ of +140 ppmV. This last finding provides further evidence of the marked changes to the water mass pH and temperature properties in the equatorial Pacific at

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the Younger Dryas- Holocene transition and the strong impact of oceanic dynamic on the atmospheric CO₂ content.

1 Introduction

The acidity of the ocean surface is increasing because of anthropogenic emission of CO₂ into the atmosphere. This increase may have severe consequences for those organisms which build their external skeleton out of calcium carbonate (CaCO₃) (Orr et al., 2005). During the last 200 years, roughly fifty percent of the anthropogenic CO₂ (IPCC, 2005) contributed to the acidification of the superficial ocean by 0.1 pH-units. Models suggest that the sea surface pH could drop by ~0.4 units before the year 2100 if the present-day trend of carbon dioxide increase continues (Caldeira and Wickett, 2003). Today, the oceanic sources and sinks of CO₂ show a high degree of spatial and annual to inter-annual variability. For instance, the wind is strengthening, and uptake of CO₂ in the Southern Ocean is reduced (Le Quéré et al., 2007). Changes in the marine biological pump control carbon removal from the upper ocean into the deep ocean (McGillicuddy et al., 2007). Changes have taken place in the oceanic circulation (Feely et al., 1999; Inoue et al., 2001), and these changes may counteract the anthropogenic acidification of the ocean. In the past, the atmospheric *p*CO₂ changed by ~80 to 100 ppmV between glacial and interglacial periods (Hönisch and Hemming, 2005; Hönisch et al., 2009; Monnin et al., 2001, 2004). Those variations are commonly ascribed to changes in the oceanic uptake of CO₂ because the ocean is the largest carbon reservoir in the atmosphere-hydrosphere-biosphere system. However, the CO₂ exchange and related mechanisms have not yet been fully understood and quantified (Archer et al., 2000; Kohfeld et al., 2005; Sigman and Boyle, 2000). Quantifying changes in the oceanic *p*CO₂ over the centennial and millennial time scales will elucidate the role of the oceanic carbon cycle and the links between the oceanic and atmospheric changes of *p*CO₂ and their consequences on ecosystems.

Ocean acidification and past sea surface pH or *p*CO₂ may be deduced from the

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



boron isotopic composition ($\delta^{11}\text{B}$) of biogenic carbonates (Hemming and Hanson, 1992; Spivack et al., 1993). During the past decade, numerous studies evaluated the “ $\delta^{11}\text{B}$ -pH” relationship using experimental studies or laboratory cultures of foraminifera and corals (Hemming et al., 1995; Hönisch et al., 2004; Reynaud et al., 2004; Sanyal et al., 1996, 2000). Only a few studies based on tropical corals focused on the past hundreds or thousands of years. These first investigations investigated the western Pacific Ocean (Pelejero et al., 2005; Wei et al., 2009) and more recently the South China Sea (Liu et al., 2009). Those studies revealed pronounced sea surface pH changes by 0.2–0.3 during the Holocene. These changes were strongly controlled by climate (monsoon) or oceanographic patterns (Pacific Decadal Oscillation).

First we describe in detail the modern “pH – Sea Surface Temperature” relationships for surface waters of the sub-Equatorial Pacific Ocean and the protocols used here to reconstruct the pH and $p\text{CO}_2$ properties of surface waters. We also present the $\delta^{11}\text{B}$ -pH and partition coefficient for boron K_D obtained from modern *Porites* collected in tropical areas of various sea surface temperatures (SSTs). Then, we use the derived relationships to investigate ancient corals recovered from submerged reefs off the Marquesas and from modern reefs near the Tahiti Islands in the central equatorial Pacific (Fig. 1) to reconstruct past pH, SST and $p\text{CO}_2$ levels of these areas. These new results are compared to previous results from modern or ancient corals of the genera *Acropora* and *Porites* (Gaillardet and Allègre, 1995) and from planktonic foraminifera (Palmer and Pearson, 2003). The deduced changes in the sea surface pH and SST are evaluated against oceanographic patterns, and the estimated oceanic $p\text{CO}_2$ values are compared to atmospheric values measured in Antarctic ice cores (Monnin et al., 2001, 2004).

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 Hydrological setting of the central tropical Pacific Ocean: “pH-SST” relationships

The pH and SST data show that in the modern equatorial Pacific, the sea surface pH and SST between 5° S and 20° S are highly negatively correlated (Fig. 2, data from <http://www.nodc.noaa.gov/OC5/SELECT/dbsearch/dbsearch.html>) (Lefèvre et al., 1994). This correlation is especially strong between 80° W and 170° W because of the cold and CO₂-rich waters coming from the major deep water upwelling along the eastern coasts of the sub-Equatorial Pacific Ocean. In the eastern Pacific, the nutrient-rich surface waters are characterised by low pH and cool SST (Fig. 2a). The pH ranges from ~7.98 to about 8.01–8.06 on the seawater scale, as reported on a modern pH scale, and the SST varies from 20.4 °C to 23.1 °C. At latitudes between 15° S and 20° S, the pH increases slightly by about 0.05 pH-units, and the SST increases by 3 °C from 80° W to 170° W. This increase is more than 0.07 pH-units between 5° S and 10° S (Fig. 2b). In the western Pacific, high pH values of 8.22±0.03 and warm SSTs are observed during the boreal summer. During the boreal winter, the pH drops by 0.1 pH-unit. The SSTs are uniformly around 29.5 °C between 5° S and 15° S and lower by about 4 °C between 15° S and 20° S. In the central Pacific between 130° W and 150° W and at latitudes higher than 15° S, warm surface waters with a SST of 28.6 °C and a pH of ~8.22 were observed during 1994, an El Niño year. These waters show pH-SST properties similar to those of the western Pacific surface waters (Fig. 2).

Most modern pH values for Marquesas and Tahiti (5–17° S) between 130° W and 150° W range from 8.10 to 8.15 (Fig. 2). This east-west pH-SSTs pattern corresponds to the well-known seasonal variation of the zonal distribution of water masses along the equator. This pattern is modified by changes in the wind strength. During El Niño years, westward water transport through the South Equatorial Current SEC decreases in the central Pacific because of eastward transport of warmer water from the western Pacific (Taft and Kessler, 1991). The sea surface pH increases under these conditions. When strong easterlies return, the westward SEC flow increases, and the equatorial

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



thermocline shoals and favours contribution of more CO₂-rich waters. Consequently, during La Niña-like conditions, acidic cold waters are advected from the east toward the central Pacific (Taft and Kessler, 1991).

3 Materials and methods

3.1 Geographical setting and regional modern SST

The “ $\delta^{11}\text{B-pH}$ ” indicator was first applied to modern *Porites* collected from New Caledonia (2001–2004), Moorea (1950) and Ishigaki Islands (Fig. 1). The New Caledonia sample used in this paper comes from outside the lagoon on the upper part of the barrier reef slope. This site is called Fausse passe d’Uitoé. The SST was measured at this site every month since 2000 and had a mean value of $24.4 \pm 1.9^\circ\text{C}$ (Table 1). The coral skeleton spans the period from 2001–2004. *Porites* off of Ishigaki Island are the coral reference material JCp-1 prepared by the Geological Survey of Japan (Inoue et al., 2004). The regional SST over the last two decades in this area is close to $26.1 \pm 2.1^\circ\text{C}$. The average SST of a sample from Moorea (MOO 3A-1-02) was $27.1 \pm 0.5^\circ\text{C}$ (Boiseau, 1998). Another modern sample is COM3 from the Red Sea. This sample grew at annually averaged SST of $22.5 \pm 1.5^\circ\text{C}$ (Gertman and Brenner, 2004; Heiss et al., 1999). According to modern observations (<http://www.nodc.noaa.gov>) (Henin, 1999), the Marquesas area has a mean annual SST of $28 \pm 1^\circ\text{C}$. The ancient *Porites* studied here were collected from submerged reefs off the Marquesas and Tahiti Islands in the Equatorial Pacific. Coral ages, which span from 20.72 ± 0.20 kyr BP to 0.25 ± 0.03 kyr BP, were obtained by mass spectrometric U-Th dating (Paterne et al., 2004; Cabioch et al., 2008).

3.2 Chemical preparation

The mean annual growth band of *Porites* is about 10 mm in our samples. About 400 mg of the coral skeleton were cut along the growth axis systematically. Each of these samples represents more than one year. The coral fragments were crushed in an agate

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mortar, and the resulting carbonate powder was washed three times with ultra-pure water in an ultrasound bath (Gaillardet and Allègre, 1995). Then, the powder was dissolved in 1 N HNO₃ before chemical purification. The solutions were purified on the anion exchange resin Amberlite IRA 743 using the method developed by Lécuyer et al. (2002). In 50 ml polypropylene corning centrifuge tubes, the solutions were neutralised to a pH of 8–9. Cleaned resin was added in sufficient quantity (500 mg) to extract 100% of the boron from solution for each standard and sample. The tubes were shaken for more than four hours before the resin was rinsed three times with MQ water. Then, the boron was eluted by three successive volumes of 5 ml of 0.1 N HNO₃. Extraction yields of close to 100% and accurate purification were verified by Inductively Coupled Plasma-Quadrupole Mass Spectrometry (ICP-QMS). Finally, the boron concentration was adjusted to 200 ppb in 2% nitric acid for MC-ICPMS analyses.

3.3 Boron isotope measurements

The boron isotope composition was determined with double-focusing sector-field multi-collector inductively coupled plasma mass spectrometers (MC-ICPMS Neptune of ThermoFisher Scientific) both in the Advanced Mass Spectrometry Laboratory of Thermo Fisher Scientific in Bremen (Germany) and at the Institut Physique du Globe in Paris (France). Common introduction in Bremen was by a quartz double-pass spray chamber, and in Paris we used a direct injection high efficiency nebuliser (d-DIHEN). Mass drift of the ¹¹B/¹⁰B ratio with time was systemically controlled by standard-sample bracketing. Most of the δ¹¹B values presented here were determined using the direct injection technique, which allows a strong reduction of the analytical blank contributions. Those contributions were lower than 0.5‰ for each isotope with a rinse time of three minutes or less (Louvat et al., 2010). Comparison of repeated analyses for four different samples (DW1281c_78a1; DW1261_68a1; DW1281_75a2; MOO 3A-1-02) using direct injection (d-DIHEN) and conventional introduction yields excellent agreement within the analytical uncertainty given here (Table 1). The reproducibility and accuracy of the ¹¹B/¹⁰B ratios were calculated from repeated analyses of the boric acid standard

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NBS-951 and the North Atlantic Seawater Standard NASS-II. The measured external reproducibility of 0.25‰ (2σ) is similar to the reproducibility that was recently published for MC-ICPMS analyses of Neptune (Foster, 2008; Louvat et al., 2010). A mean $\delta^{11}\text{B}$ of 39.9‰ ($n = 20$, 2σ) was measured for NASS-II. This value is also within the external precision values of 39.7–40.2‰ previously published for thermal ionization mass spectrometry (TIMS) or MC-ICPMS measurements (Gaillardet and Allègre, 1995; Hemming and Hanson, 1992; Hönisch et al., 2004; Wei et al., 2009; Louvat et al., 2010). The external reproducibility of 0.25‰ (2σ represents our analytical precision for each of the analyzed samples. The boron isotopic composition of seawater ($\delta^{11}\text{B}_{\text{SW}}$) used for pH calculations was 39.9‰. Boron concentrations were measured with a precision of $\pm 3\%$ (2σ) by ICP-QMS (ThermoFisher X-series) at Laboratoire des Sciences du Climat et de l'Environnement (LSCE) in Gif-sur-Yvette (France).

3.4 pH and $p\text{CO}_2$ calculations for seawater

The sea surface paleo-pH can be reliably reconstructed by measuring the boron isotopes in tropical corals. Two previous calibration exercises using laboratory-cultured corals (*Acropora* and *Porites*) demonstrate the validity of the technique and provide empirical calculation parameters (Hönisch et al., 2004; Reynaud et al., 2004). These pH calculations are based on isotopic fractionation by the coral of two boron species: borate ions $\text{B}(\text{OH})_4^-$ and boric acid $\text{B}(\text{OH})_3$. The relative proportions of these species and their isotopic composition in seawater are pH-dependent (Pagani et al., 2005; Sanyal et al., 1996, 2000). Thus borate ions $\text{B}(\text{OH})_4^-$ dissolved in seawater would be preferentially incorporated into the carbonate skeleton by substitution of bicarbonate ions (Hemming and Hanson, 1992). Consequently, the relationship (Eq. 1) between pH and $\delta^{11}\text{B}$ is as follows:

$$\text{pH}_{\text{SW}} = \text{pK}_{\text{B}} - \log \left(\frac{\delta^{11}\text{B}_{\text{SW}} - \delta^{11}\text{B}_{\text{carbonate}}}{\alpha^{-1}\delta^{11}\text{B}_{\text{carbonate}} - \delta^{11}\text{B}_{\text{SW}} + 1000(\alpha^{-1} - 1)} \right) \quad (1)$$

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where pK_B is the equilibrium constant between the boric acid $B(OH)_3$ and the borate ions $B(OH)_4^-$ in seawater (Dickson, 1990). $\delta^{11}B_{\text{carbonate}}$ ($\delta^{11}B$) is the isotopic composition of boron measured here in *Porites*. $\delta^{11}B_{\text{SW}}$ is the isotopic composition of boron measured for seawater. And the value α (α_{4-3}) is the isotopic fractionation constant of the following equilibrium (R1):



Factors such as temperature, salinity, light, food (Hönisch et al., 2004; Pagani et al., 2005), and especially the coefficient α determine the uncertainties associated with calculating the pH (Pagani et al., 2005). Until recently, α was the subject of numerous empirical or theoretical studies with the goal of precisely determining its value. The following two major trends appear today: (1) $\alpha \geq 0.980-0.981$ (Gaillardet and Allègre, 1995; Hemming et al., 1995; Hönisch et al., 2004, 2007; Kakahana et al., 1977; Palmer and Pearson, 2003; Sanchez-Valle et al., 2005; Sanyal et al., 1996, 2000; Xiao et al., 2006) and (2) $\alpha \leq 0.974-0.976$ (Foster et al., 2008; Klochko et al., 2006; Lécuyer et al., 2002; Pagani et al., 2005; Zeebe, 2005). The first trend ($\alpha \geq 0.980-0.981$) is usually applied today for paleo-reconstructions. Today, α values of 0.974–0.976 are often considered theoretical values (Klochko et al., 2006), and the calculated sea surface pH from our $\delta^{11}B$ data (described below) would range from 8.4 to 8.6. These values are 0.3 to 0.4 pH-units higher than the expected modern pH on the Sea Water Scale (pH_{SWs}). To reconstruct pH values that are more compatible with this scale, we use a value of the coefficient α of 0.981 (Kakahana et al., 1977). This value was empirically verified by calibration for both coral genera *Acropora* and *Porites* (Hönisch et al., 2004; Reynaud et al., 2004). These laboratory experiments reveal a systematic 0.6 $\delta^{11}B$ offset between these two coral types over the pH range from 7.7 to 8.2. These $\delta^{11}B$ offset may result in a systematic pH offset between the two species. This offset was taken into account by adding 0.6 δ -units to the $\delta^{11}B$ values of each ancient *Acropora* sample from Tahiti for comparison with our *Porites* $\delta^{11}B$ values (Gaillardet and Allègre, 1995).

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To estimate the oceanic $p\text{CO}_2$ from the $\delta^{11}\text{B}$ -pH values of the corals, we followed the procedure described by Hönisch and Hemming (2005). This procedure is based on the determination of two of the four parameters of the marine carbonate equilibrium ($p\text{CO}_2$, ΣCO_2 , alkalinity and pH) in addition to Sea Surface Salinity (SSS) and SST. Using the National Oceanic and Atmospheric Administration (NOAA) data for modern water masses in the sub-Equatorial Pacific Ocean, we obtain a linear correlation between the total alkalinity (A_k) and the SSS of [$A_k=0.0688\cdot\text{SSS} - 0.0484$; $R^2=0.974$; $n=233$].

We hypothesise that the present-day “ $A_k - \text{SSS}$ ” relationship was also valid in the past. Past salinities can be estimated from sea-level changes through the past (Bard et al., 1996). For example, a reduction of -60 m occurred during the Younger Dryas (YD) over a mean oceanic depth of 3800 m. This estimate takes into account the modern salinity near Tahiti and the Marquesas Islands of 35.6 ± 1 practical salinity units (psu) as a reference (Delcroix et al., 1996). From this calculation, the salinity in the past was about 36.2 psu in the central Pacific and about 34.8 psu in the eastern Pacific during the YD. We consider today’s mean seasonal salinity variability of ± 1 psu (Delcroix et al., 1996) in the calculation. The oceanic $p\text{CO}_2$ calculation was performed using the CO2SYS program by Lewis and Wallace (1998) with K_1 and K_2 from Roy et al. (1993), pH values on the seawater scale, and phosphate and silicate concentrations of $0.2\ \mu\text{M}$ and $2\ \mu\text{M}$, respectively.

4 Results and discussion

To verify the reliability of pH reconstruction from boron isotopes in corals, we first analyzed the $\delta^{11}\text{B}$ and pH data for modern *Porites* samples collected from various areas and compared our results with the modern pH_{SWS} measured in the Pacific. Those areas cover a range of SST values and allow us to further elucidate the influence of the temperature on the incorporation of boron and its isotopic composition in corals. For each area, the measured or previously published values of $\delta^{11}\text{B}$ and [B] in *Porites* are evaluated against a theoretical curve that illustrates the preferential incorporation of

borate ions in corals by substituting for the bicarbonate ions (Hemming and Hanson, 1992). This approach allows us to determine the various values of the partition coefficient K_D and to establish a link between the SST and this K_D . Similar exercises were done for the ancient corals from Marquesas and Tahiti to obtain information about the past SST. Finally, the changes in the pH and the $p\text{CO}_2$ as a function of time will be discussed in the central sub-equatorial Pacific by using the results of the “ $\delta^{11}\text{B}$ -pH” calculation for both *Porites* near Marquesas (this study) and *Acropora* near Tahiti (Gaillardet and Allègre, 1995). This analysis takes into account the observed SST data.

4.1 Reconstruction of the pH and SST from $\delta^{11}\text{B}$ and [B] data in modern *Porites*

4.1.1 pH deduced from $\delta^{11}\text{B}$ in modern *Porites*

The mean $\delta^{11}\text{B}$ values measured in modern *Porites* (Table 1) were about $25.8 \pm 0.2\%$ ($n = 6$) for Moorea (1950AD), $24.8 \pm 0.3\%$ ($n = 8$) for New Caledonia (2001–2004 AD) and $24.5 \pm 0.2\%$ ($n = 17$) for the Ishigaki Island (JCp-1). These values are similar to the modern value of 24.8% published for the Fanning Island (Hemming et al., 1998) and higher than the values of 23.6% and 23% found for the Red Sea (Gaillardet and Allègre, 1995) and Flinders Reef (~1990 AD) along the Great Barrier Reef (Pelejero et al., 2005), respectively. According to the regional SST, the deduced sea surface pH was 7.98 for the Red Sea, 8.06 for the Ishigaki area, 8.07 for both Fanning Island and New Caledonia and 8.13 and 8.18 for Moorea in 1991 AD and 1950 AD, respectively. The $\delta^{11}\text{B}$ value for Moorea was higher by + 0.5 than the previous measurement in 1991 (Gaillardet and Allègre, 1995). This change corresponds to the pH change of 0.05 pH-unit. These values from modern *Porites* off the Moorea match the pH_{SWS} values measured in this area (Fig. 2). The inter-decadal variability linked, for example, to the Pacific Decadal Oscillation (PDO) could explain the difference observed in the western Pacific (Pelejero et al., 2005), but this $\delta^{11}\text{B}$ change could also reflect the consequences of surface water acidification during the industrial era given our own

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



recent description of the last century at Fiji (Douville et al., 2009). Finally, the pH values deduced from the modern *Porites* for various areas of the Pacific Ocean match well with the modern pH map (Fig. 1) that was calculated on the Sea Water Scale (pH_{SWs}). These results corroborate the analyses that were based on empirical α values and previous lab calibrations on the tropical coral *Porites* (Hönisch et al., 2004; Reynaud et al., 2004).

4.1.2 SST and the partition constant K_D for aragonite corals

All of the $\delta^{11}\text{B}$ values versus [B] obtained here (Table 1) for modern *Porites* from Moorea, Ishigaki and New Caledonia Islands in the Pacific Ocean were compared to available published values for Moorea, Fanning Island and the Red Sea (Gaillardet and Allègre, 1995; Hemming et al., 1998) and plotted (Fig. 3a) to verify the model of a unique incorporation of boron (as borate ions) into the corals by substitution of bicarbonates ions (Hemming and Hanson, 1992; Sanyal et al., 2000; Yu and Elderfield, 2007; Foster, 2008). This also suggests that the [B] (or B/Ca) in corals increases together with the $\delta^{11}\text{B}$ and is controlled by the borate/bicarbonate abundance ratio in seawater according to the following relationship (Eq. 2):

$$[\text{B}/\text{Ca}]_{\text{corals}} = K_D \cdot [\text{B}(\text{OH})_4^- / \text{HCO}_3^-]_{\text{seawater}} \quad (2)$$

where K_D is the partition coefficient for boron. For various pH conditions, the theoretical $\delta^{11}\text{B}$ or $[\text{B}(\text{OH})_4^- / \text{HCO}_3^-]$ ratios for seawater and the [B] for corals can be calculated and plotted by setting a K_D value. This model curve can be directly compared and fitted to points that correspond to the $\delta^{11}\text{B}$ and the [B] values measured in corals for each studied area. Model curves were generated with empirical values of $\alpha=0.981$ (Kakihana et al., 1977), $\delta^{11}\text{B}_{\text{SW}}=39.9\text{‰}$ (this work) and $\text{p}K_{\text{B}}=8.56$ (Dickson, 1990) at 28°C and 35 psu. Figure 3a shows scattered points that represent various values of the partition coefficient K_D ranging between 0.0068 (Moorea) and 0.0125 (Red Sea). A similar graph for ancient corals also shows scattered dots with average K_D values of 0.0057 and 0.0068 for the Marquesas and Tahiti samples, respectively (Fig. 3b). The

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



highest values are observed for the corals that grew at the end the Younger Dryas with values up to 0.0095 (Tahiti).

To evaluate the potential effect of the SST on this coefficient, the borate/bicarbonate molar ratios for seawater $[B(OH)_4^-/HCO_3^-]_{\text{seawater}}$ were first re-calculated from the measured $\delta^{11}B\text{-pH}$ values using the regional SST and then compared to the measured boron/calcium molar ratios ($[B/Ca]_{\text{coral}}$) for all of the corals studied here (Fig. 4). This graph shows that the regional SST strongly controls the partition coefficient K_D , which is 0.0057 or less for the highest temperature (in the Marquesas area or Fanning Island during the winter season and the influence of warm and high pH waters, see below) and 0.0125 for the lowest SST (close to 22.5 °C in the Red Sea). The K_D values are plotted versus the SST for modern *Porites* in Fig. 5a. This figure shows a strong linear relationship with a regression coefficient (R^2) of 0.98. This linear dependence confirms that the boron in corals is more easily incorporated in aragonite at low temperatures, as previously observed in *Porites* (Fallon et al., 2003).

To test whether this linear temperature dependence of K_D applies (i) to ancient corals near Marquesas and (ii) to a wider temperature range that affects cold-water corals such as *Lophelia pertusa*, we extrapolated the regression towards SSTs of 28–29 °C and temperatures of 5.5 to 11 °C (Fig. 5b). This low temperature range covers the North Atlantic intermediate waters, which today have pH values approximately 7.95±0.05. For pH 7.95, the K_D deduced from boron concentrations in *Lophelia pertusa* (Douville et al., 2007) at local temperature agrees with the established regression (Fig. 5b). The average K_D of ancient Marquesas corals was 0.0057 (Fig. 3b). This value fits the linear temperature dependence described above well with deviations of no more than ±1 °C at 28 °C (Fig. 4). These reconstructed temperatures match the mean SST observed today near Marquesas. However, the lowest SSTs are estimated at the end of the Younger Dryas (YD) or during the glacial period, and higher SSTs are suggested by Figure 4 for the Marquesas corals that are about 9 and 14–15 kyr BP old, respectively.

Similar trends are observed for the old corals near Tahiti. These corals have a mean K_D of 0.0068 (SST # 27 °C) and a pronounced and unique SST decrease of 2.5 °C

**Abrupt sea surface
pH change at the end
of the Younger Dryas**

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



around 11.5 kyr BP ago. Another example of the influence of the SST on the partition coefficient K_D is from the seasonal data published at Fanning Island (Figs. 3a and 4). The seasonal SSTs in the Fanning Islands vary by 3°C. The highest temperatures are close to 28°C and correspond to surface waters of high pH recorded by the high density bands of the modern *Porites* (<http://www.nodc.noaa.gov>; Hemming et al., 1998; Hönisch et al., 2004). This amplitude of seasonal SST changes matches the temperature (24.5°C–27.5°C) described in Fig. 4. Thus, at Fanning Islands the highest SST of about 27.5°C corresponds to a low K_D close to ~0.006 (Fig. 4), and points close to a high K_D of ~0.0095 corresponding to 24.5°C.

4.2 Variations of “ $\delta^{11}\text{B-pH}$ ” as a function of time

In fossil *Porites* from Marquesas (Table 1), the $\delta^{11}\text{B}$ values vary from 26.0 to 26.6‰ during the Holocene, drastically fall to 24.7‰ at the end of the Younger Dryas, and then rise again to values between 25.6 and 26.4‰ during the beginning of the Younger Dryas and Bölling Allerød. The highest value (27.1 ‰) was observed during the last glacial period (20.7 kyr BP). During the Holocene, all of the deduced values for the sea surface pH_{SWS} from Marquesas and Tahiti are similar at 8.23 ± 0.03 (Table 1, Fig. 6). These values are equivalent to modern pH (~8.15) (Fig. 2a) when accounting for the model prediction for anthropogenic acidification on the order of 0.07 pH-units (Sabine et al., 2004). At the end of the YD at ~11.5 kyr BP, uniformly low pH values (~8.00–8.05) are observed close to both islands (Fig. 6). These low pH values are preceded by high pH values (8.15 to 8.31) during the Bölling Allerød and the last glacial/interglacial transition (Fig. 6). From foraminifera in the “western” Pacific (ERDC-92), pH reconstruction using the $\delta^{11}\text{B}$ method yields ~8.23, ~8.13, and ~8.20 for the early, middle, and late Holocene. During the YD, pH-values were ~8.16, and during the Bölling Allerød, they were between 8.1 and 8.2. During the last glacial period, the highest pH values are estimated at ~8.3 (Palmer and Pearson, 2003). The central equatorial Pacific experienced larger pH changes (0.2 pH-units) than the western Pacific (~0.1 pH-units) during the transition between the YD and the Holocene. This difference may be attributed to

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the temporal resolution obtained using corals and foraminifera. In corals, the sampling represents one or a few years of growth, and the 1cm-thick samples in the western Pacific deep-sea sediment core ERDC-92 integrate several hundred years because of the low sedimentation rate of 1.4 cm/kyr (Palmer and Pearson, 2003). Bioturbation probably smoothed the amplitude of the pH variations.

Variations of the sea surface pH are related to changes in surface water $p\text{CO}_2$, which is controlled by the CO_2 uptake from surface water through ocean-atmosphere CO_2 gas exchange. This gas exchange is modulated by the temperature, wind intensity, photosynthesis, lateral or vertical advection and mixing of different water masses. During the YD, the SSTs were slightly lower than the modern SSTs by $\sim 1^\circ\text{C}$ to $\sim 3^\circ\text{C}$ in the eastern Pacific (Kienast et al., 2006; Koutavas et al., 2002). These values were either similar (Palmer and Pearson, 2003) or lower by 3°C to 4.5°C in the western part of the Pacific (Beck et al., 1997; Corrège et al., 2004; Gagan et al., 2004) and lower by 1.5°C in the central part at Tahiti (Asami et al., 2009). If the SST dropped by $\sim 3^\circ\text{C}$ (25°C), the calculated pH would be about 8.08 rather than 8.05 because of the temperature dependence of pK_B (Dickson, 1990). As previously observed and especially for Tahiti (Fig. 4), a change in the SST of $\pm 1\text{--}3^\circ\text{C}$ has a moderate effect on the calculation of the pH.

This temperature effect cannot account for the calculated change of ~ 0.2 pH-units observed at the transition between the YD and the Holocene. Estimates of the pH from *Porites* collected on the Flinders reef system in the southwestern Pacific reveal inter-decadal changes of ~ 0.3 pH-units over the past 300 years (Pelejero et al., 2005). These changes were attributed to local effects caused by the large lagoon platform structure of the Flinders reef, the water $p\text{CO}_2$ build-up through coral calcification and the renewal rate of lagoon water by the South Equatorial current pulsed by the PDO (Pelejero et al., 2005). In the central equatorial Pacific, the impact of these local effects cannot be completely excluded. However, the probability of detecting the extreme changes of 0.2 pH-units from corals collected in the two distant islands Tahiti and Marquesas is low given that the samples only provide information for a few years in the

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Holocene, which began around 11.5 kyr BP ago. A recent sea surface pH- $\delta^{11}\text{B}$ study at the Fiji Islands showed a slight pH change because of the PDO (Douville et al., 2009). In addition, all ancient corals during the Holocene and until the last glacial period show relatively stable pH values close to 8.20–8.24, and 15 years of recent observations indicate a stable $\Delta p\text{CO}_2$ in this area (see below, Fig. 7). These observations indicate major and rapid changes in the Pacific oceanic patterns during the transition between the YD and Holocene.

The estimated pH drop near Tahiti and Marquesas at the end of the YD could be accounted for by an intensified upwelling and westward advection of cooler eastern water masses to the central Pacific. These eastern water masses are characterised today by low pHs and also low SSTs, especially for latitudes between 15° S and 20° S (Tahiti's latitudes, Fig. 2a). Sustained La Niña-like conditions operated at the end of the YD. That interpretation is supported by the presence of a strong east-west SST gradient in the equatorial Pacific around 11 500 years ago (Koutavas et al., 2006). The pH change from 8.05 to 8.23 at the YD-Holocene transition in the Marquesas area indicates a rapid change to more frequent El Niño-like climatic conditions in the equatorial south Pacific. Such these changes are also observed in the east-west Pacific SST reconstruction (Koutavas et al., 2006).

4.3 Estimates of the atmosphere-ocean $\Delta p\text{CO}_2$

In the central sub-equatorial Pacific (Tahiti and Marquesas), ten Holocene values of $p\text{CO}_2$ have a mean average of about $\sim 230 \pm 25$ ppmV for a pH of 8.23 ± 0.03 (SSS: 35.6; SST: 28 °C) (Table 1). At the end of the YD, the $p\text{CO}_2$ estimated for a pH of 8.05 was ~ 395 ppmV (SSS: 36.2; SST: 28 °C) or ~ 405 ppmV (SSS: 34.8; SST: 25 °C) when considering an intensified westward advection of the eastern Pacific water masses of low salinity and temperature. According to the past atmospheric $p\text{CO}_2$ from EPICA Dome C ice cores from Antarctica (Monnin et al., 2001, 2004), the ocean-atmosphere $\Delta p\text{CO}_2$ differences were between -67 and -11 ppmV during the Holocene, between

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



–9 and +56 ppmV before the YD with five positive values during the last deglaciation and +140 ppmV at the end of the YD ($p\text{CO}_2_{\text{atm}}=250$ ppmV). From similar calculations, the $\Delta p\text{CO}_2$ for the two modern corals (1950 AD, 1991 AD) from Tahiti are close to equilibrium (Fig. 6). This trend is in agreement with modern $\Delta p\text{CO}_2$ measurements (Fig. 7), which are either close to zero or slightly positive (Dandonneau, 1995; Feely et al., 1999, 2002; Goyet et al., 2009; Goyet and Peltzer, 1997).

Since the early Holocene until today, sea surface waters in the central sub-equatorial Pacific have mainly served as a moderate sink or source of CO_2 for the atmosphere in the vicinity of both Tahiti and the Marquesas Islands (Figs. 6 and 7). Our slight negative $\Delta p\text{CO}_2$ values agree with previous suggestions that more frequent El Niño-like oceanographic conditions occurred during the Holocene (Gagan et al., 2004; Koutavas et al., 2006). In contrast, the sub-equatorial sea surface waters in the central Pacific were a more pronounced source of CO_2 to the atmosphere during the last glacial/interglacial transition. This source of CO_2 strongly intensified around 11.5 kyr BP at the end of the YD and quickly reversed during the transition between the YD and the Holocene.

Such feature is connected first to the rapid rise of the atmospheric $p\text{CO}_2$ of ~15–20 ppmV (Fig. 6). This rise was observed quasi-synchronously in the EPICA Dome C ice core (EDC) (Monnin et al., 2001, 2004) by using refined chronology (Marchitto et al., 2007). Once the chronology of TLD and EDC ice cores is adjusted, this rise is also accompanied by a rapid increase of the atmospheric $\delta^{13}\text{CO}_2$ values. This connection was observed in the Taylor Dome ice core (TLD) in Antarctica (Smith et al., 1999) and more recently also in EPICA Dome C (Lourantou et al., 2010). Marchitto et al. (2007) showed that the ^{14}C ages of the intermediate waters in the Eastern Pacific decreased rapidly during YD-Holocene transition. These authors suggest that the aging of the intermediate waters was caused by northward penetration of the Antarctic Intermediate Water (AAIW) that was greater than the penetration occurring today (Marchitto et al., 2007), and the AAIW had old ^{14}C ages and depleted $\delta^{13}\text{C}$ values. The high $\Delta p\text{CO}_2$ at the end of the YD, that was deduced from the $\delta^{11}\text{B}$ -pH indicator provides further evidence of the marked changes in the properties of the water masses in the equa-

**Abrupt sea surface
pH change at the end
of the Younger Dryas**

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



torial Pacific and demonstrates the impact of these changes on the atmospheric CO₂ content.

5 Conclusions

The pH variations of the sea surface waters in the equatorial Pacific were determined from the isotopic composition of boron in the old *Porites* corals from reefs at Tahiti and Marquesas by MC-ICPMS analyses with an analytical uncertainty of ± 0.025 pH-units. Our $\delta^{11}\text{B}$, [B] and pH results for modern corals tend to verify the hypothesis of the unique incorporation of borate ions into the corals with the isotopic fractionation factor close to 0.981, in agreement with previous observations from laboratory calibration experiments. Modern aragonite *Porites* were sampled from different geographical areas with local SSTs ranging between 22.5°C and 27°C. The results reveal that the temperature strongly controls both the partition constant K_D and the incorporation of boron in various aragonite coral species. Boron is incorporated more in the aragonite skeleton at lower SST conditions. The $\delta^{11}\text{B}$ -pH values were close to ~ 8.23 during the Holocene in the surface waters of both Marquesas and the Tahiti Islands. This result is in agreement with the modern or pre-anthropogenic values. The $\delta^{11}\text{B}$ -pH values ranged from 8.15 to 8.31 during the last deglaciation, and the highest value measured was at 20.7 kyr BP

An abrupt pH drop to ~ 8.05 was observed at the end of the Younger Dryas around 11.5 kyr BP. The deduced $p\text{CO}_2$ values were relatively equilibrated with the atmosphere during the Holocene and the last deglaciation, but during this event, the $p\text{CO}_2$ increased to up to ~ 400 ppmV near both Tahiti and the Marquesas islands. The sea surface waters in the central equatorial Pacific (Tahiti, Marquesas) were a sink or a moderate source of CO₂ to the atmosphere during the Holocene. They were a more important source during the last glacial/interglacial transition, and they were a major source at the end of the YD with a $\Delta p\text{CO}_2$ of at least +140 ppmV. This last finding provides further evidence of the marked changes to the properties of the water masses

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in the equatorial Pacific (Marchitto et al., 2007) and demonstrates the impact of these changes on the atmospheric CO₂ content. These results highlight the great potential of the “ $\delta^{11}\text{B-pH}$ ” methodology to precisely reconstruct past oceanic fluxes of CO₂ to the atmosphere from well-dated sea surface corals such as *Porites* and *Acropora*.

5 *Acknowledgements.* We are greatly indebted to H el ene Isnard and Fr ed eric Chartier of the CEA laboratory LANIE in Gif/Yvette (France) and Claudia Bouman of the Advanced Mass Spec-

10 trometry Laboratory of Thermo Fisher Scientific in Bremen for our fruitful discussions and the use of their instruments during an early stage of the work presented here. We are also grateful to Norbert Frank and Eline Sall e for providing accurate U-series dates for the corals and to

15 Michel Fontugne for constructive discussions. This work received the financial support of the national INSU LEFE/CYBER project PHARE, the Commissariat   l’Energie Atomique (CEA), the French Centre National de la Recherche Scientifique (CNRS) and the Institut de Recherche pour le d eveloppement (IRD). Our thanks are extended to John Butscher for his help collecting the modern coral samples in New Caledonia and to the Province Sud of New Caledonia for the permit to collect these corals. We are grateful to David Varillon from IRD “IMAGO”, LEGOS laboratory, IRD Noum ea and the French Zon eco program for providing the Thermosalinograph datasets monitored at Utoe (New Caledonia). Gratitude is expressed to the captain, R. Proner and the crew of the R/V IRD “Alis” for their efficiency during the cruises in Marquesas.



The publication of this article is financed by CNRS-INSU.

References

Archer, D., Winguth, A., Lea, D., and Mahowald, N.: What caused the Glacial/Interglacial atmospheric $p\text{CO}_2$ cycles?, *Rev. Geophys.*, 38(2), 159–189, 2000.

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Asami R., Felis T., Deschamps P., Hanawa K., Iryu Y., Bard E., Durand M., and Murayama M.: Evidence for tropical South Pacific climate change during the Younger Dryas and the Bølling-Allerød from geochemical records of fossil Tahiti corals, *Earth Planet. Sci. Lett.*, 288, 96–107, 2009.

5 Bard, E., Hamelin, B., Arnold, M., Montaggioni, L., Cabioch, G., Faure, G., and Rougerie, F.: Deglacial sea-level record from Tahiti corals and the timing of global melt water discharge, *Nature*, 382, 241–244, 1996.

Beck, J. W., Récy, J., Taylor, F., Edwards, R. L., and Cabioch, G.: Abrupt changes in early Holocene tropical sea surface temperature derived from coral records, *Nature*, 385, 705–707, 1997.

10 Boiseau, M.: Etude de la variabilité climatique liée à l'ENSO dans l'Océan Pacifique Central Sud (Moorea) à partir de traceurs géochimiques contenus dans le squelette d'un Scleractiniaire à zooxanthelles (*Porites Lutea*), pH'D thesis, University of Paris VII, Paris, 202 pp., 1998.

15 Cabioch, G., Montaggioni, L., Frank, N., Seard, C., Sallé, E., Payri, C., Pelletier, B., and Paternite, M.: Successive reef depositional events along the Marquesas foreslopes (French Polynesia) since 26 ka, *Mar. Geol.*, 254, 18–34, 2008.

Caldeira, K. and Wickett, M. E.: Anthropogenic carbon and ocean pH, *Nature*, 425, p. 365, 2003.

20 Corrège, T., Gagan, M. K., Beck, J. W., Burr, G. S., Cabioch, G., and Le Cornec, F.: Interdecadal variation in the extent of South Pacific tropical waters during the Younger Dryas event, *Nature*, 428, 927–929, 2004.

Dandonneau, Y.: Sea-surface partial pressure of carbon dioxide in the eastern equatorial Pacific (August 1991 to October 1992): A multivariate analysis of physical and biological factors, *Deep-Sea Res. Pt II*, 42(2–3), 349–364, 1995.

25 Delcroix, T., Henin, C., Porte, V., and Arkin, P.: Precipitation and sea surface salinity in the tropical Pacific Ocean, *Deep-Sea Res. I*, 73(7), 1123–1141, 1996.

Dickson, A. G.: Thermodynamic of the dissociation of boric acid in synthetic seawater from 273.15 to 318.15 K, *Deep-Sea Res.*, 37, 755–766, 1990.

30 Douville, E., Copard, K., Colin, C., and Frank, N.: Major and trace elements in deep sea corals *Lophelia pertusa* from the eastern North Atlantic, ICP-9, Shanghai, China, 3–7 September 2007, P1-28, 45, 2007.

Douville, E., Juillet-Leclerc, A., Cabioch, G., Louvat, P., Gaillardet, J., Gehlen, M., Bopp, L.,

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Paterne, M.: Boron isotopes in Fiji corals and precise ocean acidification reconstruction, AGU Fall Meeting, San Francisco, California, USA, 14–18 December 2009, GC24A-04, 2009.

Fallon, S. J., McCulloch, M. T., and Alibert, C.: Examining water temperature proxies in *Porites* corals from the Great Barrier Reef: a cross-shelf comparison, *Coral Reefs*, 22, 389–404, 2003.

Feely, R. A., Boutin, J., Cosca, C. E., Dandonneau, Y., Etcheto, J., Inoue, H. Y., Ishii, M., Le Quéré, C., Mackey, D. J., McPhaden, M., Metzl, N., Poisson, A., and Wanninkhof, R.: Seasonal and interannual variability of CO₂ in the equatorial Pacific, *Deep-Sea Res. Pt II*, 49, 2443–2449, 2002.

Feely, R. A., Wanninkhof, R., Takahashi, T., and Tans, P.: Influence of El Niño on the equatorial Pacific contribution to atmospheric CO₂ accumulation, *Nature*, 398, 597–601, 1999.

Foster, G. L.: Seawater pH, pCO₂ and [CO₃²⁻] variations in the Caribbean Sea over the last 130 kyr: A boron isotope and B/Ca study of planktic foraminifera, *Earth Planet. Sci. Lett.*, 271, 254–266, 2008.

Gagan, M. K., Hendy, E. J., Haberle, S. G., and Hantoro, W. S.: Post-glacial evolution of the Indo-Pacific Warm Pool and El Niño-Southern oscillation, *Quaternary Int.*, 118–119, 127–143, 2004.

Gaillardet, J. and Allègre, C. J.: Boron isotopic compositions of corals: seawater or diagenesis record?, *Earth Planet. Sci. Lett.*, 136, 665–676, 1995.

Gertman, I. and Brenner, S.: Analysis of water temperature variability in the Gulf of Eilat, Report of Israel Oceanogr. Limnol. Res., 13 pp., 2004.

Goyet, C., Ito Gonçalves, R., and Touratier, F.: Anthropogenic carbon distribution in the eastern South Pacific Ocean, *Biogeosciences*, 6, 149–156, 2009, <http://www.biogeosciences.net/6/149/2009/>.

Goyet, C. and Peltzer, E. T.: Variation of CO₂ partial pressure in surface seawater in the equatorial Pacific Ocean, *Deep-Sea Res. Pt. I*, 44(9–10), 1611–1625, 1997.

Heiss, G. A., Dullo, W. C., Joachimski, M. M., Reijmer, J. J. G., and Schumacher, H.: Increased Seasonality in the Gulf of Aqaba, Red Sea, Recorded in the Oxygen Isotope Record of a *Porites lutea* Coral, *Senck. Marit.*, 30, 17–26, 1999.

Hemming, N. G., Guilderson, T. P., and Fairbanks, R. G.: Seasonal variations in the boron isotopic composition of coral: A productivity signal?, *Global Biogeochem. Cy.*, 12(4), 581–586, 1998.

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Hemming, N. G. and Hanson, G. N.: Boron isotopic composition and concentration in modern marine carbonates, *Geochim. Cosmochim. Ac.*, 56, 537–543, 1992.
- Hemming, N. G., Reeder, R. J., and Hanson, G. N.: Mineral-fluid partitioning and isotopic fractionation of boron in synthetic calcium carbonate, *Geochim. Cosmochim. Ac.*, 59(2), 371–379, 1995.
- Henin, C.: Surveillance Thermohaline de la ZEE en 1997 et 1998, Zoneco Report, IRD, Noumea, New Caledonia, 16 pp., 1999.
- Henin, C. and Cresswell, G. R.: Upwelling along the western barrier reef of New Caledonia, *Mar. Freshwater Res.*, 56(7), 1005–1010, 2005.
- Hönisch, B. and Hemming, N. G.: Surface ocean pH response to variations in $p\text{CO}_2$ through two full glacial cycles, *Earth Planet. Sci. Lett.*, 236, 305–314, 2005.
- Hönisch, B., Hemming, N. G., Archer, D., Siddall, M., and McManus, J. F.: Atmospheric Carbon Dioxide Concentration Across the Mid-Pleistocene Transition, *Science*, 324, 1551–1554, 2009.
- Hönisch, B., Hemming, N. G., Grottoli, A. G., Amat, A., Hanson, G. N., and Bijma, J.: Assessing scleractinian corals as recorders for paleo-pH: Empirical calibration and vital effects, *Geochim. Cosmochim. Ac.*, 68(18), 3675–3685, 2004.
- Hönisch, B., Hemming, N. G., and Loose, B.: Comment on “A critical evaluation of the boron isotope-pH proxy: The accuracy of ancient ocean pH estimates” by M. Pagani, D. Lernerhand, A. Spivack and J. Gaillardet, *Geochim. Cosmochim. Ac.*, 71, 1636–1641, 2007.
- Inoue, H. Y., Ishii, M., Matsueda, H., Saito, S., Aoyama, M., Tokieda, T., Midorikawa, T., Nemoto, K., Kawano, T., Asanuma, I., Ando, K., Yano, T., and Murata, A.: Distributions and variations in the partial pressure of CO_2 in surface waters ($p\text{CO}_{2w}$) of the central and western equatorial Pacific during the 1997/1998 El Niño event, *Mar. Chem.*, 76(1–2), 59–75, 2001.
- Inoue, M., Nohara, M., Okai, T., Suzuki, A., and Kawahata, H.: Concentrations of Trace Elements in Carbonate Reference Materials Coral JCp-1 and Giant Clam JCt-1 by Inductively Coupled Plasma-Mass Spectrometry, *Geostand. Geoanal. Res.*, 28(3), 411–416, 2004.
- IPCC: Carbon dioxide Capture and Storage, Intergovernmental Panel on Climate Change Special Report, Editors: Bert Metz, Ogunlade Davidson Heleen de Coninck, Manuela Loos, Leo Meyer, 2005.
- Kakihana, H., Kotoka, M., Satoh, S., Nomura, M., and Okamoto, M.: Fundamental studies on the ion-exchange separation of boron isotopes, *Bull. Chem. Soc. J.*, 50, 158–163, 1977.
- Kienast, M., Kienast, S. S., Calvert, S. E., Eglinton, T. I., Mollenhauer, G., François, R., and Mix,

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Abrupt sea surface
pH change at the end
of the Younger Dryas**

E. Douville et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A. C.: Eastern Pacific cooling and Atlantic overturning circulation during the last deglaciation, *Nature*, 443, 846–849, 2006.

Klochko, K., Kaufman, A. J., Yao, W., Byrne, R. H., and Tossell, J. A.: Experimental measurement of boron isotope fractionation in seawater, *Earth Planet. Sci. Lett.*, 248, 276–285, 2006.

Kohfeld, K. E., Le Quere, C., Harrison, S. P., and Anderson, R. F.: Role of marine biology in glacial-interglacial CO₂ cycles, *Science*, 38, 74–78, 2005.

Koutavas, A., deMenocal, P. B., Olive, G. C., and Lynch-Stieglitz, J.: Mid-Holocene El Niño-Southern Oscillation (ENSO) attenuation revealed by individual foraminifera in eastern tropical Pacific sediments, *Geology*, 34(12), 993–996, 2006.

Koutavas, A., Lynch-Stieglitz, J., Marchitto, T. M., and Sachs, J. P.: El Niño-like pattern in ice age tropical Pacific sea surface temperature, *Science*, 297, 226–230, 2002.

Le Quéré, C., Rödenbeck, C., Buitenhuis, E. T., Conway, T. J., Langenfelds, R., Gomez, A., Labuschagne, C., Ramonet, M., Nakazawa, T., Metzl, N., Gillett, N., and Heimann, M.: Saturation of the Southern Ocean CO₂ sink due to recent climate change, *Science*, 316, 5832, 1735–1738, doi:10.1126/science.1136188, 2007.

Lécuyer, C., Grandjean, P., Reynard, B., Albarède, F., and Telouk, P.: ¹¹B/¹⁰B analysis of geological materials by ICP-MS Plasma 54: Application to the boron fractionation between brachiopod calcite and seawater, *Chem. Geol.*, 186, 45–55, 2002.

Lefèvre, N., Andrié, C., Dandonneau, Y., Reverdin, G., and Rodier, M.: pCO₂, chemical properties, and estimated new production in the equatorial Pacific in January–March 1991, *J. Geophys. Res.*, 99, 12639–12654, 1994.

Lewis, E. and Wallace, D. W. R.: Program developed for CO₂ system calculations, ORNL/CDIAC-105, Carbon dioxide Inf. Anal. Cent. Oak Ridge Nat. Lab., U.S. Dep. of Energy, Oak Ridge, Tenn, USA, 1998.

Liu, Y., Liu, W., Peng, Z., Xiao, Y., Wei, G., Sun, W., He, J., Liu, G., and Chou, C. L.: Instability of seawater pH in the South China Sea during the mid-late Holocene: Evidence from boron isotopic composition of corals, *Geochim. Cosmochim. Ac.*, 73, 1264–1272, 2009.

Lourantou, A., Lavric, J. V., Köhler, P., Barnola, J. M., Paillard, D., Michel, E., Raynaud, D., and Chappellaz, J.: New atmospheric carbon isotopic measurements constrain the CO₂ rise during the last deglaciation, *Global Biogeochem. Cy.*, doi:10.1029/2009GB003545, in press, 2010.

Louvat, P., Bouchez, J., and Paris, G.: MC-ICP-MS isotope measurements with direct injection

nebulization (d-DIHEN): optimization and application to boron in seawater and carbonate samples, *Geostand. Geoanal. Res.*, in press, 2010.

Marchitto, T. M., Lehman, S. J., Ortiz, J. D., Flückiger, J., and vanGeen, A.: Marine radiocarbon evidence for the mechanism of deglacial atmospheric CO₂ rise, *Science*, 316, 1456–1459, 2007.

McGillicuddy, D. J., Anderson, L. A., Bates, N. R., Bibby, T., Buesseler, K. O., Carlson, C. A., Davis, C. S., Ewart, C., Falkowski, P. G., Goldthwait, S. A., Hansell, D. A., Jenkins, W. J., Johnson, R., Kosnyrev, V. K., Ledwell, J.R., Li, Q. P., Siegel, and D. A., and Steinberg, D.K.: Eddy/wind interactions stimulate extraordinary mid-ocean plankton blooms, *Science*, 316, 1021–1026, 2007.

Monnin, E., Indermühle, A., Dällenbach, A., Flückiger, J., Stauffer, B., Stocker, T. F., Raynaud, D., and Barnola, J. M.: Atmospheric CO₂ concentrations over the last glacial termination, *Science*, 291, 112–114, 2001.

Monnin, E., Steig, E. J., Siegenthaler, U., Kawamura, K., Schwander, J., Stauffer, B., Stocker, T. F., Morse, D. L., Barnola, J. M., Bellier, B., Raynaud, D., and Fischer, H.: Evidence for substantial accumulation rate variability in Antarctica during the Holocene, through synchronization of CO₂ in the Taylor Dome, Dome C and DML ice cores, *Earth Planet. Sci. Lett.*, 224, 45–54, 2004.

Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G. K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M. F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, 2005.

Pagani, M., Lemarchand, M. D., Spivack, A., and Gaillardet, J.: A critical evaluation of the boron isotope-pH proxy: The accuracy of ancient ocean pH estimates, *Geochim. Cosmochim. Ac.*, 69(4), 953–961, 2005.

Palmer, M. R. and Pearson, P. N.: A 23000 -Year record of surface water pH and pCO₂ in the Western Equatorial Pacific Ocean, *Science*, 300, 480–482, 2003.

Paterne, M., Ayliffe, L. K., Arnorld, M., Cabioch, G., Tisnérat-Laborde, N., Hatté, C., Douville, E., and Bard, E.: Paired ¹⁴C and ²³⁰Th dating of surface corals from the Marquesas and Vanuatu (Sub-Equatorial Pacific) in the 3000 to 15000 cal. yr interval, *Radiocarbon*, 46(2), 551–566, 2004.

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Abrupt sea surface
pH change at the end
of the Younger Dryas**

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Pelejero, C., Calco, E., McCulloch, M. T., Marshall, J. F., Gagan, M. K., Lough, J. M., and Opdyke, B. N.: Preindustrial to Modern interdecadal variability in Coral Reef pH, *Science*, 309, 2204–2207, 2005.

Reynaud, S., Hemming, N. G., Juillet-Leclerc, A., and Gattuso, J. P.: Effect of $p\text{CO}_2$ and temperature on the boron isotopic composition of the zooxanthellate coral *Acropora* sp., *Coral Reefs*, 23, 539–546, 2004.

Roy, R. N., Roy, L. N., Vogel, K. M., Porter-Moore, C., Pearson, T., Good, C. E., Millero, F. J., and Campbell, D. M.: The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45 °C, *Mar. Chem.*, 44, 249–267, 1993.

Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T. H., Kozyr, A., Ono, T., and Rios, A. F.: The Oceanic Sink for Anthropogenic CO_2 , *Science*, 305, 367–371, 2004.

Sanchez-Valle, C., Reynard, B., Daniel, I., Lécuyer, C., Martinez, I., and Chervin, J. C.: Boron isotopic fractionation between minerals and fluids: new insights from in situ high pressure-high-temperature vibrational spectroscopic data, *Geochim. Cosmochim. Ac.*, 69, 4301–4313, 2005.

Sanyal, A., Hemming, N. G., Broecker, W. S., Lea, D. W., Spero, H. J., and Hanson, G. N.: Oceanic pH control on the boron isotopic composition of foraminifera: Evidence from culture experiments, *Paleoceanography*, 11(5), 513–517, 1996.

Sanyal, A., Nugent, M., Reeder, R. J., and Bijma, J.: Seawater pH control on the boron isotopic composition of calcite: Evidence from inorganic calcite precipitation experiments, *Geochim. Cosmochim. Ac.*, 64(9), 1551–1555, 2000.

Sigman, D. M. and Boyle, E. A.: Glacial/interglacial variations in atmospheric carbon dioxide, *Nature*, 407, 859–869, 2000.

Smith, H. J., Fischer, H., Wahlen, M., Mastroianni, D., and Deck, B.: Dual modes of the carbon cycle since the Last Glacial Maximum, *Nature*, 400, 248–250, 1999.

Spivack, A. J., You, C. F., and Smith, H. J.: Foraminiferal boron isotope ratios as a proxy for surface ocean pH over the past 21 Myr, *Nature*, 363, 149–151, 1993.

Taft, B. A. and Kessler, W. S.: Variations of zonal currents in the central tropical Pacific during 1970 to 1987: sea-level and dynamic height measurements, *J. Geophys. Res.*, 96, 12599–12618, 1991.

Wei, G., McCulloch, M. T., Mortimer, G., Deng, W., and Xie, L.: Evidence for ocean acidification in the Great Barrier Reef of Australia, *Geochim. Cosmochim. Ac.*, 73, 2332–2346, 2009.

Xiao, Y. K., Shirodkar, P. V., Zhang, C. G., Wei, H. Z., Liu, W. G., and Zhou, W. J.: Isotopic fractionation of boron in growing corals and its palaeoenvironmental implication, *Curr. Sci. India*, 90(3), 414–420, 2006.

5 Yu, J., Elderfield, H., and Hönisch B.: B/Ca in planktonic foraminifera as a proxy for surface seawater pH, *Paleoceanography*, 22, PA2202, doi:10.1029/2006PA001347, 2007.

Zeebe, R. E.: Stable boron isotope fractionation between dissolved $B(OH)_3$ and $B(OH)_4^-$, *Geochim. Cosmochim. Ac.*, 69(11), 2753–2766, 2005.

BGD

7, 1959–1993, 2010

**Abrupt sea surface
pH change at the end
of the Younger Dryas**

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Isotopic composition of boron $\delta^{11}\text{B}$ (‰) and boron concentrations measured for coral and seawater samples. The ΔpCO_2 calculation is detailed in the text. The previously published $\delta^{11}\text{B}$ data for corals includes (a) (Gaillardet and Allègre, 1995); (b) (Hemming et al., 1998); and (c) (Pelejero et al., 2005). The $^{230}\text{Th}/\text{U}$ ages of the corals from Tahiti and Marquesas are from Bard et al., 1996; Cabioch et al., 2008; Paterne et al., 2004. * Graphically estimated from Gaillardet and Allègre, 1995. SST sources: (<http://www.nodc.noaa.gov>; Boiseau, 1998; Gertman and Brenner, 2004; Heiss et al., 1999; Henin and Cresswell, 2005).

Samples, mean SST	$^{230}\text{Th}/\text{U}$ age cal. B.P.	LSCE number	Boron (ppm)	$\delta^{11}\text{B}_{\text{seawater}}$ (‰)	Std Dev. (2 σ , n)	pH_{seaw}	pCO_2 (ppmV)	ΔpCO_2 (ppmV)
Marquesas Islands, 28 ± 1.0°C								
Nuku Hiva DR6(1)	250 ± 30	B33*	40.0	26.2	± 0.2 (2)	8.22 ± 0.02	235	-55
DW1281c_78a1	3 230 ± 30	B01, 02, 03, 04, B20*	63.2	26.6	± 0.2 (5)	8.26 ± 0.02	207	-67
DW1281_68a1	3 260 ± 30	B15, B16, B18*	60.0	26.2	± 0.2 (7)	8.22 ± 0.02	235	-39
Eiao DR16(3)	9 110 ± 130	B31*	37.4	26	± 0.3 (2)	8.20 ± 0.03	250	-11
Eiao DR16(5)	9 110 ± 130	B32*	53.4	26.3	± 0.5 (2)	8.23 ± 0.05	229	-35
Eiao DR12(1)	9 590 ± 180	B30*	43.5	26.2	± 0.2 (3)	8.21 ± 0.02	240	-24
DW1281_75a2	11,470 ± 90	B13, B14, B19**	45.3	24.7	± 0.2 (6)	8.05 ± 0.02	394	138
Hiva Oa DR10(2)	12,420 ± 100	B27*	51.9	26.2	± 0.2 (3)	8.21 ± 0.02	239	1
Eiao DR11bs(4)	13,410 ± 190	B53*	n. d.	25.6	± 0.3 (1)	8.15 ± 0.03	294	56
Eiao DR8(1)	14,560 ± 180	B29*	42.1	26.1	± 0.2 (2)	8.20 ± 0.02	251	12
Hiva Oa DR14bis(1)	15,450 ± 150	B28*	40.0	26.1	± 0.3 (3)	8.21 ± 0.03	247	23
Hiva Oa DR8bis(1)	15,460 ± 110	B26*	43.8	26.4	± 0.2 (2)	8.24 ± 0.02	224	1
Hiva Oa DR5	20,720 ± 200	B25*	63.6	27.1	± 0.3 (3)	8.31 ± 0.03	180	-9
Moorea-Tahiti Islands, 27.1 ± 0.5°C								
Porites:								
COM2 *	1991 (AD)		51.0	25.3	± 0.2 (3)	8.13 ± 0.02	309	-45
MOO 3A-1-02	1950 (AD)	B11, B12, B17**	51.9	25.8	± 0.2 (6)	8.18 ± 0.02	265	-45
Ta P8-348	12,910 ± 30	B56*	n. d.	25.9	± 0.3 (3)	8.20 ± 0.03	256	22
Ta P8-353	13,335 ± 30	B57*	n. d.	26.6	± 0.2 (3)	8.27 ± 0.02	203	-35
Acropora*:								
Ta-1	8 520 ± 40		50.0	25.7	± 0.3 (2)	8.24 ± 0.03	223	-38
Ta-2	9 260 ± 50		49.0	25.4	± 0.2 (3)	8.21 ± 0.02	244	-17
Ta-3	9 850 ± 40		51.0	25.6	± 0.2 (2)	8.23 ± 0.02	231	-33
Ta-4	10,250 ± 40		52.0	25.9	± 0.3 (1)	8.26 ± 0.03	210	-58
Ta-5	10,575 ± 50		41.0	24.6	± 0.3 (1)	8.12 ± 0.03	323	57
Ta-6	10,850 ± 50		50.0	26.6	± 0.2 (2)	8.32 ± 0.02	171	-94
Ta-7	11,000 ± 40		50.0	25.3	± 0.2 (2)	8.19 ± 0.02	259	-5
Ta-8	11,290 ± 30		41.0	24.4	± 0.2 (3)	8.09 ± 0.02	351	87
Ta-9	11,495 ± 30		41.0	23.6	± 0.2 (3)	7.99 ± 0.02	466	212
Ishigaki Island, 26.1 ± 2.1°C								
Porites JcP-1 (powder)	modern	9 samplings	47.7	24.5	± 0.1 (15)	8.05 ± 0.01		
New Caledonia, 24.4 ± 1.9°C								
NC_a	2001-2002 (AD)	B84*, B85*	55.0	24.3	± 0.2 (2)	8.04 ± 0.02		
NC_b	2002-2003 (AD)	B86*, B87*	54.6	24.5	± 0.6 (3)	8.05 ± 0.06		
NC_c	2003-2004 (AD)	B89*, B90*, B91*, B92*	54.4	24.8	± 0.5 (4)	8.10 ± 0.05		
Red Sea*, 22.5 ± 2.0°C								
COM3	1970-80 (AD)		57.0	23.7	± 0.5 (2)	7.98 ± 0.05		
Fanning Island*, 25-28°C								
Porites Lobata	mean	Seasonal	50.5	24.8	-	8.07		
	Two year range	microsampling	46.1 - 53.9	23.9 - 26.2	-	7.96-8.23		
Finders Reef*								
FLO2A	1990s (AD)		-	23	-	- 7.9		
	1950s (AD)		-	24.5	-	- 8.1		
	1700 - 2000 (AD)		-	23.0 - 25.0	-	7.9 - 8.1		

**Abrupt sea surface
pH change at the end
of the Younger Dryas**

E. Douville et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.




Fig. 1. Geographical locations of the study sites, including Moorea (close to Tahiti), Marquesas, Ishigaki, New Caledonia and the Fanning Islands. The location of the ERDC-92 box core (Palmer and Pearson, 2003) is also shown. The sea surface pH from modern corals are plotted on a seawater scale pH map modified from (Pelejero et al., 2005). The $\delta^{11}\text{B}$ data of corals: **(a)** (Gaillardet and Allègre, 1995); **(b)** (Hemming et al., 1998); **(c)** (Pelejero et al., 2005); the foraminifera $\delta^{11}\text{B}$ data: **(d)** (Palmer and Pearson, 2003).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.




Fig. 2. (a) The modern sea surface pH in the sub-Equatorial Pacific Ocean as function of the geographical location. 0.07 pH-units are added to the modern pH on the pre-industrial era pH scale (see text). The month (values in parenthesis) of the water sampling and the averaged SSTs are also indicated. (b) Sea surface pH in the sub-Equatorial Pacific Ocean plotted against SST between 15° E and 80° W. Similar pH-SST correlations were observed for surface waters between 80° W and 170° W at three different ranges of latitude (sources from <http://www.nodc.noaa.gov/OC5/SELECT/dbsearch/dbsearch.html>; measurement years: 1980–1998; depths: 0–50 m).

Abrupt sea surface
pH change at the end
of the Younger Dryas

E. Douville et al.




Fig. 3. The $\delta^{11}\text{B}$ values plotted against the boron concentrations **(a)** for modern *Porites* from various geographical areas and **(b)** for ancient *Porites* and *Acropora* near Marquesas and the Tahiti Islands. The theoretical curves for different partition coefficient K_D values were calculated assuming $T = 28^\circ\text{C}$; $S = 35\text{‰}$; $\alpha = 0.981$; and $\delta^{11}\text{B}_{\text{SW}} = 39.9\text{‰}$. Seawater values used were $[\text{B}]_{\text{seawater}} = 416 \mu\text{M}$ and $\text{TCO}_2 = 2 \text{mM}$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.




Fig. 4. B/Ca molar ratio in coral plotted against the borate/bicarbonate molar ratio in seawater. Theoretical curves for different partition coefficient K_D values were calculated assuming $T = 28^\circ\text{C}$; $S = 35\%$; $\alpha = 0.981$; and $\delta^{11}\text{B}_{\text{SW}} = 39.9\%$. Regional SSTs were used to calculate the borate/bicarbonate ratios for each of the corals. The temperature strongly influences the partition coefficient K_D . *Porites* near Marquesas showed a K_D of ~ 0.0057 . The seawater values used to calibrate the partition coefficient were $[\text{B}]_{\text{seawater}} = 416 \mu\text{M}$ and $\text{TCO}_2 = 2 \text{ mM}$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abrupt sea surface
pH change at the end
of the Younger Dryas

E. Douville et al.




Fig. 5. Linear correlation between the mean regional SST and the partition coefficient K_D obtained from modern *Porites* samples collected from various SST areas. This regression shows the influence of the temperature on K_D and is valid for various aragonite species. The tropical corals are *Porites*, *Acropora*, and the cold sea coral is *Lophelia pertusa*. The expected area for cold sea corals (7–11°C) was determined from pH 7.95 and boron concentrations measured in North Atlantic *Lophelia pertusa* samples collected between 40° N and 70° N (Douville et al., 2007).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. Comparison of the sub-Equatorial Pacific $\delta^{11}\text{B}$ -pH of marine biogenic carbonates over the last 21 000 cal. yr. Sources are (1) this study; (2) Gaillardet and Allègre (1995); and (3) Palmer and Pearson (2003). The calculation parameters for the pH are $\alpha = 0.981$; $\delta^{11}\text{B}_{\text{SW}} = 39.9\text{‰}$; $S = 35\text{‰}$; and $T = 27^\circ\text{C}$ for Tahiti (Moorea) and $T = 28^\circ\text{C}$ for the Marquesas Islands and for ERDC-92. Deduced $\Delta p\text{CO}_2$ values (Table 1, see text) and atmospheric $p\text{CO}_2$ data (Monnin et al., 2001, 2004) are also plotted to illustrate the synchronization between the abrupt changes to the sea surface pH and $\Delta p\text{CO}_2$ in the central sub-Equatorial Pacific and the atmospheric CO_2 increase of 15–20 ppmV at the end of the YD event.

1992

BGD

7, 1959–1993, 2010

Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abrupt sea surface pH change at the end of the Younger Dryas

E. Douville et al.




Fig. 7. The relative stability of modern $\Delta p\text{CO}_2$ values close to equilibrium in the Tahiti – Marquesas area between 1992 AD and 2004 AD (Dandonneau, 1995; Feely et al., 2002; Feely et al., 1999; Goyet et al., 2009; Goyet and Peltzer, 1997).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

