



This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Fueling primary productivity: nutrient return pathways from the deep ocean and their dependence on the Meridional Overturning Circulation

J. B. Palter¹, J. L. Sarmiento¹, A. Gnanadesikan², J. Simeon³, and D. Slater¹

¹Atmospheric and Oceanic Sciences Program, Princeton University, Princeton, New Jersey, USA

²Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey, USA

³Laboratoire des Sciences du Climat et de l'Environnement, UMR CEA-CNRS, CEA Saclay, 91191 Gif-sur-Yvette, France

Received: 9 May 2010 – Accepted: 11 May 2010 – Published: 1 June 2010

Correspondence to: J. B. Palter (jpalter@princeton.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

7, 4045–4088, 2010

Nutrient return pathways

J. B. Palter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

In the Southern Ocean, mixing and upwelling in the presence of heat and freshwater surface fluxes transform subpycnocline water to lighter densities as part of the upward branch of the Meridional Overturning Circulation (MOC). One hypothesized impact of this transformation is the restoration of nutrients to the global pycnocline, without which biological productivity at low latitudes would be catastrophically reduced. Here we use a novel set of modeling experiments to explore the causes and consequences of the Southern Ocean nutrient return pathway. Specifically, we quantify the contribution to global productivity of nutrients that rise from the ocean interior in the Southern Ocean, the northern high latitudes, and by mixing across the low latitude pycnocline. In addition, we evaluate how the strength of the Southern Ocean winds and the parameterizations of subgrid-scale processes change the dominant nutrient return pathways in the ocean. Our results suggest that nutrients upwelled from the deep ocean in the Antarctic Circumpolar Current and subducted in Subantarctic Mode Water support between 33 and 75% of global primary productivity between 30° S and 30° N. The high end of this range results from an ocean model in which the MOC is driven primarily by wind-induced Southern Ocean upwelling, a configuration favored due to its fidelity to tracer data, while the low end results from an MOC driven by high diapycnal diffusivity in the pycnocline. In all models, the high preformed nutrients subducted in the SAMW layer are converted rapidly (in less than 40 years) to remineralized nutrients, explaining previous modeling results that showed little influence of the drawdown of SAMW surface nutrients on atmospheric carbon concentrations.

1 Introduction

Primary productivity in the ocean's sunlit surface incorporates dissolved nutrients into organic matter, a fraction of which sinks to the ocean interior. Once in the ocean's dark interior, the organic matter is respired back to dissolved inorganic carbon and nutrients,

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the ACC's equatorward fringes that influences tracer properties throughout much of the global pycnocline (McCartney, 1982; Sarmiento et al., 2004; Hanawa and Talley, 2001). This conceptual model hinges on the upwelled nutrients being subducted with the SAMW at the time it forms, rather than being utilized and exported from it, as can be 5 the case for other mode waters (Palter et al., 2005). Indeed, the SAMW is formed in a region with highly elevated surface nutrients (Fig. 1a), and it tends to retain its nutrients during formation and subduction. Evidence supporting the dominance of the Southern Ocean nutrient upwelling pathway has been obtained by model sensitivity studies, in 10 which the strength of global primary productivity was critically reduced by completely stripping nutrients from the surface ocean in the SAMW formation region (Sarmiento et al., 2004; Marinov et al., 2006).

Additional support for the importance of a Southern Ocean upward MOC branch in influencing pycnocline nutrient concentrations comes from looking at quasi-conservative combinations of nutrient tracers. For example, in the Subantarctic Zone 15 where SAMW is formed, phosphate (PO_4^{3-}) and nitrate (NO_3^-) are abundant, while silicic acid (Si(OH)_4) is nearly depleted. Thus, a useful tracer of the SAMW is $\text{Si}^*=\text{Si(OH)}_4\text{-NO}_3^-$ (Sarmiento et al., 2004). The lowest Si^* concentrations in the global surface ocean are found in the SAMW formation region (Fig. 1b). Sarmiento et al. (2004) argue that because Si^* is approximately conserved in the ocean interior, 20 the subsurface pool of negative Si^* indicates the spreading of SAMW via advective and diffusive transport processes throughout much of the low latitude ocean (Fig. 1c).

Despite the apparent convergence of evidence from models and the Si^* tracer towards a central role for a Southern Ocean nutrient return pathway from the deep ocean to the pycnocline, several important questions still require evaluation and quantification.

25 What is the fraction of low latitude productivity sustained by the Southern Ocean nutrient return pathway and which water masses are involved? In which regions are the SAMW nutrients most important, and what are the pathways by which the SAMW arrives at these downstream locales? Here we examine these questions using a suite of ocean general circulation models (OGCMs). A second class of questions concerns

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the sensitivity of the predominant nutrient return pathway to the models' parameterizations of subgridscale processes and the prescribed atmospheric forcing that drives the ocean circulation, particularly the Southern Ocean winds. The fraction of deep upwelling that occurs in the Southern Ocean is sensitive to the strength of the winds, 5 as well as the turbulent mixing and transport due to mesoscale eddies, which are parameterized in the coarse GCMs typically used to simulate climate and biogeochemistry. The sensitivity of the modeled MOC to winds and subgridscale processes has been demonstrated in more than one idealized analytical model (Gnanadesikan, 1999; Samelson, 2004, 2009), and tested in a suite of GCMs (Gnanadesikan et al., 2007).
10 Here we quantify the sensitivity of the magnitude and distribution of nutrient supply to wind forcing and the parameterizations of subgridscale processes.

We also consider the implication of these results for non-steady behavior by investigating the lateral and temporal scales over which the preformed nutrients subducted in the SAMW layer persist in the ocean interior before fueling downstream primary productivity. 15 Preformed nutrients are those inorganic nutrients that enter the ocean interior without having fueled primary productivity while in the mixed layer. Therefore, preformed nutrients are a signature of inefficiency in the biological pump. A more efficient biological pump utilizes a greater portion of surface nutrients, converts them to organic material and increases the remineralized nutrient and carbon concentration deeper in the water column. Higher remineralized nutrient concentrations signify greater ocean 20 carbon storage and lower atmospheric carbon dioxide, while a higher preformed nutrient inventory signals a smaller ocean carbon reservoir (Archer et al., 2000; Ito and Follows, 2005; Gnanadesikan and Marinov, 2008; Marinov et al., 2008a, b). Much of the SAMW nutrients enter the ocean in their preformed state, but when they fuel productivity 25 downstream of their formation region, these inorganic nutrients are incorporated to organic matter and eventually converted to their remineralized state. Any increase in the subduction of the SAMW preformed nutrient pool would diminish the deep ocean carbon reservoir until these preformed nutrients are converted to their remineralized form. Evaluation of this potential for change in the ocean preformed nutrient reservoir

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



is especially timely given recent observations showing the intensification of Southern Hemisphere westerly wind strength (Thompson and Solomon, 2002) with implications for the stability of the ocean carbon sink (Le Quéré et al., 2007). Such an intensification is expected to increase the cross-ACC Ekman transport of preformed nutrients, 5 the preformed nutrient load of the SAMW, and the local inefficiency of the biological pump due to the subducted preformed nutrients in the mode and intermediate waters. Our model experiments shed light on the time scale that this perturbation might be expected to persist.

BGD

7, 4045–4088, 2010

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



cnal mixing, and wind forcing all determine where dense water returns to the ocean's surface, while the interplay of all three factors determines the depth and thickness of the pycnocline and the strength of the overturning cell.

For the modern ocean, the Gnanadesikan (1999) theory can be summarized as follows. Dense water forms and sinks in the high latitude North Atlantic and is exported southward in the deep limb of the MOC, a flux represented in the schematic as M_n . All along its export pathways, the downward mixing of heat transforms the dense water to lighter densities and forces an upward volume flux represented as M_u that reduces the deep southward flow as it approaches the Southern Ocean. This upward flux scales as:

$$M_u = \frac{K_v A}{D},$$

where K_v is the coefficient of diapycnal diffusion, D is the pycnocline depth that separates dense water from light, and A is the area of the low latitude pycnocline.

In the Southern Ocean, laterally divergent Ekman transport forces the upwelling of abyssal waters, a flux represented as M_s . The upwelled waters that cross the ACC into the SAMW formation region gain buoyancy along their journey, thus completing the dense-to-light transformation of the water masses formed in the Northern Hemisphere high latitudes. However, adding another layer of complexity, a volume transport due to mesoscale eddies is directed southward across the ACC, balancing some fraction of the northward Ekman transport. The eddy transport is thought to be proportional to the local slope of isopycnals, which is the source of potential energy from which the eddies grow (Gent and McWilliams, 1990). Thus, the flux of abyssal water to the light water realm in the Southern Ocean, M_s , is equal to the difference between the Ekman transport M_{ek} and the mass flux due to the eddies:

$$M_s = M_{ek} - \frac{A_i D L_x}{L_y^S}$$

where A_i is a coefficient that determines the strength of the eddy mass transport, L_y^S is

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the north-south scale over which the pycnocline shallows in the Southern Ocean, and L_x is the Earth's circumference at the Drake Passage latitude (see also Karsten and Marshall, 2002).

For the limiting case in which the eddy transport across the ACC is as large as the 5 Ekman transport, the net Southern Ocean deep water return pathway becomes negligible and the upward branch of the MOC must be accomplished exclusively through low latitude diapycnal mixing; this is the picture of the MOC hypothesized by Stommel and Arons (1961). For the alternate limiting case in which global diapycnal mixing is, on average, as small as the diapycnal mixing measured in the ocean away from topography 10 (Gregg and Sanford, 1980; Ledwell et al., 1993, 1998), the Ekman upwelling of dense waters in the Southern Ocean and their buoyancy gain provide the primary driver of meridional overturning (Toggweiler and Samuels, 1998). We note that this conceptual model does not explicitly consider spatial variability in the strength of diapycnal mixing 15 observed in the ocean (Heywood et al., 2002; Naveira Garabato et al., 2004), although the model framework outlined below allows for some exploration of this variability.

2.2 The models

The underlying architecture used to probe our questions is a suite of simulations using the Geophysical Fluid Dynamics Laboratory Modular Ocean Model version 3 (MOM3) (Pacanowski and Griffies, 1999). The MOM3 suite explores a range of mixing parameters and wind forcing, as described in several previous studies (Gnanadesikan, 20 1999; Gnanadesikan et al., 2004, 2007; Mignone et al., 2006), and summarized in Table 1. The models create variants of the MOC strength and structure by changing: a) the Southern Ocean Ekman transport, M_{ek} , via the use of two different wind re-analysis products to force the model; b) the counterbalancing eddy transport via the manipulation of A_i , the coefficient associated with the Gent and McWilliams (1990) 25 (GM) parameterization of eddy transport, and c) the diapycnal mixing across the low latitude pycnocline via variations in K_v , the coefficient of diapycnal diffusion. Here we present the results from a subset of four models from a larger MOM3 suite explored

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

in Gnanadesikan et al. (2007), chosen because they capture much of the inter-model differences of the full parameter range without being excessively cumbersome for comparison.

Two of the models are forced with wind stresses from Hellerman and Rosenstein (1983), which are nearly half as strong in the Southern Ocean as the European Centre for Medium-Range Weather Forecasts (ECMWF) climatological winds (Trenberth et al., 1989) used to force the other two models. Of these Hellerman and Rosenstein-forced models, one is run with a Low K_v and Low A_i , and is therefore referred to as LL; the other has a High K_v and High A_i , and is referred to as HH. LL-ECMWF is identical to the LL model in the ocean, but is forced with ECMWF winds. A special case we call P2A resembles the LL-ECMWF model but has been modified in several ways intended to increase the realism of the simulation: the Drake Passage was narrowed from three grid points (1165 km) to two (777 km) to be closer to the real Drake Passage, which is 800 km wide; strong vertical mixing ($5 \times 10^{-3} \text{ m}^2 \text{ s}^{-1}$) is imposed on the uppermost two grid cells (50 m) as a very crude approximation to what actually occurs in an oceanic mixed layer; and surface salinity is strongly restored to subsurface observations in four key locations at the onset of Austral winter in Southern Ocean deep water formation regions to simulate brine rejection during sea-ice formation. Without this salinity adjustment, restoring the model to observations suppresses Antarctic Bottom Water (AABW) formation, as the observations do not capture the outcropping of such waters. Finally, in P2A the coefficient of diapycnal mixing is increased in the Southern Ocean, first motivated by observations of intense ACC internal wave activity (Polzin, 1999) and later validated by observations of rapid diapycnal tracer spreading in the region (Naveira Garabato et al., 2004). Clearly the MOC strength and upward branch is controlled by the interplay of these parameter choices (Fig. 2b–c), with the clockwise part of the MOC in the HH model confined entirely to the region north of 20° N, while the LL-ECMWF overturning is actually intensified in the Southern Hemisphere. We will explore the MOC differences among the models at greater length in Sect. 3.

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A model of the biological cycling of phosphorus and carbon is coupled to the physical model, as specified by the Ocean Carbon Model Intercomparison Project-2 (OCMIP-2) protocol (Najjar et al., 2007). In this protocol, phosphorus is the only simulated nutrient and is conserved globally in all experiments. The phosphorus source/sink terms are governed by an extremely simple set of equations meant to represent the incorporation of phosphate into organic matter by photosynthesis in the euphotic zone and the sinking of this organic matter and its remineralization back to phosphate throughout the water column (Fig. 3). The simulated phosphorus appears in 2 pools: dissolved organic phosphorus (DOP) and phosphate (PO_4^{3-}). These pools are transported as passive tracers by the physical model. At depths shallower than 75 m, model PO_4^{3-} in excess of a seasonal PO_4^{3-} climatology (Louanchi and Najjar, 2000) fuels primary productivity, represented by restoring the surface model PO_4^{3-} to climatological values over a time scale of $\tau=30$ days. The resulting productivity has two fates: two thirds becomes a source term for DOP and one third fuels a parameterized sinking flux of organic phosphorus that is shunted to 75 m. Below 75 m, the sinking flux decays as a function of depth and its vertical divergence is a source term of PO_4^{3-} . DOP is remineralized to PO_4^{3-} everywhere in the model, at a rate of $\kappa=0.5 \text{ year}^{-1}$.

2.3 The experimental design

To investigate the role of the upward branch of the MOC in the supply of nutrients to the low latitude pycnocline and ultimately the euphotic zone, we created two sets of tags that are schematized in Fig. 4. The first is a simple set of dye tracers. In each tagging region (represented in black in the schematics), the dye value is set to one; in each destruction region (white in the schematics), the dye value is set to zero; in the rest of the domain (gray in the schematics), the dye tags are treated as conserved, passive tracers. The dyes are named according to their source locations and are fully described in Fig. 4 and its caption. The physical model is spun up to a steady circulation before the dyes are released. All presented results are averages between year 190 and 200

after the release of the tags, except for LL-ECMWF, which are for years 390–400. At year 190, the drift in average low latitude dye concentrations is less than 0.1% year⁻¹, very close to equilibrium, and almost all regions are fully tagged. After 200 years, only the water masses of the Bay of Bengal are not fully tagged (thus, the global sum of all tags can be slightly less than zero).

Outside the tagging and destruction regions (i.e. in and above the low latitude pycnocline), the dye tracers directly measure the fractions of water in and above the main thermocline that derive from each tagging region. The abyssal ocean is tagged only in low latitudes and only at potential densities greater than 27.4 kg m⁻³, which is beneath the main thermocline in all models. The Northern Hemisphere High latitudes are tagged everywhere north of 40° N, and further designated as stemming from the North Atlantic and North Pacific basins. In the Southern Ocean, the tags are divided into density classes in order to approximate the contribution of SAMW and AAIW to the low latitude pycnocline. By design, the southernmost tag is trapped in the Southern Ocean, and is used primarily to ensure that all water masses are tagged and a water mass budget can be balanced.

In the Southern Ocean, AAIW is most commonly identified as a salinity minimum along the 27.3 isopycnal. Because the SAMW is formed in large part by convective processes, it can be identified as a pycnostad, or region of low vertical density stratification. A comparison of our simulations with a well-studied salinity and density section along 100° E (Sloyan and Kamenkovich, 2007) (Fig. 5), reveals that the models do a good job simulating the salinity and stratification minima that characterize the AAIW and SAMW, respectively. The model density structure in the Southern Ocean resembles the observations, giving confidence that our Southern Ocean water mass definitions are appropriate to address the question at hand. It is also reassuring that the SAMW dye tracer on the 26.8 isopycnal shows a similar spatial distribution to the Si* tracer from data (compare Fig. 6 with Fig. 1c). We note that the SAMW penetrates into the Northern Hemisphere in all of the models, regardless of whether there is net destruction or creation of light waters in the Southern Ocean. The differences between

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the models are largely seen in tropical latitudes where the HH model (Fig. 6b) has much lower concentrations of SAMW dye than the others.

The second set of tags is designed to trace nutrients from the same source regions as the dyes. In each source region, all model PO_4^{3-} and DOP are tagged, or named

5 according to their region, such that the total pools of PO_4^{3-} and DOP everywhere in the model domain are the sum of the regionally-tagged pools. As an illustrative example, PO_4^{3-} and DOP are endowed with a SAMW tag between the 26.5 and 27.1 isopycnals in the Southern Hemisphere, southward of where the 26.5 isopycnal intersects 200 m (as for the SAMW dye tracer, shown in Fig. 4c).

10 As the SAMW-tagged DOP (hereafter *SAMW DOP*) is transported in the flow field from its tagging region, it is resupplied to a separate pool of *remineralized SAMW phosphate*. The same is true for the *SAMW phosphate* that fuels primary productivity downstream of its tagging domain: the remineralization of the resultant DOP and sinking flux provides a source of *remineralized SAMW phosphate*. Thus, although the phosphorus

15 maintains its regional identity until it enters a destruction region, it is marked as either coming directly from the tagging region, which we will refer to as *preformed*, or being created by remineralization away from the tagging region, which we refer to as *remineralized*. We note that, because the tags are not exclusively introduced in the mixed layer, *preformed* nutrients in our experiments differ from a traditional definition

20 of preformed nutrients as those that have been cycled through the mixed layer and returned to the ocean interior without having fueled primary productivity. This distinction becomes particularly apparent in our discussion of the *deep low latitude phosphate*, which is tagged only in the interior ocean, and may therefore bear no resemblance to the traditional definition of preformed nutrients. Thus, throughout this work, references

25 to regional tags and tagged nutrients and their *preformed* and *remineralized* components will appear in *italics* for clarity. Combining the equations for the source/sink terms of PO_4^{3-} and DOP in the OCMIP-2 biogeochemistry model (Eqs. 2–7 in Fig. 3), allows an estimate of the primary productivity that is fueled by each pool of *preformed* and *remineralized* tagged phosphate.

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



3 Results and discussion

The simulated MOC in the different models behaves as predicted by the simple analytical model presented in Sect. 2.1. In all models, the interplay of the chosen parameter values creates a pycnocline with a realistic depth and density and an MOC at 48° N similar to the 16–19 Sv inferred from observations (Table 2) (Lumpkin and Speer, 2007; Talley, 2008; Forget, 2009). However, the upward branch of the MOC occurs at notably different locations among the models (Table 2). The LL model with its weak Southern Ocean winds and low mixing creates an MOC of 13.6 Sv, with more than half of the upward branch supplied in the Southern Ocean; the HH model creates 17.9 Sv of overturning, but its upward branch is realized entirely at low latitudes; the P2A and LL-ECMWF models yield more than 18 Sv of overturning, but with upward branches primarily in the Southern Ocean. Some observationally-driven inverse models (Lumpkin and Speer, 2007) and, more recently, the results from data-assimilating forward model used in Estimating the Circulation and Climate of the Ocean (ECCO) (Wunsch and 10 Heimbach, 2007; Forget, 2009) have suggested that the upward branch of the MOC is accomplished primarily in the Southern Ocean (Table 2). Yet, another observationally-driven estimate suggests a more important role for low latitude upwelling and implies values of low latitude diapycnal mixing as high as in our HH model (Talley, 2008). Thus, the relative importance of these two possible upwelling branches lacks a definitive solution at present, providing a justification for including in our analysis models that cover 15 the range of possibilities. Furthermore, the modeled MOC has a direct influence on the large-range of air-sea CO₂ fluxes estimated from coupled models (Cao et al., 2009).

Radiocarbon, a useful tracer of ocean circulation, offers evidence consistent with the upwelling of deep waters in the Southern Ocean. We briefly compare the radiocarbon results from our four models with observations to offer a context with which to judge the various circulations. Radiocarbon enters the ocean only at the air-sea interface and then decays in the ocean interior with a half-life of 5730 years. Thus, the most-recently ventilated ocean waters are most enriched in radiocarbon, and the 20

BGD

7, 4045–4088, 2010

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

tributes 67% of the pycnocline waters but only 48% of the phosphate (Table 3). These differences arise because the phosphate concentration in a tagged water mass may be highly elevated above the phosphate concentration in the pycnocline. Thus, when the water mass is mixed or advected into the pycnocline it can contribute a greater percentage of the resultant phosphate concentration than the corresponding contribution of the dye fraction. Such is the case for the highly phosphate-enriched deep ocean.

5 Each of the models demonstrates some skill at reproducing the large-scale features of the nutricline: simulated PO_4^{3-} concentrations arch upward from the deep, depleted bowls of the subtropical gyres to the shallow nutricline of the equator and the high latitudes, just as they do in the observational data (Figs. 8a–b and 9a–b). For brevity, we have chosen to show the zonally-averaged sections only for the HH and P2A models in 10 Figs. 8 and 9; though HH is quite different from the other three models, LL, LL-ECMWF, and P2A are qualitatively similar, at least as far as the zonally-averaged nutrient sections are concerned.

15 The Southern Ocean is notable for its elevated near-surface nutrient concentrations (Figs. 8 and 9), which result from strong vertical and lateral advection and mixing of nutrients to the surface (Sloyan and Rintoul, 2001; Naveira Garabato et al., 2004; Sarmiento et al., 2004) in the presence of weak, iron- and light-limited biological drawdown (Coale et al., 2004). The simplified biogeochemical model from the OCMIP-2 20 protocol does not simulate an iron cycle, nor does it evaluate the irradiance available for photosynthesis. Nonetheless, the biological model does represent the sluggish biological PO_4^{3-} drawdown in the Southern Ocean that results from iron and light limitation, because it allows productivity to remove surface nutrients only when model PO_4^{3-} concentrations exceed observed climatological mean surface PO_4^{3-} concentrations. 25 Because there are high surface PO_4^{3-} concentrations in the Subantarctic Zone where SAMW is formed, the modeled SAMW is subducted with some of the highest preformed nutrient concentrations in the global pycnocline. This result is independent of the physical model; rather, it is set by the PO_4^{3-} observations in this region. The degree to which these nutrients spread from the SAMW source region to fuel downstream

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

primary productivity, on the other hand, is highly dependent on model physics.

This intermodel difference can clearly be seen in tropical nutrient cycling (Figs. 8 and 9). In the HH model compared to the P2A model, the spreading of the *SAMW phosphate* is restricted to a narrow band near its source region, largely because the high GM coefficient (A_i) creates a parameterized southward eddy transport that largely compensates the northward Ekman transport of SAMW (Figs. 8f and 9f). Similarly, this same GM transport causes a more restricted *North Pacific phosphate* spreading in the HH run relative to the runs with a lower GM coefficient (Figs. 8c and 9c). Furthermore, the *deep low latitude phosphate* diffuses strongly across the low latitude pycnocline due to the choice of high diapycnal diffusion (Figs. 8e and 9e). This strong diffusion of nutrients across the low-latitude pycnocline also causes the HH model to have lower concentrations of phosphate in the dense waters underlying the tropics, the highest low-latitude productivity, and the one that disagrees the most with observationally-based estimates (Gnanadesikan et al., 2002).

Winds also affect the extent to which the spreading of the SAMW transports nutrients away from the Southern Ocean (Fig. 10). In the models forced with sluggish Southern Ocean winds (LL and HH), the *preformed SAMW phosphate* comprises the majority of the total pycnocline nutrients only in a narrow band south of 30° S near the tagging region (Fig. 10) and represents less than 10% of the nutrients in the Northern Hemisphere gyres. Spreading of the *preformed SAMW phosphate* is stronger in the P2A and LL-ECMWF models, with a noticeable contribution to total PO_4^{3-} above the 26.8 isopycnal in the subtropical North Atlantic (Fig. 10, top row). This density horizon is chosen to represent the central pycnocline, but the qualitative results are not sensitive to the density horizon used, as long as it is above the 27.4 isopycnal below which the *deep low latitude* tag is assigned.

The importance of SAMW supply is more apparent in the spreading of its *remineralized* component (Fig. 10, bottom row). Whereas the *preformed SAMW phosphate* is largely confined to the Southern Ocean, the *remineralized SAMW phosphate* can propagate far downstream of its tagging region, contributing a significant phosphate

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

small contribution of the northern high latitudes can be attributed to the experimental design, which re-tags all nutrients that enter the low latitude abyss with the *deep low latitude* tag. A portion of the *North Atlantic DOP* and *phosphate* sinks and spreads to the *deep low latitude* tagging region (where the North Atlantic tag is reset to zero) without first fueling productivity.

How do nutrients subducted in the SAMW layer at densities greater than those in the subtropical euphotic zone come to influence productivity throughout the low latitudes? Originally tagged beneath the 26.5 isopycnal, the SAMW rises to shallower depths and lighter densities after spreading from its tagging region (Fig. 12). First, in the down-welling domain just the north of the SAMW tagging region, advective and diffusive fluxes of the SAMW at the depth of the 26.5 isopycnal are downward. This downward flux is caused by the tagging routine in which the SAMW tag is destroyed beneath the 27.4 isopycnal, but only diluted by mixing above this layer, so that the SAMW concentration maximum moves to densities slightly above the 26.5 isopycnal just to the north of the tagging region. Downstream of this downwelling domain, the SAMW is transported upwards. The upward transport is primarily achieved by vertical advection in tropical regions, along the eastern fringes of the Southern Hemisphere subtropical basins, and in the South Atlantic western boundary current off Brazil (Fig. 12). Once advected and mixed to shallower depths, the SAMW is transformed by surface buoyancy forcing and diapycnal mixing and may no longer carry the characteristic temperature, salinity and potential vorticity signature it had upon formation and subduction. However, our tags trace the water mass fraction and its nutrients downstream of the tagging regions even as the water mass itself is destroyed by such transformations.

The SAMW nutrients that are upwelled into the euphotic zone in the tropics and along the eastern boundaries of the subtropics are consumed and converted to organic particles that sink through the water column. These sinking particles are partially remineralized on layers with lighter densities and shallower depths than the SAMW had upon subduction (Toggweiler and Carson, 1995; Williams et al., 2006). Thus, a portion of the nutrients originally subducted in the SAMW remains available to support

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



subtropical productivity as it spreads along these lighter surfaces to the neighboring subtropical gyres.

BGD

7, 4045–4088, 2010

5 Our model results suggest that nutrients slowly lost to depths beneath the pycnocline throughout the low latitudes return to lighter layers primarily in the Southern Ocean. This picture emerges from all models but HH and is summarized in Fig. 13. In the Southern Ocean where biological export does not efficiently compete with the advection and mixing of nutrients to the surface layers, nutrients are upwelled and mixed vertically in the ACC and then subducted in the SAMW. Therefore, much of the water masses formed in the Southern Ocean enter the ocean interior with elevated nutrient 10 concentrations. Although much of these nutrients are subducted to depths well below the subtropical euphotic zone, their upwelling along the eastern and western fringes of the subtropics and in the tropics allows these nutrients to sustain subtropical primary productivity.

15 Not far from the SAMW source region, the SAMW *preformed* nutrients are consumed and converted to their *remineralized* counterparts within the pycnocline; further downstream the *remineralized* nutrients are re-incorporated into organic matter wherever they are mixed and advected into the sunlit surface ocean. In these downstream locales, the slow leakage of organic phosphorus past the base of the seasonally-accessible layers is almost entirely balanced by the lateral and vertical supply of nutrients 20 that were cycled through Southern Ocean water masses. In the HH model, the nutrient budget achieves a slightly more one-dimensional balance: a majority of the sinking flux of nutrients downward across the low latitude pycnocline is restored locally via the mixing of nutrients upwards across the low latitude pycnocline. However, even in the HH model, *SAMW phosphate* sustains more than a third of low latitude productivity; in the other models the percentage is between 57% and 75% (Fig. 13). Each of 25 the other tagging regions besides *SAMW* and *deep low latitude* contributes less than 10% of the phosphate that fuels global low latitude productivity.

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



4 Conclusions, implications, and connections

Our model framework has allowed us to identify the dominant nutrient return pathways from the dense ocean interior to depths that are accessible to the sunlit euphotic zone. Between 30° S and 30° N, the upper ocean nutrient reservoir is dominated by those nutrients that were returned from the deep ocean to the pycnocline in the SAMW formation/subduction region of the Southern Ocean (Fig. 13). Only in the model where the coefficient of vertical mixing is set to levels higher than those observed in the pycnocline of the open ocean does mixing directly across the low latitude nutricline provide the dominant return pathway of nutrients to the low latitude euphotic zone. The Northern Hemisphere subpolar regions are not the primary suppliers of nutrients to the low latitude euphotic zones; nor are the regions of Southern Ocean south of the SAMW formation region.

These results are in line with previous modeling studies (Marinov et al., 2006) and with inferences from the distribution of tracers such as Si* (Sarmiento et al., 2004). But there are two major new insights afforded by this study. The first stems from our focus on the sensitivity of nutrient pathways to the winds, vertical mixing, and the strength of lateral eddy transport (via changes in the GM coefficient). Our results suggest that in models with high vertical mixing, high eddy transport, and low Southern Ocean winds, the dominant nutrient balance is more one-dimensional, i.e. organic nutrients lost by downward sinking at low latitudes are restored through the upward mixing of inorganic nutrients across the pycnocline at these latitudes. At the other extreme, in models with low vertical mixing, weak GM transport, and stronger Southern Ocean winds, organic nutrients lost across the low latitude pycnocline are restored to the pycnocline primarily in the Southern Ocean and depend on the full three-dimensional ocean circulation for their delivery to the low latitude euphotic zone.

The second major insight afforded by the study is an approximation of the spatial and temporal scale over which nutrients that enter the ocean interior in the SAMW formation region persist before being utilized in downstream locations. In less than 40 years

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



on average, phosphate entering the ocean interior in the SAMW region is utilized to fuel downstream productivity. In large part, the SAMW nutrients first fuel primary productivity south of 20° S. This short spatial and temporal scale separating the subduction of SAMW nutrients and their downstream utilization helps explain previous modeling results that tested the influence of the SAMW region on atmospheric carbon concentrations. In the modeling study of Marinov et al. (2006), the SAMW nutrient source was “turned off” by forcing nutrients towards zero at the Southern Ocean surface north of the outcropping 27.3 isopycnal, the equivalent of increasing nutrient drawdown by enhanced export production. Although this forced drawdown decimated low latitude productivity by starving these regions of their nutrients, the experiment had little impact on ocean carbon storage or atmospheric carbon concentrations on millennial time scales. The reason for the lack of impact on atmospheric carbon is that the biological pump had been made much more efficient locally in the SAMW formation region, but the global efficiency was unchanged once the model returned to equilibrium (Marinov et al., 2006). In experiments without the forced nutrient drawdown, the SAMW nutrients were already being efficiently consumed downstream of the SAMW formation region on decadal time scales.

Our model experiments suggest a paradigm for returning nutrients to the surface ocean that contextualizes a number of phenomena. For example, it has been hypothesized that biological nitrogen fixation in the North Atlantic is sustained by excess phosphate (phosphate in excess of nitrate relative to their Redfield ratio) advected in the shallow limb of the MOC (Moore et al., 2009). The excess phosphate signature in the shallow MOC layers stems from the SAMW formation region, where high-phosphate waters from the Indo-Pacific Basin are mixed with the water masses advected across the ACC. Iron limits biological nitrogen fixation in the SAMW formation region and throughout the subtropical South Atlantic, so the excess phosphate signature is preserved during formation and subduction and is exported to the North Atlantic. Because nitrogen fixation in the relatively iron-rich North Atlantic appears to be phosphate-limited (Sañudo-Wilhelmy et al., 2001; Moore et al., 2009), the SAMW

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

models are configured with one vertical mixing profile for the global ocean, which was altered only in P2A in order to simulate higher Southern Ocean mixing. Such uniformity in mixing does not capture the observed enhancement of mixing in regions of rough topography, such as tidal mixing (St. Laurent and Simmons, 2006) which is parameterized in more recent GCMs (Simmons et al., 2004). Finally, the OCMIP-2 biology is not fully prognostic because it is forced towards observations at the surface. Thus, the models cannot represent changes to biology at the surface forced by the models' differing physics, a challenge that can be tackled with fully prognostic biological models. Future efforts could also move beyond our steady state solution and address temporal variability at time scales other than seasonal, incorporate ocean-atmosphere coupling, and examine the hypothesized alterations in the nutrient return pathways under different climate scenarios.

Acknowledgements. The authors gratefully acknowledge helpful comments from Robbie Togweiler and Stephanie Downes.

15 References

Anderson, R. F., Ali, S., Bradtmiller, L. I., Nielsen, S. H. H., Fleisher, M. Q., Anderson, B. E., and Burckle, L. H.: Wind-Driven Upwelling in the Southern Ocean and the Deglacial Rise in Atmospheric CO₂, *Science*, 323, 1443–1448, doi:10.1126/science.1167441, 2009.

Archer, D. E., Eshel, G., Winguth, A., Broecker, W., Pierrehumbert, R., Tobis, M., and Jacob, R.: Atmospheric $p\text{CO}_2$ sensitivity to the biological pump in the ocean, *Global Biogeochem. Cy.*, 14, 1219–1230, 2000.

Cao, L., Eby, M., Ridgwell, A., Caldeira, K., Archer, D., Ishida, A., Joos, F., Matsumoto, K., Mikolajewicz, U., Mouchet, A., Orr, J. C., Plattner, G.-K., Schlitzer, R., Tokos, K., Totterdell, I., Tschumi, T., Yamanaka, Y., and Yool, A.: The role of ocean transport in the uptake of anthropogenic CO₂, *Biogeosciences*, 6, 375–390, doi:10.5194/bg-6-375-2009, 2009.

Coale, K. H., Johnson, K. S., Chavez, F. P., Buesseler, K. O., Barber, R. T., Brzezinski, M. A., Cochlan, W. P., Millero, F. J., Falkowski, P. G., Bauer, J. E., Wanninkhof, R. H., Kudela, R. M., Altabet, M. A., Hales, B. E., Takahashi, T., Landry, M. R., Bidigare, R. R., Wang, X.,

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Gnanadesikan, A., De Boer, A. M., and Mignone, B. K.: A simple theory of the pycnocline and overturning - revisited., *Ocean Circulation: Mechanisms and Impacts*, Geophys. Monogr. Ser., 173, 335–349, 2007.

5 Gnanadesikan, A. and Marinov, I.: Export is not enough: nutrient cycling and carbon sequestration, *Mar. Ecol. Prog. Ser.*, 364, 289–294, doi:10.3384/meps07550, 2008.

Gregg, M. C. and Sanford, T. B.: Signatures of mixing from the Bermuda Slope, the Sargasso Sea and the Gulf Stream, *J. Phys. Oceanogr.*, 10, 105–127, 1980.

10 Hallberg, R. and Gnanadesikan, A.: The Role of Eddies in Determining the Structure and Response of the Wind-Driven Southern Hemisphere Overturning: Results from the Modeling Eddies in the Southern Ocean (MESO) Project, *J. Phys. Oceanogr.*, 36, 2232–2252, 2006.

Hanawa, K. and Talley, L.: Mode Waters, in: *Ocean Circulation and Climate*, edited by: Siedler, G., and Church, J., International Geophysics Series, Academic Press, 373–386, 2001.

15 Hellerman, S. and Rosenstein, M.: Normal monthly wind stress over the world ocean with error estimates, *J. Phys. Oceanogr.*, 13, 1093–1104, 1983.

Heywood, K. J., Naveira Garabato, A. C., and Stevens, D. P.: High mixing rates in the abyssal Southern Ocean, *Nature*, 415, 1011–1014, 2002.

Ito, T. and Follows, M. J.: Preformed phosphate, soft tissue pump and atmospheric CO₂, *J. Marine Res.*, 63, 813–839, 2005.

20 Karsten, R. H. and Marshall, J.: Constructing the Residual Circulation of the ACC from Observations, *J. Phys. Oceanogr.*, 32, 3315–3327, 2002.

Key, R. M., Kozyr, A., Sabine, C., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy, C., and Peng, T.-H.: A global ocean carbon climatology: Results from GLODAP, *Global Biogeochem. Cy.*, 18, GB4031, doi:10.1029/2004gb002247, 2004.

25 Kuhlbrodt, T., Griesel, A., Montoya, M., Levermann, A., Hoffman, M., and Rahmstorf, S.: On the driving processes of the Atlantic Meridional Overturning Circulation, *Rev. Geophys.*, 45, RG2001, doi:10.1029/2004rg000166, 2007.

Le Quéré, C., Rödenbeck, C., Buitenhuis, E. T., Conway, T. J., Langenfelds, R., Gomez, A., Labuschagne, C., Ramonet, M., Nakazawa, T., Metzl, N., Gillett, N., and Heimann, M.: Saturation of the southern ocean CO₂ sink due to recent climate change, *Science*, 316, 1735–1738, 2007.

30 Ledwell, J. R., Watson, A. J., and Law, C. S.: Evidence for slow mixing across the pycnocline from an open-ocean tracer-release experiment, *Nature*, 364, 701–703, 1993.

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Ledwell, J. R., Watson, A. J., and Law, C. S.: Mixing of a tracer in the pycnocline, *J. Geophys. Res.*, 103, 21499–21529, 1998.

Levitus, S., Burgett, R., and Boyer, T. P.: World Ocean Atlas, NOAA Atlass NESDIS 3, National Oceanic and Atmospheric Administration, Silver Spring, MD, 1994.

5 Louanchi, F. and Najjar, R. G.: A global monthly climatology of phosphate, nitrate and silicate in the upper ocean: Spring-summer export production and shallow remineralization, *Global Biogeochem. Cy.*, 14, 957–977, 2000.

Lumpkin, R. and Speer, K.: Global Ocean Meridional Overturning, *J. Phys. Oceanogr.*, 37, 2550–2562, 2007.

10 Marinov, I., Gnanadesikan, A., Toggweiler, J. R., and Sarmiento, J. L.: The Southern Ocean biogeochemical divide, *Nature*, 441, 964–967, 2006.

Marinov, I., Follows, M. J., Gnanadesikan, A., Sarmiento, J. L., and Slater, R. D.: How does ocean biology affect atmospheric $p\text{CO}_2$?, Theory and models, *J. Geophys. Res.-Oceans*, 113, C07032, doi:10.1029/2007jc004598, 2008a.

15 Marinov, I., Gnanadesikan, A., Sarmiento, J. L., Toggweiler, J. R., Follows, M., and Mignone, B. K.: Impact of oceanic circulation on biological carbon storage in the ocean and atmospheric $p\text{CO}_2$, *Global Biogeochem. Cy.*, 22, GB3007, doi:10.1029/2007gb002958, 2008b.

McCartney, M. S.: The subtropical recirculation of Mode Waters, *J. Marine Res.*, 40, 427–464, 1982.

20 Mignone, B. K., Gnanadesikan, A., Sarmiento, J. L., and Slater, R. D.: Central role of Southern Hemisphere winds and eddies in modulating the oceanic uptake of anthropogenic carbon, *Geophys. Res. Lett.*, 33, L01604, doi:10.1029/2005gl024464, 2006.

Moore, C. M., Mills, M. M., Achterberg, E. P., Geider, R. J., LaRoche, J., Lucas, M. I., McDonagh, E. L., Pan, X., Poulton, A. J., Rijkenberg, M. J. A., Suggett, D. J., Ussher, S. J., and Woodward, E. M. S.: Large-scale distribution of Atlantic nitrogen fixation controlled by iron availability, *Nature Geosci.*, 12, 867–871, doi:10.1038/ngeo667, 2009.

25 Munk, W. and Wunsch, C.: Abyssal recipes II: Energetics of tidal and wind mixing, *Deep-Sea Res. Pt. I*, 45, 1977–2010, 1998.

Najjar, R. G., Jin, X., Louanchi, F., Aumont, O., Caldeira, K., Doney, S. C., Dutay, J. C., Follows, M., Gruber, N., Joos, F., Lindsay, K., Maier-Reimer, E., Matear, R. J., Matsumoto, K., Monfray, P., Mouchet, A., Orr, J. C., Plattner, G. K., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Weirig, M. F., Yamanaka, Y., and Yool, A.: Impact of circulation on export production, dissolved organic matter, and dissolved oxygen in the ocean: Results from Phase II of the

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Ocean Carbon-cycle Model Intercomparison Project (OCMIP-2), *Global Biogeochem. Cy.*, 21, GB3007, doi:10.1029/2006gb002857, 2007.

Naveira Garabato, A. C., Polzin, K. L., King, B. A., Heywood, K. J., and Visbeck, M.: Widespread Intense Turbulent Mixing in the Southern Ocean, *Science*, 303, 210–213, 2004.

5 Pacanowski, R. C. and Griffies, S. M.: The MOM3 manual, Alpha version., Geophysical Fluid Dynamics Laboratory, Princeton, NJ, 580, 1999.

Palter, J. B., Lozier, M. S., and Barber, R. T.: The effect of advection on the nutrient reservoir in the North Atlantic subtropical gyre, *Nature*, 437, 687–692, 2005.

10 Polzin, K. L.: A rough recipe for the energy balance of quasi-steady internal lee waves, 'Aha Huliko'a: Dynamics of oceanic internal gravity waves II., Hawaii, 1999, 117–128, 1999.

Ridgway, K. R., Dunn, J. R., and Wilkin, J. L.: Ocean Interpolation by Four-Dimensional Weighted Least Squares-Application to the Waters around Australasia, *J. Atmos. Ocean. Tech.*, 19, 1357–1375, 2002.

15 Samelson, R. M.: Simple mechanistic models of middepth meridional overturning, *J. Phys. Oceanogr.*, 34, 2096–2103, 2004.

Samelson, R. M.: A simple dynamical model of the warm-water branch of the middepth meridional overturning cell, *J. Phys. Oceanogr.*, 39, 1216–1230, 2009.

20 Sañudo-Wilhelmy, S. A., Kustka, A. B., Gobler, C. J., Hutchins, D. A., Yang, M., Lwiza, K., Burns, J., Capone, D. G., Raven, J. A., and Carpenter, E. J.: Phosphorus limitation of nitrogen fixation by *Trichodesmium* in the central Atlantic Ocean, *Nature*, 411, 66–69, 2001.

Sarmiento, J. L., Gruber, N., Brzezinski, M. A., and Dunne, J. P.: High-latitude controls of thermocline nutrients and low latitude biological productivity, *Nature*, 427, 56–60, 2004.

25 Sarmiento, J. L., Simeon, J., Gnanadesikan, A., Gruber, N., Key, R. M., and Schlitzer, R.: Deep ocean biogeochemistry of silicic acid and nitrate, *Global Biogeochem. Cy.*, 21, GB1S90, doi:10.1029/2006gb002720, 2007.

Simmons, H. L., Jayne, S. R., Laurent, L. C. S., and Weaver, A. J.: Tidally driven mixing in a numerical model of the ocean general circulation, *Ocean Modell.*, 6, 245–263, 2004.

Sloyan, B. M. and Rintoul, S. R.: Circulation, renewal, and modification of Antarctic mode and intermediate water, *J. Phys. Oceanogr.*, 31, 1005–1030, 2001.

30 Sloyan, B. M. and Kamenkovich, I. V.: Simulation of Subantarctic Mode and Antarctic Intermediate Waters in Climate Models, *J. Climate*, 20, 5061–5080, 2007.

St. Laurent, L. and Simmons, H.: Estimates of Power Consumed by Mixing in the Ocean Interior, *J. Climate*, 19, 4877–4890, 2006.

Nutrient return pathways

J. B. Palter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 1. Description of the four ocean GCMs. Model acronyms are described in the text. Differences from LL are highlighted in bold.

Configuration:	LL	HH	LL-ECMWF	P2A
Diapycnal diffusivity (tropics) (m ² s ⁻¹)	1.5×10 ⁻⁵	6×10⁻⁵	1.5×10 ⁻⁵	1.5×10 ⁻⁵
Diapycnal diffusivity (Southern Ocean) (m ² s ⁻¹)	1.5×10 ⁻⁵	6×10⁻⁵	1.5×10 ⁻⁵	13×10⁻⁵
Enhanced diffusivity in top 50 m	No	No	No	Yes
Along isopycnal diffusivity AND coefficient for GM thickness diffusion (m ² s ⁻¹)	1000	2000	1000	1000
Applied surface moisture fluxes	(Da Silva et al., 1994)	(Da Silva et al., 1994)	(Da Silva et al., 1994)	(Da Silva et al., 1994) plus correction globally
Temperature and salinity toward which restoring occurs	(Levitus et al., 1994)	(Levitus et al., 1994)	(Levitus et al., 1994)	(Levitus et al., 1994) plus correction factor near Antarctica in winter
Wind stress	Hellerman	Hellerman	ECMWF	ECMWF
Topography	Wide Drake Passage	Wide Drake Passage	Wide Drake Passage	Narrow Drake Passage

Title Page

Abstract

Introduction

Conclusion

References

Tables

Figures

1

71

1

5

Page 1

81

Full Screen / Esc



Table 2. Comparison of the pycnocline and MOC characteristics of the four models from the MOM3 suite and with estimates from observations.

Metric:	LL	HH	LL- ECMWF	P2A	Estimates from observations (in- cluding inverse models and a data- assimilating forward model)
Pycnocline depth (m) ¹	862	881	980	915	834 estimated from Conkright et al. (2001)
$D = \frac{\int_0^{2.5 \text{ km}} (\sigma_{1\max} - \sigma_1) dz}{\int_0^{2.5 \text{ km}} (\sigma_{1\max} - \sigma_1) dz}$					
σ_θ at pycnocline depth ¹	27.31	27.25	27.32	27.28	27.24 (Conkright et al., 2001)
M_n (Sv): max northward transport summed vertically from the surface at 48° N	13.6	17.9	18.4	18.8	19 (Talley 2008) ² 18 (Lumpkin and Speer 2007) ² 16 (ECCO) ³
M_s (Sv): transport at 32° S above the 27.4 isopycnal	6.7	-3.7	15.9	12.2	-1.1 (Talley 2008) 7.9 (Lumpkin and Speer, 2007) 15.8 (ECCO) ³
$M_u = M_n - M_s$ (Sv)	6.9	21.6	2.5	6.6	20.1 (Talley 2008) ² 10.1 (Lumpkin and Speer 2007) ² 0.2 (ECCO) ³

¹ Averaged from 30° S–40° N.² Estimated from inverse models at approximately 48° N (M_n) and 32° S (M_s) using hydrographic data; details in Lumpkin and Speer (2007) and Talley (2008).³ Maximum global overturning at 48° N and 32° S calculated from the Ocean Comprehensive Atlas (OCCA), produced at the Massachusetts Institute of Technology (MIT) by the Estimating the Circulation and Climate of the Ocean (ECCO) group. The OCCA Atlas is publicly available (<http://www.ecco-group.org/>) for the data-rich Argo period from 2004 to 2006 (Forget, 2009). OCCA is an ocean climatology produced by calculating the least squares fit of a global full-depth-ocean and sea-ice configuration of the MIT general circulation model (MITgcm) to satellite and in situ data.**Nutrient return pathways**

J. B. Palter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nutrient return pathways

J. B. Palter et al.

Table 3. Percentage contribution of each tagged water mass to the upper ocean volume (above the 26.8 isopycnal) between 30° S and 30° N in bold lettering. Listed in parentheses are the percentage contributions of *preformed* tagged phosphate (first number) and the *remineralized* tagged phosphate (second number) to the total phosphate inventory above the 26.8 isopycnal. The columnwise sum of the preformed and remineralized contributions (both numbers in parentheses) should be close to 100% of the total phosphate reservoir. However, some columns may sum to less than 100% due to roundoff error and because part of the Bay of Bengal is only partially tagged.

	LL	HH	LL-ECMWF	P2A
North Atlantic	5 (2, 3)	4 (1, 4)	5 (2, 2)	6 (3, 5)
North Pacific	24 (10, 22)	14 (5, 11)	16 (8, 16)	15 (6, 15)
Deep Low latitude	8 (9, 9)	33 (22, 21)	6 (6, 7)	5 (6, 5)
SAMW	60 (18, 21)	45 (12, 13)	68 (26, 28)	67 (24, 24)
AAIW	3 (2, 3)	4 (2, 4)	3 (2, 3)	2.5 (3, 4)
Southern	0 (0, 1)	0 (0, 2)	0 (0, 1)	0 (1, 3)

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)



Nutrient return pathways

J. B. Palter et al.




Fig. 1. Global nutrients at the ocean's surface and in the SAMW layer. **(a)** Surface phosphate from WOA01 (Conkright et al., 2001). Surface $\text{Si}^* = \text{Si(OH)}_4 - \text{NO}_3^-$ from WOA01. **(c)** Si^* on the 26.8 isopycnal, at the center of the SAMW layer. In panels (a) and (b), black hatching indicate regions between 65°S and 65°N where annual maximum mixed layer depths exceed 400 m, where mixed layer depths are taken from de Boyer Montégut et al. (2004). Panel (c) is recreated after Sarmiento et al. (2004). The white area represents the region where the annual mean surface density exceeds 1026.8 kg m^{-3} .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.




Fig. 2. Schematic of the simple analytical model of the meridional overturning circulation (MOC) and the model results of the MOC. **(a)** Adapted from Gnanadesikan (1999), the schematic of the flows (gray arrows) which comprise the MOC and their driving mechanisms (boxed text). The black line schematizes the global pycnocline, below which nutrient-rich water is found, represented by green shading. The sinking of water masses in the Northern Hemisphere is balanced by the sum of the Southern Ocean upwelling (M_s) and low latitude upwelling (M_u). The rate of Southern Ocean upwelling is set by the wind stress over the Southern Ocean (τ_s) and the eddy return flow, which is determined largely by the GM coefficient (A_i). The low latitude upwelling is driven by the downward mixing of heat, which is set by the vertical mixing coefficient (κ_v). **(b)** The zonally integrated meridional transport above the 27.4 isopycnal (northward is positive) for each of the four models described in Table 1. **(c)** The same as panel (b), but below the 27.4 isopycnal. By design, panels (b) and (c) are approximately mirror images of one another because the total transport sums to zero for the volume-conserving ocean of the GCMs. A meridional increase in the meridional transport above (below) the 27.4 signals the upwelling (downwelling) of water across the 27.4 isopycnal. The four models are described in Sect. 2.2 of the text and in Table 1.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.




Fig. 3. A schematic showing the governing equations of the OCMIP-2 biological model. Terms that begin with the letter J are source/sink terms (black text), while terms that begin with the letter F are flux terms (grey text). At depths shallower than 75 m , model PO_4^{3-} (PO_4^{model}) in excess of a seasonal climatology (PO_4^{Obs}) (Louanchi and Najjar, 2000) fuels primary productivity (J_{Prod}), thus restoring the surface model to climatological values over a time scale of $\tau=30$ days (Eq. 1). The resulting productivity has two fates: two thirds becomes a source term for DOP (J_{DOP} , Eq. 2), and one third fuels a parameterized sinking flux of organic phosphorus, that is shunted to 75 m (F_{75} , Eq. 3). Below 75 m , the sinking flux decays as a function of depth (Eq. 4). The vertical divergence of this sinking flux is a source term of PO_4^{3-} (Eq. 5). DOP is remineralized to PO_4^{3-} everywhere in the model, at a rate of $\kappa=0.5 \text{ year}^{-1}$ (Eqs. 5 and 6). This DOP to PO_4^{3-} decay is a sink term for DOP ($-\kappa[DOP]$, Eqs. 2 and 7).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.




Fig. 4. A schematic of the experimental design used to examine the nutrient return pathways to the low latitude surface ocean. Each panel represents the deployment of a regional tag in zonal cross-section. The curvy lines represent selected isopycnals, as labeled at the top of each panel. In the black area, the dye tag, phosphate, and DOP are tagged according to the following conventions: **(A)** the Northern tag (which was further subdivided into North Atlantic and North Pacific) set on all water north of 40° N. **(B)** The Deep Low latitude tag, set for all water with potential densities greater than 27.4 between 40° N and a southern boundary defined by the latitude where the 26.5 isopycnal intersects 200 m. **(C)** The SAMW tag, set between that same boundary and the 27.1 isopycnal. **(D)** AAIW tag, set between the 27.1 and 27.4 isopycnals southward of the same boundary. **(E)** Southern tag for all water denser than AAIW and southward of the boundary. In the white areas, the tag is destroyed and set to its new region. In the gray areas, all tags behave according to their conservation equation.

Nutrient return pathways

J. B. Palter et al.




Fig. 5. Salinity (colors) and density (contours) along 100°E in (a) observations and (b–e) the 4 models, as labeled above each panel. The SAMW layer is visible as a pycnostad between the 26.5 and 27.1 isopycnals just north of the ACC, and the AAIW is characterized by a salinity minimum on the dense side of the SAMW. Observations taken from the CARS2006 database (Ridgway et al., 2002). Vertical lines show the boundary used to separate the SAMW tag region from the low-latitudes in the models.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.




Fig. 6. SAMW dye on the 26.8 isopycnal for the 4 models described in Table 1 and labeled at the top of each panel.

[Title Page](#) | [Discussion Paper](#) | [Discussion Paper](#)

[Abstract](#) | [Introduction](#)
[Conclusions](#) | [References](#)
[Tables](#) | [Figures](#)

[◀](#) | [▶](#)
[◀](#) | [▶](#)
[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)
[Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.




Fig. 7. Zonal and depth-average $\delta^{14}\text{C}$ above **(a)** and below **(b)** the 27.4 isopycnal for the observations from the GLODAP database (Key et al., 2004) and 3 of the 4 ocean models described in Table 1. The LL-ECMWF model is excluded, because it was not run with $\delta^{14}\text{C}$ as a tracer.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.




Fig. 8. Zonally-averaged PO_4^{3-} in (a) the HH model and (b) observations. (c–h) show the percentage of total PO_4^{3-} comprised by each regional phosphate tag in the HH model, as labeled at the top of each panel. The colorbar applies to all panels, but is in mmol m^{-3} for panels (a–b) and percentage of total PO_4^{3-} for (c–h).

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.




Fig. 9. Same as Fig. 8, but for the P2A model.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.




Fig. 10. The fraction of total phosphate above the 26.8 isopycnal comprised by *preformed SAMW phosphate* (top panels) and *remineralized SAMW phosphate* (bottom panels) in the 4 models described in Table 2, as labeled at the top of each map. Note that we refer to *preformed SAMW phosphate* as phosphate that comes directly from the tagging region without ever fueling productivity, while *remineralized SAMW phosphate* derives either from DOP tagged in the SAMW source region and remineralized to phosphate or SAMW phosphate that has fueled primary productivity at least once (see Sect. 2.3).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.




Fig. 11. The zonally-averaged fraction of primary productivity fueled by phosphate (*pre-formed+remineralized*) from each of the tagging regions (left column) and by the *remineralized* components of each tagged phosphate between 30° S and 30° N (right column) for the four models described in Table 1. Note the panels on the left show the full latitude range of the model ocean, while the panels on the right show only the low latitudes. The contribution of the *pre-formed* components of tagged phosphate to sustaining low latitude productivity is the difference between the contributions of the total tagged phosphate (left) and the *remineralized* component of the tagged phosphate (right). Negative values reflect tagged DOP that is respired to phosphate without having fueled primary productivity before entering a destruction region. The lines refer to phosphate from the North Pacific (red solid), North Atlantic (red dotted), Deep low latitude (green solid), SAMW (black solid), AAIW (black dotted), and Southern (magenta solid) regions (refer to Fig. 4 for tagging protocol).

[Title Page](#)
[Abstract](#) [Introduction](#)
[Conclusions](#) [References](#)
[Tables](#) [Figures](#)

[◀](#) [▶](#)
[◀](#) [▶](#)
[Back](#) [Close](#)
[Full Screen / Esc](#)

[Printer-friendly Version](#)
[Interactive Discussion](#)

Nutrient return pathways

J. B. Palter et al.




Fig. 12. Vertical fluxes of SAMW dye (in fractional concentration per $m^2 s$) at the depth of the 26.5 isopycnal. Upper row for the HH model and lower row for P2A. The left hand panels are the vertical advective flux (wC , where w is the vertical velocity and C is the concentration of the SAMW dye tag). The right hand panels are the vertical diffusive flux ($\kappa_v dC/dz$, where κ_v is the coefficient of diapycnal diffusion). The colorbar applies to all panels, and positive numbers represent an upwards flux. The line plots are the zonal means of the maps (black for vertical advection and blue for vertical diffusion).

Nutrient return pathways

J. B. Palter et al.




Fig. 13. Percentage of low latitude (30° S– 30° N) primary productivity fueled by nutrients from different tagged regions, separated by *preformed* and *remineralized* components and by model configuration. The salient feature of the graph is that a majority (57–75%) of low latitude productivity is sustained by SAMW nutrients, except in the HH model configuration (33%). Roughly half of the SAMW nutrients sustaining low latitude productivity are from the preformed pool. Negative values for *remineralized* tags indicate that *preformed* DOP tagged in a given source region is respiration to phosphate without fueling primary productivity. Such is the case for almost all of the *Southern DOP* and much of the *North Atlantic DOP*.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)