

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# CO<sub>2</sub> maximum in the oxygen minimum zone (OMZ)

**A. Paulmier<sup>1</sup>, D. Ruiz-Pino<sup>2</sup>, and V. Garçon<sup>1</sup>**

<sup>1</sup>LEGOS, UMR 5566, Toulouse, France

<sup>2</sup>LOCEAN, Paris, France

Received: 3 August 2010 – Accepted: 4 August 2010 – Published: 25 August 2010

Correspondence to: A. Paulmier (aurelien.paulmier@legos.obs-mip.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Oxygen minimum zones (OMZs), known as suboxic layers mainly localized in the Eastern Boundary Upwelling Systems, are expanding since the 20th “high CO<sub>2</sub>” century, probably due to the global warming. OMZs are also known to contribute significantly to the oceanic production of N<sub>2</sub>O, a greenhouse gas (GHG) more efficient than CO<sub>2</sub>. However, the contribution of the OMZs on the oceanic sources and sinks budget of CO<sub>2</sub>, the main GHG, still remains to be established.

We present here the dissolved inorganic carbon (DIC) structure, associated locally with the Chilean OMZ and globally with the main most intense OMZs (O<sub>2</sub> < 20 μmol/kg) in the open ocean. To achieve this, we jointly examine simultaneous DIC and O<sub>2</sub> data collected off Chile during 4 cruises and a monthly monitoring (2000–2002) in one of the shallowest OMZ, along with international DIC and O<sub>2</sub> databases for other OMZs.

High DIC concentrations (>2225 μmol/kg, up to 2350 μmol/kg) have been reported over the whole OMZ thickness, allowing to define for all studied OMZs a Carbon Maximum Zone (CMZ). The CMZs-OMZs constitute the largest carbon reserves of the ocean in subsurface waters and could induce a positive feedback for the atmosphere during upwelling activity, as potential direct local sources of CO<sub>2</sub>. The CMZ paradoxically presents a slight “carbon deficit” in its core, meaning a DIC increase from the oxygenated ocean to the OMZ lower than the corresponding O<sub>2</sub> decrease (assuming classical C/O molar ratios). This “carbon deficit” would be related to thermal mechanisms affecting faster O<sub>2</sub> than DIC (due to the carbonate buffer effect) and occurring upstream in warm waters (e.g., in the Equatorial Divergence), where the CMZ-OMZ core originates. The “carbon deficit” in the CMZ core would be mainly compensated locally at the oxycline, by a “carbon excess” induced by a specific remineralization. Indeed, a possible co-existence of bacterial heterotrophic and autotrophic processes usually occurring at different depths could stimulate an intense aerobic-anaerobic remineralization, inducing deviation of C/O molar ratio from the canonical Redfield ratios. Further studies to confirm these results for all OMZs are

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



required to understand the OMZ effects on both climatic feedback mechanisms and marine ecosystem perturbations.

## 1 Introduction

The most intense ( $O_2 < 20 \mu\text{mol/kg}$ ) and largest oxygen minimum zones (OMZs), known as suboxic layers, are mainly localized in subsurface of the upwelling regions in the Eastern Pacific and Northern Indian open oceans (Paulmier and Ruiz-Pino, 2008). OMZs can be considered as analogs of the primitive anoxic ocean associated with a high  $CO_2$  atmosphere ( $> 1000 \text{ ppmv}$ : Royer et al., 2004), because of comparable reduced chemical conditions and similarities between ancient bacteria and those living in the OMZs (e.g., Planctomycetales: Gribaldo and Philippe, 2002). OMZs appear as a refuge of abundant specific microbes capable of chemolithoautotrophic carbon assimilation (Walsh et al., 2009) and organisms adapted to low  $O_2$  (e.g., zooplankton as specific copepods and euphausiids: Wishner et al., 2008; Antezana, 2009). These biological specificities and perturbations, associated with shift in animals distribution and changes in ecosystems structure (e.g., towards gelatinous species as jellyfish and salps: Childress and Siebel, 1998), may potentially impact the carbon export. In addition, OMZs are submitted to climate changes (lower ventilation by stratification and  $O_2$  solubility decrease) and natural or anthropogenic fertilization (remineralization increase) through nutrient or metal inputs by upwelling, rivers or dust (Fuhrman and Capone, 1991; Naqvi et al., 2006; Rabalais et al., 2010). In the past, OMZs have probably extended and contracted in warm (interglacial) and cold (glacial) periods (e.g., Cannariato and Kennett, 1999; Rogers, 2000), associated with high (300 ppmv) and low (180 ppmv) atmospheric  $CO_2$  (Jansen et al., 2007), respectively. In the present and according to the last decades observations associated with high anthropogenic  $CO_2$  uptake by the ocean (Sabine et al., 2004), OMZs would increase or intensify (Stramma et al., 2008), or even new ones appear locally and episodically (Chan et al., 2008).

## **$CO_2$ maximum in the OMZ**

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**CO<sub>2</sub> maximum in the OMZ**

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



OMZs are known for playing an essential role in the nitrogen cycle, in which different autotrophic and heterotrophic bacterial processes intervene (e.g., Arrigo, 2005) such as ammonification, nitrogen assimilation, anammox, DNRA (Dissimilatory nitrate reduction to ammonia), denitrification and nitrification. As a result of these two last processes, OMZs would produce up to 50% of the oceanic N<sub>2</sub>O (e.g., Bange, 2008), a greenhouse gas, ~300 times more efficient than CO<sub>2</sub> (Jain et al., 2000; Ramaswamy et al., 2001). So far, no study has investigated the direct link between OMZs and the most important greenhouse gas, CO<sub>2</sub>.

However, OMZs may play an indirect role on the carbon reservoir through denitrification which converts nitrate, one of the main oceanic nutrients, into gaseous nitrogen (e.g., N<sub>2</sub>, N<sub>2</sub>O) lost towards the atmosphere. Indeed, according to Falkowski's (1997) hypothesis, the denitrification process would decrease the oceanic nitrogen reserve, and consequently total primary production reduction will limit atmospheric CO<sub>2</sub> sequestration by the ocean. Locally, Paulmier et al. (2006) suggested from a study in the OMZ off Chile a specific remineralization at the oxycline up to 5 times more intense than in the oxygenated ocean, which consequently produces more CO<sub>2</sub> than classically. In addition, Koeve and Kähler (2010) show theoretically that the combined effects of the processes in the OMZ core turn out to be clearly heterotrophic, suggesting net OMZ CO<sub>2</sub> release.

The main objective of this study is thus to determine and discuss the total CO<sub>2</sub> or DIC (dissolved inorganic carbon) structures which are forming in the OMZs, from a study focused in the OMZ off Chile and extended to the global ocean.

## 2 Methodology

To characterize the DIC structure associated with an OMZ, a high vertical resolution sampling (5–10 m), adapted to low O<sub>2</sub> concentrations (<20 μmol/kg), has been performed simultaneously for DIC and O<sub>2</sub> in the OMZ off Chile to capture the intense biogeochemical gradients. This sampling is based on 4 cruises between 2000 and

2002 in non El Niño years (at 21° S with a cross-shore transect and at 30° S) and a monthly monitoring at 36° S off a semi-enclosed bay (S14 station), to document illustrative spatio-temporal variations detailed in Paulmier et al. (2006).

Total dissolved inorganic CO<sub>2</sub> (DIC) samples were analyzed by potentiometry (Bradshaw and Brewer, 1988) with the National CO<sub>2</sub> measurement Service (SNAPCO<sub>2</sub>, Paris, France) protocols (e.g., Corbière et al., 2007). From 15 certified Dickson standards (Batches #44 and #49, SCRIPPS, USA), an accuracy of ~1.7 µmol/kg and, a reproducibility of ~2.7 µmol/kg, based on 16 replicates, have been determined (Paulmier et al., 2008). O<sub>2</sub> samples were analyzed according to the Winckler method, improved and adapted to low concentration detection (O<sub>2</sub> < 40 µmol/kg) following Broenkow and Cline (1969), with an accuracy of 1 µmol/kg and reproducibility of 1.8 µmol/kg (Paulmier et al., 2006).

To validate results obtained off Chile for the most intense OMZs of the world ocean, the international data bases GLODAP (GLObal Ocean Data Analysis Project; Sabine et al., 2005) for DIC and WOA2005 (World Ocean Atlas, 2005; Boyer et al., 2006) for O<sub>2</sub> have been used at the global scale. For a more consistent comparison between DIC and O<sub>2</sub> data in terms of spatio-temporal coverage and sampling protocols, the analysis has been focused on the same cruises carried out during the WOCE program (World Ocean Circulation Experiment, www.coaps.fsu.edu/woce; >300 transects between 1992 and 1998). Indeed, the horizontal coverage by WOCE cruises measuring both DIC and O<sub>2</sub> (between 1992 and 1995, also in non El Niño years) is very similar in the main OMZ regions of the Pacific and Indian Oceans. Both accuracy and reproducibility of the data, respecting WOCE standards quality (WOCE Operation Manual, 1994), are: ~4 µmol/kg for DIC; ~2 µmol/kg (with a detection limit of ~10 µmol/kg) for O<sub>2</sub>.

For comparing DIC increase versus O<sub>2</sub> decrease due to remineralization, which has been considered as the main mechanism responsible of the existence of all O<sub>2</sub> minima observed in the ocean (Wyrтки, 1962), an index is proposed:

$$DIC^* = DIC + 0.69O_2 \quad (1)$$

0.69=117/170 is the stoichiometric carbon to oxygen ratio C/O of Anderson and

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sarmiento (1994). It corresponds to the lowest and most commonly used C/O ratio value in the data analysis literature.

DIC\* is defined as the sum of the observed DIC and DIC produced during remineralization by O<sub>2</sub> consumption. DIC\* can be considered as an analogue of nitrogen deficit index (e.g.: NO of Broecker, 1974; N\* of Gruber and Sarmiento, 1997) or P\* (Deutsch et al., 2007). DIC\* should be conservative if no physico-chemical and/or biological mechanisms are affecting DIC and O<sub>2</sub> concentrations other than remineralization and/or photosynthesis with the same C/O stoichiometry. The accuracy and reproducibility of DIC and O<sub>2</sub> measurements lead to an accuracy and reproducibility for DIC\* lower than 4.5 µmol/kg.

A water mass analysis based on an inverse method is performed to estimate the biogeochemical contributions for DIC production and O<sub>2</sub> consumption locally within the OMZ off Chile. This hydrological method (e.g., Minas et al., 1986) has been already described for O<sub>2</sub> in Paulmier et al. (2006) and detailed in Paulmier (2005). The contribution due to mixing processes has been determined with a correlation coefficient  $R^2$  of ~0.9, using potential temperature and salinity as semi-conservative parameters. The measured DIC and O<sub>2</sub> concentrations can be expressed as the sum of a contribution due to mixing processes (DIC<sub>mixing</sub> and O<sub>2 mixing</sub>) and a contribution due only to biogeochemical processes called ΔDIC and ΔO<sub>2</sub>. Positive or negative ΔDIC and ΔO<sub>2</sub> indicate (DIC and O<sub>2</sub>) production or consumption, respectively.

### 3 Results

DIC and O<sub>2</sub> data sets collected off Chile (Fig. 1) show a specific DIC structure associated with an OMZ already described in Paulmier et al. (2006), mainly characterized by an upper O<sub>2</sub> gradient (oxycline) and a core (O<sub>2</sub><20 µmol/kg, down to <1 µmol/kg). In particular, the oxycline is: i) very intense (>1 µmol/kg/m) and stratified; ii) associated with an O<sub>2</sub> continuum from oxic (O<sub>2</sub>>200 µmol/kg) to suboxic (O<sub>2</sub><20 µmol/kg) conditions; iii) shallow (from 5–20 m), then intercepting the euphotic

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



layer depth. Then, in order to determine if the description of the DIC structure detailed for the OMZ off Chile can be extended to the global scale, an analysis of the representative DIC structure in the main most intense OMZs ( $O_2 < 20 \mu\text{mol/kg}$ ; Paulmier and Ruiz-Pino, 2008) is carried out. The DIC structure was determined in the permanent OMZs in the Tropical Pacific ocean (North – ETNP: Costa Rica, Mexico; South – ETSP: Peru) and Northern Indian ocean (Arabian sea – AS; Bay of Bengal – BB), in the seasonal OMZs in the West Bering Sea (WBS) and the Gulf of Alaska (GA), and in the Eastern Boundary Upwellings of the Atlantic Ocean (Canary and Benguela/Namibia).

### 3.1 Carbon maximum zone (CMZ) along the OMZ off Chile

The DIC structure, observed at  $21^\circ \text{S}$  from the shelf to off-shore (Iquique: Fig. 1a and c) but also at  $30^\circ \text{S}$  (Coquimbo: Fig. 1c; stations location in Fig. 1f), corresponds to a permanent structure off Northern and Central Chile. The DIC structure at  $21^\circ \text{S}$  shows low values ( $\text{DIC} < 2140 \mu\text{mol/kg}$ ) in surface, and a strong increase with depth from 20 m called carboncline (by analogy with the oxycline) up to maximal DIC concentrations ( $> 2300 \mu\text{mol/kg}$ ) on the continental shelf. The continental slope induces a 30–40 m shoaling of the carboncline (cf., DIC isolines between 2200 and  $2275 \mu\text{mol/kg}$ ) and of the maximal DIC ( $> 2250 \mu\text{mol/kg}$ : in red-white, Fig. 1a). This shoaling follows the oxycline and OMZ core shoaling (cf., the  $20 \mu\text{M } O_2$  isoline, Fig. 1d). The DIC structure at  $21^\circ \text{S}$  ( $\sim 2300 \mu\text{mol/kg}$  in average on the core) presents higher concentrations than at  $30^\circ \text{S}$  by  $\sim 50 \mu\text{mol/kg}$  (Fig. 1c), also associated with a less intense OMZ at Coquimbo from  $\sim 40 \mu\text{mol/kg}$  (Fig. 1f). At  $36^\circ \text{S}$  on the shelf (Concepcion Bay), DIC concentrations are much more variable than at  $21^\circ \text{S}$  and  $30^\circ \text{S}$ , with the highest concentrations reported off Chile ( $> 2500 \mu\text{mol/kg}$  in spring 2000; Fig. 1b and c). The highest DIC concentrations ( $> 2250 \mu\text{mol/kg}$ : in red, Fig. 1b) appear from the bottom ( $\sim 50 \text{ m}$  depth) up to 10 m depth in austral spring-summer (October–March), associated with the OMZ establishing ( $O_2 < 20 \mu\text{M}$  from 20 to 50 m in white: Fig. 1e). The DIC maximum ( $> 2250 \mu\text{mol/kg}$ ) starts to settle in spring during the high upwelling activity period,

## **CO<sub>2</sub> maximum in the OMZ**

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





simultaneously with the OMZ core. In fall (April–May), the 2175  $\mu\text{mol/kg}$  DIC isoline extends from 5 m near the surface to the bottom (Fig. 1b), indicative of a homogenized water column with lower DIC. It is coincident with the complete OMZ destruction (Fig. 1e). In the OMZ off northern and central Chile, the DIC maximum appears spatially and temporally collocated with the  $\text{O}_2$  minimum of the OMZ core, despite differences in the OMZ intensity and shape between each site (Fig. 1f).

Considering all sampled data in the OMZ off Chile (Fig. 1), the whole DIC and  $\text{O}_2$  profiles confirm a same DIC structure associated with the OMZ: low concentrations in surface waters and an increase with depth, as in the oxygenated ocean (cf., classical profile out of the OMZ: Fig. 1c). However, in average, the observed increase off Chile is  $\sim 5$  times stronger than the usual increase. DIC concentrations reach values as high as  $2330 \pm 60 \mu\text{mol/kg}$  at 50–100 m. Usually for the same area but out of the OMZ, these high DIC are observed from 1000 m and deeper (WOCE, 1992–1995; Takahashi and Goddard, 1998). In addition, DIC concentrations in the OMZ off Chile remain high ( $>2250 \mu\text{mol/kg}$ ) on the whole core thickness ( $\text{O}_2$  minimum between 30 and 750 m). Thus, as compared with a representative classical DIC profile out of the OMZ, the OMZ off Chile presents a relative maximum of DIC (hatched in red: Fig. 1c). This DIC maximum, defined by  $\text{DIC}_{\text{CMZ}} = \text{DIC}_{\text{Average in OMZ}} - \text{DIC}_{\text{Classical out of OMZ}}$ , is called, by analogy with the OMZ, a carbon maximum zone (CMZ).

Although the CMZ defined here and the OMZ off Chile are relatively well correlated in terms of position and thickness, some differences appear, especially between the carboncline and the oxycline. The carbonclines start from the surface, at shallower depths than the oxyclines (from 5–10 m). In addition, the carbonclines are associated with some extreme DIC peaks (up to  $>2350 \mu\text{mol/kg}$ ), whereas smooth low  $\text{O}_2$  profiles can be observed at the oxycline.

### 3.2 Towards a validation of the CMZ existence at the global scale

The hypothesis of the CMZs existence at the global scale for all the most intense OMZs is tested with GLODAP and WOA2005 datasets.

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Vertically, the DIC distributions in the OMZs in the Eastern Tropical South and North Pacific (ETSP and ETNP), in the Arabian Sea (AS) and in the Bay of Bengal (BB), are low in surface (between 1900 and 2100  $\mu\text{mol/kg}$ ) and increase with depth up to 2300  $\mu\text{mol/kg}$  (Fig. 2a). As observed in the OMZ off Chile, all these OMZs present high DIC concentrations (always  $>2225 \mu\text{mol/kg}$ , Fig. 2a) over their core depths in average between  $\sim 175$  and  $\sim 950$  m depth (Fig. 2b). These high DIC concentrations in the OMZ cores are associated with a large DIC relative maximum, higher than 100  $\mu\text{mol/kg}$  as compared to the global averaged DIC profile (continuous profile in Fig. 2a). These large DIC relative maxima allow to suggest the existence of CMZs in all main OMZs, as defined for the OMZ off Chile.

The CMZs are the most intense in ETNP ( $\text{DIC} > 2300 \mu\text{mol/kg}$ ) and the least intense in AS and BB, up to 150  $\mu\text{mol/kg}$  and  $\sim 100 \mu\text{mol/kg}$  higher than the average for the global ocean, respectively (Fig. 2a). This is likely a consequence of the higher accumulation of DIC along the Conveyor Belt in the ETNP CMZ than for the CMZs in the Northern Indian Ocean. In particular, the CMZ in BB presents the lowest DIC concentrations ( $\text{DIC} \approx 1800 \mu\text{mol/kg}$ ) near the surface, probably due to the advection of important riverine freshwaters (e.g., Brahmaputra, Gange).

Horizontally, the analysis has been illustrated on DIC and  $\text{O}_2$  fields over the 175 and 950 m layer, intercepting the core of all main tropical OMZs (Fig. 2b and d). The highest DIC concentrations ( $>2225 \mu\text{mol/kg}$ : in red-orange, Fig. 2b) are found in the Tropical Eastern North (ETNP) and South (ETSP) Pacific and Northern Indian Oceans (AS; BB), and at high latitude around  $60^\circ \text{N}$  in the Northern Pacific. These regions correspond to the most intense ( $\text{O}_2 < 20 \mu\text{mol/kg}$ ) OMZs and to the seasonal OMZs of the West Bering Sea and the Gulf of Alaska, respectively (in blue-purple, Fig. 2d). The Canary and Benguela upwelling systems and the China Sea, corresponding to the least intense  $\text{O}_2$  minima ( $\text{O}_2 \geq 40 \mu\text{mol/kg}$ : Karstensen et al., 2008; Chen et al., 1995; respectively) also correspond to high DIC (maxima around 2225  $\mu\text{mol/kg}$ ), albeit lower concentrations than for the Pacific and Indian OMZs. All main OMZs appear thus as potential CMZs. The reciprocal is not true, since the high DIC concentrations averaged

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



between 175 and 950 m depth south of 50° S in the Southern Ocean ( $\sim 2250 \mu\text{mol/kg}$ ) are not correlated with an OMZ. These high DIC concentrations are mainly associated with the deepest maximal mixed layer depths ( $\geq 250 \text{ m}$ : De Boyer Montégut et al., 2004).

The existence of CMZ has been validated in all main OMZs on both vertical mean profiles for each OMZs (Fig. 2a and c) and horizontal distribution (Fig. 2b and d). In addition, sensitivity tests on DIC and  $\text{O}_2$  averages have been performed, taking into account the lower and upper vertical levels of the GLODAP and WOA2005 grids for each OMZ limit: 137.5 and 225 m for the upper limit (175 m); 850 and 1050 m for the lower limit (950 m). These tests have been summarized in Table 1 for the total OMZs, each OMZ and the global ocean. In general, the average DIC are more sensitive to the choice of the lower CMZ limit and in the ETNP ( $\pm 8 \mu\text{mol/kg}$ ), where the DIC concentrations are the highest. On the contrary, the average  $\text{O}_2$  are more sensitive to the choice of the upper CMZ-OMZ limit, corresponding to the highest  $\text{O}_2$  concentrations close to the saturation values. They are less sensitive in AS, with the lowest  $\text{O}_2$  concentration on the thickest OMZ core. The maximal differences for both DIC and  $\text{O}_2$ , lower than to the data accuracy + reproducibility (cf., Methodology), suggest that the choice of the upper and lower CMZ-OMZ limits does not have a significant effect on results of the analysis proposed here.

## 4 Discussion

The CMZs-OMZs, because of the accumulation of high DIC ( $> 2225 \mu\text{mol/kg}$ ) in subsurface, constitute the largest reserves of carbon near the surface in the global ocean,  $\sim 570 \text{ Pg C}$ , i.e. a comparable stock to the total atmospheric  $\text{CO}_2$  content. This simple calculation is based on a mean DIC concentration of  $\sim 2225 \mu\text{mol/kg}$ , reported over 940 m of the OMZ thickness between 10 and 950 m, multiplied by the total area of the tropical shallow OMZs ( $22.2 \times 10^6 \text{ km}^2$ : from Paulmier and Ruiz-Pino, 2008). The high DIC concentrations of  $\sim 2225 \mu\text{mol/kg}$  are considered on both the

**BGD**

7, 6353–6385, 2010

## **$\text{CO}_2$ maximum in the OMZ**

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



carbonline-oxycline (10–175 m, as reported from local cruises: cf. Fig. 1c) and CMZ cores (175–950 m, as reported from global data bases: cf. Fig. 2a). The existence of such CMZ explains why the OMZ can be an intense GHG oceanic local source of N<sub>2</sub>O but also CO<sub>2</sub> (e.g., off Chile: Paulmier et al., 2008).

The mechanisms responsible for the carbon maximum structure (CMZ) associated with the OMZ are discussed. We will first examine the physico-chemical mechanisms inducing a differential CO<sub>2</sub> accumulation and deoxygenation in the CMZ-OMZ core. We will then focus our attention on more local and biogeochemical processes, associated with remineralization and carbon to oxygen molar ratio (C/O) at the carbonline-oxycline of the Chilean OMZ.

#### 4.1 Differential regional physico-chemical CO<sub>2</sub> accumulation and deoxygenation in the CMZ-OMZ core

It is in no way surprising to find DIC maxima associated with O<sub>2</sub> minima in subsurface, since remineralization degrading organic matter into CO<sub>2</sub> occurs mainly in subsurface layer and is classically aerobic, i.e. consuming O<sub>2</sub>. CMZ in the OMZ is thus a logical consequence of remineralization end-products accumulation, agreeing with subsurface fields of nitrate, phosphate and silicate at the depth of the OMZ core. This accumulation of remineralization end-products is not distributed uniformly horizontally (Fig. 2b) and vertically (Fig. 2a). It mainly occurs in privileged zones, in particular the OMZs in the top 1000 m. The same dynamical mechanisms acting at different scales, already proposed for the OMZ formation contribution (e.g., Paulmier and Ruiz-Pino, 2008; Karstensen et al., 2008), contribute to the CMZ formation. At the global scale, DIC accumulation occurs in the oldest waters, following the Conveyor Belt (Broecker and Peng, 1982) with the highest concentrations in Northern Pacific (the seasonal West Bering Sea and Gulf of Alaska OMZs). Low subsurface ventilation of shadow zones (Luyten et al., 1983) and confinement in regions surrounded by continents explain why CMZ-OMZs are located in the Eastern Pacific and in the Northern Indian Ocean rather than in the Western Pacific and in the Southern Indian, respectively. The location

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of tropical shadow zones with older waters (10<100 years: Karstensen et al., 2008) explains why the DIC maximum and O<sub>2</sub> minimum are found in the equatorwards subsystems of the eastern boundary upwelling in both the Atlantic and Pacific Oceans. The upwelling activity (intensity; typology of cells: Endoh, 1976) and undercurrents transport (intensity, depth), as well as mixed-layer depth, affects the position of the CMZ-OMZs in subsurface (Paulmier and Ruiz-Pino, 2008). In particular, the upwelling explains why the DIC maximum starts from shallower (<175 m) in the CMZ-OMZs than the rest of the ocean in average (Fig. 2b and d).

However, despite of similarities and co-localization of CMZ and OMZ, some differences appears. Let's assume that the dynamical transport and accumulation of DIC produced by remineralization is the only cause explaining the CMZ existence in an OMZ, then the DIC increase should be proportional to the O<sub>2</sub> decrease. The proportionality factor should be the C/O stoichiometric coefficient, if the molar C/O ratio of remineralization is constant. In this case, values of DIC\* (=DIC+0.69O<sub>2</sub>; Eq. 1) should be similar in the oxygenated ocean and in the OMZs. But, as illustrated in Fig. 3a and b, DIC\* averages in OMZs core (DIC\*<sub>OMZ</sub>) are slightly lower than the average in the global ocean (DIC\*<sub>Global</sub>) for the same depth layer, by a significant difference of ~10–20 μmol/kg ranging from 7 (ETNP) up to 25 (AS) μmol/kg. This difference is higher than DIC\* uncertainty (cf. Methodology). This difference suggests that the CMZ core exhibits a slight DIC deficit compared to the expected DIC maximum, i.e. the DIC produced by a remineralization (Re) consuming the entire O<sub>2</sub> amount necessary to form the OMZ and following the chosen constant C/O stoichiometry. Estimating a DIC<sub>Re</sub> of ~93 μmol/kg (=0.69\*(157–22) from O<sub>2</sub><sub>OMZ</sub>=22 μmol/kg and O<sub>2</sub><sub>Global ocean</sub>=157 μmol/kg; Table 1), the DIC deficit is accounting around >10% of the total DIC produced by remineralization. Note that for all DIC\* calculations presented here, sensitivity tests were performed, especially with a higher C/O value (the canonical C/O=106/138 from Redfield et al., 1963). These tests did not change the conclusions.

Theoretically, the history of the water masses which give origin to the CMZ-OMZ will affect the DIC and O<sub>2</sub> values in addition to the strict biological effect. Water

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**CO<sub>2</sub> maximum in the OMZ**

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



masses forming the CMZ-OMZs cores come from surface equatorial and/or tropical warm waters. In the Pacific, CMZs-OMZs waters mainly originate from the Pacific equatorial waters advected through zonal equatorial undercurrents (e.g., Tsuchiya, 1981; Karstensen et al., 2008): i) from the Equatorial SubSurface Water through the Peru-Chili undercurrent (PCU) for the ETSP (Strub et al., 1998; Paulmier et al., 2006); ii) from the SubTropical Underwater Water formed near the Equator for the ETNP (O'Conner et al., 2002; Maske et al., 2010). The AS CMZ-OMZs waters originate from the Red Sea and Persian Gulf (Morrison et al., 1999) and the BB subsurface waters come from the surface Equatorial Indian Ocean (Rao et al., 1996; Sarma et al., 1998). The equatorial regions are net O<sub>2</sub> sinks, despite a thermal O<sub>2</sub> source effect (Najjar and Keeling, 2000), which tends to diminish the surface O<sub>2</sub> concentration thereby affecting DIC\*. Since DIC\* = DIC + 0.69O<sub>2</sub>, a decrease of DIC\* is expected in the warm regions due to the higher temperature lowering the gas solubility. Indeed, an increase of temperature from 15° to 25°C, corresponding to the mean subsurface and surface temperatures for tropical and equatorial regions, and for a standard mean salinity of 35 (CARS database, 2009), would decrease the O<sub>2</sub> solubility by >10% (Weiss, 1970). A similar additional effect is occurring on dissolved CO<sub>2</sub> (Weiss, 1974), but it remains negligible because of the buffer effect due to the CO<sub>2</sub> dissociation (<1%; Lewis and Wallace, 1998).

The global dataset analysis confirms that the differential CO<sub>2</sub> accumulation and deoxygenation is occurring upstream in the regions where the CMZs-OMZs water are pre-formed, mainly in the equatorial regions (Fig. 4). In the subsurface layer (175–950 m) corresponding to the CMZs-OMZs depths, DIC\* is decreasing from the global ocean to the equatorial regions by ~10 μmol/kg (Fig. 4c), whereas DIC\* is similar in the CMZs-OMZs and in equatorial regions. For Chile, OMZ DIC and O<sub>2</sub> concentrations in PCU confirm this result, since PCU is advecting waters poorer in O<sub>2</sub> (Paulmier et al., 2006) than richer in DIC, inducing a lower DIC\*. The differential thermal effect on DIC and O<sub>2</sub> is needed to explain why DIC\* decreases from the global ocean (GO) to Equatorial regions (ER), from a DIC\* value of 2311 μmol/kg to a DIC\* value of

2300  $\mu\text{mol/kg}$  (Fig. 4c). Let's use the  $\text{DIC}^*$  definition and the decomposition of  $\text{DIC}^*$ ,  $\text{DIC}$  and  $\text{O}_2$  into the physical and biogeochemical components. Let's assume that the thermal effect (Th) and remineralization (Re) can be associated to the physical and biogeochemical components in subsurface waters, respectively.

$$\text{DIC}_{\text{GO}}^* - \text{DIC}_{\text{ER}}^* = (\text{DIC}_{\text{GO Th}}^* + \text{DIC}_{\text{GO Re}}^*) - (\text{DIC}_{\text{ER Th}}^* + \text{DIC}_{\text{ER Re}}^*)$$

Because  $\text{DIC}_{\text{GO Re}}^* - \text{DIC}_{\text{ER Re}}^* = 0$  by definition,

$$\begin{aligned} \text{DIC}_{\text{GO}}^* - \text{DIC}_{\text{ER}}^* &= \text{DIC}_{\text{GO Th}}^* - \text{DIC}_{\text{ER Th}}^* \\ &= (\text{DIC}_{\text{GO Th}} + 0.69\text{O}_{2 \text{ GO Th}}) - (\text{DIC}_{\text{ER Th}} + 0.69\text{O}_{2 \text{ ER Th}}) \\ &= (\text{DIC}_{\text{GO Th}} - \text{DIC}_{\text{ER Th}}) + 0.69(\text{O}_{2 \text{ GO Th}} - \text{O}_{2 \text{ ER Th}}) \end{aligned}$$

Considering here  $\text{DIC}_{\text{GO Th}} - \text{DIC}_{\text{ER Th}} \approx 0.01(\text{CO}_{2 \text{ GO Th}} - \text{CO}_{2 \text{ ER Th}})$  with a constant  $\text{CO}_2$  buffer effect of 1% and a negligible temperature effect on this Revelle factor of 0.01 (Lewis and Wallace, 1998), and assuming  $\text{CO}_{2 \text{ GO Th}} - \text{CO}_{2 \text{ ER Th}} \approx \text{O}_{2 \text{ GO Th}} - \text{O}_{2 \text{ ER Th}}$  (similar solubility variation for  $\text{CO}_2$  and  $\text{O}_2$ ):

$$\begin{aligned} \text{DIC}_{\text{GO}}^* - \text{DIC}_{\text{ER}}^* &= 0.01(\text{O}_{2 \text{ GO Th}} - \text{O}_{2 \text{ ER Th}}) + 0.69(\text{O}_{2 \text{ GO Th}} - \text{O}_{2 \text{ ER Th}}) \\ &= 0.70(\text{O}_{2 \text{ GO Th}} - \text{O}_{2 \text{ ER Th}}) \end{aligned}$$

with  $\text{O}_{2 \text{ GO Th}} - \text{O}_{2 \text{ ER Th}} = (\text{O}_{2 \text{ GO}} - \text{O}_{2 \text{ ER}}) - (\text{O}_{2 \text{ GO Re}} - \text{O}_{2 \text{ ER Re}})$  and

$$(\text{O}_{2 \text{ GO Re}} - \text{O}_{2 \text{ ER Re}}) = -(\text{DIC}_{\text{GO Re}} - \text{DIC}_{\text{ER Re}})/0.69 \approx -(\text{DIC}_{\text{GO}} - \text{DIC}_{\text{ER}})/0.69 \quad (2)$$

$$\begin{aligned} \text{O}_{2 \text{ GO Th}} - \text{O}_{2 \text{ ER Th}} &= (\text{O}_{2 \text{ GO}} - \text{O}_{2 \text{ ER}}) + (\text{DIC}_{\text{GO}} - \text{DIC}_{\text{ER}})/0.69 \\ &= 77 - 61/0.69 = 16 \mu\text{mol/kg} \text{ (from the dashed arrows: Fig. 4a} \end{aligned}$$

and b), and

$$\text{DIC}_{\text{GO}}^* - \text{DIC}_{\text{ER}}^* = 0.70 \cdot 16 \approx 11 \mu\text{mol/kg}$$

as expected from the direct calculation.

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Therefore, the temperature effect on the solubility appears to be the mechanism decreasing the  $O_2$  concentration in the CMZs-OMZs originating regions, and mainly driving and explaining the slightly lower  $DIC^*$  observed in the CMZs-OMZs (Fig. 3). In particular, the faster air-sea exchanges for  $O_2$  than for  $CO_2$  should explain why carbonclines start slightly shallower (by 5–10 m) than oxyclines, in addition to the photosynthesis effect in surface depleting more DIC than the supersaturating  $O_2$  (e.g., Minas et al., 1986). Our approach only demonstrates that a DIC deficit ( $DIC^* < DIC^*_{Global}$ ) could be due to physico-chemical mechanisms prior to the complete CMZ-OMZ formation, since dynamical connection between Equator waters and OMZ cores is complex, indirect and still not very well known.

## 4.2 Local remineralization and biogeochemical C/O variations at the carboncline-oxycline

In addition to regional physico-chemical contribution to the DIC deficit of the CMZ-OMZ core, specific local mechanisms could affect C/O in OMZs, and thus help to explain why  $DIC^*_{OMZ}$  is different from  $DIC^*_{Global}$ . Off Chile (Fig. 3c), the CMZ-OMZ core also presents a  $DIC^*$  ( $\sim 2225 \mu\text{mol/kg}$ ) lower than in oxygenated waters, by a difference up to  $\sim 60 \mu\text{mol/kg}$  and  $\sim 20 \mu\text{mol/kg}$  in average similar to those discussed from the global dataset (Fig. 3a and b). On the contrary, above the core, at the carboncline-oxycline depths between 10 and 175 m depending on the site,  $DIC^*_{Chile}$  is much higher by  $\sim 80 \mu\text{mol/kg}$  in average than in oxygenated waters and in the core (Fig. 3c). The highest variability (up to  $\sim 750 \mu\text{mol/kg}$ ) and concentrations ( $DIC^* > 2500 \mu\text{mol/kg}$ ) correspond to  $36^\circ \text{S}$  (Fig. 3c), the shallowest site on the continental margin associated with a seasonal CMZ-OMZ (Fig. 1b and e). The variation of  $DIC^*$  can be due to the local deviation of the C/O in the CMZ-OMZ from the fixed value chosen for the global scale analysis. Indeed, a plot of  $[DIC]$  over  $[O_2]$  for the global ocean (85 650 data points from the WOCE cruises; Fig. 5a) indicates that a linear regression leads to a slope of 0.7 very close to  $C/O=0.69$  (Anderson and Sarmiento, 1994) chosen for our  $DIC^*$  computation. This plot presents strongly higher and lower

**BGD**

7, 6353–6385, 2010

## **CO<sub>2</sub> maximum in the OMZ**

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





deviations from a linear regression, indicated by a low coefficient  $R^2=0.25$ . We thus investigated for local C/O deviations, especially at the carboncline-oxycline. To study the potential influence of the high remineralization on DIC production contributing to the CMZ formation, we evaluated the biogeochemical DIC production. Results of the hydrological method provide generally positive  $\Delta\text{DIC}=\text{DIC}_{\text{measured}}-\text{DIC}_{\text{mixing}}$ , indicating a net “new” DIC production by biogeochemical processes (Fig. 1c), in opposition with an “old” DIC advected in the CMZ-OMZ. Note that this approach provides a time-integrated information for the biogeochemical contribution to DIC concentration:  $\Delta$  are expressed in terms of concentration ( $\mu\text{mol/kg}$ ), and not in terms of rate.

In this study, no direct in situ measurement of any bacterial activity has been performed along the Chile OMZ allowing to identify microbial organisms and to prove the existence of biogeochemical processes. However, from theoretical stoichiometric equations for the known biogeochemical processes, although under-constrained and taking into account the uncertainties in the inverse method, it is possible to detect the “signature” of different dominant processes on both DIC and  $\text{O}_2$ . For example, aerobic remineralization produces DIC ( $\Delta\text{DIC}>0$ ) and consumes  $\text{O}_2$  ( $\Delta\text{O}_2<0$ ), and inversely for photosynthesis. Anaerobic remineralization (e.g., denitrification, or even DNRA) produces DIC ( $\Delta\text{DIC}>0$ ) without consuming  $\text{O}_2$  ( $\Delta\text{O}_2\approx 0$ ) by definition. Bacteria involved in the autotrophic processes such as nitrification and anammox should consume DIC (carbon source for assimilation;  $\Delta\text{DIC}<0$ ), but do ( $\Delta\text{O}_2<0$ ) and do not ( $\Delta\text{O}_2\approx 0$ ) consume  $\text{O}_2$  as aerobic and anaerobic process, respectively. All computed ( $\Delta\text{DIC}$ ,  $\Delta\text{O}_2$ ) for our sampling off Chile are reported in the carboncline-oxycline (Fig. 5b) where  $\text{DIC}_{\text{Chile}}^*$  appears the highest and most variable. Here, the  $\Delta\text{DIC}-\Delta\text{O}_2$  diagram (Fig. 5b) can provide qualitative information only, since each point corresponds to the predominance of a given process, and a large spatio-temporal variability is expected (e.g., between shelf and open ocean). Each situation corresponding theoretically to one dominant identified biogeochemical process (Fig. 5b) has been already observed off Chile and/or in other global OMZs. Indeed, for the same sites and/or periods off the Chilean OMZ, denitrification (Graco, 2002), anammox (Thamdrup et al., 2006; Galán

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2009) and nitrification (Molina et al., 2007) were observed to occur and tend to confirm the results of our simple biogeochemical approach. The reported ( $\Delta\text{DIC}$ ,  $\Delta\text{O}_2$ ) correspond to a net biogeochemical contribution, which is the net result of different superimposed processes. In particular, the situations ( $\Delta\text{DIC}>0$ ,  $\Delta\text{O}_2>0$ ), because of unknown biogeochemical process, could be interpreted as the superimposition of remineralization processes and photosynthesis. One needs to be very cautious concerning the indirect anammox detection, because anammox would correspond to two groups of ( $\Delta\text{DIC}$ ,  $\Delta\text{O}_2$ ) points:  $\Delta\text{DIC}\approx-70\text{ }\mu\text{mol/kg}$ , and  $\approx-300\text{ }\mu\text{mol/kg}$ . Note also that DIC and  $\text{O}_2$  can be affected not only from biogeochemical processes occurring in the water column, but also from sedimentary mechanisms (respiration), particularly active off the semi-enclosed Bay of Concepción as early diagenesis (e.g., Graco, 2002). Especially in non El Niño periods, bottom waters are influenced by a benthic organic remineralization (e.g., Graco et al., 2006) and then could produce DIC.

Net heterotrophy (remineralization) appears to be coexistent with net autotrophy (e.g., mainly nitrification but also anammox and photosynthesis: Fig. 5b), as suggested by chemolithoautotrophic evaluations for same sampling location than this study ( $36^\circ\text{S}$ ; Farias et al., 2009). However, net heterotrophy ( $\Delta\text{DIC}>0$ ) corresponds to most situations ( $>50\%$  of the points in Fig. 5b), compared to net autotrophy ( $\Delta\text{DIC}<0$ ). In aerobic configurations ( $\Delta\text{O}_2<0$ ), net heterotrophy would induce higher DIC production versus  $\text{O}_2$  consumption than expected from reference  $\text{C}/\text{O}_{\text{Re}}$  ratio, with mainly  $110>\Delta\text{DIC}>40\text{ }\mu\text{mol/kg}$ , whereas net autotrophy correspond mainly to  $-40<\Delta\text{DIC}<0\text{ }\mu\text{mol/kg}$ . This suggests a potential predominance of heterotrophy versus autotrophy for  $\Delta\text{DIC}$ , which would lead to higher C/O at the oxycline, and explaining the high  $\text{DIC}^*_{\text{ChileOxycline}}$  compared to the oxygenated ocean reported in Fig. 3c. This heterotrophy over autotrophy predominance would also be associated with predominance of net aerobic (the highest  $\Delta\text{DIC}$  correspond to the most negative  $\Delta\text{O}_2$ :  $<20\text{ }\mu\text{mol/kg}$ ) over net anaerobic (anammox and denitrification with low  $\Delta\text{DIC}$ :  $<40\text{ }\mu\text{mol/kg}$  on axis  $\Delta\text{O}_2=0$ ) activity (Fig. 5b). Referring to Paulmier et al. (2006), heterotrophy and autotrophy would correspond to active and passive OMZ phases, with

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



high ( $|\Delta O_2| > 20 \mu\text{mol/kg}$ ) and low ( $|\Delta O_2| < 20 \mu\text{mol/kg}$ )  $O_2$  consumption, respectively. The transition from active to passive phases suggests the importance of a potential coupling between mechanisms for C/O variations: heterotrophic-autotrophic (e.g., remineralization-photosynthesis); aerobic-anaerobic (e.g., aerobic remineralization-denitrification; nitrification-anammox, cf. Lam et al., 2007); but also photic-aphotic (photosynthesis-nitrification). Local high DIC production contributing to CMZ formation ( $\Delta\text{DIC}$  up to  $110 \mu\text{mol/kg}$ ) would be associated with high  $O_2$  consumption, stimulated by  $O_2$  availability in the oxycline rather than surface biomass availability (Paulmier et al., 2006). This DIC production would be encouraged by specific characteristics of the shallow oxycline (e.g., large range of  $O_2$  concentrations; well-lit; stratified) allowing a potential co-existence of different bacterial processes (Fig. 5b) which usually do not occur at the same depth. As for  $O_2$  consumption, intense DIC productions estimated at the oxycline agree with previous works in configurations of oxic-anoxic oscillation (e.g., Abril et al., 1999, 2010) and other experimental works (e.g., Sun et al., 2002). In addition to the net heterotrophy of the suboxic OMZ core, demonstrated by Koeve and Kähler (2010), the oxycline would highly contribute to the CMZ formation.

DIC production (heterotrophy) and consumption (autotrophy) at the oxycline can affect C/O molar ratios but also C/N ratios. Here,  $\Delta\text{DIC}/\Delta O_2$  corresponds to the drawdowns from a superimposition of several processes with different molar biomass ratio.  $\Delta\text{DIC}/\Delta O_2$  do not exactly correspond to strict C/O Redfield ratios (RR) determined in organic matter, but provide a first C/O approximation according to the Redfield definition linking molar ratios to biological processes (Redfield et al., 1963). In the oxycline, we have  $|\Delta\text{DIC}|_{\text{max}}/|\Delta O_2|_{\text{max}}$  varying from 0.6 to 1.5, and  $|\Delta\text{DIC}|_{\text{max}}/|\Delta\text{NO}_3^-|_{\text{max}}$  varying from  $\sim 0$  to 16, quite concomitantly (data not shown).  $|\Delta\text{DIC}|_{\text{max}}/|\Delta O_2|_{\text{max}}$  and  $|\Delta\text{DIC}|_{\text{max}}/|\Delta\text{NO}_3^-|_{\text{max}}$  differ from the classical canonical RR of  $106/138 \approx 0.8$  and  $106/16 \approx 6.6$ , respectively. In average for the whole dataset in the oxycline off Chile,  $|\Delta\text{DIC}|_{\text{max}}/|\Delta O_2|_{\text{max}}$  ( $1.1 \pm 0.3$ ) are twice as high as the RR, and  $|\Delta\text{DIC}|_{\text{max}}/|\Delta\text{NO}_3^-|_{\text{max}}$  ( $8.4 \pm 4.5$ ) is 20% higher than RR.

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In OMZs, anomalous low N/P ( $<10$  instead of 16) have been classically reported as for example off Peru (Codispoti and Christensen, 1985). Abnormally high C/O of  $\sim 1.1$  (Millero et al., 1998) and C/N of 7–8 (Graco, 2002) have also already been observed, but in iron fertilization experiment (IRONEX II, Indian Ocean) and anoxic sediments (on the shelf off Chile), respectively. The OMZ specific remineralization associated with the coexistence of different bacterial processes should be a key-element to understand why  $|\Delta\text{DIC}|_{\text{max}}/|\Delta\text{O}_2|_{\text{max}}$  ratio in the carboncline-oxycline is abnormally variable. High  $\Delta\text{DIC}$  and  $|\Delta\text{DIC}|_{\text{max}}/|\Delta\text{O}_2|_{\text{max}}$  could theoretically and locally result from higher rates of DIC production but also from more complete carbonated organic matter (OM) degradation and different OM stoichiometries (cf., Van Mooy et al., 2002; Paulmier et al., 2009). Abnormality of  $|\Delta\text{DIC}|_{\text{max}}/|\Delta\text{O}_2|_{\text{max}}$  molar ratios reported here in the OMZ off Chile suggests a perturbation of the ecosystem as compared to the oxygenated ocean, and could provide some information about molar ratios of the primitive anoxic ocean.

## 5 Conclusions

This study has showed that the OMZ off Chile is associated with very high DIC concentrations (mean of  $2330 \pm 60 \mu\text{mol/kg}$ ) at depths from  $\sim 175$ – $950$  m. These high DIC concentrations, corresponding to a local DIC maximum as compared to the oxygenated ocean, allow to define a carbon maximum zone (CMZ) over the whole OMZ thickness, including a carboncline associated with the oxycline. At the global scale, all OMZs form CMZs ( $\text{DIC} > 2225 \mu\text{mol/kg}$ ), with comparable vertical and horizontal structures and similar biogeochemical characteristics.

The existence of CMZs associated with all OMZs suggests that marine remineralization and/or accumulation of remineralized end-products are not uniformly distributed, but mainly occur in the OMZs. CMZ and OMZ are probably forming simultaneously with the same dynamical (low ventilation and upwelling) and

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



biogeochemical (remineralization producing DIC and consuming O<sub>2</sub>) mechanisms. However and paradoxically, CMZs exhibit a “carbon deficit” (DIC\*) in its core of ~10% of the remineralized DIC. This “carbon deficit” can be related to the physico-chemical (thermal effect) genesis of CMZ-OMZ core from upstream (e.g., equatorial) waters. Locally, the “carbon deficit” in the CMZ core appears to be compensated by a “carbon excess” at the carboncline-oxycline, induced by an intense aerobic-anaerobic remineralization. This specific remineralization is due to a potential net heterotrophic co-existence of different bacterial processes usually occurring at different depths. This co-existence contributes to a shallow and strong CO<sub>2</sub> production easy to be released toward the atmosphere, and may induce C/O ratio deviation from canonical Redfield ratios. However, if OMZs are associated with high remineralization activity, phytoplankton in surface could benefit from an “OMZ-regenerated” primary production after an upwelling episode, inducing a negative feedback effect on the oceanic CO<sub>2</sub> release. Indeed, OMZs highly recycle main nutrients such as phosphates and silicates, much less nitrate due to denitrification and anammox. Reduced conditions created in OMZs could be also favourable to more bio-availability for metals such as iron.

The existence of the CMZs-OMZs would imply anomalous molar ratios, which remain to be confirmed for all main OMZs. High DIC, associated with relatively low alkalinity (~2325 µeq/kg) and predominance of aerobic remineralization and nitrification, would largely acidify the OMZ core waters. This acidification would induce perturbations of marine biogeochemistry and ecosystems in addition of the respiratory barrier due to the O<sub>2</sub> deficit. CMZs-OMZs can be considered as “natural laboratories” of a low O<sub>2</sub> and already “high CO<sub>2</sub>” ocean. They are ideally suited for studies looking back in the past and aiming to predict the future, as planned in international programs, such as SOLAS (Surface Ocean Lower Atmosphere Study) mid-term strategies.

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



*Acknowledgements.* This study was supported by a French CNRS Ph.D. Fellowship to A. Paulmier and financial support provided by the ECOS Sur program (Foreign Office Ministry), by EU contract N° SIP3-CT-2003-502885 to LEGOS/CNRS (IP-MERSEA project), and IRD. We thank to O. Ulloa and L. Farias for making possible the DIC sampling during the cruises, and to the R/V *Vidal Gormaz*, *Carlos Porter* and *Kay-Kay* crews. We also thank C. Duarte, H. J. Minas and M. Graco for previous fruitful discussions.



The publication of this article is financed by CNRS-INSU.

## References

- Antezana, T.: *Euphausia mucronata*: a keystone herbivore and prey of the Humboldt Current System, *Deep-Sea Res. Pt. II*, 57(7–8), 652–662, doi:10.1016/j.dsr2.2009.10.014, 2009.
- Anderson, L. and Sarmiento, J.: Redfield ratios of remineralization determined by nutrient data analysis, *Global Biogeochem. Cy.*, 8, 65–80, 1994.
- Arrigo, K. R.: Marine microorganisms and global nutrient cycles, *Nature*, 437, 349–355, 2005.
- Abril, G., Etcheber, H., Le Hir, P., Bassoullet, P., Boutier, B., and Frankignoulle, M.: Oxic/anoxic oscillations and organic carbon mineralization in an estuarine maximum turbidity zone (The Gironde, France), *Limnol. Oceanogr.*, 44(5), 1304–1315, 1999.
- Abril, G., Commarieu, M.-V., Etcheber, H., Deborde, J., Deflandre, B., Zivacinovic, M. K., Chaillou, G., and Anschutz, P.: In vitro simulation of oxic/suboxic diagenesis in an estuarine fluid mud subjected to redox oscillations, *Estuar. Coast. Shelf S.*, 88(2), 279–291, doi:10.1016/j.ecss.2010.04.003, 2010.
- Bange, H. W.: Gaseous Nitrogen Compounds (NO, N<sub>2</sub>O, N<sub>2</sub>, NH<sub>3</sub>) in the ocean, in: *Nitrogen in Marine Environment*, edited by: Capone, D. G., Bronk, D. A., Mulholland, M. R., and Carpenter, E. J., Chapt. 2, Elsevier, 51–94, 2008.

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Boyer, T. P., Antonov, J. I., Garcia, H. E., Johnson, D. R., Locarnini, R. A., Mishonov, A. V., Pitcher, M. T., Baranova, O. K., and Smolyar, I. V.: World Ocean Database 2005, edited by: Levitus, S., NOAA Atlas NESDIS 60, Washington, DC, 190 pp., DVDs, 2006.

Bradshaw, A. L. and Brewer, P. G.: High precision measurements of alkalinity and total carbon dioxide in seawater by potentiometric titration, *Mar. Chem.*, 23, 69–86, 1988.

Broecker, W. S.: “NO”, a conservative water-mass tracer, *Earth Planet. Sci. Lett.*, 23, 100–107, 1974.

Broecker, W. S. and Peng, T. H.: *Tracers in the Sea*, Eldigio Press, 1982.

Broenckow, W. W. and Cline, J. D.: Colorimetric determination of dissolved oxygen at low concentrations, *Limnol. Oceanogr.*, 14, 450–454, 1969.

Cannariato, K. and Kennett, J.: Climatically related millennial-scale fluctuations in strength of California margin oxygen-minimum zone during the past 60 k.y., *Geology*, 27, 975–978, 1999.

CARS: CSIRO Atlas of Regional Seas, a <http://www.marine.csiro.au/~dunn/cars2009/#refs>, 2010.

Chan, F., Barth, J., Lubchenko, J., Kirincich, A., Weels, H., Perterson, W., and Menge, A.: Emergence of anoxia in the California current large marine ecosystem, *Science*, 319, p. 920, 2008.

Chen, C. T. A., Ruo, R., Pai, S. C., Liu, C. T., and Wong, G. T. F.: Exchange of water masses between the East China Sea and the Kuroshio off Northeastern Taiwan, *Cont. Shelf Res.*, 15, 19–39, 1995.

Childress, J. J. and Siebel, B. A.: Life at stable low oxygen levels: adaptations of animals to oceanic oxygen minimum layers, *J. Exp. Biol.*, 201, 1223–1232, 1998.

Codispoti, L. A. and Christensen, J. P.: Nitrification, denitrification, and nitrous oxide cycling in the Eastern Tropical Southern Ocean, *Mar. Chem.*, 16, 277–300, 1985.

Corbière, A., Metzl, N., Reverdin, C., and Takahashi, T.: Interannual and decadal variability of the oceanic carbon sink in the North Atlantic subpolar gyre, *Tellus B*, 59(2), 168–178, 2007.

De Boyer Montégut C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: an examination of profile data and a profile-based climatology, *J. Geophys. Res.*, 109, 1–20, 2004.

Deutsch, C., Sarmiento, J. L., Sigman, D. L., Grüber, N., and Dunne, J. P.: Spatial coupling of nitrogen inputs and losses in the ocean, *Nature*, 445, 163–167, 2007.

Endoh, M.: Double-celled circulation in coastal upwelling, *J. Oceanogr.*, 33, 30–37, 1976.

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





- Falkowski, P. G.: Evolution of the nitrogen cycle and its influence on the biological sequestration of CO<sub>2</sub> in the ocean, *Nature*, 387, 272–275, 1997.
- Farías, L., Fernández, C., Faúndez, Cornejo, M., and Alcaman, M. E.: Chemolithoautotrophic production mediating the cycling of the greenhouse gases N<sub>2</sub>O and CH<sub>4</sub> in an upwelling ecosystem, *Biogeosciences*, 6, 3053–3069, doi:10.5194/bg-6-3053-2009, 2009.
- Fuhrman, J. A. and Capone, D. G.: Possible biogeochemical consequences of ocean fertilization, *Limnol. Oceanogr.*, 36, 1951–1959, 1991.
- Galán, A., Molina, V., Thamdrup, B., Woebken, D., Lavik, G., Kuypers, M. M. M.: Anammox bacteria and the anaerobic oxidation of ammonium in the oxygen minimum zone off northern Chile, *Deep-Sea Res. Pt. II*, 56(16), 1021–1031, doi:10.1016/j.dsr2.2008.09.016, 2009.
- Graco, M.: Variación temporal de la producción de amonio en un área de surgencias de Chile Central (36° S), Factores y procesos involucrados en su reciclaje, PhD Thesis, UPMC Paris VI-France, Universidad de Concepción-Chile, 2002.
- Graco, M., Gutiérrez, D., and Farías, L.: Interannual variability of the pelagic-benthic coupling in the upwelling system off central Chile, *Adv. Geosci.*, 6, 127–132, 2006, <http://www.adv-geosci.net/6/127/2006/>.
- Gribaldo, S. and Philippe, H.: Ancient Phylogenetic Relationships, *Theor. Popul. Biol.*, 61, 391–408, 2002.
- Gruber, N. and Sarmiento, J. L.: Global patterns of marine nitrogen fixation and denitrification, *Global Biogeochem. Cy.*, 11(2), 235–266, 1997.
- Ivanoff, A.: Introduction à l'océanographie, Propriétés physiques et chimiques des eaux de mer, Vuibert, Paris, 2, 208 pp., 1972.
- Jain, A. K., Briegleb, B. P., Minschwaner, K., and Wuebbles, D. J.: Radiative forcing and global warming potentials of 39 greenhouse gases, *J. Geophys. Res.*, 105(D16), 20773–20790, 2000.
- Jansen, E., Overpeck, J., Briffa, K. R., Duplessy, J.-C., Joos, F., Masson-Delmotte, V., Olago, D., Otto-Bliesner, B., Peltier, W. R., Rahmstorf, S., Ramesh, R., Raynaud, D., Rind, D., Solomina, O., Villalba, R., and Zhang, D.: Palaeoclimate, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK and New York, 2007.
- Karstensen, J., Stramma, L., and Visbeck, M.: Oxygen minimum zones in the Eastern Tropical

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Atlantic and Pacific Oceans, *Prog. Oceanogr.*, 77, 331–350, 2008.
- Koeve, W. and Kähler, P.: Heterotrophic denitrification vs. autotrophic anammox – quantifying collateral effects on the oceanic carbon cycle, *Biogeosciences*, 7, 2327–2337, doi:10.5194/bg-7-2327-2010, 2010.
- 5 Lam, P., Jensen, M. M., Lavik, G., McGinnis, D. F., Müller, B., Schubert, C. J., Amann, R., Thamdrup, B., and Kuypers, M. M. M.: Linking crenarchaeal and bacterial nitrification to anammox in the Black Sea, *P. Natl. Acad. Sci. USA*, 104(17), 7104–7109, 2007.
- Lewis, E. and Wallace, D. W. R.: Program developed for CO<sub>2</sub> system calculations, ORNL/CDIAC-105, Carbon Dioxide Information Analysis Center, Oak Ridge National
- 10 Laboratory, US Department of Energy, 1998
- Luyten, J. L., Pedlosky, J., and Stommel, H.: The ventilated thermocline, *J. Phys. Oceanogr.*, 13, 292–309, 1983.
- Maske, H., Medrano, R. C., Castro, A. T., Mercado, A. J., Jauregui, C. O. A., Castro, G. G., and Ochoa, J.: Inorganic carbon and biological oceanography above a shallow oxygen minimum
- 15 in the entrance to the Gulf of California in the Mexican Pacific, *Limnol. Oceanogr.*, 55(2), 481–491, 2010.
- Millero, F. J., Degler, A. A., O'Sullivan, D. W., Goyet, C., and Eiseheid., G.: The carbon dioxide system in the Arabian Sea, *Deep-Sea Res.*, 45, 2225–2252, 1998.
- Minas H. J., Minas, M., and Packard, T. T.: Productivity in upwelling areas deduced from hydrographic and chemical fields, *Limnol. Oceanogr.*, 31(6), 1182–1206, 1986.
- 20 Molina, V., Ulloa, O., Farias, L., Urrutia, H., Ramirez, S., Junier, P., and Witzel, K. P.: Ammonia-oxidizing  $\beta$ -proteobacteria from the oxygen minimum zone off Northern Chile, *Appl. Environ. Microb.*, 73, 3547–3555, 2007.
- Morrison, J. M., Codispoti, L. A., Smith, S. L., Wishner, K., Flagg, C., Gardner, W. D., Gaurin, S., Naqvi, S. W. A., Manghnani, V., Prosperie, L., and Gundersen, J. S.: The oxygen minimum
- 25 zone in the Arabian Sea during 1995, *Deep-Sea Res.*, 46, 1903–1931, 1999.
- Najjar, R. G. and Keeling, R. F.: Mean annual cycle of the air-sea oxygen flux: a global view, *Global Biogeochem. Cy.*, 14(2), 573–584, 2000.
- Naqvi, S. W. A., Naik, H., Pratihary, A., D'Souza, W., Narvekar, P. V., Jayakumar, D. A., Devol, A. H., Yoshinari, T., and Saino, T.: Coastal versus open-ocean denitrification in the Arabian
- 30 Sea, *Biogeosciences*, 3, 621–633, doi:10.5194/bg-3-621-2006, 2006.
- O'Conner, B. M., Fine, R. A., Maillet, K. A., and Olson, D. B.: Formation rates of subtropical underwater in the Pacific Ocean, *Deep-Sea Res. Pt. I*, 49, 1571–1590, 2002.

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Paulmier, A.: Zones de Minimum d'Oxygène (OMZs) de l'océan moderne, Ph.D. thesis, Pierre et Marie Curie University, Paris, France, 248 pp., 2005 (in Fench).
- Paulmier, A., Ruiz-Pino, D., Garçon, V., and Farías, L.: Maintaining of the East South Pacific Oxygen Minimum Zone (OMZ) off Chile, *Geophys. Res. Lett.*, 33, L20601, doi:10.1029/2006GL026801, 2006.
- Paulmier, A. and Ruiz-Pino, D.: Oxygen Minimum Zones (OMZs) in the modern ocean, *Prog. Oceanogr.*, 80(3–4), 113–128, doi:10.1016/j.pocean.2008.05.001, 2008.
- Paulmier, A., Ruiz-Pino, D., and Garçon, V.: The Oxygen Minimum Zone (OMZ) off Chile as intense source of CO<sub>2</sub> and N<sub>2</sub>O, *Cont. Shelf Res.*, 28(20), 2746–2756, doi:10.1016/j.csr.2008.09.012, 2008.
- Paulmier, A., Kriest, I., and Oschlies, A.: Stoichiometries of remineralisation and denitrification in global biogeochemical ocean models, *Biogeosciences*, 6, 923–935, doi:10.5194/bg-6-923-2009, 2009.
- Rabalais, N. N., Díaz, R. J., Levin, L. A., Turner, R. E., Gilbert, D., and Zhang, J.: Dynamics and distribution of natural and human-caused hypoxia, *Biogeosciences*, 7, 585–619, doi:10.5194/bg-7-585-2010, 2010.
- Ramaswamy, V., Boucher, O., Haigh, J., Hauglustaine, D., Haywood, J., Myhre, G., Nakajima, T., Shi, G. Y., and Solomon, S.: Radiative forcing of climate change, in: “Climate CHANGE 2001: The scientific basis. Contribution of working group I to the third assessment report of the intergovernmental panel on climate change”, edited by: Houghton, J. T., Ding, Y., Griggs, D. J., Noguer, M., Van der Linden, P. J., Dai, X., Maskell, K., and Johnson, C. A., Cambridge University Press, Cambridge, UK, 349–416, 2001.
- Rao, D. P., Sarma, V. V., Subba Rao, V. Sudhakar, U., and Gupta, G. V. M.: On watermass mixing ratios and regenerated silicon in the Bay of Bengal, *Indian J. Mar. Sci.*, 25, 56–61, 1996.
- Redfield, A. C., Ketchum, B. H., Richards, P. A.: The influence of organisms on the composition of seawater, in: *The Sea*, Vol. 2, edited by: Hill, M. N., Interscience Publishers, NY, 26–77, 1963.
- Rogers, A. D.: The role of the oceanic oxygen minima in generating biodiversity in the deep-sea, *Deep-Sea Res.*, 47(1–2), 119–148, 2000.
- Royer, D. L., Bemmer, R. A., Montañez, I. P., Tabor, N. J., and Beerling, D. J.: CO<sub>2</sub> as a primary driver of Phanerozoic climate, *GSA Today*, 14(3), 4–10, doi:10.1130/1052-5173, 2004.

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T. H., Kozyr, A., Ono, T., and Rios, A. F.: The oceanic sink for anthropogenic CO<sub>2</sub>, *Science*, 305, 367–371, 2004.
- Sabine, C. L., Key, R. M., Kozyr, A., Feely, R. A., Wanninkhof, R., Millero, F. J., Peng, T. H., Bullister, J. L., and Lee, K.: Global Ocean Data Analysis Project: Results and Data. ORNL/CDIAC-145, NDP-083, Carbon Dioxide Information Analysis Centre, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, unpublished, 110 pp., 2005.
- Sarma, Y. V. B., Rama Rao, E. P., Saji, P. K., and Sarma, V. V. S. S.: Hydrography and circulation of the bay of Bengal during withdrawal phase of the Southwest Monsoon, *Oceanol. Acta*, 22, 453–473, 1998.
- Stramma, L., Johnson, G. C., Sprintall, J., and Mohrholz, V.: Expanding oxygen-minimum zones in the tropical oceans, *Science*, 320, 655–658, 2008.
- Strub, P., Mesías, J., Montecino, V., Rutllant, J., and Salinas, S.: Coastal ocean circulation off Western South America, in: *The Sea*, Vol. 11, Regional Studies and Synthesis, edited by: Robinson, A. R. and Brink, K. H., John Wiley, Hoboken, NJ, 273–313, 1998.
- Sun, M. Y., Aller, R. C., Lee, C., and Wakeham, S. G.: Effects of oxygen and redox oscillation on degradation of cell-associated lipids in surficial marine sediments, *Geochim. Cosmochim. Ac.*, 66, 2003–2012, 2002.
- Takahashi, T. and Goddard, J.: Measurements of total CO<sub>2</sub> Concentration and total alkalinity in seawater collected during the Sonne Cruise 102, Final Tech. Report, 15 March 1998.
- Thamdrup, B., Dalsgaard, T., Jensen, M. M., Ulloa, O., Farías, L., and Escobano, R.: Anaerobic ammonium oxidation in the oxygen deficient waters off Northern Chile, *Limnol. Oceanogr.*, 55, 2145–2156, 2006.
- Tsuchiya, M.: The origin of the Pacific equatorial 13 °C water, *J. Phys. Oceanogr.*, 11, 794–812, 1981.
- Van Mooy, B., Keil, R., and Devol, A.: Impact of suboxia on sinking particulate organic carbon: Enhanced carbon flux and preferential degradation of amino acids via denitrification, *Geochim. Cosmochim. Ac.*, 66, 457–465, 2002.
- Viecelli, J. A.: The atmospheric carbon dioxide response to oceanic primary productivity fluctuations, *Climatic Change*, 6, 203–215, 1984.
- Walsh, D. A., Zaikova, E., Howes, C. G., Song, Y. C., Wright, J. J., Tringe, S. G., Tortell, P. D., and Hallam, S. J.: Metagenome of a versatile chemolithoautotroph from expanding oceanic

**BGD**

7, 6353–6385, 2010

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- dead zones, *Science*, 326, 578–582, 2009.
- Weiss, R. F.: The solubility of nitrogen, oxygen and argon in water and seawater, *Deep Sea Res.*, 17, 721–735, 1970.
- 5 Weiss, R. F.: Carbon dioxide in water and seawater: The solubility of a nonideal gas, *Mar. Chem.*, 2, 203–215, 1974.
- Wishner, K. F., Gelfman, C., Gowing, M. M., Outram, D. M., Rapien, M., Williams, R. L.: Vertical zonation and distributions of calanoid copepods through the lower oxycline of the Arabian Sea oxygen minimum zone, *Prog. Oceanogr.*, 78, 163–191, 2008.
- 10 World Ocean Circulation Experiment: WOCE Operation Manual, Vol. 3, The Observational Programme: Part 3.1.3 – WHP Operations and Methods, 68/91, Geneva, 1994.
- Wyrski, K.: The oxygen minima in relation to ocean circulation, *Deep-Sea Res.*, 9, 11–23, 1962.

## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.

Title Page

## Abstract

## Introduction

## Conclusions

## References

## Tables

## Figures



▶

1

▶

[Back](#)

Close

Full Screen / Esc


[Printer-friendly Version](#)

## Interactive Discussion



**Table 1.** Averaged DIC and O<sub>2</sub> concentrations in a mean layer intercepting the core of all main OMZs (175–950 m), and sensitivity tests (in **bold**; commented in Sect. 3.2) on upper (for 137.5 and 225 m) and lower limits (for 850 and 1050 m) from GLODAP and WOA2005 databases. Averages are given for tropical OMZs together (ETSP+ETNP+AS+BB) and separately, and for the global ocean. ETNP: Eastern Tropical North Pacific; ETSP: Eastern Tropical South Pacific; AS: Arabian Sea; BB: Bay of Bengal.

Averaged layer Parameters ( $\mu\text{mol/kg}$ )	175–950 m		137.5–950 m		225–950 m		175–850 m		175–1050 m		Maximal differences in the 175–950 m layer	
	DIC	O <sub>2</sub>	DIC	O <sub>2</sub>	DIC	O <sub>2</sub>	DIC	O <sub>2</sub>	DIC	O <sub>2</sub>	DIC	O <sub>2</sub>
Total OMZs	2284	22	<b>2279</b>	<b>24</b>	<b>2289</b>	<b>20</b>	<b>2277</b>	<b>21</b>	<b>2289</b>	<b>23</b>	<b>±7</b>	<b>±2</b>
ETSP	2285	22	<b>2283</b>	<b>22</b>	<b>2287</b>	<b>22</b>	<b>2282</b>	<b>19</b>	<b>2288</b>	<b>26</b>	<b>±3</b>	<b>±4</b>
ETNP	2286	24	<b>2280</b>	<b>26</b>	<b>2292</b>	<b>22</b>	<b>2278</b>	<b>24</b>	<b>2292</b>	<b>25</b>	<b>±8</b>	<b>±2</b>
AS	2276	12	<b>2273</b>	<b>13</b>	<b>2279</b>	<b>12</b>	<b>2272</b>	<b>12</b>	<b>2280</b>	<b>13</b>	<b>±4</b>	<b>±1</b>
BB	2278	20	<b>2274</b>	<b>20</b>	<b>2281</b>	<b>20</b>	<b>2274</b>	<b>18</b>	<b>2282</b>	<b>22</b>	<b>±4</b>	<b>±2</b>
Global ocean	2203	157	<b>2199</b>	<b>159</b>	<b>2208</b>	<b>155</b>	<b>2196</b>	<b>159</b>	<b>2209</b>	<b>155</b>	<b>±7</b>	<b>±2</b>
Averages on the whole water column for:												
	DIC: 2255						O <sub>2</sub> : 176					




**Fig. 1.** Vertical sections along an east-west cross-shore transect on 0–100 m (DIC: (a);  $O_2$ : (d)) and versus time on 0–50 m (DIC: (b);  $O_2$ : (e)). Points indicate sampling locations. Profiles of DIC (c) and  $O_2$  (f) on 0–1000 m for all stations sampled in the OMZ off Chile in 2000–2002. Continuous thin profiles correspond to classical profiles out of the OMZ in the Eastern Equatorial Pacific (for DIC on (c): Viecelli, 1984; for  $O_2$  on (f): Ivanoff, 1972). Continuous thick black profiles indicate representative profiles for DIC on (c) and for  $O_2$  on (f). On (c) and (f), data from Iquique at 21° S (squares), Coquimbo at 30° S (diamonds) and Concepción at 36° S (triangles). On (c): the hatched red area indicates  $DIC_{CMZ}$  ( $DIC_{CMZ} = DIC_{Average \text{ in OMZ}} - DIC_{Classical \text{ out of OMZ}}$ ) corresponding to the carbon maximum zone (CMZ); continuous thick red profile indicates the representative profile for the sum of  $DIC_{Classical \text{ out of OMZ}}$  and local DIC production contributing to the CMZ ( $DIC_{Classical \text{ out of OMZ}} + \Delta DIC$ ), with  $\Delta DIC$  the biogeochemical contribution to DIC. The points on the map in (f) indicate the three sampled latitudes (21° S; 30° S; 36° S).



# CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.



**Fig. 2.** Vertical profiles of DIC (**a**) and O<sub>2</sub> (**c**) on 0–5500 m for the main most intense OMZs in the open ocean from GLODAP and WOA2005 databases. The continuous profile in (a) and (c) corresponds to a mean global profile. Vertical dashed line in (c) corresponds to O<sub>2</sub>=20 μmol/kg (upper O<sub>2</sub> concentrations of the OMZs core), and horizontal continuous lines in (a) and (c) correspond to the mean upper (175 m) and lower (950 m) depths of the OMZs core. Mean DIC (**b**) and O<sub>2</sub> (**d**) distribution average in μmol/kg on the 175–950 m, a layer intercepting the core of all main tropical OMZs (cf. Table 1). For ETNP, ETSP, AS and BB, same abbreviations than in Table 1 caption.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



# CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.



**Fig. 3.** Vertical profiles of DIC\* (=DIC+0.69O<sub>2</sub>; cf. Methodology and the comments provided for Eq. 1) for the main OMZs from GLODAP and WOA2005 databases on 0–5500 m **(a)**, with a zoom in the layer intercepting the core of all considered OMZs **(b)**, and from all stations sampled in the OMZ off Chile in 2000–2002 **(c)**. Continuous profile in (a) and (c) corresponds to a mean profile for the global ocean and for the oxygenated Pacific out of the OMZ, respectively. Rectangles in (a) and (b) indicate the OMZ core. For ETNP, ETSP, AS and BB, same abbreviations than in Table 1 caption.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



CO<sub>2</sub> maximum in the OMZ

A. Paulmier et al.



**Fig. 4.** DIC (a), O<sub>2</sub> (b) and DIC\* (c) mean concentrations for global ocean, equatorial regions (Indian and Pacific only, where the main intense OMZs are located; between 10° N and 10° S; between 175 and 950 m, in the same depth layer than the subsurface OMZ core), and CMZs-OMZs. In dashed arrows, the remineralization effect from the global ocean towards the Equatorial regions and the CMZs-OMZs. In thick grey arrows, the temperature effect decreasing mainly the solubility, not visible on DIC (a) because of the buffer effect, but on O<sub>2</sub> (b) and DIC\* (c) only.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 5.** (a) DIC- $\text{O}_2$  diagram for the global ocean (all WOCE data with both DIC and  $\text{O}_2$  available). (b)  $\Delta\text{DIC}-\Delta\text{O}_2$  diagram (from all stations sampled in the OMZ off Chile in 2000–2002, mainly focused in the oxycline). For (a), the grey line corresponds to classical linear regression, with  $N=85\,650$  data and a regression coefficient  $R^2$  of 0.25:  $\text{DIC} = -0.7\text{O}_2 + 2330$ . For (b),  $\Delta\text{DIC}$  and  $\Delta\text{O}_2$  correspond to the biogeochemical contribution to DIC and  $\text{O}_2$ , respectively, using an inverse method (cf. Methodology). Each symbol corresponds to one dominating biogeochemical process, which has been suggested and identified indirectly from theoretical chemical reactions: aerobic remineralization (small squares); nitrification (triangles); anaerobic remineralization (diamonds, mainly near  $\Delta\text{DIC}-\Delta\text{O}_2 \approx (0,0)$ ); photosynthesis (large squares); anammox (circles). Crosses correspond to non determined process. Accuracy and reproducibility on DIC and  $\text{O}_2$  induce an uncertainty on the  $\Delta$  of  $\sim 4 \mu\text{mol/kg}$  (cf., Paulmier et al., 2006).  $(\text{C/O})_{\text{Re}}$  corresponds to the reference molar ratio for this study of 117/170 (Anderson and Sarmiento, 1994).