

**Quality or
decomposer
efficiency**

J. Å. M. Wetterstedt and
G. I. Ågren

Quality or decomposer efficiency – which is most important in the temperature response of litter decomposition? A modelling study using the GLUE methodology

J. Å. M. Wetterstedt and G. I. Ågren

Department of Ecology, Swedish University of Agricultural Sciences, P.O. Box 7044,
750 07 Uppsala, Sweden

Received: 3 November 2010 – Accepted: 5 November 2010 – Published: 30 November 2010

Correspondence to: G. I. Ågren (goran.agren@ekol.slu.se)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Effects of temperature history on litter decomposition were evaluated using the GLUE modelling framework together with the Q-model calibrated to a needle litter incubation experiment. The needle litter incubation was a full factorial design with initial and final temperatures 5, 15 and 25 °C. Samples were moved from the initial to the final temperature when approximately 12% of the initial carbon had been respired. We used four variations of the Q-model: one or two initial litter quality values and fixed or temperature-dependent decomposer efficiency. The model was calibrated to the constant temperature data subset. Evaluation against temperature shift subsets gave good results, except just after the change in temperature where the model predicted a smaller response than observed. Using one or two initial litter quality values and fixed decomposer efficiency had little effect on final litter quality and respiration at the final incubation temperature. Allowing decomposer efficiency to vary with temperature showed that decomposer efficiency should decrease between 5 to 15 °C but with no change between 15 and 25 °C. A flexible decomposer efficiency resulted also in substantial differences in litter quality at the end of the initial incubation in response to incubation temperature. The results suggests that it is important to consider other factors than the variation in temperature sensitivity with quality when evaluating effects of temperature changes on soil organic matter stability.

1 Introduction

In view of the expected future climatic change (Solomon, 2007) the temperature dependence of decomposition of litters and soil organic matter (SOM) has since long attracted much interest because a strong positive temperature dependence would create a strong positive climatic feedback. However, both in laboratory incubations and field studies, temperature history and not only current temperature have been shown to affect respiration rates, such that SOM with different temperature histories will have

BGD

7, 8699–8722, 2010

Quality or decomposer efficiency

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

different decomposition rates at the same temperature. These effects can be short term or long term, and be a result of factors like substrate depletion, changes in decomposer community composition and abundance, and changes in quality composition (Kirschbaum, 2006). Quality changes have been challenged as a major factor because the temperature dependence of the rate of utilisation might not respond strongly enough to explain observations as there are several processes involved with sometimes counteracting temperature responses (Davidson and Janssens, 2006; Ågren and Wetterstedt, 2007). It is thus necessary to also consider the temperature dependence of the other factors regulating decomposition. Decomposer community composition and decomposer biomass are two important factors that may be affected by temperature. For example, it has been demonstrated (Devêvre and Horwath, 2000; Steinweg et al., 2008) that the carbon use efficiency of decomposers decreases with temperature, probably as a result of higher maintenance costs. Allison et al. (2010) suggest that the decomposer biomass should go down with increasing maintenance cost and as a result less extracellular enzymes are produced and decomposition slows down.

Most models, e.g. Century (Parton et al., 1987), G'Day (Comins and McMurtrie, 1993), RothC (Coleman and Jenkinson, 1995), Q (Ågren and Bosatta, 1998) dealing with soil organic carbon (SOC) conform to the same generic structure. SOC is described as consisting of carbon of different pools or qualities with three main processes driving the changes in SOC quantity and quality: (i) a decomposer community feeding on SOC at some rate (*growth rate*); (ii) when doing so, part of the carbon they use is respired as carbon dioxide and part remains as SOC; we call the fraction remaining efficiency (*decomposer efficiency*); (iii) the fraction remaining undergoes changes in quality. We call this transfer between qualities (pools) *dispersion*, on average SOC increases in recalcitrance/decreases in quality with time.

To address the question of how the factors quality and decomposer efficiency affect the temperature response we have used the Q-model (Bosatta and Ågren, 2003; Ågren and Bosatta, 1998) to see how well we can predict respiration of needle litter from an incubation experiment, and analyse the importance of those factors (quality and

BGD

7, 8699–8722, 2010

Quality or decomposer efficiency

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



decomposer efficiency). The Q-model was chosen because the fate of carbon and the decomposition processes are relatively easy to follow in it. Data from a recently published temperature variation experiment (Wetterstedt et al., 2010) is used, both for calibration and evaluation. We will use the model to explore the consequences of having one or two initial litter qualities in combination with fixed or flexible (with regard to temperature) decomposer efficiency.

We have chosen to use the GLUE (Generalised Likelihood Uncertainty Estimation, Beven, 2006) framework for model calibration and evaluation. GLUE can be used as a modelling protocol and is well suited to give uncertainty estimations in model output. It also provides criteria for complete model rejection, i.e. the model structure needs to be changed if the model fails to predict empirical data well enough.

2 Materials and methods

2.1 The Q-model

The Q-model describes litter or SOC as consisting of a continuous spectrum of carbon qualities or decomposabilities instead of being partitioned into a small number of discrete pools. The Q-model has certain advantages over discrete models. Firstly, there are analytical solutions, making it easier to understand and explain model behaviour. Approximate solutions, which are similar in their behaviour, are also available (Bosatta and Ågren, 2003). The approximate solutions are much less computationally demanding and are therefore preferred when doing large model runs, for example during calibration. They substitute the complete distribution of litter qualities in the exact solution with one average quality. Secondly, there are also fewer parameters in a continuous formulation, as opposed to pool models, and the model formulation enforces consistency between them (Bruun et al., 2010). Parameters estimated with the approximate solutions can also be used in the exact solution, possibly with some slight recalibration.

BGD

7, 8699–8722, 2010

Quality or decomposer efficiency

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Quality or decomposer efficiency

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In the Q-model, the growth rate of the decomposers depends on carbon quality (q) and temperature (T) as well as the base rate parameter u_0 , which describes growth rate. The temperature response of the growth rate is an Arrhenius function with activation energy ΔG^0 . Decomposer efficiency (e) is set to be either temperature independent (fixed) or allowed to vary with temperature (flexible). Transfer of carbon to lower qualities, dispersion, is expressed through the parameter η_{12} , which is assumed to be temperature independent. Table 1 lists the parameters used in the model. When running the model with two initial litter qualities, the two q_0 's will be selected from behavioural models (see below) where the qualities are somewhat separated. The reason we are using two initial q_0 's in this experiment is to explore the effect of how the different temperature sensitivity of different q_0 's translates into a differential quality evolution.

Four versions of the model were run with combinations of *one* or *two* initial qualities combined with *fixed* or *flexible* decomposer efficiencies. In the case of two initial litter qualities, the initial amount of substrate was partitioned equally between the qualities.

2.2 Observational data

We have chosen to use the spruce (*Picea abies*) needle litter data from the temperature experiment by Wetterstedt et al. (2010) (see Figs. 1 and 2). The data consists of time series (four replicates) of litter respiration rates at different temperatures. In some time series the samples have been shifted from one temperature to another when approximately 12% of the initial carbon had been respired. We will denote temperature combinations as initial temperature + final temperature, e.g. 5 + 15 °C, meaning that the sample was first exposed to 5 and then 15 °C. The data used for calibration were from needles stored at three temperatures without shifts in temperature (5 + 5 °C, 19 data points; 15 + 15 °C, 14 data points; and 25 + 25 °C, 16 data points). To reduce the variability in data between measurement points, we used a running mean of three consecutive points to smooth the curve (the first and last points were averaged from two points). We also normalised the variance at each measurement point by averaging over the whole measurement period for each temperature, i.e. the variances used when

calibrating the model are

$$\bar{O}_i \sum_i \frac{\text{Var}(\bar{O}_i)}{\bar{O}_i} / n,$$

where O_i is the observation at point i and n is the number of points in the measurement series. These steps were taken to obtain a more robust calibration process. However, when the calibrated models are evaluated using the least square method, R^2 are calculated with respect to non-smoothed data (Figs. 1 and 2).

2.3 GLUE

The GLUE methodology introduced by Beven och Binley (1992) is a framework for calibrating and using models in predictions. It includes criteria and methods for model rejection and sensitivity analysis of model parameters. A “model” in GLUE terminology is the combination of the “model structure”, e.g. the Q-model as opposed to some other model, and the parameter values used to run the model. A “behavioural model” is a model that can simulate real data “good enough”. It follows that a non-behavioural model should not be used to forecast data; instead, it would need better parameterisation or a change in model structure. In this text we will however use the term “model” with the meaning “model structure”. The use of GLUE includes the following steps (Beven, 2009).

2.3.1 Likelihood measure

Decide on an informal (or formal) likelihood measure or measures (LM) for use in evaluating each model run, including the rejection criteria, which for a non-behavioural model run will be given a likelihood of zero. Ideally this should be done before running the model, taking into account possible input and observational errors: since calibration data contain means as well as standard deviations, we used a triangular shaped

BGD

7, 8699–8722, 2010

**Quality or
decomposer
efficiency**

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



likelihood measure:

$$I(M, O_{\min}, O_{\text{mean}}, O_{\max}) = \begin{cases} \frac{M - O_{\min}}{O_{\text{mean}} - O_{\min}} & M \leq O_{\text{mean}} \\ \frac{O_{\max} - M}{O_{\max} - O_{\text{mean}}} & M > O_{\text{mean}} \end{cases}$$

If model output equals the average measured value (O_{mean}) the function returns 1, at ± 1.96 standard deviations (O_{\max}, O_{\min}) zero, and negative values when model output deviates more from the observed mean value. These values were summed for each temperature and divided by the number of observed values, $n_{O,T}$, and then averaged over the three temperatures

$$L(M(\Theta), O) = \sum_{T=5,15,25^{\circ}\text{C}} \left(\sum_{O_T} \frac{I(M_{O,T}, O_{\min,T}, O_{\text{mean},T}, O_{\max,T})}{n_{O,T}} \right) / 3.$$

Θ stands for the parameter set used. This likelihood measure takes advantage of the variability in the observed data and is less influenced than the least square method by outliers. Ideally, the model together with its parameters, should predict all observed data points within their 95% error bounds; i.e. for all $L(M, O_{\min}, O_{\text{mean}}, O_{\max})$ observations.

2.3.2 Model parameters

Decide which model parameters and input variables are to be considered uncertain: all model parameters were considered uncertain (Table 1).

2.3.3 Parameter distributions

Decide on prior distributions from which the uncertain parameters and variables can be sampled: we have chosen uniform initial distributions for all parameters except u_0 , for which a logarithmic one was used (Table 1). To further narrow the sampling space, initial sample runs were made to localise parts of the parameter space that were more likely to generate good fits.

2.3.4 Random realisations of the model

Decide on a method to generate random realisations of models consistent with the assumptions in steps 1 and 2: twenty thousand parameter sets were drawn from uniform distributions for all parameters except u_0 , which was drawn from a log-distribution, (Table 1) and used as initial points in the Simulated annealing algorithm (Mathematica 7.0.0 Ubuntu/Linux). This procedure results in only one “optimum” set of parameters. The procedure was therefore repeated 28 000 times and the optimum sets together with their resulting likelihood values were stored. Calibration was made simultaneously against samples that had been kept at 5 + 5 °C, 15 + 15 °C and 25 + 25 °C.

2.3.5 Dotty plots

There exist a number of methods to assess sensitivity in non-linear models. The method most often used within the GLUE framework is to make a scatter-plot/dotty plot of each parameter (on the x-axis) versus the likelihood measure (y-axis) obtained during the calibration/conditioning process. From the resulting swarm of points, one can find trends showing for example that certain parameters are present in only a short interval of the initially sampled points, whereas others have a uniform density along the x-axis. If only a small segment of the initially sampled parameter space is found among the behavioural model runs, restricting that parameter to a smaller range will probably improve the number of behavioural model runs. On the contrary, if behavioural runs are equally distributed along the parameter axis, extending the parameter range might disclose/unravel areas of the parameter space which are more likely to prove behavioural.

2.4 Using the model

Behavioural parameter sets are used in ensemble runs to generate a mean output value and likely error bounds. An ensemble run is obtained when running the same

BGD

7, 8699–8722, 2010

**Quality or
decomposer
efficiency**

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



model with many parameter sets (as is the case in this article) or running different models to obtain a distribution of results. The likelihood measure, LM, or any other performance measure, can then be used to weigh the different outputs to create a weighted mean. Error bounds can be generated from the max and min from the model runs, or at any preferred significance level obtained from a cumulative density curve. In this article we will simply use max and min of the selected models as bounds.

3 Results

3.1 One initial quality, fixed decomposer efficiency

With one initial quality (q_0) and fixed decomposer efficiency (e_0) the dot plots (data not shown) showed more or less evenly distributed LM's, except for u_0 for which fits tended to be better with increasing u_0 . This indicates that the upper boundary for u_0 might have been too small. The generally even distribution of all parameters means that, within the used ranges, different parameters compensate for each other, making the model rather insensitive to changes in single parameters.

Ideally, all of modelled points should have been within the error bounds of the calibration data but that was not possible. Therefore, we decided to use all parameter sets with positive LM in the ensemble model runs. We then got 257 parameter sets that were behavioural. The best fit yielded a LM of 0.243, and was within boundaries at 37 out of 48 data points in the calibration set (Fig. 1).

When validated against experiments with shifts in temperature, the model follows the data well during the initial temperature phase; this is not surprising because it was calibrated on similar data (Fig. 1). During the final incubation after a temperature increase, the model underestimates the increase in respiration during the first days when going from 5 to 15 or 25 °C. When shifting downwards in temperature the model predicts initially slightly higher values than observed.

With fixed decomposer efficiency, temperature history has negligible effect on future respiration rates (Fig. 3a). The respiration at the final temperature after the shift for the

BGD

7, 8699–8722, 2010

**Quality or
decomposer
efficiency**

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 + 25 °C treatment is the same as the respiration after 12% C loss in the 25 + 25 °C treatment. The same holds for the 25 + 5 °C and 5 + 5 °C treatment.

3.2 One initial quality, flexible decomposer efficiency

With one initial quality (q_0) and flexible decomposer efficiency (e_0) there are few points in the dotted plots at the extremes of the x-axis for q_0 , meaning that high and low q_0 where unlikely to give good fits. u_0 , ΔG^0 and e_0 's are fairly evenly distributed (Fig. 4). η_{12} is skewed towards the lower end of the spectrum. The best fit yielded a LM of 0.284, and was within boundaries at 37 out of 48 points (Fig. 2). 33 sets were found behavioural, i.e. with LM > 0.

The model with flexible decomposer efficiency fits the data slightly better than the fixed decomposer efficiency version when validated against the experiment with temperature shifts, as well as bracketing more of the data points due to the wider uncertainty bounds (Fig. 2). When going down in temperature, the model seems to overshoot slightly, at least initially (15 + 5 °C, 25 + 5 °C) and when going up (5 + 25 °C and possibly 15 + 25 °C) the model misses the initial respiratory peak.

To search for trends in how e_0 varied with temperature we reran the simulations with the one q_0 flexible e_0 model to obtain a larger number (160) of behavioural parameter sets (LM > 0). Decomposer efficiencies were plotted in pairs, i.e. e_{0-5} versus e_{0-15} , e_{0-15} versus e_{0-25} , and e_{0-5} versus e_{0-25} (Fig. 5). The plots show that the e_0 's are highly correlated. Average e_0 decreased by 0.03 units when going from 5 to 15 °C (left), increased with 0.02 when going from 15 to 25 °C (middle), resulting in an overall decrease of 0.01 from 5 to 25 °C (right).

With temperature dependent decomposer efficiency, respiration responded strongly to temperature history (Fig. 3b). For example, the sample initially at 25 °C respired substantially more than the one initially at 5 °C when both were at 25 °C. Similarly, the sample initially at 25 °C respired more than the one initially at 5 °C when both are at 5 °C.

Quality or decomposer efficiency

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3 Two initial qualities, fixed and flexible decomposer efficiency

In the model runs with two initial qualities, the contribution (both at 0 and at 12% carbon loss) of the lower quality (q_{0-1}) to respiration is much lower (1/4700 and 1/2700 at 5 °C and 25 °C, respectively) than the respiration of the higher quality (q_{0-2}). Therefore the respiration from the high quality totally dominated the respiration and the model behaved qualitatively the same as with a single initial quality but with different “optimal” parameters.

4 Discussion

4.1 Model behaviour

The calibration data showed considerable variation in the variability between days. Also, the respiration did not always decrease monotonically as expected. We do not know whether this variability in input data comes from short-time biological variation or from measurement errors. We had, therefore, to relax the condition that, for each behavioural parameter set, predictions should be within error bounds for all points in each temperature series. Despite that, calibration to the constant temperature subsets worked well with R^2 values in the range of 0.83–0.96. However, even though the ensemble runs mostly covered all calibration points, at 5 and 15 °C the data points might have a more concave pattern than what the model can predict (Figs. 1 and 2 at 5 + 5 °C, 15 + 15 °C). When the model is validated against the temperature shift experiments, experiments tend to respond more strongly just at the temperature shift than the model.

4.2 Choice of likelihood measure (LM)

Our choice of likelihood measure, LM, is subjective. Ultimately, the objective should be to acquire parameters “useful in model prediction” (Beven, 2009, p. 124), and the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



LM should be chosen to help in doing so. One way of interpreting “useful in model prediction” is that the model should be able to bracket our observations, which it did in most of the cases (Figs. 1 and 2). Since the primary objective of this paper is not to model decomposition in general, but rather to highlight the effect of efficiency and quality coupled with temperature, the likelihood measure used is of lesser importance; see Beven (2009, pp. 165) for further discussion of choice of likelihood measure.

4.3 Mechanisms for temperature history to influence current respiration

4.3.1 One or several initial qualities

We have considered two main ways in which temperature history can affect current respiration rates. The first is that different qualities have different temperature dependencies, which should lead to a difference in quality composition at different temperatures and equal carbon loss (for a more detailed discussion, see Wetterstedt et al. (2010)). However, with our choice of two different initial qualities ($q_{0-1}=1.80$, $q_{0-2}=2.50$), the lower quality decomposed at only about 1/4700 at 5 °C and 1/2700 at 25 °C of the rate of the higher one. Together with the relatively small difference in temperature sensitivity between the two q_0 's (fixed e_0 , two q_0 model: $Q_{10-1} = 2.9$, $Q_{10-2} = 2.1$; flexible e_0 , two q_0 model: $Q_{10-1} = 2.9$, $Q_{10-2} = 2.3$) this did not in this short-term experiment translate into a sufficiently large difference in quality evolution between the temperatures; it is essentially only the highest quality that decomposes. Choosing a larger q_{0-1} resulted in more use also of the lower quality, but at the expense of a smaller difference in Q_{10} between the two qualities. However, whatever q_{0-1} is chosen, the effect on the temperature response is small.

4.3.2 Fixed or flexible decomposer efficiency

The second mechanism, letting decomposer efficiency (e_0) vary with temperature, resulted in a clear effect on quality distributions and thus temperature sensitivity and

BGD

7, 8699–8722, 2010

**Quality or
decomposer
efficiency**

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Quality or decomposer efficiency

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

respiration rates (Figs. 3 and 6). The reason for this is two-fold. Most importantly, with higher efficiency, when carbon is taken from the initial quality, a smaller part is lost by respiration and a larger part is converted to lower qualities. Thus, to obtain the same mass loss more of the initial quality has to be processed. Secondly, the dispersed, lower-quality, carbon will for the same reason persist for longer which means that yet more initial carbon needs to be processed before reaching the same cumulative respiration as at the lower decomposer efficiency. As a result, at equal mass losses, the higher efficiency produces a lower quality.

The flexible model is better fitted to the initially more rapid decrease at the beginning of the experiment as well as after the shift in temperature. Having a temperature dependent e_0 also leads to a model that simulates differences in respiration rates at the same final temperature when comparing samples from different initial incubation temperatures (Fig. 3). Surprisingly, having flexible decomposer efficiency resulted in fewer behavioural parameter sets. This is surprising because it adds two extra parameters which should increase the possibility of finding better fits. It seems however that the two extra parameters decreased the probability of finding good parameter sets and because the calibration was run with the same number of trial parameter sets, this resulted in fewer behavioural parameter sets.

The behaviour of e_0 points in the direction that decomposer efficiency might decrease with increasing temperature. This could be one of the explanations to why respiration is so strongly correlated to temperature. However, it can be difficult to compare e_0 between different models, between models and experiment, or indeed, between different experiments (cf. Devêvre and Horwath, 2000; Steinweg et al., 2008). In experiments e_0 is not measured directly and a number of more or less explicit assumptions are introduced when calculating e_0 from measurable quantities such as consumed substrate and respiration; such assumptions may or may not distort the relation between conceptual and observed values. In models, we also simplify the system; simplifications that differ between models.

4.3.3 Other temperature effects

One part where the model has difficulties in reproducing the experiment is directly after temperature shifts, where respiration is underestimated and overestimated after shifts upward and downward, respectively. These deviations between predictions and observations are similar to those observed when extrapolating respiration rates obtained at constant temperatures to temperature shifts in the study by Wetterstedt et al. (2010). We propose that these deviations over a few days represent transient adjustments in decomposer properties to new conditions. A possible interpretation is that these transients result from decomposer adaptation to new temperatures and that previously cold-adapted organisms respond more strongly than previously warm-adopted (Bradford et al., 2008). It remains an open question how important such transients may be under field conditions where temperatures are changing continuously, albeit less rapidly than in most experimental studies.

It should also be born in mind that the temperature response we find in e_0 depends on the assumptions we have made about the temperature dependence of the other factors. For example, we are assuming that the dispersion function is temperature independent although the rate of decomposition is highly sensitive to the strength of dispersion (Hyvönen et al., 2005). This is a simplifying assumption but we are not aware of any experiments demonstrating temperature sensitivity of dispersion. Likewise, although there are theoretical arguments for the effects of quality on the temperature dependence of the rate of carbon utilisation (Bosatta and Ågren, 1999), this has not been tested rigorously experimentally. Allison et al. (2010) point out another complication from temperature dependent decomposer efficiency. If decomposer efficiency goes down with temperature, decomposers assimilating the same amount of carbon will produce less biomass, which in turn should lead to a lower production of extracellular enzymes that can release assimilable carbon. In our terminology this should correspond a positive coupling between e_0 and u_0 . The increased loss of carbon caused by a temperature increase resulting from lowered decomposer efficiency would then be

BGD

7, 8699–8722, 2010

Quality or decomposer efficiency

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

counteracted by a lower use of carbon. It is possible that we need to consider whether there are additional such couplings between processes.

5 Conclusions

When fitting complex models with many adjustable parameters it is a common situation that many different parameter combinations are almost equally good and there is no additional information to be used for discriminating between them (Hyvönen et al., 2005; Beven, 2006). The strength of the GLUE method in a context like this one is that it does not select just one optimal set of parameters but allows all the possible parameters that match preselected conditions. What we learn in this study is that for almost all parameter combinations, e_{0-5} is larger than e_{0-15} and e_{0-25} . This is a strong suggestion that decomposer efficiency is, indeed, temperature dependent at least in the range 5 – 15 °C; above that range the results are less clear. This is a key result of this study.

A temperature-sensitive decomposer efficiency was shown to have a much stronger influence on quality differentiation, and thus respiration, than the temperature sensitivity of utilisation of different qualities. The difficulties in capturing changes in respiration rates at rapid temperature changes should caution us about extrapolating short term effects to longer time periods (cf. Wetterstedt et al., 2010); understanding the rate at which a microbial community can adjust requires more investigations. Our results show also that it is necessary to more carefully consider the temperature dependence of other processes than those directly coupled to the rate of substrate utilisation.

Acknowledgements. We would like to thank Keith Beven for help with the GLUE methodology. This work was supported by the Swedish Research Council.

BGD

7, 8699–8722, 2010

Quality or decomposer efficiency

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Ågren, G. I. and Bosatta, E.: Theoretical Ecosystem Ecology – Understanding Element Cycles, 234 pp., 1998.
- Ågren, G. I. and Wetterstedt, J. Å. M.: What determines the temperature response of soil organic matter decomposition?, *Soil Biol. Biochem.*, 39(7), 1794–1798, 2007.
- Allison, S. D., Wallenstein M. D., and Bradford, M. A.: Soil-carbon response to warming dependent on microbial physiology, *Nat. Geosci.*, 3, 336–340, 2010.
- Beven, K.: A manifesto for the equifinality thesis, *J. Hydrol.*, 320(1–2), 18–36, doi:10.1016/j.jhydrol.2005.07.007, 2006.
- Beven, K.: Environmental modelling: An uncertain future?, Routledge, London, 310 pp., 2009.
- Beven, K. and Binley, A.: The future of distributed models: Model calibration and uncertainty prediction, *Hydrol. Process.*, 6(3), 278–298, doi:10.1002/hyp.3360060305, 1992.
- Bosatta, E. and Ågren, G.: Soil organic matter quality interpreted thermodynamically, *Soil Biol. Biochem.*, 31(13), 1889–1891, 1999.
- Bosatta, E. and Ågren, G. I.: Exact solutions to the continuous-quality equation for soil organic matter turnover, *J. Theor. Biol.*, 224(1), 97–105, 2003.
- Bradford, M. A., Davies, C. A., Frey, S. D., Maddox, T. R., Melillo, J. M., Mohan, J. E., Reynolds, J. F., Treseder, K. K., and Wallenstein, M. D.: Thermal adaptation of soil microbial respiration to elevated temperature, *Ecol. Lett.*, 11(12), 1316–1327, doi:10.1111/j.1461-0248.2008.01251.x, 2008.
- Bruun, S., Ågren, G. I., Christensen, B. T., and Jensen, L. S.: Measuring and modeling continuous quality distributions of soil organic matter, *Biogeosciences*, 7, 27–41, doi:10.5194/bg-7-27-2010, 2010.
- Coleman, K. and Jenkinson, D. S.: RothC-26.3 – a model for the turnover of carbon in soil: Model description and users guide, Lawes Agricultural Trust, Harpenden, 45 pp., 1995.
- Comins, H. N. and McMurtrie, R. E.: Long-term response of nutrient-limited forests to CO₂ enrichment; Equilibrium behavior of plant-soil models, *Ecol. Appl.*, 3(4), 666–681, 1993.
- Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440(7081), 165–173, doi:10.1038/nature04514, 2006.
- Devèvre, O. C. and Horwath, W. R.: Decomposition of rice straw and microbial carbon use efficiency under different soil temperatures and moistures, *Soil Biol. Biochem.*, 32(11–12), 1773–1785, doi:10.1016/S0038-0717(00)00096-1, 2000.

Quality or decomposer efficiency

J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



BGD

7, 8699–8722, 2010

**Quality or
decomposer
efficiency**J. Å. M. Wetterstedt and
G. I. Ågren

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Hyvönen, R., Ågren, G. I., and Dalias, P.: Analysing temperature response of decomposition of organic matter, *Glob. Change Biol.*, 11, 770–778, doi:10.1111/j.1365-2486.2005.00947.x, 2005.
- 5 Kirschbaum, M. U. F.: The temperature dependence of organic-matter decomposition—still a topic of debate, *Soil Biol. Biochem.*, 38(9), 2510–2518, 2006.
- Parton, W. J., Schimel, D. S., Cole, C. V., and Ojima, D. S.: Analysis of Factors Controlling Soil Organic Matter Levels in Great Plains Grasslands, *Soil Sci. Soc. Am. J.*, 51(5), 1173–1179, 1987.
- 10 Solomon, S.: *Climate Change 2007, The physical science basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, 996 pp., 2007.
- Steinweg, J. M., Plante, A. F., Conant, R. T., Paul, E. A., and Tanaka, D. L.: Patterns of substrate utilization during long-term incubations at different temperatures, *Soil Biol. Biochem.*, 40(11), 2722–2728, 2008.
- 15 Wetterstedt, J. Å. M., Persson, T., and Ågren, G. I.: Temperature sensitivity and substrate quality in soil organic matter decomposition: results of an incubation study with three substrates, *Glob. Change Biol.*, 16, 1806–1819, 2010.

Quality or decomposer efficiency

J. Å. M. Wetterstedt and G. I. Ågren

Table 1. List of parameters. Parameter values are those corresponding to the highest *LM*. Range is the range used in the calibration procedure. Sampling was done uniformly for all parameters but u_0 which was sampled from a log-distribution. With two initial qualities the q_0 's (q_{0-1} and q_{0-2}) were set to values obtained with one initial quality (q_0). Fixed and Flex. in the table heading refer to fixed or flexible e_0 . Initial parameter range was greater to find suitable parameter space.

Parameter	Range	One q_0		Two q_0		Comment
		Fixed	Flex.	Fixed	Flex.	
q_0	1.5–4.5	2.50	2.5	–	–	Initial litter quality (unit-less)
q_{0-1}	fixed	–	–	1.80	1.80	1st initial litter quality in the two q_0 version
q_{0-2}	fixed	–	–	2.50	2.50	2nd initial litter quality in the two q_0 version
ΔG^0	a	127	93	127	99.5	Gibbs' free energy (kJ)
e_0	0.15–0.40	0.377	–	0.220	–	Decomposer efficiency in the fixed version (unit-less)
e_{0-5}	– : –	–	0.369	–	0.393	e_0 at 5 °C in the flexible version
e_{0-15}	– : –	–	0.327	–	0.336	e_0 at 15 °C – : –
e_{0-25}	– : –	–	0.321	–	0.330	e_0 at 25 °C – : –
u_0	10^7 – 10^9	7.63×10^7	2.46×10^7	65.7×10^7	1.21×10^7	Decomposer growth rate regulating parameter ($\text{g C (g C)}^{-1}\text{d}^{-1}$)
η_{12}	0.1–0.45	0.120	0.199	0.0663	0.0476	Rate of quality decrease, approximate solution (unit-less)
Q_{10}	1.4–4.0					Only used to create plausible parameter range for ΔG^0

^a For ΔG^0 the range is $q_0 R^{\frac{T_5 - T_{15}}{10}} [-\log [Q_{10 \max}] \dots -\log [Q_{10 \min}]]$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. Model predictions of respiration rates for the one initial quality, fixed decomposer efficiency model and observed respiration rates for all combinations of initial (T_i) and final (T_f) temperatures (5, 15, and 25°C.) Weighted ensemble run predictions (solid black line) with max/min curves (blue dashed lines) for the behavioural parameter sets. The yellow fields show 95% error bounds around measured data points (dots). Least square R^2 -values are shown in top right corner of each sub-graph.




Fig. 2. Model predictions of respiration rates for the one initial quality, flexible decomposer efficiency model and observed respiration rates for all combinations of initial (T_i) and final (T_f) temperatures (5, 15, and 25°C.) Weighted ensemble run predictions (solid black line) with max/min curves (blue dashed lines) for the behavioural parameter sets. The yellow fields show 95% error bounds around measured data points (dots). Least square R^2 -values are shown in top right corner of each sub-graph.

Quality or decomposer efficiency


J. Å. M. Wetterstedt and
G. I. Ågren

Fig. 3. Respiration rates of samples as a function of fraction of respired carbon, predicted by the flexible and fixed decomposer efficiency models with constant temperatures (solid lines) and with a switch between 5 and 25 °C at 12% respired carbon (broken lines) using the parameter set with the highest LM (Table 1). For clarity, the transitions between temperatures are slightly displaced from exactly 12%. **(a)** Fixed e_0 . **(b)** Flexible e_0 . Note, these simulations do not give the same result as the weighted ensemble predictions (Figs. 2 and 3).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Dotty plots for the one initial quality, flexible decomposer efficiency. Each dot represents one model run. The likelihood measure, LM, or “goodness of fit”, is plotted against parameter values. The x-scales cover the allowed ranges of the parameters. Parameter sets resulting in a $LM > 0$ where used in ensemble simulation runs. Only simulations with $LM > -4$ are shown.

Quality or decomposer efficiency

J. Å. M. Wetterstedt and G. I. Ågren

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Quality or decomposer efficiency

J. Å. M. Wetterstedt and
G. I. Ågren




Fig. 5. Correlations between efficiencies at 5, 15, and 25°C from behavioural parameter sets in the one initial quality, flexible decomposer efficiency model. Solid line: linear regression of data. Broken line: 1:1 line.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Quality or decomposer efficiency


J. Å. M. Wetterstedt and
G. I. Ågren

Fig. 6. Distributions of qualities for combinations of one or two initial qualities and fixed or flexible decomposer efficiency (e_0) when 12% carbon has been lost for sample incubated at 5 or 25°C. Solid lines and black bars are for samples at 5°C. Dashed lines and grey bars are for samples at 25°C. The bars have been shifted slightly leftwards and rightwards from their value to visually separate them. Bars show the amount of carbon that has not been used by decomposers so far (remaining at the initial qualities). The lines show the distributions of carbon that the decomposers have converted into new qualities. With flexible decomposer efficiency, more carbon has been converted (lines) and less remains at the initial quality (bar). With two initial qualities the losses have essentially only occurred from the highest quality.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion