

Interactive  
Comment

***Interactive comment on “Carbon input control  
over soil organic matter dynamics in a temperate  
grassland exposed to elevated CO<sub>2</sub> and warming”  
by Y. Carrillo et al.***

**Y. Carrillo et al.**

dcarril1@uwyo.edu

Received and published: 25 March 2010

Response to anonymous reviewer # 1

Comment 1

We appreciate the opportunity to address Reviewer 1's concerns about the application of controlled laboratory incubations to assess soil organic matter (SOM) decomposition. This topic has generated considerable debate in the literature, including several highly cited papers in Biogeosciences (Conen et al., 2006; Reichstein et al., 2005). We contend that laboratory incubation is a valuable approach which is sensitive to changes in SOM decomposability that cannot be detected using other methods. The present

C315

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



study demonstrates (for the first time in the literature) interactive effects of experimental warming and elevated carbon dioxide concentrations on the proportion of soil C in labile form and its decomposability, despite no apparent effect in the total C pool. We also discovered an interesting and novel relationship between resistant SOM decomposition rate and biomass and dissolved organic carbon, indicated the importance of substrate availability of labile C in resistant C turnover. This priming effect is likely to play a strong role in regulating climate change effects on resistant SOM decomposition rates and is a unique and important finding of this work.

We agree that actual field decomposition rates will be dependent on field temperature and moisture conditions. However, our work aimed at determining whether the exposure to the elevated carbon dioxide and warming treatments had generated changes in the intrinsic decomposability of the labile and resistant pools. That is precisely an advantage of incubations at standard conditions: they allow to assess whether experimental treatments have modified the characteristics of C pools that affect their susceptibility to degradation, while removing the environmental constraints and confounding effects at play in a field situation (Davidson and Janssens, 2006). The temperature and moisture levels chosen for our study are near optimal, and also relevant to natural field conditions.

We strongly disagree with the idea that laboratory incubations cannot be used to estimate kinetic parameters. Most ecosystem and global scale biogeochemistry models use decomposition constants derived from laboratory incubations, such as CENTURY (Parton et al., 1987; Parton et al., 1993) and ROTHAMSTED-C (Jenkinson et al., 1991). In addition, many previous studies have used laboratory incubation approaches to evaluate responses in the kinetic parameters of organic matter pools to experimental field treatments (e.g. Collins et al., 2000; Dijkstra and Cheng, 2007; Fissore et al., 2008; Pendall and King, 2007; Taneva and Gonzalez-Meler, 2008; Dijkstra et al., 2006) as well as to compare processes rates under standard conditions as affected by previous exposure to experimental treatments (e.g. Carney et al., 2007; Langley

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

et al., 2009). Attempts to assess kinetic decomposition parameters from intact soils in the field are unfortunately confounded by contributions from root respiration, unless mechanical or isotopic partitioning techniques are used (Hanson et al., 2000), and by moisture limitations (Davidson and Jannsens, 2006). Laboratory incubations therefore complement field studies.

#### Comment 2

We are happy to clarify the potential contribution of carbonates to the evolved CO<sub>2</sub> in the laboratory incubations. Carbonate contents in our field site are highly stratified and the concentrations in top soil are very low. Average carbonate content by weight in our soils is 0.0045 % at 0-5 cm (S.D. = 0.062 as in most cases carbonates were not detectable) and 0.2056 % (S.D.= 0.61) at 5-15 cm (Sherrod et al., 2002). Further, the isotopic composition of CO<sub>2</sub>-C released from these soil incubations ranges from -18 to -25‰ consistent with SOM values in the different treatments (CO<sub>2</sub> and SOM  $\delta^{13}\text{C}$  data not included in this manuscript but available on request). Thus, we are confident that C from carbonates did not cause a bias in our results. All of our soils are routinely processed for carbonate removal before C and  $\delta^{13}\text{C}$  analyses for consistency, because carbonate content below 15 cm is higher.

#### Comment 3

Our study is unique because it utilizes soil exposed to the combined effects of elevated carbon dioxide and ecosystem warming in an intact native grassland. We agree that the early responses measured in the first few years are likely to contrast longer-term responses (Melillo et al., 2002). Our study is valuable because it documents early responses in sensitive (labile) pools, supporting the hypothesis that labile C may be more susceptible to initial impacts of experimental climate change than resistant C. Part of the reason why responses of soil C to climate change treatments are difficult to detect in the short term is that most studies do not assess individual C pools. One of the contributions of our study results from the fact that individual pools were assessed.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Additional ongoing experiments are following the longer-term responses of SOM pool dynamics in the PHACE experiment.

## References

Carney, K. M., Hungate, B. A., Drake, B. G., and Megonigal, J. P.: Altered soil microbial community at elevated CO<sub>2</sub> leads to loss of soil carbon, *Proc. Natl. Acad. Sci. U. S. A.*, 104, 4990-4995, 2007.

Collins, H. P., Elliott, E. T., Paustian, K., Bundy, L. C., Dick, W. A., Huggins, D. R., Smucker, A. J. M., and Paul, E. A.: Soil carbon pools and fluxes in long-term corn belt agroecosystems, *Soil Biol. Biochem.*, 32, 157-168, 2000.

Conen, F., Leifeld, J., Seth, B., and Alewell, C.: Warming mineralises young and old soil carbon equally, *Biogeosciences*, 3, 515-519, 2006.

Davidson, E. A., and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165-173, 2006.

Dijkstra, F. A., Hobbie, S. E., and Reich, P. B.: Soil processes affected by sixteen grassland species grown under different environmental conditions, *Soil Sci. Soc. Am. J.*, 70, 770-777, 2006.

Dijkstra, F. A., and Cheng, W. X.: Interactions between soil and tree roots accelerate long-term soil carbon decomposition, *Ecol. Lett.*, 10, 1046-1053, 2007.

Fissore, C., Giardina, C. P., Kolka, R. K., Trettin, C. C., King, G. M., Jurgensen, M. F., Barton, C. D., and McDowell, S. D.: Temperature and vegetation effects on soil organic carbon quality along a forested mean annual temperature gradient in North America, *Global Change Biology*, 14, 193-205, 2008.

Hanson, P. J., Edwards, N. T., Garten, C. T., and Andrews, J. A.: Separating root and soil microbial contributions to soil respiration: A review of methods and observations, *Biogeochemistry*, 48, 115-146, 2000.

**BGD**

7, C315–C320, 2010

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Jenkinson, D. S., Adams, D. E., and Wild, A.: Model Estimates of CO<sub>2</sub> Emissions from Soil in Response to Global Warming, *Nature*, 351, 304-306, 1991.

Langley, J. A., McKinley, D. C., Wolf, A. A., Hungate, B. A., Drake, B. G., and Megonigal, J. P.: Priming depletes soil carbon and releases nitrogen in a scrub-oak ecosystem exposed to elevated CO<sub>2</sub>, *Soil Biol. Biochem.*, 41, 54-60, 2009.

Melillo, J. M., Steudler, P. A., Aber, J. D., Newkirk, K., Lux, H., Bowles, F. P., Catri-  
cala, C., Magill, A., Ahrens, T., and Morrisseau, S.: Soil warming and carbon-cycle  
feedbacks to the climate system, *Science*, 298, 2173-2176, 2002.

Parton, W. J., Schimel, D. S., Cole, C. V., and Ojima, D. S.: Analysis of Factors Con-  
trolling Soil Organic-Matter Levels in Great-Plains Grasslands, *Soil Sci. Soc. Am. J.*,  
51, 1173-1179, 1987.

Parton, W. J., Scurlock, J. M. O., Ojima, D. S., Gilmanov, T. G., Scholes, R. J., Schimel,  
D. S., Kirchner, T., Menaut, J. C., Seastedt, T., Moya, E. G., Kamnalrut, A., and Kinya-  
mario, J. I.: Observations and Modeling of Biomass and Soil Organic-Matter Dynamics  
for the Grassland Biome Worldwide, *Global Biogeochemical Cycles*, 7, 785-809, 1993.

Pendall, E., and King, J. Y.: Soil organic matter dynamics in grassland soils under  
elevated CO<sub>2</sub>: Insights from long-term incubations and stable isotopes, *Soil Biol.*  
*Biochem.*, 39, 2628-2639, 2007.

Reichstein, M., Katterer, T., Andren, O., Ciais, P., Schulze, E. D., Cramer, W., Papale,  
D., and Valentini, R.: Temperature sensitivity of decomposition in relation to soil organic  
matter pools: critique and outlook, *Biogeosciences*, 2, 317-321, 2005.

Sherrod, S. K., Belnap, J., and Miller, M. E.: Comparison of methods for nutrient mea-  
surement in calcareous soils: Ion-exchange resin bag, capsule, membrane, and chem-  
ical extractions, *Soil Sci.*, 167, 666-679, 2002.

Taneva, L., and Gonzalez-Meler, M. A.: Decomposition kinetics of soil carbon of differ-  
ent age from a forest exposed to 8 years of elevated atmospheric CO<sub>2</sub> concentration,

**BGD**

7, C315–C320, 2010

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Soil Biol. Biochem., 40, 2670-2677, 2008.

---

Interactive comment on Biogeosciences Discuss., 7, 1575, 2010.

**BGD**

7, C315–C320, 2010

---

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C320

