

Interactive comment on “Synoptic relationships quantified between surface Chlorophyll-a and diagnostic pigments specific to phytoplankton functional types” by T. Hirata et al.

Anonymous Referee #1

Received and published: 12 October 2010

Error-quantified, synoptic-scale relationships between satellite based chlorophyll-a (Chla) and phytoplankton pigment groups at the sea surface are presented. A total of nine pigment groups were considered to represent nine phytoplankton functional types (PFTs) including microplankton, nanoplankton, picoplankton, diatoms, dinoflagellates, green algae, picoeukaryotes, prokaryotes and Prochlorococcus sp. The present paper applies the approach of Uitz et al. (2006) to get from satellite (SeaWiFS) chl-a directly information on the fractional contribution of various phytoplankton classes by a parameterization based on a large data set on the distribution of diagnostic pigments obtained from global HPLC data sets. The paper extends the Uitz et al. 2006 approach to get more phytoplankton groups besides the three size classes, but it limits the approach to

C3292

get only surface information instead of information on the vertical profile.

The paper contributes with its error estimates and a large (and extended as compared to Uitz et al. 2006) data base to the understanding of phytoplankton group dynamics for marine ecosystem and biogeochemical modelling and therefore fits well in the scope of the paper it has been submitted to. However, still a major revision of the paper is necessary to improve the results and discussion section: The discussion of the results in comparison to other methods has not been done and also the shortcoming of the presented method are not discussed in detail. One fundamental limitation of the method is that it is only able to retrieve from satellite chl-a phytoplankton groups dynamics which are expected. Unexpected shifts in PFT composition will not be detected by this method because no optical information measured by the ocean color satellite sensor is taken into account. In addition, the approach uses diagnostic pigments to infer the different phytoplankton groups globally, but this has not been verified globally by other techniques such as microscopy, particle size counter and flowcytometry. Since this is probably beyond the scope of the paper, this should at least be discussed as a limitation to the method. In addition, the paper should be extended by using the results of the fractional contribution of the various phytoplankton groups to the total chl-a to calculate their absolute chl-a and the dominant phytoplankton group for each pixel. In addition, it will be of great value to see the seasonal variation of relative and/or absolute phytoplankton composition. By that the results can directly be compared to other approaches to derive phytoplankton groups from satellite measurements and data are useful for further modelling studies. Also some citations in the text are not appropriate and should be changed.

In detail the following should be changed: - why write e.g. on page 6677 line 18 and ff. 5.8[% Chla] instead of 5.8% Chla? The brackets do not make sense. - Abstract, page 6677, line 25 ff. The sentence is unclear? If nanoplankton is the background globally with only 44%, why picoplankton with 47% globally is only the background for the subtropical gyres - this has to be expressed more clearly? -Page 6678, line 11-14 the

C3293

citation of Nair et al. (2008) for the DMSP production in the ocean is not appropriate! Better use Sunda et al. (2002), Nature 418, p317-320 -Page 6678, line 21: the model is called "NOBM" not "NOMB" -Page 6680, line 3-11ff: you should also add for completeness to this sentence that another method to get information on the distribution of phytoplankton groups is the method by Bracher et al. (2009) which uses hyperspectral satellite data to get quantitative information on the distribution of phytoplankton groups (diatoms and cyanobacteria). -Page 6688, Line 15: three words are missing after ("instead of % Chla")- read again and change it -Page 6688, last paragraph- in order to judge if the different size classes but also other phytoplankton groups play a dominant role overall or in certain regions, it would be helpful to calculate the dominant groups globally. Also add to the results the chl-a conc. of the different phytoplankton groups. Both will enable the comparison of results to other methods deriving PFT or size classes from satellite ocean color data. -Page 6689: as said above, improve the discussion by discussing the limitations to the presented method, e.g. that unexpected shifts in PFT composition will not be detected by this method and that to use diagnostic pigments in order to get certain various phytoplankton groups from HPLC data needs further verification. - Page 6682: give size classes for micro-, nano- and picoplankton in μm in the text in order to make clear what does this classification mean discuss the seasonal coverage of the supplied HPLC data

Interactive comment on Biogeosciences Discuss., 7, 6675, 2010.

C3294