

Interactive  
Comment

## ***Interactive comment on “Soil-atmosphere exchange of nitrous oxide, methane and carbon dioxide in a gradient of elevation in the coastal Brazilian Atlantic forest” by E. Sousa Neto et al.***

**E. Sousa Neto et al.**

eraklito@gmail.com

Received and published: 19 January 2011

1. The effect of increasing temperature on CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes The authors suggest that increasing temperatures will result in a consequent increase in soil CO<sub>2</sub> and N<sub>2</sub>O fluxes and CH<sub>4</sub> consumption. Considering the limitations of their observed data (i.e. once a month frequency and missing data) and provided statistical results (i.e. no clear correlation between soil temperature with N<sub>2</sub>O and CH<sub>4</sub> fluxes through the gradient of elevation), the suggestion may not be robustly supported by the results of this study.

Author’s comment: We accept the reviewer’s comment. Temperature increase alone

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



may be insufficient to cause increases in N<sub>2</sub>O and CO<sub>2</sub> emissions and in CH<sub>4</sub> uptake rates. The variation of soil gas emissions with altitude responds to combinations of factors including climatic conditions, species composition and structure, nutrient supply and soil physical and chemical properties. We reformulated the abstract and the conclusions to reflect greater caution. See lines 35-38 and lines 352-364 in the manuscript.

2. Estimate cumulative annual CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes for different altitudes The authors provide annual mean flux of N<sub>2</sub>O and CH<sub>4</sub> for different altitudes and statistical significance in the difference of fluxes by altitudes. The cumulative annual flux of N<sub>2</sub>O and CH<sub>4</sub> could also be calculated by linear interpolation and numerical integration of observed fluxes between sampling times (i.e. area under the flux curves) and the authors can test the significance in differences of cumulative annual flux by altitudes. Additional efforts will provide clearer information. In the case of CO<sub>2</sub> flux, the Q10 model (relationship between soil temperature and CO<sub>2</sub> flux) can be developed with currently available soil temperature and CO<sub>2</sub> flux data and the authors can then estimate the missing CO<sub>2</sub> flux (Oct. 2006 to Dec. 2006) using the Q10 model with observed soil temperature (Oct. 2006 to Dec. 2006). After filling the gap of CO<sub>2</sub> flux, the authors can estimate cumulative annual CO<sub>2</sub> flux.

Author's comment: We accept the reviewer's comment. We calculated the cumulative annual flux of N<sub>2</sub>O and CH<sub>4</sub> and tested the differences of cumulative annual flux by altitudes. This calculation approach is now described in the methods and results sections (See lines 175-177, 220-224) and is summarized for N<sub>2</sub>O and CH<sub>4</sub> in Table 5. The statistical outcome does not change for N<sub>2</sub>O. For CH<sub>4</sub>, the cumulative results show no significant difference among altitudes. This is because the test has fewer degrees of freedom. We do not believe that the cumulative results for N<sub>2</sub>O or CH<sub>4</sub> change any of our conclusions. They may be useful to compare our sites to other sites. (See lines 175-177, 220-224 and Table 5). For CO<sub>2</sub>, we estimated the missing CO<sub>2</sub> values by using the exponential model and estimated the cumulative annual CO<sub>2</sub> fluxes as suggested. See lines 177-180, 225-236.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

3. Add a rainfall figure and check possibility of peak N<sub>2</sub>O emission caused by rewetting of dry soil. Beyond the significantly higher annual mean flux of N<sub>2</sub>O at 100 m, two very high N<sub>2</sub>O peak emissions occurred in Dec. 2006 and Jun. 2007 in 100 m. Since the soil temperatures of both months were not particularly high (Fig. 1), the peak emissions may not be caused by high soil microbial activity influenced by soil temperature. Looking at the WFPS figure (Fig. 2), there is an interesting common point: WFPS abruptly increased in both months. Considering the well-matched timing of N<sub>2</sub>O peak emissions and WFPS changes, it is possible the peak emissions may be caused by rewetting events (rewetting of dry soils). Studies have reported increased soil N<sub>2</sub>O emission following the wetting of dry soil in various ecosystems, including forest (e.g., Groffman Tiedje, 1988; Vitousek et al., 1989; Garcia-Mendez et al., 1991; Davidson et al., 1993; Nobre et al., 2001). If there are available rainfall data for the sites (i.e. from the nearest weather station) plot them with WFPS and check whether the changes of WFPS were associated with rainfall events. If they are well matched, further discussion related to N<sub>2</sub>O peak emissions and rewetting events will contribute to our understanding of N<sub>2</sub>O fluxes. It would be very interesting if the authors could discuss why high N<sub>2</sub>O peak emissions occurred only in the 100-m site.

Author's comment: We followed the referee's interesting suggestion but we found no evidence for a rewetting event as the cause of the increased N<sub>2</sub>O fluxes. In our text (lines 298-306) we discuss the coincidence of three factors that promote N<sub>2</sub>O production, soil moisture, soil temperature and organic matter decomposition at the beginning of the Austral summer. This explanation may explain the December peak in N<sub>2</sub>O emissions but does not explain the June peak.

Please also note the supplement to this comment:

<http://www.biogeosciences-discuss.net/7/C4797/2011/bgd-7-C4797-2011-supplement.pdf>

Interactive comment on Biogeosciences Discuss., 7, 5227, 2010.

C4799

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

