



Received: 6 December 2011 – Accepted: 7 December 2011 – Published: 21 December 2011

Correspondence to: O. Duteil (oduteil@ifm-geomar.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**BGD**

8, 12423–12449, 2011

---

**Preformed and  
regenerated PO<sub>4</sub> in  
OGCMs**

O. Duteil et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Phosphate distributions simulated by seven state-of-the-art biogeochemical ocean circulation models are evaluated against observations of global ocean nutrient distributions. The biogeochemical models exhibit different structural complexities, ranging from simple nutrient-restoring to multi-nutrient NPZD type models. We evaluate the simulations using the observed volume distribution of phosphate. The errors in these simulated volume class distributions are significantly larger when preformed phosphate (or regenerated phosphate) rather than total phosphate is considered. Our analysis reveals that models can achieve similarly good fits to observed total phosphate distributions for a very different partitioning into preformed and regenerated nutrient components. This has implications for the strength and potential climate sensitivity of the simulated biological carbon pump. We suggest complementing the use of total nutrient distributions for assessing model skill by an evaluation of the respective preformed and regenerated nutrient components.

## 1 Introduction

The distribution of phosphate ( $\text{PO}_4$ ) in the ocean is controlled by a wide range of processes. In the absence of major external sources, internal processes are the main drivers, namely the physical circulation and the marine biology.  $\text{PO}_4$  is incorporated into organic matter by photosynthetic phytoplankton in the euphotic zone. A fraction of this organic matter escapes recycling in the surface layer and is exported into the ocean interior. Remineralisation of organic matter by bacteria and zooplankton releases nutrients such as  $\text{PO}_4$  and consumes oxygen. This phosphate is called “regenerated phosphate”. Biotically unutilized surface phosphate can also enter the ocean by subduction during water mass formation. According to observations, this so-called “preformed phosphate” makes up for about 64 % of the global ocean’s phosphate inventory, whereas regenerated phosphate derived from remineralisation of organic matter makes up for the remaining 36 % (Ito and Follows, 2005).

**BGD**

8, 12423–12449, 2011

## Preformed and regenerated $\text{PO}_4$ in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Preformed and regenerated PO<sub>4</sub> in OGCMs**

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Patterns of PO<sub>4</sub> concentrations as well as the partitioning into preformed and regenerated components are sensitive to ocean circulation, export production and remineralisation length scale. A more active biology increases the transfer of phosphate from surface to intermediate depth by remineralisation and consequently depletes surface waters, decreasing the amount of phosphate subducted at high latitudes and thus decreasing preformed PO<sub>4</sub> and increasing regenerated PO<sub>4</sub> concentrations. An accelerated circulation results in higher supply rates of nutrients from the deep ocean to the euphotic zone, often increasing surface nutrient levels and thereby tending to increase the preformed PO<sub>4</sub> and decrease the regenerated PO<sub>4</sub> pool. A model that correctly simulates biogeochemical and physical processes should represent not only the total phosphate distribution, but also reproduce its preformed and regenerated pools in a realistic way.

The global PO<sub>4</sub> distribution is characterised by generally low PO<sub>4</sub> values in surface waters and high concentrations in the deep ocean, with maximum concentrations often found in intermediate waters associated with low oxygen concentrations. This picture is usually well reproduced by ocean general circulation models, as depicted by model intercomparison studies (Najjar et al., 2007), which have been conducted for instance in the context of the Ocean Carbon-cycle Model Intercomparison Project (OCMIP). OCMIP compared models with a simplistic representation of marine biogeochemistry, i.e. restoring of nutrients at the sea surface and no explicit representation of phytoplankton. More complex biological models were used in the Coupled Carbon Cycle Climate Model Intercomparison Project (C4MIP) that focussed on the coupling of climate and the carbon cycle (Friedlingstein et al., 2006). Schneider et al. (2008) assessed surface PO<sub>4</sub>, export and primary production simulated by three C4MIP models. Carbon fluxes at the ocean atmosphere interface have been assessed using a suite of fully coupled climate models (Roy et al., 2011).

All these studies focused on total nutrient concentrations. The partitioning between regenerated and preformed pools has, to our knowledge, not been considered in previous model intercomparison studies. Previous studies aimed at a better understanding

of the interactions of climate and biogeochemistry have found preformed  $\text{PO}_4$  to be an excellent indicator of atmospheric  $p\text{CO}_2$ , with a positive correlation between the atmospheric  $\text{CO}_2$  levels and the relative amount of preformed phosphate in the ocean interior (Marinov et al., 2008). By contrast, the total surface nutrient inventory is not necessarily correlated with atmospheric  $p\text{CO}_2$  and oceanic circulation changes.

The objective of this study is to illustrate that additional information can be gained by explicitly considering preformed and regenerated nutrients pools in the evaluation of biogeochemical models. Without the intention to perform a complete model intercomparison, we here exploit 7 state-of-the-art coupled biogeochemical-circulation models to demonstrate how the separation of the phosphate into preformed and regenerate concentrations can improve the assessment of the simulations.

## 2 Datasets

Seven global biogeochemical-circulation models used in previous studies are considered here (see Table 1 for details): OPA-PISCES (Aumont and Bopp, 2006), MPIOM-HAMOCC (Maier-Reimer, 1993; Maier-Reimer et al., 2005), CCSM3-BEC (Krishnamurthy et al., 2009), UVIC2.8 (Oschlies et al., 2008), om1p7-BLINGv0 (Galbraith et al., 2010), CSIRO (Matear and Hirst, 2003), MOM P2A (Gnanadesikan et al., 2004).

Given that the partitioning into preformed and regenerated nutrients is likely sensitive to the models' parameter choices and the forcing used, the behaviour of the models shown here should not be extrapolated to other model configurations.

All models were forced by prescribed atmospheric conditions in an attempt to obtain realistic pre-industrial circulation fields. The pelagic ecosystem part of the biogeochemical models ranges from relatively simple nutrient restoring-type models (CSIRO, MOM P2A) to intermediate complexity nutrient-phytoplankton-zooplankton-detritus (NPZD) (UVIC2.8, om1p7-BLINGv0) and more complex models with multiple plankton functional types (OPA-PISCES, MPIOM-HAMOCC, CCSM3-BEC). All models were integrated for at least 2000yr to reach a steady state quasi-equilibrium.

**BGD**

8, 12423–12449, 2011

## Preformed and regenerated $\text{PO}_4$ in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Annual mean output from these models is compared against objectively analysed annual means of the World Ocean Atlas (WOA) 2005 (Garcia et al., 2006a,b).

Prior to analysis, all model output fields were regridded onto the 33 levels 1x1 degree WOA grid.

The distribution of the regenerated phosphate can be expressed in a general way as  $P_{\text{Regenerated}} = \sum P_{\text{Regenerated (Oxidant)}}$  where possible oxidants are oxygen, nitrate, manganese oxides, iron oxides, sulfate and eventually  $\text{CO}_2$ . Only the first one or two terms are globally relevant and hence usually considered in OGCMs.

For the main analysis of this paper, we assume that  $P_{\text{Regenerated}}$  can be approximated by  $P_{\text{Regenerated (O}_2)}$ . Impacts of denitrification, i.e.  $P_{\text{Regenerated (NO}_3)}$ , are estimated for models that include both a nitrogen and phosphorus cycle (see Appendix). However, due to the limited validity of this estimate and the relatively small impact of phosphorus regeneration by denitrification on our results (see Fig. A1), we focus on aerobic processes in the main part of this study. Regenerated phosphate is computed by multiplying the Apparent Oxygen Utilisation (AOU) with the oxidation ratio  $R_{\text{P}:-\text{O}_2}$

$$P_{\text{Regenerated}} = R_{\text{P}:-\text{O}_2} \text{AOU} \quad (1)$$

Apparent Oxygen Utilisation (AOU) evaluates integral oxygen consumption by biological activity and is computed as the difference between the oxygen saturation concentration, which depends on thermohaline properties (Weiss, 1970), and the actually observed oxygen concentration. The  $R_{\text{P}:-\text{O}_2}$  ratio is assumed to be a fixed and constant stoichiometric ratio between phosphate production and oxygen consumption.

This ratio has been set to 1/172 in OPA-PISCES, MPIOM-HAMOCC, 1/170 in CCSM3-BEC, CSIRO, MOM P2A, UVIC2.8 and 1/150 in om1p7-BLINGv0. We use these model values when analyzing the respective models outputs. A common  $R_{\text{P}:-\text{O}_2}$  value of 1/170 is used for the analysis of WOA data independently of the value considered in the different models. Indeed, generally accepted values derived from ocean data are close to 1/170 (Takahashi et al., 1985; Kortzinger et al., 2001; Anderson and Sarmiento, 1994; Li et al., 2000).

**Preformed and regenerated  $\text{PO}_4$  in OGCMs**

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Preformed phosphate is computed as the difference between total and regenerated phosphates

$$P_{\text{Preformed}} = P - P_{\text{Regenerated}} \quad (2)$$

We note that, despite the rapidity of the air-sea gas exchange, equilibrium of surface water with atmospheric oxygen is not perfect and may be prevented by physical processes, such as surface heat loss or entrainment of oxygen depleted deep water. Computed AOU might be higher than the true oxygen utilisation, especially for waters with high-latitude outcrop regions (Ito et al., 2004). While the processes controlling the saturation state of oxygen in the surface waters are not well constrained by available observations, we pragmatically use the AOU metric in this study. We acknowledge the limitation of the AOU approach, and we assume that all our datasets, model outputs as well as observations, are affected similarly.

A more detailed analysis of the sensitivity of the simulated undersaturation in the surface waters to biogeochemical and physical model components is beyond the current study and will be presented elsewhere for a subset of the models used here.

### 3 Means and volume distributions

In WOA, total  $\text{PO}_4$  concentrations range from close to 0 to maximum values of  $4.9 \text{ mmol m}^{-3}$ , with only very small parts of the global ocean volume exhibiting concentrations greater than  $3.5 \text{ mmol m}^{-3}$  (Fig. 1). The global mean value is  $2.26 \text{ mmol m}^{-3}$ . The volume associated with concentrations lower than  $1 \text{ mmol m}^{-3}$  is also small and represents only 5 % of the ocean volume. This low concentrations range corresponds to the surface layer and the upper thermocline (Fig. 3a,b). The total oceanic phosphate pool is composed by 64 % of preformed and 36 % of regenerated  $\text{PO}_4$ .

The observed (WOA) total phosphate volume distribution is bimodal (Fig. 1). A first mode is characterised by concentrations ranging from 1 to  $1.8 \text{ mmol m}^{-3}$ , with an associated volume of 20 % of the total ocean. This first mode corresponds mainly to Atlantic

**BGD**

8, 12423–12449, 2011

## Preformed and regenerated $\text{PO}_4$ in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Preformed and regenerated PO<sub>4</sub> in OGCMs**

O. Duteil et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

waters (Fig. 3), which are characterised by low phosphate values (around  $1 \text{ mmol m}^{-3}$ ) in northern subduction regions, intermediate values ( $1.5 \text{ mmol m}^{-3}$ ) in tropical regions and large values (above  $2 \text{ mmol m}^{-3}$ ) in the Southern Atlantic. Subduction regions are characterised by a high fraction (0.75) of preformed over total phosphate, as the sinking water mass is young and thus little biological activity occurred since its last contact with the atmosphere. This fraction is much lower in the tropical regions of the Atlantic, where biological production is high and subsurface waters are older. Preformed phosphate is low in the northern high latitude region ( $0.8 \text{ mmol m}^{-3}$ ), high in the Southern Atlantic (above  $1.3 \text{ mmol m}^{-3}$ ) and intermediate in tropical waters ( $1 \text{ mmol m}^{-3}$ ), where mixing occurs between waters of northern and southern origin.

The second mode of the total PO<sub>4</sub> volume distribution (Fig. 1, WOA) ranges from 2.2 to  $3.5 \text{ mmol m}^{-3}$  and represents 70 % of the ocean volume. This range of concentrations is typical for subsurface waters of the Southern Ocean and the Pacific Ocean. Phosphate concentrations increase when water gets older, e.g., from the Southern Ocean ( $2.2 \text{ mmol m}^{-3}$ ) towards the North Pacific, where values reach  $3.0 \text{ mmol m}^{-3}$  at intermediate depths, as seen in Fig. 3. Preformed phosphate concentrations are relatively constant in the Pacific with values of 1.4 to  $1.6 \text{ mmol m}^{-3}$  (Fig. 3b), because in contrast to the Atlantic Ocean, deep waters of the Pacific originate exclusively from the Southern Ocean. In contrast, the ratio of preformed over total phosphate decreases from the Southern Ocean (0.75) northwards and reaches minimum values of 0.5 in the North Pacific, where regenerated PO<sub>4</sub> and total PO<sub>4</sub> concentrations are highest along the Pacific transect.

Global mean phosphate concentrations are similar between observations and all the studied models. Values range from  $2.17 \text{ mmol m}^{-3}$  (MPIOM-HAMOCC) to  $2.35 \text{ (MOM P2A)}$ . This similarity is not surprising, since all models were initialised with phosphate fields derived from WOA. Differences reflect to a large extent the different topographies and resolution used in the different models. In order to quantify differences in the PO<sub>4</sub> distribution between models and data, we computed a “Volume Class Error”, which is the percentage of water simulated in the wrong concentration class compared to the

WOA.

$$VCE = \left( \sum_i |V_{PO_4(i)_{model}} - V_{PO_4(i)_{WOA}}| \right) / (2 \cdot V_{ocean}) \quad (3)$$

where  $V_{PO_4(i)}$  is the volume of water for a given modeled or observed phosphate concentration class  $i$  and  $V_{ocean}$  is the total volume of the ocean.

When measured in terms of total phosphate concentration classes of  $0.1 \text{ mmol m}^{-3}$  width (Fig. 1), for our set of models the volume class error varies from 9% for the UVIC2.8 model to 32% in the MPIOM-HAMOCC model, with OPA-PISCES (19%), CCSM3-BEC (20%) and MOM P2A (21%), om1p7-BLINGv0 (31%), CSIRO (31%), displaying intermediate model-data differences (Fig. 1).

The partitioning into preformed and regenerated phosphate displays large variations among the different models. In the om1p7-BLINGv0 model, which assumes  $R_{P:-O_2} = 1/150$ , the global ratio of preformed over total is only 0.5, indicating a larger share of regenerated phosphate and hence a more intensive biological carbon pump comparing to the physical one. The oceanic phosphate field simulated by the OPA-PISCES model, on the other hand, is determined to a larger extent by circulation-driven preformed phosphate, with a global ratio of preformed-to-total phosphate of 0.68. Preformed-to-total phosphate ratios of other models lie between these two extremes (om1p7-BLINGv0: 0.5, CCSM3-BEC: 0.55; CSIRO: 0.6, MPIOM-HAMOCC: 0.62, WOA: 0.64, MOM P2A: 0.65, UVIC2.8: 0.67, OPA-PISCES: 0.68) and are all relatively close to the WOA ratio of 0.64.

The “Volume Class Error” computed for the preformed pool distribution (Fig. 2) spans from 17% for the CSIRO model to 69% for om1p7-BLINGv0 (35% MPIOM-HAMOCC, 44% UVIC2.8, 47% CCSM3-BEC, 48% OPA-PISCES, 49% MOM P2A). For all models but CSIRO, the error in preformed phosphate is greater than the error in the total phosphate. It demonstrates that the mechanisms, whether dynamical or biological, responsible for the setting of preformed phosphate concentrations are not as well represented as suggested by just the total phosphate concentrations. The good

**Preformed and regenerated  $PO_4$  in OGCMs**

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



performance of CSIRO might be due to the use of flux adjustments at ocean surface and particularly in the Southern Ocean (Matear and Hirst, 2003), leading to a more realistic winter mixed layer, convection depth and subduction of water in the Southern Ocean.

## 4 Water masses

In order to better understand what causes the differences in the  $\text{PO}_4$  distributions, we analyse the models' representations of the main water masses in the Atlantic and the Pacific Ocean.

Deep waters of the Atlantic Ocean are formed at high latitudes, both in the north (North Atlantic Deep Water: NADW) and in the Southern Ocean (Antarctic Bottom Water: AABW). In the North Atlantic interior (below 500 m and from 40 to 60° N) (Fig. 3), concentrations of total phosphate range from around 0.8 (CSIRO) to 1.5 (MOM P2A)  $\text{mmol m}^{-3}$  and concentrations of preformed phosphate from 0.4 (om1p7-BLINGv0), 0.6 (CSIRO) to 1  $\text{mmol m}^{-3}$  (MOM P2A). Some models display preformed values in the NADW substantially lower than the WOA preformed  $\text{PO}_4$  of about 0.8  $\text{mmol m}^{-3}$  (CCSM3-BEC, om1p7-BLINGv0, CSIRO).

This apparent depletion of nutrients in the high-latitude water mass formation areas could be caused by too intense biological activity or too sluggish circulation.

Simulated total  $\text{PO}_4$  concentrations in the Southern Ocean (80 to 60° S) show relatively small variations among the different models ranging from around 1.9 (MPIOM-HAMOCC) to 2.5  $\text{mmol m}^{-3}$  (MOM P2A) (Fig. 3a). However, the partitioning into preformed and regenerated phosphate varies substantially among the different models (Fig. 3b). Simulated preformed phosphate concentrations in the Southern Ocean range from around 1.1 (MPIOM-HAMOCC) to 1.8  $\text{mmol m}^{-3}$  (OPA-PISCES). For relatively simple nutrient-restoring biogeochemistry models it has been shown that low preformed Southern Ocean phosphate concentrations can be related to sluggish circulation and weak overturning (Marinov et al., 2008). Significant differences between the

**BGD**

8, 12423–12449, 2011

## Preformed and regenerated $\text{PO}_4$ in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



end member values of preformed  $\text{PO}_4$  in the different models suggest that an identical concentration of preformed  $\text{PO}_4$  is achieved through a different combination of NADW and AABW in the deep ocean, which is likely to have biogeochemical implications, e.g. on oxygen concentration.

In all of the models investigated, the simulated tongue of NADW with low preformed phosphate concentrations occupies a depth range of about 2000–3000 m and is thinner than in the World Ocean Atlas (Fig. 3b). AABW, generally rich in preformed nutrients, extends northward throughout the Atlantic in most models, whereas it is more mixed with NADW in observations in tropical regions. A tongue of relatively high phosphate concentrations is seen at intermediate depths in the tropical Central Atlantic Ocean. This feature is due both to the subduction of Antarctic Intermediate Water, characterised by high preformed nutrient values, and high remineralisation in the tropics. In some models (MPIOM-HAMOCC, MOM P2A, CSIRO) the role of remineralisation is dominant in generating this feature, indicated by a meridional increase in total, but not in preformed, phosphate concentrations.

Waters found in the Pacific Ocean originate both from the North Atlantic and from the Southern Ocean (Broecker et al., 1991).  $\text{PO}_4$  concentrations progressively increase from south to north due to remineralisation processes (Fig. 3a, WOA). Simulated total phosphate concentrations are highest in MPIOM-HAMOCC, om1p7-BLINGv0, UVIC2.8 and CSIRO, where phosphate concentrations exceed  $3 \text{ mmol m}^{-3}$  at intermediate depths (1000 to 2000 m) from  $20^\circ \text{ S}$  to  $50^\circ \text{ N}$ . These high concentrations are associated with a preformed-to-total phosphate ratio of less than 0.5 (Fig. 3b). Simulated  $\text{PO}_4$  fields are dominated by the remineralised fraction in the om1p7-BLINGv0 over most of the Pacific Ocean. OPA-PISCES and MOM-P2A do not display  $\text{PO}_4$  concentrations higher than  $3 \text{ mmol m}^{-3}$ .

In the case of OPA-PISCES, remineralisation occurs at shallow depths and thus in regions where residence times are shorter, limiting the accumulation of regenerated phosphate. This shallow remineralisation could be due to too slow sinking speed or too fast remineralisation. Inversely in MPIOM-HAMOCC, remineralisation occurs in

**BGD**

8, 12423–12449, 2011

## Preformed and regenerated $\text{PO}_4$ in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



deeper and older water, where preformed values are already high. The apparent non-conservative behavior of simulated preformed phosphate concentrations in MPIOM-HAMOCC and CCSM3 is an artefact that can be linked to high denitrification rates (see Appendix), as only aerobic respiration has been taken in account in the computation of the regenerated phosphate pool.

Having described the main differences and similarities between simulated and observed basin-scale  $\text{PO}_4$  distributions, a quantitative measure of the respective model's fit to observations can, for example, be obtained by the absolute  $L1$  misfit (Kriest et al., 2010)

$$L1_{\text{abs}} = \sum_i \sum_j \sum_k |\text{PO}_{4\text{Model}} - \text{PO}_{4\text{Obs}}| \quad (4)$$

where  $\text{PO}_{4\text{Model}}$  and  $\text{PO}_{4\text{Obs}}$  are modeled and observed (WOA) phosphate concentrations and  $i, j, k$  denote the dimensions of the spatial domain. The sums of the norm are weighted by the relative volume of the corresponding grid boxes  $\frac{V_{ijk}}{V_{\text{total}}}$ . Absolute  $L1$  norms are computed for total, preformed and regenerated  $\text{PO}_4$ , respectively.

For the model configurations studied here, total  $\text{PO}_4$  misfits as well as misfits of the preformed and remineralised components are shown in Fig. 4. Absolute total  $\text{PO}_4$  misfits are generally smallest in the Southern Ocean, higher in the Pacific Ocean (Fig. 4c) and largest in the Atlantic Ocean (Fig. 4d). The different magnitudes of the misfits are correlated with the complexity of the composition of the different water masses in the different basins. For instance, the Southern Ocean presents a relatively simple vertical water mass structure, whereas the Atlantic ocean is characterised by different deep and intermediate waters. Any vertical misplacement of this structure by the models can generate high misfits in total phosphate.

Several models display absolute misfits in total phosphate that are smaller than the individual misfits of preformed phosphate or regenerated phosphate, indicated by the symbols located to the left of the red lines in Fig. 4. This indicates some compensation of errors in preformed phosphate by errors in remineralised phosphate, e.g., by too low

**BGD**

8, 12423–12449, 2011

## Preformed and regenerated $\text{PO}_4$ in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



preformed phosphate concentrations being accompanied by too high concentrations of regenerated phosphate (and vice versa). A compensation is also suggested by the fact that models with large errors in preformed phosphate often display large errors in regenerated phosphate, as seen by the relatively close mirror symmetry of the different panels of Fig. 4 with respect to the horizontal zero line. If no compensation of errors in preformed and regenerated phosphate existed, and if total amounts of preformed and regenerated phosphate misfits were identical, the model solutions would lie on the green lines displayed in Fig. 4. This is not the case, and essentially all model solutions are located to the left of the green lines.

A compensation of errors in preformed and regenerated phosphate is largest in the Southern Ocean, where model-data misfits of total phosphate are for most models smaller than total phosphate misfits in the Pacific or Atlantic Ocean as well as in the global average. Model-data misfits in both preformed and regenerated phosphate, in contrast, tend to be higher in the Southern Ocean than in the other ocean basins. This suggests that particularly in the Southern Ocean, a relatively good fit to observed total phosphate concentrations can be reached by a very unrealistic partitioning into preformed and regenerated phosphate. Given the close correlation of changes in preformed phosphate and atmospheric CO<sub>2</sub> observed for some models (Marinov et al., 2008), a model evaluation based on preformed (regenerated) rather than total phosphate concentrations may reveal more details about the quality of the model. In particular, Southern Ocean biogeochemistry is, in many models, not as well simulated as one might think when looking at simulated total PO<sub>4</sub> distributions only.

Using the root-mean-square norm to measure model-data misfits yields very similar results (see Appendix Fig. A1). Also, a correction for denitrification impacts on the estimated regenerated phosphate (see Appendix) for the models including an explicit nitrogen cycle (MPIOM-HAMMOCC and CCSM3-BEC) does not change the results significantly as indicated by the yellow and purple symbols included in Fig. 4.

**BGD**

8, 12423–12449, 2011

## Preformed and regenerated PO<sub>4</sub> in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 5 Conclusions

Using a set of state-of-the-art coupled biogeochemistry-circulation models of the global ocean, this study has shown that a good representation of annual mean phosphate distributions in the ocean does not automatically imply that also the preformed and regenerated phosphate components are reproduced at a similar level of accuracy. A realistic representation of preformed and regenerated nutrient distributions is, however, relevant for assessing the sensitivity of marine biogeochemical cycles and marine CO<sub>2</sub> uptake to environmental changes.

The underlying notion is that preformed (regenerated) phosphate reflects the amount of phosphate returning to the ocean interior by physical (biological) processes.

The partitioning into preformed and regenerated phosphate has been based on the canonical AOU concept (Redfield et al., 1963; Weiss, 1970), which assumes that a subducted water mass is fully saturated with respect to atmospheric oxygen at its temperature and salinity values. However, in reality as well as in an ocean model, the degree to which surface saturation is reached before the water is transported away from the surface layer depends on several factors. These factors include the intensity of air-sea gas exchange (e.g. wind speed), cooling of surface water, entrainment of low-oxygen waters from deeper layers as well as in situ biological processes (respiration or production of oxygen). At high latitudes, ice-cover can further slow down air-sea exchange. Namely the processes of cooling, ice-cover and entrainment act in the same direction and lead to an undersaturation of surface waters during winter at high latitudes when and where most of the water masses inhabiting the deep ocean are formed. High latitudes are sparsely sampled during the winter season (Koeve, 2001, 2006). However, winter-time undersaturation has been observed, for example in the Wedell Sea, one prominent region of deep water formation, of up to about 20 % (Weiss et al., 1979; Gordon and Huber, 1990; Hopema et al., 1995; Russell and Dickson, 2003) and is in good agreement with a model study that applied an explicit tracer of preformed oxygen (Ito et al., 2004). Oxygen undersaturation in newly formed deep waters translates into an overestimation of oxygen use when the canonical AOU concept is applied, as

### Preformed and regenerated PO<sub>4</sub> in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



recognized from early on (Redfield et al., 1963). The published observations and model studies available so far suggest that this bias, however, is comparable in models and the real ocean.

The models we used here may differ in the degree and realism of simulated under-saturation of oxygen at high latitudes and, in consequence, the associated bias in computed AOU. However, lacking information about preformed oxygen from these models, and in particular from the real ocean, we pragmatically assume that inter-model variations in this bias are of second-order importance for data-based model evaluation. A more detailed analysis of this subject is under way for a smaller number of ocean models.

Despite the limits of the canonical AOU concept, the use of regenerated and preformed phosphate pools in global scale “skill” metrics provides a diagnostic that is more sensitive to differences in circulation and biology than metrics based on total phosphate alone.

For the models investigated here, it turned out that low (high) values of preformed tend to be compensated by high (low) values of remineralised phosphate, suggesting some corrections of deficiencies in the physics by a tuned biogeochemical model component. This indicates that models may fit observed phosphate distributions for the wrong reasons. For the models analyzed here, this is particularly the case in the Southern Ocean.

As a conclusion, we re-iterate the need of establishing a set of validation tools for biogeochemical models. In addition to the common inspection of total nutrient concentrations, we recommend considering also preformed and regenerated nutrient components. This comes at little additional cost, but can provide additional information about mechanisms responsible for the generation of biogeochemical tracer distributions. Using the concepts of preformed and regenerated nutrients, which reflect the interplay between marine biology and ocean circulation, should help to better constrain the ability of coupled biogeochemical models to reproduce the processes relevant for global biogeochemical tracer distributions.

**BGD**

8, 12423–12449, 2011

## Preformed and regenerated PO<sub>4</sub> in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### Estimated impact of denitrification on regenerated phosphate

The denitrification process has been estimated in the computation of the regenerated phosphate pool. We assume that denitrification has occurred wherever there is a nitrate deficit with respect to the Redfield equivalent of total phosphate, computed as  $\text{NO}_3 - 16\text{PO}_4$ . Remineralisation of 1 mol of phosphate consumes  $4/5 \cdot \text{ratio}(-\text{N}/\text{O}_2)$  moles of nitrate (Paulmier et al., 2009). However, the limitation of this approach is the assumption that waters with  $\text{NO}_3 - 16\text{PO}_4 > 0$  have not experienced denitrification and that all denitrification leaves a trace in negative  $\text{NO}_3 - 16\text{PO}_4$ . Applying this correction, preformed phosphate conservative properties are significantly improved in MPIOM-HAMOCC and CCSM-BEC (Figs. 4 and A2). At basin scale, model-data differences computed following the rationale of Kriest et al. (2010) display slight improvements; however, the general picture is virtually unchanged (Fig. 4).

*Acknowledgements.* We acknowledge financial support to O.D. and A.O. from the Deutsche Forschungsgemeinschaft (SFB 754), to W.K. from the German Federal Ministry of Education and Research (FKZ 03FO608A, BIOACID), to E.G from Compute Canada and to J.S. from the EU FP7 project CARBOCHANGE (Grant no. 264879). The research of D.B. was supported by the Office of Science (BER), US Department of Energy, Award DE-FG02-07ER64467. The work of E.G was supported in part by BP and Ford Motor Company through the Carbon Mitigation Initiative at Princeton University and under award NA17RJ2612 and NA08OAR4320752 from the National Oceanic and Atmospheric Administration, US Department of Commerce. The statements, findings, conclusions, and recommendations are those of the authors and do not necessarily reflect the views of the National Oceanic and Atmospheric Administration, or the US Department of Commerce.

### Preformed and regenerated $\text{PO}_4$ in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## References

- Anderson, L. and Sarmiento, J.: Redfield ratios of remineralization determined by nutrient data analysis, *Global Biogeochem. Cy.*, 8, 65–80, 1994. 12428
- Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global Biogeochem. Cy.*, 20(2), GB2017, doi:10.1029/2005GB002591, 2006. 12427
- Broecker, W., Blanton, S., Smethie Jr., W., and Ostlund, G.: Radiocarbon decay and oxygen utilization in the deep Atlantic Ocean, *Global Biogeochem. Cy.*, 5(1), 87–117, 1991. 12433
- Broecker, W. S., Peacock, S. L., Walker, S., Weiss, R., Fahrbach, E., Schroeder, M., Mikolajewicz, U., Heinze, C., Key, R., Peng, T. H., and Rubin, S.: How much deep water is formed in the Southern Ocean? *J. Geophys. Res.*, 103(C8), 15833–15843, 1998.
- Dutay, J., Bullister, J., Doney, S., Orr, J., Najjar, R., Caldeira, K., Campin, J., Drange, H., Follows, M., Gao, Y., Gruber, N., Hecht, M., Ishida, A., Joos, F., Lindsay, K., Madec, G., Maier-Reimer, E., Marshall, J., Matear, R., Monfray, P., Mouchet, A., Plattner, G., Sarmiento, J., Schlitzer, R., Slater, R., Totterdell, I., Weirig, M., Yamanaka, Y., and Yool, A.: Evaluation of ocean model ventilation with CFC-11: comparison of 13 global ocean models, *Ocean Modelling*, 4(2), 89–120, 2002.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K., Schnur, R., Strassmann, K., Weaver, A., Yoshikawa, C., and Zeng, N.: Climate-carbon cycle feedback analysis: results from the C4MIP model intercomparison, *J. Climate*, 19(14), 3337–3353, 2006. 12426
- Galbraith, E. D., Gnanadesikan, A., Dunne, J. P., and Hiscock, M. R.: Regional impacts of iron-light colimitation in a global biogeochemical model, *Biogeosciences*, 7, 1043–1064, doi:10.5194/bg-7-1043-2010, 2010. 12427
- Garcia, H. E., Locarnini, R. A., Boyer, T. P., and Antonov, J. I.: *World Ocean Atlas 2005, Vol. 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Oxygen Saturation*, edited by: Levitus, S., NOAA Atlas NESDIS 63, US Government Printing Office, Washington, DC, 342 pp., 2006a. 12428
- Garcia, H. E., Locarnini, R. A., Boyer, T. P., and Antonov, J. I.: *World Ocean Atlas 2005, Vol. 4: Nutrients (Phosphate, Nitrate, Silicate)*, edited by: Levitus, S., NOAA Atlas NESDIS 64, US Government Printing Office, Washington, DC., 396 pp., 2006b. 12428

---

**Preformed and regenerated PO<sub>4</sub> in OGCMs**

O. Duteil et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Preformed and regenerated PO<sub>4</sub> in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gnanadesikan, A., Dunne, J., Key, R., Matsumoto, K., Sarmiento, J., Slater, R., and Swathi, P.: Oceanic ventilation and biogeochemical cycling: Understanding the physical mechanisms that produce realistic distributions of tracers and productivity, *Global Biogeochem. Cy.*, 18(4), GB4010, doi:10.1029/2003GB002097, 2004. 12427

5 Gordon, A. and Huber, B.: Southern ocean mixed layer, *J. Geophys. Res.-Oceans*, C7(95), 11655–11672, 1990. 12436

Hoppema, M., Fahrbach, E., Schröder, M., Wisotzki, A., and De Baar, H. J. W.: Winter–summer differences of carbon dioxide and oxygen in the Weddell Sea surface layer, *Mar. Chem.*, 51, 177–192, 1995. 12436

10 Ito, T., and Follows, M.: Preformed phosphate, soft tissue pump and atmospheric CO<sub>2</sub>, *J. Mar. Res.*, 63(4), 813–839, 2005. 12425

Ito, T., Follows, M., and Boyle, E.: Is AOU a good measure of respiration in the oceans?, *Geophys. Res. Lett.*, 31(17), L17305, doi:10.1029/2004GL020900, 2004. 12429, 12436

Koeve, W.: Wintertime nutrients in the North Atlantic – new approaches and implications for estimates of seasonal new production, *Mar. Chem.*, 74, 245–260, 2001. 12436

15 Koeve, W.: Stoichiometry of the biological pump in the North Atlantic – constraints from climatological data, *Global Biogeochem. Cy.*, 20, GB3018, doi:10.1029/2004GB002407, 2006. 12436

Kortzinger, A., Hedges, J., and Quay, P.: Redfield ratios revisited: Removing the biasing effect of anthropogenic CO<sub>2</sub>, *Limnol. Oceanogr.*, 46, 964–970, 2001. 12428

Kriest, I., Khatiwala, S., and Oschlies, A.: Assessment of simple global marine biogeochemical models of increasing complexity, *Prog. Oceanogr.*, 86, 336–360, 2010. 12434, 12438

Krishnamurthy, A., Moore, J., Mahowald, N., Luo, C., Doney, S., Lindsay, K., and Zender, C.: Impacts of increasing anthropogenic soluble iron and nitrogen deposition on ocean biogeochemistry, *Global Biogeochem. Cy.*, 23, GB3016, doi:10.1029/2008GB003440, 2009. 12427

25 Li, Y. H., Karl, D. M., Winn, C. D., Mackenzie, F. T., and Gans, K.: Remineralization ratios in the subtropical north Pacific gyre, *Aquatic Geochem.*, 6, 65–86, 2000. 12428

Maier-Reimer, E.: Geochemical cycles in an ocean general circulation model: preindustrial tracer distributions, *Global Biogeochem. Cy.*, 7, 645–677, 1993. 12427

30 Maier-Reimer, E., Kriest, I., Segschneider, J., Wetzol, P.: The Hamburg Ocean Carbon Cycle Model HAMOCC5.1 – Technical Description Release 1.1 – Berichte zur Erdsystemforschung 14/2005, Max-Planck-Institut für Meteorologie, Hamburg, Germany, available at: <http://www.mpimet.mpg.de/wissenschaft/publikationen/berichteerdsystemforschung.html>, last ac-

cess: December 2011, 2005. 12427

Marinov, I., Gnanadesikan, A., Sarmiento, J., Toggweiler, J., Follows, M., and Mignone, B.: Impact of oceanic circulation on biological carbon storage in the ocean and atmospheric  $p\text{CO}_2$ , *Global Biogeochem. Cy.*, 22, GB3007, 2008. 12427, 12432, 12435

5 Matear, R. and Hirst, A.: Long-term changes in dissolved oxygen concentrations in the ocean caused by protracted global warming, *Global Biogeochem. Cy.*, 17(4), doi:10.1029/2002GB001997, 2003. 12427, 12432

Najjar, R., Jin, X., Louanchi, F., Aumont, O., Caldeira, K., Doney, S., Dutay, J., Follows, M., Gruber, N., Joos, F., Lindsay, K., Maier-Reimer, E., Matear, R., Matsumoto, K., Monfray, P., 10 Mouchet, A., Orr, J., Plattner, G., Sarmiento, J., Schlitzer, R., Slater, R., Weirig, M., Yamanaka, Y., and Yool, A.: Impact of circulation on export production, dissolved organic matter, and dissolved oxygen in the ocean: Results from phase II of the Ocean Carbon-cycle Model Intercomparison Project (OCMIP-2), *Global Biogeochem. Cy.*, 21(3), GB3007, doi:10.1029/2006GB002857, 2007. 12426

15 Oschlies, A., Schulz, K., Riebesell, U., and Schmittner, A.: Simulated 21st century's increase in oceanic suboxia by  $\text{CO}_2$ -enhanced biotic carbon export, *Global Biogeochem. Cy.*, 22(4), GB4008, doi:10.1029/2007GB003147, 2008. 12427

Paulmier, A., Kriest, I., and Oschlies, A.: Stoichiometries of remineralisation and denitrification in global biogeochemical ocean models, *Biogeosciences*, 6, 923–935, doi:10.5194/bg-6-923-2009, 2009. 12438

20 Redfield, A. C., Ketchum, B. H., and Richards, F. A.: The influence of organisms on the composition of seawater, in: *The Sea*, vol. 2, edited by: Hill, M. N., John Wiley and Sons, New York, 26–77, 1963. 12436, 12437

Roy, T., Bopp, L., Gehlen, M., Schneider, B., Cadule, P., Frolicher, T., Segschneider, J., Tjiputra, J., Heinze, C., and Joos, F.: Regional impacts of climate change and atmospheric  $\text{CO}_2$  25 on future ocean carbon uptake: a multimodel linear feedback analysis, *J. Climate*, 24, 2300–2318, 2011. 12426

Russell, J. L. and Dickson, A. G.: Variability in oxygen and nutrients in South Pacific Antarctic intermediate water, *Global Biogeochem. Cy.*, 17, 1033, doi:10.1029/2000GB001317, 2003. 30 12436

Schneider, B., Bopp, L., Gehlen, M., Segschneider, J., Frölicher, T. L., Cadule, P., Friedlingstein, P., Doney, S. C., Behrenfeld, M. J., and Joos, F.: Climate-induced interannual variability of marine primary and export production in three global coupled climate carbon cycle

**BGD**

8, 12423–12449, 2011

---

**Preformed and regenerated  $\text{PO}_4$  in OGCMs**

O. Duteil et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- models, *Biogeosciences*, 5, 597–614, doi:10.5194/bg-5-597-2008, 2008. 12426
- Takahashi, T., Broecker, W. S., and Langer, S.: Redfield ratio based on chemical data from isopycnal surfaces, *J. Geophys. Res.-Oceans*, C4(90), 6907–6924, 1985. 12428
- 5 Weiss, R.: The solubility of nitrogen, oxygen, and argon in water and seawater, *Deep Sea Res.*, 17, 721–756, 1970. 12428, 12436
- Weiss R. F., Ostlund, H. G., and Craig, H.: Geochemical studies of the Weddell sea, *Deep Sea Res. Pt. I*, 26(10), 1093–1120, 1979. 12436

---

**Preformed and  
regenerated PO<sub>4</sub> in  
OGCMs**

O. Duteil et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Preformed and regenerated PO<sub>4</sub> in OGCMs


O. Duteil et al.

**Table 1.** General description of the models used in this study.

Model	OPA-PISCES	MPIOM-HAMOCC	CCSM3-BEC	UVIC	om1p7-BLINGv0	CSIRO	MOM-P2A
Resolution	0.5 (eq.) to 2 × 2	0.5(NA) to 1.5 × 1.5	0.9 (eq.) to 2 × 3.6	1.8 × 3.6	0.6 (eq.) to 2 × 3	3.2 × 5.6	3.75 × 4
Levels	31	40	25	19	28	21	24
Mixed Layer	TKE	TKE	KPP	–	KPP	–	–
Lateral diffusion	Iso-GM	Iso-GM	Iso-GM	Iso-GM	Iso-GM	Iso-GM	Iso-GM
Wind stress	ERS	NCEP-NCAR	NCEP-NCAR	NCEP-NCAR	CORE	Previous coupled exp.	ECMWF
Biology Boxes	NPZD 24 2 phy., 2 zoo., NO <sub>3</sub> , PO <sub>4</sub> , Detritus, O <sub>2</sub> , Carb., Iron, Sil.	NPZD 18 1 phy.*, 1 zoo., NO <sub>3</sub> , PO <sub>4</sub> , Detritus, O <sub>2</sub> , Carb., Iron, Sil	NPZD 18 4 phy., 1 zoo., NO <sub>3</sub> , PO <sub>4</sub> , Detritus, O <sub>2</sub> , Carb., Iron, Sil.	NPZD 9 2 phy., 1 zoo., NO <sub>3</sub> , PO <sub>4</sub> , Detritus, O <sub>2</sub> , DIC, TALK	N 4 implicit PZD, PO <sub>4</sub> , DOP, O <sub>2</sub> , Iron	N 4 implicit PZD, PO <sub>4</sub> , O <sub>2</sub> , DIC, TALK	OCMIP (N) 5 implicit PZD, PO <sub>4</sub> , DOP, O <sub>2</sub> , DIC, TALK
Ref.	Aumont and Bopp (2006)	Maier-Reimer et al. (2005)	Krishnamurty et al. (2009)	Oschlies et al. (2008)	Galbraith et al. (2010)	Matear and Hirst (2003)	Gnanadesikan et al. (2004)

\* Implicitly partitioned into diatoms and coccolithophores depending on available silicate.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Fig. 1.** Total phosphate ( $\text{mmol m}^{-3}$ ) volume distribution in WOA, OPA-PISCES, MPIOM-HAMOCC, CCSM3-BEC, UVIC2.8, om1p7-BLINGv0, CSIRO, MOM P2A for bin sizes of  $0.1 \text{ mmol m}^{-3}$ . The numbers indicate the global mean fraction of preformed over total phosphate. The blue (red) bars indicate the mean fraction of preformed (regenerated) phosphate in each bin.

**Preformed and regenerated  $\text{PO}_4$  in OGCMs**

O. Duteil et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 2.** Preformed phosphate ( $\text{mmol m}^{-3}$ ) volume distribution in WOA, OPA-PISCES, MPIOM-HAMOCC, CCSM3-BEC, UVIC2.8, om1p7-BLINGv0, CSIRO, MOM P2A.

Preformed and regenerated  $\text{PO}_4$  in OGCMs

O. Duteil et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Fig. 3.** Total **(a)** and preformed **(b)** phosphate concentrations ( $\text{mmol m}^{-3}$ ) along an Atlantic meridional section ( $30^\circ \text{W}$ ) and a Pacific meridional section ( $150^\circ \text{W}$ ) in different datasets: WOA, OPA-PISCES, MPIOM-HAMOCC, CCSM3-BEC, UVIC2.8, om1p7-BLINGv0, CSIRO, MOM P2A.

## Preformed and regenerated PO<sub>4</sub> in OGCMs

O. Duteil et al.



**Fig. 4.** Absolute misfits ( $\text{mmol m}^{-3}$ ) for regenerated, preformed (ordinate) and total phosphate (abscissa) in **(a)** global, **(b)** Southern Ocean, **(c)** Pacific and **(d)** Atlantic Ocean. In a model where no compensation between preformed and regenerated pool exists, one should expect that the total phosphate misfit is the sum of the preformed and regenerated misfits. In this case, and considering that total phosphate misfits are, at each location, caused by either the preformed or regenerated pool, with equal amounts of preformed and regenerated phosphate misfits, the models should be located along the green lines. Several models present lower total misfits than the sum of preformed and regenerated misfits. This is particularly the case for models reaching the red line (misfits in preformed or regenerated are at least equal to the total misfit). Yellow/purple symbols visible for the models MPIOM-HAMMOCC and CCSM3-BEC include a correction for the estimated contribution of denitrification to the regenerated phosphate pool (see Appendix).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Preformed and regenerated $\text{PO}_4$ in OGCMs

O. Duteil et al.



**Fig. A1.** Absolute misfits ( $\text{mmol m}^{-3}$ ) for regenerated, preformed (ordinate) and total phosphate (abscissa) in **(a)** global, **(b)** Southern Ocean, **(c)** Pacific and **(d)** Atlantic Ocean. Root square of the L2 absolute misfit has been used:  $L2_{\text{abs}} = \sum_i \sum_j \sum_k (\text{PO}_{4\text{Model}} - \text{PO}_{4\text{Obs}})^2$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. A2.** Preformed phosphate concentration ( $\text{mmol m}^{-3}$ ) taking into account the impact of denitrification along an Atlantic meridional section ( $30^\circ \text{W}$ ) and a Pacific meridional section ( $150^\circ \text{W}$ ) in different datasets: WOA, OPA-PISCES, MPIOM-HAMOCC, CCSM3-BEC, UVIC2.8, om1p7-BLINGv0, CSIRO, MOM P2A.