

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Fate of mercury in tree litter during decomposition

A. K. Pokharel and D. Obrist

Desert Research Institute, Division of Atmospheric Sciences, Reno, Nevada, USA

Received: 16 February 2011 – Accepted: 26 February 2011 – Published: 15 March 2011

Correspondence to: D. Obrist (dobrist@dri.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

We performed a controlled laboratory litter incubation study to assess changes in dry mass, carbon (C) mass and concentration, mercury (Hg) mass and concentration, and stoichiometric relations between elements during decomposition. Twenty-five surface litter samples each, collected from four forest stands, were placed in incubation jars open to the atmosphere, and were harvested sequentially at 0, 3, 6, 12, and 18 months. Using a mass balance approach, we observed significant mass losses of Hg during decomposition (5 to 23% of initial mass after 18 months), which we attribute to gaseous losses of Hg to the atmosphere through a gas-permeable filter covering incubation jars. Percentage mass losses of Hg generally were less than observed dry mass and C mass losses (48% to 63% Hg loss per unit dry mass loss), although one species showed similar losses. A field control study using the same litter types exposed at the original collection locations for one year showed that field litter samples were enriched in Hg concentrations by 8 to 64% compared to samples incubated for the same time period in the laboratory, indicating strong additional sorption of Hg in the field. Solubility of Hg, assessed by exposure of Hg to water upon harvest, was very low ($< 0.22 \text{ ng Hg g}^{-1}$ dry mass) and decreased with increasing stage of decomposition for all litter types. Our results indicate large gaseous emissions, or re-emissions, of Hg originally associated with plant litter upon decomposition. Results also suggest that Hg accumulation in litter and surface layers in the field is driven mainly by sorption of Hg – such as from atmospheric deposition – with minor contributions from “internal” accumulation due to preferential loss of C over Hg. Litter types showed highly species-specific differences in Hg levels during decomposition – suggesting that emissions, retention, and sorption of Hg are dependent on litter type.

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Atmospheric deposition of mercury (Hg), a potent environmental pollutant, enters remote ecosystems primarily through atmospheric deposition (Fitzgerald et al., 1998; Mason and Sheu, 2002). Terrestrial ecosystems serve as important receptors of Hg and sources to aquatic systems (Lorey and Driscoll, 1999; Harris et al., 2007), and present large storage pools for atmospheric Hg depositions which accumulate in surface litter and soil pools (Grigal, 2003). There are several important deposition pathways for atmospheric Hg, including wet deposition resulting from uptake of reactive gaseous Hg (RGM) and precipitation scavenging of particulate Hg (PHg; Lindberg et al., 1992), plus dry deposition including direct surface deposition of gaseous elemental and reactive Hg as well as uptake of Hg by particulates (Zhang et al., 2009; Fu et al., 2010). Vegetation also plays an important role in deposition of atmospheric Hg (Lindberg, 1996; Grigal, 2003; Driscoll et al., 2007), both through uptake of atmospheric Hg to leaves and other plant tissues and consecutive input as plant detritus (e.g., litterfall deposition), plus by wash-off of Hg previously deposited on plant surfaces (i.e., throughfall deposition). A series of field investigations have shown that litterfall alone accounts for 30% to 60% of total atmospheric Hg inputs in forests (Schwesig and Matzner, 2000; St. Louis et al., 2001; Rea et al., 2002; Sheehan et al., 2006; Demers et al., 2007) and that leaf litterfall inputs often exceed direct wet deposition inputs by a factor of two or more (Iverfeldt, 1991; Lindberg, 1996; Grigal et al., 2000b; St. Louis et al., 2001). Laboratory studies confirm that the main source of Hg in above-ground biomass is from atmospheric uptake (Ericksen et al., 2003; Frescholtz et al., 2003; Millhollen et al., 2006a, b; Rutter et al., 2011). Field studies show uptake of atmospheric gaseous Hg during peak vegetation periods, particularly when leaf areas are at maximum expansion (Obrist et al., 2006; Fritsche et al., 2008a). Uptake of Hg by vegetation may occur both through stomatal processes (Ericksen et al., 2003; Stamenkovic and Gustin, 2009; Rutter et al., 2011) and non-stomatal sorption to plant surfaces (Rea et al., 2000; Stamenkovic and Gustin, 2009).

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



39° 23' N, longitude of 119° 50' W, and at an elevation of 1821 m – shows 241 mm of annual precipitation. The dominant tree species here is *Populus tremuloides*. The pine forest near Georgetown, California, is located at 38° 54' N and 120° 39' W, at an elevation of 1302 m, and with annual precipitation averaging 1660 mm. Dominant tree species include ponderosa pine (*Pinus ponderosa*) and sugar pine (*Pinus lambertiana*), with lesser contributions of white fir (*Abies concolor*), incense cedar (*Calocedrus decurrens*), Douglas fir (*Pseudotsuga menziesii*), and California black oak (*Quercus kelloggii*). The oak forest in California, dominated by blue oak (*Quercus douglasii*), is situated at 39° 15' N, and 121° 17' W, at an elevation of 193 m and with total annual precipitation of 775 mm. Other species on the site include interior live oak, (*Q. wislizenii*) and foothill pine (*Pinus sabiniana*).

All sites are considered remote locations not affected by known, specific point sources of Hg. All litter samples were collected within days to a few weeks after the occurrence of leaf litterfall in late summer and fall of 2008. Litter samples were comprised predominantly of leaves from the dominant tree species, with smaller contributions from other trees and from understory vegetation. Woody litter components were removed from samples, and the remaining leaf litter was mixed well prior to use in the laboratory and field decomposition studies.

2.2 Controlled laboratory incubation study

For the controlled laboratory incubation study, we prepared 100 glass jars of 960 ml volume (wide-mouth, clear USP Type III soda-lime glass with PTFE-lined lids). All glass jars were cleaned with chelating soap and dilute nitric acid (5%), dried, and weighed. A 2.7 cm diameter hole was drilled in the lids of the jars and covered with a Teflon® filter membrane (pore size 0.2 µm). This allowed air exchange between the jars and the atmosphere, and, at the same time, avoided deposition of dust (including particulate-bound Hg) while likely minimizing transfer and deposition of reactive gaseous Hg to jars. Each of 25 glass jars was filled with homogenized litter samples from the four

BGD

8, 2593–2627, 2011

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

5 sites, with an initial starting weight (fresh mass) of 30 g of litter (15 g for blue oak litter). Immediately after sample preparation, litter from the first five of the 25 replicate jars was harvested for determination of dry mass and initial C, N, and Hg concentrations (i.e., time: $t = 0$ month). The 20 remaining jars of each species were placed
10 in an environmentally-controlled chamber at the Desert Research Institute (EcoPODS at DRI) at a constant temperature of 25 °C throughout the experiment. Walls and the ceiling of the chamber were covered by black cloth to allow decomposition under dark conditions. Five replicate glass jars from each species were harvested after 3, 6, 12, and 18 months of incubation (i.e., 4 litter types \times 5 harvest times \times 5 replications = 100 incubation samples total). Litter samples were kept moist by additions of ultra-purified Millipore water every three weeks.

Litter was incubated for 0 (initial starting point), 3, 6, 12, and 18 months, respectively. Upon harvest, we analyzed all samples for dry mass and concentrations of C, nitrogen (N), and total Hg (Table 1). We calculated total mass of the respective elements through
15 time. Harvested litter samples also were used for determination of Hg solubility: ultra-purified Millipore water (500 g) was added to half of the harvested litter samples, and after one hour of exposure in water, 26.5 ml aliquots of solution were extracted for measurement of soluble Hg. Aliquots were filtered with 0.45 μm pore size filters (PTFE membrane; PP housing), preserved by adding 1% concentrated hydrochloric acid, and
20 stored at 5 °C until analysis.

2.3 Field study

We conducted a field component using the same litter types to assess changes of Hg, C, and N concentrations plus their stoichiometric relationships after one year of decomposition. Samples for field decomposition were packed in mash bags made
25 from nylon screen. All bags were washed prior to use with a chelating soap bath for 48 h and were tested for Hg concentrations, which averaged $15.0 \pm 7.0 \mu\text{g Hg kg}^{-1}$. Hence, mash Hg concentrations were lower than all litter concentrations observed (see results), and hence litter bags were unlikely a significant source of Hg to litter in the

field. We measured litter concentrations of C, N, and Hg prior to field exposure. Litter bags were exposed in the field for one full year through placement in the surface Oi litter horizons of the respective collection sites. Hg, C, and N concentrations and ratio were re-measured after collection to assess concentration changes. Mass balances of litter bags were not quantified in the field study due to potential losses of small litter fractions from bags during field exposure and transport.

2.4 Sample analyses for C, N, total Hg, and soluble Hg

Upon harvests, litter samples were freeze dried using a freeze dryer (Model Micro Modulyo-115, Thermo Scientific, Waltham, MA, USA) for 48 h. After freeze drying, dry weights of each litter sample contained in glass jars were determined. Half of the well-homogenized litter samples were then milled using stainless steel coffee mills, while the other portions were used for determination of soluble Hg. Milled samples were analyzed for total Hg using a Model MA-2000 Total Mercury Analyzer (Nippon Inc., Takatsuki, Japan) according to US EPA Method 7473. The analyzer was calibrated using 0.1 ppm and 0.01 ppm Hg stock solutions made from 1000 ppm HgCl₂ standard in 0.001% L-cysteine solution according to the manufacturer guidelines. National Institute of Standards and Technology (NIST) solid standard reference materials (# 1575: Pine leaves: 39.9 μg Hg kg⁻¹; and # 1515: Apple Leaves: 44.4 μg Hg kg⁻¹) were measured at the beginning of each analytical run and repeated after every six samples. When analysis of NIST standards deviated more than 5% from their values, the analyzer was recalibrated and all samples were re-run. Across all sample analyses, NIST standard samples averaged 39.7 ± 1.2 μg Hg kg⁻¹ (mean ± standard deviation; *n* = 27; 99% recovery) for pine leaves, and 44.8 ± 0.9 μg Hg kg⁻¹ (mean ± standard deviation; *n* = 41; 101% recovery) for apple leaves. All samples were analyzed in duplicates, and analyses were repeated when the coefficient of variability of samples exceeded 10%. Total C and N in litter samples were analyzed using a Leco Turspec carbon/nitrogen analyzer (LECO, St. Joseph, Michigan, USA) at the Soil Forage and Water Analysis Laboratory at Oklahoma State University (<http://www.soiltesting.okstate.edu/>), which

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



guarantees accuracy and precision of test results through daily analysis of quality control samples. All instruments used for analysis were calibrated with certified standards and maintained in accordance with specification. Using standards every 10 samples, specific quality control for C/N analysis included frequent blank test and sample checks.

Water samples for determination of soluble Hg were analyzed according to US EPA Method 1631 for Total Mercury in Water using dual stage gold pre-concentration. We used a Tekran 2600 Mercury Analyzer (Tekran Inc., Toronto, Canada). Bromine monochloride (0.5% BrCl) was added to each sample 12 h prior to analysis for sample digestion. Immediately prior to sample analysis, we added 0.25% hydroxylamine hydrochloride (HH) to samples to ensure full destruction of the free halogens. We used a 3% solution of freshly prepared stannous chloride, purged with a slow flow of ultra-high purity (UHP) argon at 20 ml min^{-1} , to reduce Hg in samples to gaseous Hg, which was separated by a liquid-gas separator and loaded on gold traps of the analyzer. After thermal desorption, Hg was analyzed using atomic fluorescence spectroscopy. The analyzer was calibrated with standards of 0.5, 5, 10, 25, and 50 ppt of Hg in solution of 0.5% BrCl, and 0.25% HH. Quality control included three calibration blanks (consisting of 0.5% BrCl and 0.25% HH reagent), Ongoing Precision and Recovery (OPR) of 5 ppt Hg after every six samples (average recovery $97 \pm 9\%$, $n = 21$), and matrix spikes of at least two samples per analytical run. Detection limit of the system was estimated at < 0.5 ppt, based on $3 \times$ standard deviation of the reagent blanks.

2.5 Statistical analyses

All numbers in text, figures, and tables (Table 1) are shown as mean ± 1 standard deviation of five replicate samples for the laboratory study, and mean ± 1 standard deviation of 15 replicate samples for the field study. Statistical analyses of laboratory samples (Table 2) were performed using analysis of variance (ANOVA) and Bonferroni post-hoc comparison tests to analyze for statistical effects of variables: *Time*, *Species*, and *Time* \times *Species* interactions. Dependent variables tested were dry mass, C mass, C concentrations, Hg Mass, Hg concentrations, Hg/C ratio, C/N ratios, and total soluble

BGD

8, 2593–2627, 2011

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



182 ng, 42 ng, and 104 ng in mixed deciduous, aspen, pine, and oak litter, respectively. These Hg mass losses accounted for 5 to 23% of the initial Hg mass present at the start of the incubation. ANOVA tests including all data showed significant effects of *Time*, *Species*, and *Time* × *Species* interactions, indicative of significant mass losses of Hg across all litter samples. Hg mass losses were species-specific, and Bonferroni post-hoc tests showed that the significant effect of *Time* was driven by significant Hg mass losses in mixed deciduous litter, aspen litter, and to a smaller degree in oak litter (at the 10% significance level). No *Time* effect, based on Bonferroni post-hoc tests, was observed for Hg mass in pine litter.

3.2 Stoichiometric relationships between Hg, C, and N

Similar to Hg concentrations, Hg/C ratios showed only minor changes during the 18 months of laboratory decomposition (Fig. 2; Table 1). Overall, ANOVA tests showed significant effects of *Time*, *Species*, and *Time* × *Species* interaction on Hg/C ratios. As with Hg concentrations, the significant increase in Hg/C ratios was driven entirely by aspen as indicated by Bonferroni post-hoc tests, while no significant *Time* effect was evident for the other species. In aspen species, Hg/C ratios increased from $106.1 \pm 3.3 \mu\text{g kg}^{-1} \text{ C}$ at the start of the experiment to $135.6 \pm 12.2 \mu\text{g kg}^{-1} \text{ C}$ at the end of the experiment.

In regards to C/N ratios, ANOVA tests showed significant effects of *Time*, *Species*, and *Time* × *Species* interaction, in support of decreases in C/N ratios across all laboratory samples. Post-hoc Benferroni tests showed that the *Time* effects were significant in all four litter types, although occurring to various degrees. Regression analyses between Hg concentrations and C/N ratios in laboratory samples (Fig. 3) showed significant increase in Hg concentrations as a function of decreasing C/N ratios in aspen and pine litter samples (significant negative slope of linear regression; $r^2 = 0.77$ and $r^2 = 0.22$, respectively), while the two other species showed no significant regression slopes.

BGD

8, 2593–2627, 2011

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.3 Field litter decomposition patterns

Paired Student *t*-tests of Hg concentrations of samples prior to and after exposure for one full year in the field showed that Hg concentration increased significantly across all species. When each species was analyzed independently, three of four showed significant Hg concentration increases after field exposure. Only in pine litter was the effect not statistically significant. Increases in concentrations during the one-year field exposure accounted for 23 $\mu\text{g kg}^{-1}$ in mixed deciduous (+67%), 15 $\mu\text{g kg}^{-1}$ in aspen (+28%), and 21 $\mu\text{g kg}^{-1}$ in oak (+27%). Compared with samples that were exposed in the laboratory for 12 months, Hg concentrations in field samples incubated for the same time period were enriched in Hg by +64% (mixed deciduous), +62% (oak), +10% (aspen), and +8% (pine) (Fig. 3; Table 1). Given that field decomposition likely proceeded at different speed than decomposition in the laboratory (see discussion), we also analyzed Hg concentrations as a function of C/N ratios (to standardize to the degree of decomposition). Figure 3 shows that for field samples of mixed deciduous and oak litter, Hg concentrations fall above the regression line between Hg concentrations and C/N ratios for laboratory samples. In fact, for these two species, field sample concentrations were enriched by 22 $\mu\text{g kg}^{-1}$ (mixed deciduous) and 24 $\mu\text{g kg}^{-1}$ (oak). In aspen and pine field concentrations also were higher than those observed in the laboratory after one year (by 6 $\mu\text{g kg}^{-1}$ and 3 $\mu\text{g kg}^{-1}$, respectively), but the data points follow a similar regression line as observed between Hg and C/N ratios in laboratory samples. As with Hg concentrations, Hg/C ratios of field samples were significantly enriched after field exposure, and individual analysis showed that the increase was significant in all species but pine.

C/N levels decreased significantly during exposure in the field in all litter types, with the strongest decrease observed in mixed deciduous, aspen, pine, and to a lesser degree in oak. C/N ratios of field samples after 12 months of exposure were significantly lower than C/N ratios observed in laboratory samples after 12 months of exposure, with field C/N ratios decreased by 23 to 42% compared to those exposed in the laboratory (Fig. 3; Table 1).

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.4 Solubility of Hg in decomposing litter

Concentrations of soluble Hg – measured during one-hour exposure of half the harvested samples to 500 ml of Hg-free water and standardized to the amount of dry mass – were highly variable across different litter types (Fig. 4). For example, the initial amount of Hg per unit of dry litter mass accounted for $0.22 \pm 0.02 \text{ ng g}^{-1}$ in oak litter, but only for $0.07 \pm 0.01 \text{ ng g}^{-1}$ in pine litter. The amount of soluble Hg decreased strongly in all species throughout the experiment (Fig. 4). After 18 months, concentrations of soluble Hg were negligible and mainly below the detection limit of the water analyzer ($\sim 0.5 \text{ ng l}^{-1}$, based on three times standard deviation of reagent blanks), except in mixed deciduous litter. Overall, ANOVA tests showed significant effects of *Time* and *Species* on soluble Hg per unit of dry litter mass (both when only samples above the detection limit were considered and when samples below the detection limit were considered as zero values; Table 2). ANOVA tests indicated no *Time* \times *Species* interaction when only samples above the detection limits were considered, but indicated species-specific responses when all samples were considered (i.e., samples below detection limit set as zero), with Post-hoc Benferroni tests showing significant *Time* effects in mixed deciduous, aspen, and oak litter, but not in pine litter (Table 2).

4 Discussion

4.1 Changes in dry mass as well as concentration and mass changes of C and N

Dry mass losses, with relatively consistent temporal decreases in all four litter types through time (Fig. 1), show that sampling replications and time of exposure (i.e., 18 months) yielded substantial and detectable mass losses during the laboratory incubation study. Rates of mass decreases were different among litter types, and aspen leaf litter showed the strongest dry mass decrease (31% after 18 months) while pine litter

BGD

8, 2593–2627, 2011

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

showed only small losses (8%). Many field studies have shown that different litter is subject to different decomposition rates. For example, studies have shown that deciduous litter decomposed faster than coniferous litter (Kaneko and Salamanca, 1999; Hall and St. Louis, 2004), while others found the opposite (Demers et al., 2007). Some studies observed inconsistent trends or time-dependent trends where initial differences between litter types disappeared during longer time (Moore et al., 1999; Sundarapandian and Swamy, 1999; Prescott et al., 2000). In general, these field studies showed higher mass losses (e.g., up to full mass loss after one year; Sundarapandian and Swamy, 1999) than we observed during our controlled laboratory study. Possible reasons for the slower decomposition rates in the laboratory study include an artificial environment without presence of underlying soil substrates, which may have reduced diversity and mass of microbial communities (Couteaux et al., 1995). Although we regularly watered samples every 2–3 weeks, the moisture conditions for mineralization (Horner et al., 1988) may have been unfavorable compared to a field environment. Associated with dry mass losses, we observed similar losses of C mass, although C mass losses were less consistent in time. C concentrations did not significantly change throughout the experiment, and remained within 97 to 99% of the original C levels after 18 months of decomposition.

Lower C/N ratios in the field control samples (Fig. 3) compared to laboratory samples after one year of decomposition suggest that litter samples experienced higher decomposition rates under field conditions than during laboratory incubation. C/N ratios reflect the degree of decomposition with high C/N ratios associated with fresh and undecomposed organic C, while lower C/N ratios are indicative of older and more decomposed fractions (Paul and Clark, 1989). In our study, the highest C/N ratios were observed in pine litter and may be linked to its correspondingly slow decomposition rate. High C/N ratios have been considered “poor” for litter quality due to low N availability to microorganisms (Webster and Benfield, 1986; Bryant et al., 1998).

4.2 Concentration and mass changes of Hg plus stoichiometric relationships to C and N

Across all litter types, litter incubated under controlled laboratory conditions exhibited significant Hg mass losses through time (Fig. 1), with losses after 18 months ranging from 5 to 23% of initial Hg mass present at the start of the experiment. We ensured that all dry mass was harvested from sample jars and was well homogenized prior to analysis; even if small fractions of dry mass were lost during harvests, any losses would have been constant throughout the experiment (i.e., including at the initial harvest at time = 0 months) and would not explain mass losses in time. Hg losses were relatively consistent and linear in time. At the level of individual species (post-hoc test after ANOVA), significant mass losses were observed in the three deciduous species but were not significant in pine litter. The lack of mass loss in pine may be associated with its correspondingly slow rate of decomposition.

Because litter samples were kept in glass jars that only allowed mass losses by air exchange through a 0.2 μm Teflon® filter membrane in the lid, we attribute the observed Hg mass losses from litter to gaseous evasion to the atmosphere, most likely in elemental form. We originally hypothesized (Hypothesis 1) that C mass loss due to mineralization (as CO_2) would result in corresponding losses of Hg associated with organic C, and our experimental results confirm this hypothesis. Percentage Hg mass losses, however, were generally smaller than observed dry mass and C losses; for example, Hg mass losses after 18 month were 12%, 15%, and 5%, compared to dry mass losses of 19%, 31%, and 8% (percentage loss of original mass). The percentage Hg mass lost per unit C loss in these three litter types was between 48% to 63%. Only in oak litter did we observe almost identical mass losses of dry mass and Hg mass (i.e., 100% Hg lost per dry mass loss). Differences among species indicate that ratios of Hg subject to evasion versus retention is species-specific, a notion also supported by observed Hg concentrations through time: ANOVA results indicate a significant *Time* and *Time* \times *Species* effect on Hg concentration, and post-hoc tests showed that the

BGD

8, 2593–2627, 2011

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

with forest litter – ranging from 48% to full loss and depending on species – may be subject to evaporation losses once litter is subject to decomposition, at least under the conditions implemented in our controlled laboratory study.

Our observations of significant Hg mass losses, and relatively constant Hg concentrations and Hg/C ratios, during decomposition contrast with observations made in the field, both from previous studies and our field control. Demers et al. (2007) observed that total litter Hg increased by 134% and 128% of its initial mass in deciduous and coniferous litter after two years of exposure in a mixed deciduous and coniferous forest, while at the same time litter dry mass decreased by 33% and 43%, respectively. Similarly, Hall and St. Louis (2004) reported that Hg mass in deciduous and coniferous litter exposed in forests increased by 147%, and 37% compared to initial Hg mass after the 798 days of decomposition in a boreal, upland forest, while dry mass declined by 57% to 46%, respectively. Hayes et al. (1998) showed no Hg mass change in spruce litter in the field during 2.5 years, despite 20 to 40% dry mass losses. Along with Hg mass increases, Hall and St. Louis (2004) observed Hg concentration increases during field exposure (151 to 474% after 798 days of field exposure), and Demers et al. (2007) observed Hg concentration increases of 109 to 127% after two years of field exposure. In our study, field control samples exposed at the respective sampling locations showed consistently higher Hg concentrations and Hg/C ratios after one year compared to levels observed prior to field exposure. Also, litter Hg concentrations were enhanced by 8 to +64% compared to samples incubated for the same time period in the laboratory (Fig. 3). We graphed Hg concentrations versus respective C/N ratios (Fig. 3) to quantify Hg concentration changes per unit change C/N ratio. By doing this, we account for potentially different decomposition rates of field and laboratory samples, and basically “standardize” Hg concentrations to the degree of decomposition. Results show that two species (mixed deciduous and oak litter) showed particularly pronounced Hg concentration enhancements in the field compared to the laboratory (as evident by data points above the linear trendline drawn for laboratory samples). Aspen litter, however – the species that already showed Hg concentration enhancement in the laboratory –

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



showed similar concentration increases in the field, while pine litter showed inconsistent and variable patterns.

The above patterns suggest pronounced Hg sorption in litter exposed in the field (Hypothesis 3), leading to 60% concentration enhancements in the field compared to litter exposed in the laboratory where atmospheric Hg inputs were minimized by 0.2 μm Teflon[®] membranes covering litter jars and addition of Hg-free water. Our results indicate that Hg sorption in the field is dependent on litter types or on location of field exposure. We did not directly measure Hg deposition at the field sites; but based on National Atmospheric Deposition Program data, wet deposition in the area of our field sites ranged between 4 to 6 $\mu\text{g Hg m}^{-2} \text{yr}^{-1}$ (NADP, 2009). Additional Hg deposition can be expected in forest sites due to throughfall deposition (Iverfeldt, 1991; St Louis et al., 2001; Lindberg, 1996) and dry deposition (Driscoll et al., 2007; Lyman et al., 2007; Selin et al., 2007). Previous litter studies have attributed increases in Hg concentrations in the field to significant sorption of Hg from the surrounding environment (e.g., Hall and St. Louis, 2004; Demers et al., 2007), although these studies have not looked at the potential for “internal” Hg accumulation due to preferential losses of C over Hg in decomposing litter. Our field and laboratory comparisons support the notion that sorption of Hg in the field is the main reason for Hg accumulation in litter but suggest that preferential release of C over Hg also leads to “internal” Hg accumulation in some species. In addition to sorption of atmospheric Hg deposition, it is possible that tissues in the field equilibrate Hg concentrations with those of the surrounding substrates: Hall and St. Louis (2004) observed that litter with initial Hg concentration of less than 30 ng g^{-1} showed increases in Hg concentration, while litter with Hg concentrations more than 30 ng g^{-1} generally showed a decrease compared to initial concentration. In our study, we did not find indication for such a “compensation” point. The notion of significant sorption of Hg in decomposing litter is in good agreement with many studies that report strong accumulation of Hg in litter and surface soil layers in the field (Mierle, 1990; Aastrup et al., 1991; Nater and Grigal, 1992; Munthe et al., 1998; Grigal, 2003; Friedli et al., 2007; Obrist et al., 2011).

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.3 Solubility of Hg in litter during decomposition

When harvested litter was submerged in water for quantification of Hg solubility, the amount of soluble Hg (filtered by 0.45 μg membrane) per litter dry mass strongly decreased in time. At the start of the experiment, soluble Hg averaged between 0.07 and 0.22 ng Hg g^{-1} dry mass, depending on species. This amount, although highly variable, decreased to close to the detection limit (about 0.05 ng Hg g^{-1} dry mass) to 0.13 ng Hg g^{-1} dry mass after 12 months. After 18 months, most litter samples showed no soluble Hg concentrations above the detection limit anymore. These results indicate that the fraction of Hg available for mobilization as soluble Hg decreases with increasing time of decomposition.

Comparisons of these results with other studies are challenging since we are not aware of studies that also addressed Hg solubility of variously decomposed litter substrates. In a controlled litter incubation study, Tsui et al. (2008) submerged forest litter samples in different stream water for 66 days and observed that soluble Hg (both total Hg and methylated Hg) in water samples increased significantly with time of exposure. The Tsui et al. study demonstrated that litter in streams and lakes decomposing under hypoxic conditions contribute Hg to these water bodies and showed that the contributions are highly dependent on stream water chemistry. The experimental conditions between Tsui et al.'s study and ours are not comparable – given that we decomposed litter under aerobic conditions and assessed Hg solubility at the time of harvests, while Tsui et al.'s study assessed the cumulative amount of soluble Hg released under anaerobic decomposition. In the field, Hall and St. Louis (2004) reported Hg concentration increases in tree litter submerged in flooded reservoirs whereby litter seemed to reach equilibrium conditions with the surrounding water. In sphagnum tissues, Heyes et al. (1998) and Hall and St. Louis (2004) reported Hg mass losses from litter submerged in wetland and flooded reservoirs, respectively.

Possible reasons for declines in soluble Hg release with increasing stage of decomposition include gaseous losses of Hg (see above) that may have depleted Hg that was

BGD

8, 2593–2627, 2011

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

initially present in soluble form. Other reasons include the possibility that decomposition may have led to increased surface area as well as increased presence of humic and fulvic substances, which may result in more efficient sorption of Hg in decomposing litter tissues. Allard et al. (1991) showed that humic and fulvic acids have strong sorption capacity for Hg, thereby reducing its mobility; and humic matter plays a major role in controlling solubility and mobilization of Hg in freshwater systems (Mierle and Ingram, 1991; Weber, 1993). It is unclear if soluble Hg extracted from litter was of external or internal origin (i.e., surface wash-off or internal leaching; Rea et al., 2000). Ericksen et al. (2003) showed that about 1.5 to 3% of total Hg present in fresh leaves is water soluble and easily rinsed from the leaves. In our study, the fraction of soluble Hg, as a percentage of total Hg, only ranged between 0.2 and 0.6% at the beginning of the experiment and further declined with time. Reasons for reduced solubility in our study may include tissue types, leaf age, and stage of decomposition (e.g., fresh leaves versus surface Oi litter), and that litter was subject to precipitation wash-off prior to collection in the field while leaves used by Ericksen et al. (2003) were grown in environmentally controlled growth chambers without experiencing rain.

5 Summary

The results of our laboratory incubation study showing substantial Hg mass losses in decomposing litter are important in regard to emissions of Hg from terrestrial ecosystems. Such releases back into the atmosphere can be considered re-emission fluxes of Hg that was originally deposited by leaf litterfall. The issue of re-emission processes is of high interest since thousands of tons of Hg are contained and stored in surface litter and surface soils (e.g., Grigal et al., 2003; Obrist, 2007), including substantial amounts of past anthropogenic emissions due to increased atmospheric Hg pollution during the last 150 years (Fitzgerald et al., 1998; Schuster et al., 2002). Substantial re-emission of Hg contained in terrestrial storage pools, particularly during mineralization processes, could have important consequences for global environmental change

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

due to predicated losses in C storage (Grigal, 2003; Obrist, 2007). Recent studies using stable isotopes (Hintelmann et al., 2002; Ericksen et al., 2005; Harris et al., 2007; Graydon et al., 2009) showed that most of newly deposited Hg, when applied via wet deposition, is retained in watersheds and soils and that atmospheric re-emission may account for less than 10% in the first year after application (Ericksen et al., 2005; Harris et al., 2007). Our study points toward potentially stronger re-emission, up to 23% of Hg within 18 months, of Hg that is originally associated with plant litter. Our study, however, cannot address the degree to which re-emission will reach the atmosphere in the field. For example, we found that in the field, gaseous Hg/C ratios in emissions only accounted for a few percent of the Hg/C ratios of the soil pool (Obrist et al., 2009), while in the laboratory, we observed up to equivalent loss of Hg and C upon decomposition in litter. Our results also indicate that litter types show highly species-specific behaviors. Evasion losses of Hg were variable by species (ranging from 5 to 23%), and so was sorption of Hg litter observed in the field, indicating that different plant tissues show inherently different behaviors regarding sorption, retention, and re-emission of Hg.

Acknowledgements. We would like to thank Rebecca Wenk, Andrew Richardson, Scott Ollinger, Michelle Day, and Rob York for help with litter collection and placement of litter bags in the field. Thanks also to Johnny Dagget, Bill Coulomb, Russ Bergin, So Lee, and Xavier Fäin for the experimental set-up and analytical support. We appreciate editorial comments by Roger Kreidberg. Funding was provided by the US Environmental Protection Agency through a Science-To-Achieve-Results grant (# RD833378010) and through DRI Internal Project Assignment funding.

References

- Aastrup, M., Johnson, J., Bringmark, E., Bringmår, I., and Iverfeldt, A.: Occurrence and transport of mercury within a small catchment area, *Water Air Soil Poll.*, 56, 155–167, 1991.
- Allard, B. and Arsenie, I.: Abiotic reduction of mercury by humic substances in aquatic system – an important process for the mercury cycle, *Water Soil Air Poll.*, 56, 457–464, 1991.

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bryant, D. M., Holland, E. A., Seastedt, T. R., and Walker, M. D.: Analysis of litter decomposition in an alpine tundra, *Can. J. Bot.*, 76, 1295–1304, 1998.
- Carpi, A. and Lindberg, S. E.: Sunlight-mediated emission of elemental mercury from soil amended with municipal sewage sludge, *Environ. Sci. Technol.*, 31, 2085–2091, 1997.
- 5 Choi, H.-D., Sharac, T. J., and Holsen, T. M.: Mercury deposition in the Adirondacks: a comparison between precipitation and throughfall, *Atmos. Environ.*, 42, 1818–1827, 2007.
- Couteaux, M. M., Bottner, P., and Berg, B.: Litter decomposition, climate and litter quality, *Trends Ecol. Evol.*, 12, 63–66, 1995.
- Demers, J. D., Triscoll, T. C., Timothy, J. F., and Joseph, B. Y.: Mercury cycling in litter and soil in different forest types in the Adirondack region, New York, *Ecol. Appl.*, 17, 1341–1351, 2007.
- 10 Driscoll, C. T., Blette, V., Yan, C., Schofield, C. L., Munson, R., and Holsapple, J.: The role of dissolved organic carbon in the chemistry and bioavailability of mercury in remote Adirondack lakes, *Water Air Soil Poll.*, 80, 499–508, 1995.
- 15 Driscoll, C. T., Han, Y.-J., Chen, C. Y., Evers, D. C., Lambert, K. F., Holsen, T. M., Kamman, N. C., and Munson, R. K.: Mercury contamination in forest and freshwater ecosystems in the Northeastern United States, *Bioscience*, 57, 17–28, 2007.
- Engle, M., Gustin, M. S., and Zhang, H.: Quantifying natural source mercury emissions from the Ivanhoe mining district, North-Central Nevada, USA, *Atmos. Environ.*, 35, 3987–3997, 2001.
- 20 Ericksen, J. and Gustin, M. S.: Foliar exchange of mercury as a function of soil and air mercury concentrations, *Sci. Total Environ.*, 324, 271–279, 2004.
- Ericksen, J. A., Gustin, M. S., Schorran, D. E., Johnson, D. W., Lindberg, S. E., and Coleman, J. S.: Accumulation of atmospheric mercury in forest foliage, *Atmos. Environ.*, 37, 1613–1622, 2003.
- 25 Ericksen, J. A., Gustin, M. S., Lindberg, S. E., Olund, S. D., and Krabbenhoft, D. P.: Assessing the potential for re-emission of mercury deposited in precipitation from arid soils using a stable isotope, *Environ. Sci. Technol.*, 39, 8001–8007, 2005.
- Fitzgerald, W. F., Engstrom, D. R., Mason, R. P., and Nater, E. A.: The case for atmospheric mercury contamination in remote areas, *Environ. Sci. Technol.*, 32, 1–7, 1998.
- 30 Frescholtz, T. F., Gustin, M. S., Schorran, D. E., and Fernandez, G. C. J.: Environmental toxicology – assessing the source of mercury in foliar tissue of quaking aspen, *Environ. Toxicol. Chem.*, 22, 2114–2119, 2003.

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Friedli, H. R., Radke, L. F., Payne, N. J., McRae, D. J., Lynham, T. J., and Blake, T. W.: Mercury in vegetation and organic soil at an upland boreal forest site in Prince Albert National Park, Saskatchewan, Canada, *J. Geophys. Res.*, 112, G01004, doi:10.1029/2005JG000061, 2007.
- 5 Fritsche, J., Obrist, D., Zeeman, M. J., Conen, F., Wugster, W., and Alewell, C.: Elemental mercury fluxes over a sub-alpine grassland with two micrometeorological methods, *Atmos. Environ.*, 42, 2922–2933, 2008a.
- Fritsche, J., Obrist, D., and Alewell, C.: Evidence of microbial control of Hg-0 emissions from uncontaminated terrestrial soils, *Journal of Plant Nutrition and Soil Science*, 171, (2), 200–209, 2008.
- 10 Fu, X., Feng, X., Zhu, W., Rothenberg, S., Yao, H., and Zhang, H.: Elevated atmospheric deposition and dynamics of mercury in a remote upland forest of Southwestern China, *Environ. Pollut.*, 158, 2324–2333, 2010.
- Graydon, J. A., St. Louis, V. L., Hintelmann, H., Lindberg, S. E., Sandilands, K. A., Rudd, J. W. M., Kelly, C. A., Tate, M. T., Krabbenhoft, D. P., and Lehnher, I.: Investigation of uptake and retention of atmospheric Hg (II) by boreal forest plants using stable Hg isotopes, *Environ. Sci. Technol.*, 43, 4960–4966, 2009.
- 15 Grigal, D. F.: Inputs and outputs of mercury from terrestrial watersheds: a review, *Environ. Rev.*, 10, 1–39, 2002.
- 20 Grigal, D. F.: Mercury sequestration in forests and peatlands: a review, *J. Environ. Qual.*, 32, 393–405, 2003.
- Grigal, D. F., Kolka, R. K., Fleck, J. A., and Nater, E. A.: Mercury budget of an upland-peatland watershed, *Biogeochemistry*, 50, 95–109, 2000b.
- Gustin, M. S., Taylor, G. E., and Maxey, R. A.: Effect of temperature and air movement on the flux of elemental mercury from substrate to the atmosphere, *J. Geophys. Res.-Atmos.*, 102, 3891–3898, 1997.
- 25 Hall, B. D. and St. Louis, V. L.: Methylmercury and total mercury in plant litter decomposing in upland forests and flooded landscapes, *Environ. Sci. Technol.*, 38, 5010–5021, 2004.
- Hanson, P. J., Lindberg, S. E., Tabberer, T. A., Owens, J. G., and Kim, K. H.: Foliar exchange of mercury vapor: evidence for a compensation point, *Water Air Soil Poll.*, 80, 373–82, 1995.
- 30 Harris, R. C., Rudd, J. W. M., Amyot, M., Babiarz, C., Beaty, K. G., Blanchfield, P. J., Boddaly, R. A., Branfireun, B. A., Gilmour, C. C., Graydon, J. A., Heyes, A., Hintelmann, H., Hurley, J. P., Kelly, C. A., Krabbenhoft, D. P., Lindberg, S. E., Mason, R. P., Paterson, M. J.,

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Podemski, C. L., Robinson, A., Sandilands, K. A., Southworth, G. R., St. Louis, V. L., and Tate, M. T.: Whole-ecosystem study shows rapid fish-mercury response to changes in mercury deposition, *P. Natl. Acad. Sci. USA*, 104, 16586–16591, 2007.

Heyes, A., Moore, T. R., and Rudd, J. W. M.: Mercury and methylmercury in decomposing vegetation of a pristine and impounded wetland, *J. Environ. Qual.*, 27, 591–599, 1998.

Hintelmann, H., Harris, R., Heyes, A., Hurlley, J., Kelly, C., Krabbenhoft, D., Lindberg, S., Rudd, J., Scott, K., and St. Louis, V. L.: Reactivity and mobility of new and old mercury deposition in a boreal forest ecosystem during the first year of the METAALICUS study, *Environ. Sci. Technol.*, 36, 5034–5040, 2002.

Horner, D. J., Gosz, J. R., and Cates, R. G.: The role of carbon-based plant secondary metabolites in decomposition in terrestrial ecosystems, *Am. Nat.*, 132, 869–883, 1988.

Iverfeldt, A.: Mercury in forest canopy throughfall water and its relation to atmospheric deposition, *Water Air Soil Poll.*, 56, 553–564, 1991.

Johansson, K. and Iverfeldt, Å.: The relation between mercury content in soil and transport of mercury from small catchments in Sweden, in: *Mercury Pollution: Integration and Synthesis*, edited by: Watras, C. J. and Huckabee, J. W., Lewis Publishers, Monterey, California, USA, 323–328, 1994.

Joslin, J. D.: Regional differences in mercury levels in aquatic ecosystems: a discussion of possible causal factors with implications for the Tennessee River system and the Northern Hemisphere, *Environ. Manage.*, 18, 559–567, 1994.

Kaneko, N. and Salamanca, E.: Mixed leaf litter effects on decomposition rates and soil microarthropod communities in an oak-pine stand in Japan, *Ecol. Res.*, 14, 131–138, 1999.

Kolka, R. K., Grigal, D. F., Verry, E. S., and Nater, E. A.: Mercury and organic carbon relationships in streams draining forested upland/peatland watersheds, *J. Environ. Qual.*, 28, 766–775, 1999b.

Kuiken, T., Zhang, H., Gustin, M. S., and Lindberg, S.: Mercury emission from terrestrial background surfaced in the Eastern USA, part I: Air/surface exchange of mercury within a South-eastern deciduous forest (Tennessee) over one year, *Appl. Geochem.*, 23, 345–55, 2008a.

Kuiken, T., Gustin, M. S., Zhang, H., Lindberg, S. E., and Sedinge, B.: Mercury emission from terrestrial background surfaces in the Eastern USA, Part II: Air/surface exchange of mercury within forests from South Carolina to New England, *Appl. Geochem.*, 23, 356–368, 2008b.

Lee, Y. H., Bishop, K. H., Munthe, J., Iverfeldt, Å., Verta, M., Parkman, H., and Hultberg, H.: An examination of current Hg deposition and export in Fenno-Scandinavian catchments,

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Biogeochemistry, 40, 125–135, 1998.

Lindberg, S. E.: Forest and the global biogeochemical cycle of mercury: the importance of understanding air/vegetation exchange processes, in: *Global and Regional Mercury Cycles: Sources, Fluxes and Mass Balances*, NATOASI Series 21, edited by: Baeyens, W., Ebinghaus, R., and Vasilie, O., Kluwer Academic Publishers, Dordrecht, The Netherlands, 359–380, 1996.

Lindberg, S. E. and Harris, R. C.: Mercury enrichment in estuarine plant detritus, *Mar. Pollut. Bull.*, 5, 93–95, 1974.

Lindberg, S. E., Hanson, P. J., Meyers, T. P., and Kim, K. H.: Atmosphere-surface exchange of mercury in a forest: results of modeling and gradient approaches, *J. Geophys. Res.-Atmos.*, 97, 2519–2528, 1992.

Lorey, P. and Driscoll, C. T.: Historical trends of mercury deposition in Adirondack lakes, *Environ. Sci. Technol.*, 33, 718–722, 1999.

Lyman, S. N., Gustin, M. S., Prestbo, E. M., and Marsik, F. J.: Estimation of dry deposition of atmospheric mercury in Nevada by direct and indirect methods, *Environ. Sci. Technol.*, 41, 1970–1976, 2007.

Mason, R. P. and Sheu, G. R.: Role of the ocean in the global mercury cycle, *Global Biogeochem. Cy.*, 16, 1093, 2002.

Meili, M.: The coupling of mercury and organic matter in the biogeochemical cycle – towards a mechanistic model for the boreal forest zone, *Water Air Soil Poll.*, 56, 333–347, 1991.

Millhollen, A. G., Gustin, M. S., and Obrist, D.: Mercury accumulation in grass and forb species as a function of atmospheric carbon dioxide concentrations and mercury exposures in air and soil, *Chemosphere*, 65, 889–897, 2006a.

Millhollen, A. G., Obrist, D., and Gustin, M. S.: Foliar mercury accumulation and exchange for three tree species, *Environ. Sci. Technol.*, 40, 6001–6006, 2006b.

Mierle, G.: Aqueous inputs of mercury to Precambrian shield lakes in Ontario, *Environ. Toxicol. Chem.*, 9, 843–851, 1990.

Mierle, G. and Ingram, R.: The role of humic substances in the mobilization of mercury from watersheds, *Water Air Soil Poll.*, 56, 349–357, 1991.

Moore, T. R., Trofymow, J. W., Taylor, B., Prescott, C., Camire, C., Duschene, L., Fyles, J., Kozak, L., Kranavetter, M., Morrison, I., Siltanesn, M., Smith, S., Titus, B., Visser, S., Wein, R., and Zoltai, S.: Litter decomposition rates in Canadian forests, *Glob. Change Biol.*, 5, 75–82, 1999.

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Munthe, J., Lee, Y. H., Hultberg, H., and Iverfeldt, Å., Borg, G. C., and Andersson, I.: Cycling of mercury and methylmercury in the Gårdsjön catchments, in: *Experimental Reversal of Acid Rain Effects, The Gårdsjön Roof Project*, edited by: Hultberg, H. and Skeffington, R., John Wiley & Sons, New York, USA, 261–276, 1998.

5 National Atmospheric Deposition Program: Mercury Deposition Program (NADP), available at: <http://nadp.sws.uiuc.edu/mdn/>, last access: January 2011, 2009.

Nater, E. A. and Grigal, D. F.: Regional trends in mercury distribution across the Great Lakes states, North Central USA, *Nature*, 358, 139–141, 1992.

Obrist, D.: Atmospheric mercury pollution due to losses of carbon pools?, *Biogeochemistry*, 85, 119–123, 2007.

10 Obrist, D., Conen, F., Vogt, R., Siegwolf, R., and Alewell, C.: Estimation of Hg^0 exchange between ecosystems and the atmosphere using ^{222}Rn and Hg^0 concentration changes in the stable nocturnal boundary layer, *Atmos. Environ.*, 40, 856–866, 2006.

Obrist, D., Johnson, D. W., and Lindberg, S. E.: Mercury concentrations and pools in four Sierra Nevada forest sites, and relationships to organic carbon and nitrogen, *Biogeosciences*, 6, 765–777, doi:10.5194/bg-6-765-2009, 2009.

Obrist, D., Fäin, X., and Berger, C.: Relationships between gaseous elemental mercury emissions and CO_2 respiration rates in terrestrial soils under aerobic and anaerobic conditions, *Sci. Total Environ.*, 408, 1691–1700, 2010.

20 Obrist, D., Johnson, D. W., Lindberg, S. W., Luo, Y., Hararuk, O., Bracho, R., Battles, J. J., Dail, D. B., Edmonds, R. L., Monson, R. K., Ollinger, S. V., Pallardy, S. G., Pregitzer, K. S., and Todd, D. E.: Mercury distribution across 14 US forests, Part I: Spatial patterns of total Hg concentrations in biomass, litter, and soils, *Environ. Sci. Technol.*, submitted, 2011.

Paul, E. A. and Clark, F. E.: *Soil Microbiology and Biochemistry*, Academic Press, San Diego, California, USA, ISBN 0125468059, 1989.

25 Poissant, L. and Casimir, A.: Water-air and soil-air exchange rate of total gaseous mercury measured at background sites, *Atmos. Environ.*, 32, 883–893, 1998.

Prescott, C. E., Zabek, L. M., Staley, C. L., and Kabzems, R.: Decomposition of broad leaf and needle litter in forests of British Columbia: influences of litter type, forest type, and litter mixtures, *Can. J. Forest Res.*, 30, 1742–1750, 2000.

30 Rea, A. W., Keeler, G. J., and Scherbatskoy, T.: The deposition of mercury in throughfall and litterfall in the Lake Champlain watershed: a short-term study, *Atmos. Environ.*, 30, 3257–3263, 1996.

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Rea, A. W., Lindberg, S. E., and Keeler, G. J.: Assessment of dry deposition and foliar leaching of mercury and selected trace elements based on washed foliar and surrogate surfaces, *Environ. Sci. Technol.*, 34, 2418–2425, 2000.
- 5 Rea, A. W., Lindberg, S. E., Scherbatskoy, T., and Keeler, G. J.: Mercury accumulation in foliage over time in two northern mixed hardwood forests, *Water Air Soil Poll.*, 133, 49–67, 2002.
- Rogers, R. D. and McFarlane, J. C.: Factors influencing the volatilization of mercury from soil, *J. Environ. Qual.*, 8, 255–260, 1979.
- Rutter, A. P., Schauer, J. J., Shafer, M. M., Creswell, J. E., Olson, M. R., Robinson, M., Collins, R. M., Parman, A. M., Katzman, T. L., and Mallek, J. L.: Dry deposition of gaseous
10 elemental mercury to plants and soils using mercury stable isotopes in a controlled environment, *Atmos. Environ.*, 45, 848–855, 2011.
- Schuster, P. F., Krabbenhoft, D. P., and Naftz, D. L.: Atmospheric mercury deposition during the last 270 years: a glacial ice core record of natural and anthropogenic sources, *Environ. Sci. Technol.*, 36, 2303–2310, 2002.
- 15 Schwesig, D. and Matzner, E.: Pools and fluxes of mercury and methylmercury in two forested catchments in Germany, *Sci. Total Environ.*, 260, 213–223, 2000.
- Selin, N. E., Jacob, D. J., Park, R. J., Yantosca, R. M., Strode, S., Jaegle, L., and Jaffe, D.: Chemical cycling and deposition of atmospheric mercury: global constraints from observations, *J. Geophys. Res.*, 112, D02308, doi:10.1029/2006JD007450, 2007.
- 20 Sheehan, K. D., Fernandez, I. J., Kahl, J. S., and Amirbahman, A.: Litterfall mercury in two forested watersheds at Acadia National Park, Maine, USA, *Water Air Soil Poll.*, 170, 249–265, 2006.
- Stamenkovic, J. and Gustin, M. S.: Non-stomatal versus stomatal uptake of atmospheric mercury, *Environ. Sci. Technol.*, 43, 1367–1372, 2009.
- 25 Stamenkovic, J., Gustin, M. S., Arnone, J. A., Johnson, D. W., Larsen, J. D., and Verburg, P. S. J.: Atmospheric mercury exchange with a tallgrass prairie ecosystem housed in mesocosms, *Sci. Total Environ.*, 406, 227–238, 2008.
- St. Louis, V., Rudd, J. W. M., Kelly, C. A., Hall, B. D., Rolfhus, K. R., Scott, K. J., Lindberg, S. E., and Dong, W.: Importance of the forest canopy to fluxes of methyl mercury and total mercury to boreal ecosystems, *Environ. Sci. Technol.*, 35, 3089–3098, 2001.
- 30 Sundarapandian, S. M. and Swamy, P. S.: Litter production and leaf litter decomposition of selected tree species in tropical forest at Kodayar in the Western Ghats, India, *Forest Ecol. Manag.*, 123, 231–244, 1999.

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Tsui, M. T. K., Finlay, J. C., and Nater, E. A.: Effects of stream water chemistry and tree species on release and methylation of mercury during litter decomposition, *Environ. Sci. Technol.*, 42, 8692–8697, 2008.

Weber, J. H.: Review of possible paths for abiotic methylation of mercury(II) in the aquatic environment, *Chemosphere*, 26, 2063–2077, 1993.

Webster, J. R. and Benfield, E. F.: Vascular plant breakdown in freshwater ecosystems, *Annu. Rev. Ecol. Syst.*, 17, 567–594, 1986.

Wickland, K., Krabenhof, D., and Olund, S.: Evidence for a link between soil respiration and mercury emissions from organic soils, Eighth International Conference on Mercury as a Global Pollutant, Madison, Wisconsin, 6–11 August 2006.

Xin, M. and Gustin, M. S.: Gaseous elemental mercury exchange with low mercury containing soils: investigation of controlling factors, *Appl. Geochem.*, 22, 1451–66, 2007.

Zhang, H., Lindberg, S. E., Marsik, F. J., and Keeler, G. J.: Mercury air/surface exchange kinetics of background soils of the Tahquamenon River watershed in the Michigan Upper Peninsula, *Water Air Soil Poll.*, 126, 151–69, 2001.

Zhang, L., Wright, L. P., and Blanchard, P.: A review of current knowledge concerning dry deposition of atmospheric mercury, *Atmos. Environ.*, 43, 5853–5864, 2009.

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Table 1. Dry mass, C concentration, C mass, Hg concentration, Hg mass, N concentration, N mass, C/N ratio, Hg/C ratio, and Hg/N ratio of laboratory samples.

Time (months)	Dry mass (g)				C concentration (%)				C mass (g)			
	Mixed deciduous	Aspen	Pine	Oak	Mixed deciduous	Aspen	Pine	Oak	Mixed deciduous	Aspen	Pine	Oak
0	24.1 ± 0.4	23.9 ± 0.3	21.3 ± 0.6	11.1 ± 1.2	48.56 ± 0.67	47.28 ± 0.53	50.80 ± 0.48	47.90 ± 0.14	11.70 ± 0.30	11.29 ± 0.99	10.80 ± 0.29	5.33 ± 0.58
3	22.2 ± 0.5	22.5 ± 0.9	20.9 ± 0.4	10.6 ± 1.1	49.72 ± 0.50	45.54 ± 0.81	50.65 ± 0.43	48.16 ± 0.55	11.01 ± 0.13	10.24 ± 0.40	10.59 ± 0.28	5.11 ± 0.55
6	21.8 ± 0.5	21.3 ± 0.4	20.6 ± 0.4	10.4 ± 1.2	50.26 ± 0.40	47.66 ± 0.23	44.98 ± 0.52	42.50 ± 0.58	10.94 ± 0.24	10.15 ± 0.20	9.25 ± 0.20	4.41 ± 0.54
12	19.9 ± 0.5	16.8 ± 1.0	19.9 ± 0.5	9.5 ± 0.6	47.74 ± 0.40	45.70 ± 0.31	48.88 ± 0.30	46.22 ± 0.42	9.54 ± 0.20	7.70 ± 0.52	9.70 ± 0.24	4.38 ± 0.29
18	19.4 ± 0.3	16.4 ± 0.9	19.7 ± 0.6	8.6 ± 0.7	49.58 ± 0.43	45.87 ± 0.61	51.27 ± 0.48	47.40 ± 0.79	9.64 ± 0.20	7.52 ± 0.47	10.08 ± 0.25	4.09 ± 0.34

Time (months)	Hg Concentration ($\mu\text{g kg}^{-1}$)				Hg mass (ng)				N Concentration (%)			
	Mixed deciduous	Aspen	Pine	Oak	Mixed deciduous	Aspen	Pine	Oak	Mixed deciduous	Aspen	Pine	Oak
0	32.43 ± 4.17	50.16 ± 1.60	39.28 ± 2.52	40.67 ± 1.53	782 ± 97	1198 ± 30	834 ± 57	452 ± 48	0.56 ± 0.05	0.67 ± 0.06	0.45 ± 0.03	0.81 ± 0.05
3	35.52 ± 1.67	54.60 ± 1.51	37.85 ± 2.97	40.81 ± 1.21	788 ± 53	1228 ± 52	792 ± 61	430 ± 37	0.64 ± 0.03	0.74 ± 0.04	0.39 ± 0.03	0.83 ± 0.09
6	35.00 ± 3.39	55.08 ± 2.89	38.36 ± 2.27	36.08 ± 1.84	764 ± 71	1174 ± 44	790 ± 46	378 ± 55	0.5 ± 0.04	0.77 ± 0.07	0.41 ± 0.05	0.80 ± 0.05
12	34.94 ± 2.14	61.82 ± 3.08	37.87 ± 1.83	38.99 ± 2.67	700 ± 62	1036 ± 52	754 ± 42	372 ± 24	0.65 ± 0.06	1.00 ± 0.09	0.43 ± 0.02	0.86 ± 0.06
18	35.53 ± 1.55	62.13 ± 4.95	40.30 ± 3.84	40.25 ± 3.82	690 ± 26	1016 ± 31	792 ± 84	348 ± 36	1.07 ± 0.03	1.40 ± 0.12	0.83 ± 0.05	1.35 ± 0.19

Time (months)	N mass (g)				C/N ratio				Hg/C ratio ($\mu\text{g kg}^{-1}\text{C}$)			
	Mixed deciduous	Aspen	Pine	Oak	Mixed deciduous	Aspen	Pine	Oak	Mixed deciduous	Aspen	Pine	Oak
0	0.134 ± 0.011	0.161 ± 0.013	0.096 ± 0.007	0.090 ± 0.012	87.76 ± 8.68	70.55 ± 5.66	113.25 ± 6.72	59.21 ± 3.43	66.8 ± 8.9	106.1 ± 3.3	77.4 ± 5.6	84.9 ± 3.3
3	0.142 ± 0.008	0.166 ± 0.013	0.081 ± 0.006	0.088 ± 0.010	77.61 ± 4.38	62.09 ± 4.50	130.81 ± 8.76	58.69 ± 6.93	71.5 ± 4.0	119.9 ± 5.0	74.8 ± 6.3	84.8 ± 2.6
6	0.123 ± 0.010	0.164 ± 0.012	0.084 ± 0.009	0.083 ± 0.010	89.40 ± 7.71	62.37 ± 5.18	111.22 ± 13.75	53.03 ± 4.14	69.7 ± 7.3	115.6 ± 6.3	85.3 ± 5.0	84.9 ± 3.9
12	0.130 ± 0.015	0.167 ± 0.012	0.085 ± 0.004	0.081 ± 0.005	73.82 ± 6.54	45.94 ± 3.89	114.36 ± 5.31	54.05 ± 3.48	73.2 ± 5.0	135.3 ± 7.4	77.5 ± 3.7	84.4 ± 5.2
18	0.210 ± 0.007	0.227 ± 0.011	0.164 ± 0.011	0.116 ± 0.020	46.23 ± 1.40	33.20 ± 3.00	61.77 ± 3.28	35.86 ± 5.81	71.7 ± 3.5	135.6 ± 12.2	78.6 ± 7.5	84.9 ± 9.0

Time (months)	Hg/N ratio ($\mu\text{g kg}^{-1}\text{N}$)			
	Mixed deciduous	Aspen	Pine	Oak
0	5808.8 ± 378.4	7480.8 ± 591.9	8753.8 ± 700.6	5027.0 ± 334.7
3	5541.9 ± 340.2	7434.3 ± 364.1	9748.9 ± 582.3	4963.0 ± 463.7
6	6195.4 ± 383.3	7187.9 ± 167.8	9433.8 ± 632.4	4497.7 ± 348.1
12	5385.5 ± 336.6	6192.7 ± 214.9	8849.5 ± 374.6	4552.1 ± 307.1
18	3313.4 ± 185.1	4474.9 ± 187.2	4836.1 ± 219.5	3018.7 ± 334.2

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist

Table 2. Results of statistical Analysis Of Variance (ANOVA) and Bonferroni post-hoc tests to test for effects of incubation time (continuous variable), species (categorical variable), and interactions for the dependent variables: dry mass, Hg mass, Hg concentration, Hg/C ratio, C mass, C concentration, C/N ratio, N mass, N concentration, and soluble Hg.

	Dry mass (g)	Hg mass (ng)	Hg concentration ($\mu\text{g kg}^{-1}$)	Hg/C ratio	Dependent variables		C/N ratio	N mass (g)	N concentration (%)	Soluble Hg (ng g^{-1} dry mass)
					C mass (g)	C concentration (%)				
ANOVA test	Variable Time: DF = 1, $P < 0.01$	Variable Time: DF = 1, $P < 0.01$	Variable Time: DF = 1, $P < 0.01$	Variable Time: DF = 1, $P < 0.01$	Variable Time: DF = 1, $P < 0.01$	Variable Time: DF = 1, $P = 0.64$	Variable Time: DF = 1, $P < 0.01$	Variable Time: DF = 1, $P < 0.01$	Variable Time: DF = 1, $P < 0.01$	Variable Time: DF = 1, $P < 0.01$
	Variable Species: DF = 3, $P < 0.01$	Variable Species: DF = 3, $P < 0.01$	Variable Species: DF = 3, $P < 0.01$	Variable Species: DF = 3, $P < 0.01$	Variable Species: DF = 3, $P < 0.01$	Variable Species: DF = 3, $P < 0.01$	Variable Species: DF = 3, $P < 0.01$	Variable Species: DF = 3, $P < 0.01$	Variable Species: DF = 3, $P < 0.01$	Variable Species: DF = 3, $P < 0.01$
	Interaction Time \times Species: DF = 3, $P < 0.01$	Interaction Time \times Species: DF = 3, $P < 0.01$	Interaction Time \times Species: DF = 3, $P < 0.01$	Interaction Time \times Species: DF = 3, $P < 0.01$	Interaction Time \times Species: DF = 3, $P < 0.01$	Interaction Time \times Species: DF = 3, $P = 0.64$	Interaction Time \times Species: DF = 3, $P < 0.01$	Interaction Time \times Species: DF = 3, $P = 0.03$	Interaction Time \times Species: DF = 3, $P < 0.01$	Interaction Time \times Species: DF = 3, $P < 0.01$
Bonferroni comparison	Significant effect of Time in: Mixed deciduous Aspen Pine Oak	Significant effect of Time in: Mixed deciduous Aspen Oak*	Significant effect of Time in: Aspen	Significant effect of Time in: Aspen	Significant effect of Time in: Mixed deciduous Aspen Pine Oak	N/A	Significant effect of Time in: Mixed deciduous Aspen Pine Oak	Significant effect of Time in: Mixed deciduous Aspen Pine Oak	Significant effect of Time in: Mixed deciduous Aspen Oak	Significant effect of Time in: Mixed deciduous Aspen Oak

* Significant post-hoc test at significance level = 0.10, but not at 0.05.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist




Fig. 1. Development of dry mass, carbon (C) mass, and C concentrations throughout 18 months of litter decomposition in a controlled laboratory study. Bars represent means \pm standard deviations of five replicate samples. P -values in title of figures are effects of *Time* based on ANOVA results using all species, and P -values on top of bars are tests for individual species based on Bonferroni post-hoc analyses (Table 2).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Fate of mercury in tree litter during decomposition

A. K. Pokharel and
D. Obrist




Fig. 2. Development of Hg concentration, Hg mass, and Hg/C ratio throughout the 18 months of litter decomposition in the controlled laboratory study. Bars represent means \pm standard deviations of five replicate samples. P -values in title of figure are effects of *Time* based on ANOVA results using all species, and P -values on top of bars are tests for individual species based on Bonferroni post-hoc analyses (Table 2).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. Scatter plots of Hg concentration and C/N ratios of all litter samples in the laboratory decomposition study (black symbols). Samples at different harvest times are marked with different symbols. Regression line marks linear trendline (solid line when $P < 0.05$) between Hg and C/N ratios in laboratory samples. Red symbols are Hg concentrations, and C/N ratios of samples exposed in the field. In two species, points fall clearly above trendlines for laboratory samples.

Fate of mercury in tree litter during decomposition

A. K. Pokharel and D. Obrist

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soluble Hg : $P < 0.05$




Fig. 4. Development of soluble Hg per unit dry mass throughout the 18 months of litter decomposition in the controlled laboratory study. Bars represent means \pm standard deviations of five replicate samples. P -value in title of figure is effect of *Time* based on ANOVA results using all species, and P -values on top of bars are tests for individual species based on Bonferroni post-hoc analyses (Table 2).