



This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Short term changes in methanol emission and pectin methylesterase activity are not directly affected by light in *Lycopersicon esculentum*

P. Y. Oikawa¹, L. Li¹, M. P. Timko¹, J. E. Mak², and M. T. Lerdau^{1,3}

¹Department of Biology, University of Virginia, P.O. Box 400328, Charlottesville, 22904-4328 VA, USA

²School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, 11794-5000, NY, USA

³Department of Environmental Sciences, University of Virginia, P.O. Box 400123, Charlottesville, 22904-4123 VA, USA

Received: 5 November 2010 – Accepted: 23 December 2010 – Published: 17 January 2011

Correspondence to: P. Y. Oikawa (pyo6n@virginia.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

8, 415–434, 2011

Short term changes
in methanol emission

P. Y. Oikawa et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Plants are an important source of atmospheric methanol (MeOH), the second most abundant organic gas after methane. Factors regulating phytogenic MeOH production are not well constrained in current MeOH emission models. Previous studies have indicated that light may have a direct influence on MeOH production. As light is known to regulate cell wall expansion, it was predicted that light would stimulate MeOH production through the pectin methylesterase (PME) pathway. MeOH emissions normalized for stomatal conductance (g_s) did not, however, increase with light over short time scales (20–30 min). After experimentally controlling for g_s and temperature, no light activation of PME activity or MeOH emission was observed. The results clearly demonstrate that light does not directly influence short-term changes in MeOH production and emission. Our data suggest that substrate limitation may be important in regulating MeOH production over short time scales. Future investigation of the long-term impacts of light on MeOH production may increase understanding of MeOH emission dynamics at the seasonal time scale.

1 Introduction

Plants play a dominant role in the global production of methanol (MeOH), an important feature of atmospheric chemistry. Annual global emissions of MeOH range from 100–300 Tg (teragrams, 10^{12} g) yr^{-1} , with plants contributing approximately 75% of this flux, and the remainder's coming from industrial processes, plant decay, biomass burning, and in situ atmospheric production (Singh et al., 2000; Galbally and Kirstine, 2002; Heikes et al., 2002; Tie et al., 2003; von Kuhlmann et al., 2003a, b; Jacob et al., 2005). In contrast to other volatile organic compounds (VOCs), such as isoprene, which have atmospheric lifetimes on the order of tens of minutes, the lifetime of MeOH is approximately ten days (Jacob et al., 2005). This long lifetime allows MeOH to move into the upper troposphere where it can substantially lower hydroxyl radical concentrations as well as increase concentrations of ozone and formaldehyde (Singh et al., 2000, 2001; Tie et al., 2003).

BGD

8, 415–434, 2011

Short term changes in methanol emission

P. Y. Oikawa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Short term changes in methanol emission

P. Y. Oikawa et al.

The dominant biosynthetic pathway for MeOH production in leaves is believed to be the demethylation of pectin by the enzyme pectin methylesterase (PME) (Nemecek-Marshall et al., 1995; Fall and Benson, 1996; Frenkel et al., 1998; Rose and Bennett, 1999; Galbally and Kirstine, 2002; Keppler et al., 2004). Rapidly growing cells export highly methylesterified chains of galacturonic acid (GA; the backbone of pectin) to cell walls (Goldberg et al., 1996). In an ester hydrolysis reaction, GA is demethylated by PME, allowing chains of GA to cross-link and stabilize the cell wall. The methyl groups cleaved from the GA chain form MeOH, which accumulates in the leaf and is released through stomatal openings. As a by-product of this growth-related process, cumulative daily MeOH flux is known to strongly correlate with leaf expansion (Hüve et al., 2007) and studies have consistently shown that young expanding leaves emit greater amounts of MeOH than mature leaves (Macdonald and Fall, 1993; Hüve et al., 2007). Although expanding leaves are associated with higher rates of MeOH production, it is important to note that PMEs are also active in mature tissue (Bordenave and Goldberg, 1994). Plant PMEs are known to belong to large multigene families yielding numerous isoforms of PME, each of which has activities that are regulated by pH (Bordenave and Goldberg, 1994; Goldberg et al., 1996; Willats et al., 2001; Pelloux et al., 2007). The regulation of PME isoforms helps maintain a more flexible cell wall in expanding cells and a rigid cell wall in mature non-expanding cells. Therefore, both mature and immature leaves have active PMEs that produce MeOH at different rates.

Unlike other VOCs such as monoterpenes and isoprene that are directly linked to photosynthesis, less than 10% of MeOH is produced from recently assimilated carbon (Folkers et al., 2008). Although the majority of MeOH is not immediately derived from light-dependent photosynthesis, light may regulate MeOH production through the stimulation of leaf expansion. Light is known to lower the pH of growing cell walls, thereby increasing extensibility of the wall and allowing expansion (Van Volkenburgh, 1999). The process of light-induced apoplastic acidification and cellular growth occurs on the scale of seconds to minutes (Elzenga et al., 1997; Stahlberg and Van Volkenburgh, 1999). Light-induced acidification of the cell wall may influence the activity of certain

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Short term changes in methanol emission

P. Y. Oikawa et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

on MeOH production is an important step in understanding and modeling emission behavior.

In this study, we look at the effects of light on short term changes in MeOH production and emission in *Lycopersicon esculentum* under conditions where potential confounding factors such as temperature variation and changes in g_s are limited. In this manner, we can determine the direct relationship between light and MeOH production.

2 Materials and methods

2.1 Study species

All *Lycopersicon esculentum* Mill. individuals were Micro Tom clones, a dwarf variety

10 of tomato (Meissner et al., 1997). *L. esculentum* was chosen as a model plant due to its rapid growth and high MeOH emission behavior. Plants were grown in the greenhouse at the University of Virginia in Charlottesville (38° N, 78° W). Pots were placed in flats filled with 1 inch of water and were illuminated during a 16 h period with natural light supplemented with high-pressure sodium lamps. Plants were fertilized every
15 two weeks (Scotts 20% N, 20% P, 20% K; Scotts Miracle-Gro Company, Marysville, OH, USA) and kept insect-free using a variety of insecticides. Immature leaves were sampled three weeks past germination and mature leaves six weeks past germination. Leaf size was measured regularly with calipers to ensure that immature leaves were rapidly expanding and mature leaves were fully expanded.

20 2.2 Gas exchange measurements

Leaf-level gas exchange measurements were made with a LI-COR LI-6400 portable gas exchange system (LI-COR Inc., Lincoln, NE, USA). Temperature was regulated within the cuvette using thermoelectric (Peltier) coolers (LI-COR). Photosynthetic photon flux density (PPFD) within the cuvette was controlled with a set of red and blue light-emitting diodes (LI-COR). Depending on the treatment, leaves were exposed

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Short term changes in methanol emission

P. Y. Oikawa et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

to five light levels ranging from low to high light conditions (50, 300, 650, 900, 1150 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Leaf surface area enclosed in the cuvette was measured using a LI-COR Leaf Area Meter (LI-COR Inc., Lincoln, NE, USA). Photosynthetic rates (P_s) and stomatal conductance (g_s) are expressed on a per unit leaf area basis ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ and $\text{mol H}_2\text{O m}^{-2} \text{ s}^{-1}$ for P_s and g_s , respectively). Relative humidity and leaf temperature were also recorded during gas exchange measurements.

2.3 MeOH emission measurements

Leaf-level MeOH emissions were quantified with a LI-COR LI-6400 portable gas exchange system (LI-COR Inc., Lincoln, NE, USA) coupled with a proton-transfer-reaction mass spectrometer (High sensitivity PTR-MS; Ionicon Analytik, Innsbruck, Austria). PTR-MS has been described in detail elsewhere (Lindinger et al., 1998). PTR-MS requires no pre-concentration or chromatography of VOC. Instead, the air flows directly to the drift tube where VOCs undergo chemical ionization via proton-transfer reaction with H_3O^+ . Protonated VOCs are then counted by the ion detector and can be measured down to the ppt level. Air exiting the LI-6400 cuvette was routed to the PTR-MS inlet via 1/4 inch Teflon tubing with a T-fitting in order to release extra flow. Flow rates through the cuvette ranged from 150 to 350 $\mu\text{mol s}^{-1}$. Despite typically stable concentrations of MeOH in ambient air throughout the sampling periods, empty cuvette measurements were coupled with each leaf measurement in order to control for fluctuations in background MeOH. All measurements were taken between 1000 and 1600 h. Leaves were allowed to stabilize at each light level for 20–30 min prior to taking the MeOH emission measurement. PTR-MS measurements were recorded for 20 cycles for each sample. MeOH emission rates are expressed on a per unit leaf area basis ($\text{nmol m}^{-2} \text{ s}^{-1}$). MeOH emissions were divided by g_s in order to normalize emission rates for changes in g_s . Normalized MeOH emissions are expressed in nmol MeOH per mol of H_2O . Four point calibrations were made regularly throughout the sampling period with dilutions of a gravimetrically prepared MeOH gas standard provided by the Riemer lab (University of Miami) containing 3 ppmv (3 $\mu\text{l/l}$) \pm 2% MeOH.

Short term changes in methanol emission

P. Y. Oikawa et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

in nitrogen gas. Accuracy of MeOH measurements was estimated to be around 20% (based on accuracy of calibration measurements) and reproducibility of around 10%.

Three sets of MeOH emission measurements were conducted. All measurements were taken under steady-state conditions. First, repeated MeOH emission measurements were conducted on four immature and four mature *L. esculentum* in which each leaf was exposed to five light levels while keeping temperature constant and allowing g_s to change. Second, repeated MeOH emission measurements were conducted on five immature and five mature *L. esculentum* in which each leaf was exposed to five light levels while temperature and g_s were held constant. Using 12 g LI-COR CO₂ cartridges, g_s was kept relatively constant by changing CO₂ concentrations within the cuvette. The third set of MeOH emission measurements were made on leaves that were destructively sampled for enzyme assay analysis directly following the emission measurement (see Sect. 2.4). Emission measurements were collected at five light levels, where a different group of plants ($n = 5$) were sampled at each level totaling 25 immature and 25 mature *L. esculentum*. Temperature and g_s were held constant.

2.4 PME enzyme activity assay

For the third set of MeOH emission measurements, sampled leaves were excised and frozen in liquid nitrogen directly following the emission measurement. Frozen samples were assayed for PME enzyme activity via a titration technique previously developed for *L. esculentum* (Harriman et al., 1991). Plant tissue was ground in a mortar and pestle to a fine powder, weighed, and mixed in equal weight with a solution composed of 50% 2 M NaCl and 50% 10 mM phosphate buffer (pH 7.5). Samples were then centrifuged at 8000 g for 5 min. 25 µl of plant supernatant was added to 2.5 ml of pectin solution containing 0.5% pectin, 0.2 M NaCl, and 0.1 mM phosphate buffer (pH 7.5). Sample solution pH was adjusted to 7.5 using small amounts of 0.1 M NaOH (in 1–5 µl). Once the solution dropped back down to pH 7, 1–5 µl of 0.1 M NaOH was added until solution pH reached 7.3. Time for solution to drop back down to pH 7 was recorded. PME activity is expressed in µmol g fwt⁻¹ min⁻¹ based on the change in pH for a given

amount of fresh tissue over time. Measuring change in pH over time is a proxy for PME activity and not a direct measurement of enzyme activity, but this change in pH has been shown to be a highly repeatable proxy for enzyme activity (Harriman et al., 1991). A total of 25 immature and 25 mature plants were assayed for PME enzyme activity.

5 2.5 Statistical analysis

Repeated measurements collected in the first MeOH emission dataset, where leaves were exposed to five light levels under constant temperature, were analyzed using a mixed model ANOVA (Proc MIXED, SAS 9.1; SAS Institute Inc., Cary, NC, USA). Strength of association between MeOH emission and PPFD was determined by the 10 significance of the slope of the mixed model. Normalized MeOH emission data in the first dataset exhibited a decreasing exponential relationship with PPFD and were examined with a generalized linear model with a negative binomial distribution (Proc GENMOD, SAS 9.1; SAS Institute Inc., Cary, NC, USA). The relationship between g_s and PPFD was analyzed using a repeated-measures ANOVA model (Proc GLM, SAS 15 9.1; SAS Institute Inc., Cary, NC, USA). Linear and non-linear regression lines were fit using SigmaPlot 9.0 (Systat software, Inc. Point Richmond, California, USA).

The second set of repeated MeOH emission measurements, where leaves were exposed to five light levels under constant temperature and g_s , was analyzed using a mixed model ANOVA with MeOH emission data normalized for g_s (Proc MIXED, 20 SAS 9.1; SAS Institute Inc., Cary, NC, USA). Again, strength of association between normalized MeOH emission and PPFD was determined by the significance of the slope of the mixed model.

In the third set of MeOH emission measurements, leaves were destructively sampled for PME activity assays under conditions of constant temperature and g_s . This third 25 dataset was examined in a two-way MANOVA where the effect of plant type and PPFD on MeOH emission and PME activity was analyzed (Proc GLM, SAS 9.1; SAS Institute Inc., Cary, NC, USA). Data used in the analysis were Log and square root transformed to meet normality and homogeneity of variance assumptions.

3 Results

In the first set of measurements, the relationship between g_s and MeOH emission was investigated under conditions where light varied and temperature was held constant. MeOH emission and g_s were measured at five light levels. Stomatal conductance ranged from 0.03–0.17 mol H₂O m⁻² s⁻¹ and 0.04–0.22 mol H₂O m⁻² s⁻¹ for mature and immature leaves, respectively. Measurements of mature and immature leaves were taken under steady-state conditions at leaf temperature 29 ± 1 °C, photosynthetic rates 6.7 ± 4 µmol CO₂ m⁻² s⁻¹ mature and 10 ± 4 µmol CO₂ m⁻² s⁻¹ immature, and relative humidity $55 \pm 5\%$ mature and $58 \pm 6\%$ immature (means \pm SD). MeOH emissions increased with light ($F = 16.69$ $df = 1, 15$ $P = 0.001$ for mature; $F = 110.59$ $df = 1, 15$ $P < 0.0001$ for immature; Fig. 1a). Slopes were positive and significant for both leaf types (slope = 0.0015 $t = 4.09$ $P = 0.001$ for mature; slope = 0.0044 $t = 10.52$ $P < 0.0001$ for immature). Light also had an overall significant effect on g_s for both leaf types (Fig. 1b; Wilk's lambda $F = 77.78$ $df = 4, 3$ $P = 0.002$). When MeOH flux values were normalized by g_s , the influence of light disappeared. The relationship between normalized MeOH and light was not positive (regression coefficient = -0.0006, $z = -5.53$, $P < 0.0001$ for mature leaves; regression coefficient = -0.0008, $z = -50.20$, $P < 0.0001$ for immature leaves; Fig. 1c), suggesting that g_s is responsible for apparent short-term MeOH emission responses to light. It is important to note that the calculated negative relationship between normalized MeOH and PPF is almost certainly an artifact of the normalization of MeOH and not a biological phenomenon.

As an extension to the first set of measurements, g_s and temperature were experimentally controlled before making repeated MeOH flux measurements across five light levels. Measurements of mature and immature leaves were taken under steady state conditions at leaf temperature 29 ± 1 °C, $g_s 0.08 \pm 0.02$ mol H₂O m⁻² s⁻¹, and relative humidity $58.3 \pm 7\%$ (means \pm SD). Under constant temperature and g_s no significant effect of light on MeOH emissions for mature ($F = 2.96$ $df = 1, 19$ $P = 0.10$) or immature leaves ($F = 0.6$ $df = 1, 19$ $P = 0.45$) was observed (Fig. 2). The results agreed with

Short term changes in methanol emission

P. Y. Oikawa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



those from the first set of measurements in that light did not directly stimulate MeOH emission.

In the third set of MeOH emission measurements, the direct influence of light on MeOH production was investigated by assaying PME activity at each light level. Measurements of mature and immature leaves were taken under steady state conditions at leaf temperature 29 ± 1 °C, g_s 0.09 ± 0.02 mol H₂O m⁻² s⁻¹, and relative humidity $54.5 \pm 9\%$ (means \pm SD). Immature leaves had significantly greater MeOH emission and PME activity overall (Wilk's lambda $F = 45.47$ $df = 2, 38$ $P < 0.0001$). Light had no significant effect on either PME activity or MeOH emission under constant temperature and g_s (Figs. 3 and 4; Wilk's lambda $F = 0.57$ $df = 8, 76$ $P = 0.80$). Light was therefore not observed to stimulate MeOH production through the PME pathway on the time scale of 0–30 min.

4 Discussion

Factors regulating MeOH production in plants are not well constrained in current MeOH emission models (Galbally and Kirstine, 2002; Karl et al., 2003; Harley et al., 2007). Previous studies have suggested that light directly stimulates MeOH production (Harley et al., 2007). As light is known to regulate cell wall expansion on the scale of minutes, we predicted that short-term increases in light would stimulate PME activity and MeOH emission in immature leaves. The results, however, did not bear out these predictions, and MeOH emissions normalized for g_s demonstrated that light did not have a direct influence on MeOH emission. Our data also demonstrated that short-term changes in light do not influence MeOH emission if g_s and temperature are experimentally controlled. In accord with previous studies, immature leaves had significantly higher MeOH emission (Nemecek-Marshall et al., 1995; Hüve et al., 2007), as well as higher PME activity, than mature leaves. Despite being in a rapidly expanding growth phase, PME activity in immature leaves was not stimulated by light over time scales of 20–30 min. It is clear that MeOH emission and production do not behave similarly to other VOCs

BGD

8, 415–434, 2011

Short term changes in methanol emission

P. Y. Oikawa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



such as isoprene which are known to increase production and emission in response to short term changes in light. Light does not stimulate PME activity or MeOH emission in either mature or immature leaves on short time scales.

BGD

8, 415–434, 2011

MeOH production is regulated not only by PME activity, but also by multiple interacting factors, such as cell wall pH, substrate availability, and PME gene expression (Goldberg et al., 1996; Pelloux et al., 2007). The acidification of the cell wall in response to light-induced growth is known to occur on the scale of seconds to minutes (Elzenga et al., 1997; Stahlberg and Van Volkenburgh, 1999), and we hypothesized that light-induced changes in cell wall pH and cell expansion would influence MeOH production. The results, however, suggested that the fast-acting effects of light on the acidification of the cell wall did not significantly alter overall PME activity. As PME activity is dependent on cell wall pH, it is surprising that PME activity did not change across a wide spectrum of light intensity. This lack of change in PME activity and MeOH emission response to light suggests that PME may, at times, be substrate limited. Studies have shown that cell expansion is not immediately accompanied by cell wall synthesis. Rapidly growing cells are known to stretch cell walls thin due to the lack of cell wall components, such as pectin polysaccharides (Refregier et al., 2004). The addition of methylated pectin to the cell wall during 20–30 min of light-induced growth can therefore not be assumed. Although pectin synthesis is generally known to be under developmental control and also induced by mechanical damage, pathogenesis, and cell-cell interactions, understanding of the transcriptional control of cell wall polysaccharides is incomplete (Somerville et al., 2004). In addition to pectin synthesis, the regulation of PME gene expression could also limit MeOH production. Unfortunately, little is known concerning the gene expression of pectin and PMEs and how they regulate MeOH production in plants. Despite the lack of knowledge, future research should focus on measuring PME substrate limitation and gene regulation, with the possibility of linking these processes with MeOH production.

Investigation of long-term changes in MeOH production and emission in response to light may be valuable for modeling efforts. Factors such as PME activity, PME

Short term changes in methanol emission

P. Y. Oikawa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Short term changes in methanol emission

P. Y. Oikawa et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

References

Bordenave, M. and Goldberg, R.: Immobilized and free apoplastic pectinmethyl esterases in mung bean hypocotyl, *Plant Physiol.*, 106, 1151–1156, 1994.

Ciccioli, P., Fabozzi, C., Brancaleoni, E., Cecinato, A., Frattoni, M., Loreto, F., Kesselmeier, J., Schafer, L., Bode, K., Torres, L., and Fugit, J. L.: Use of the isoprene algorithm for predicting the monoterpene emission from the Mediterranean holm oak *Quercus ilex* L.: performance and limits of this approach, *J. Geophys. Res.-Atmos.*, 102, 23319–23328, 1997.

Elzenga, J. T. M., Staal, M., and Prins, H. B. A.: Calcium-calmodulin signalling is involved in light-induced acidification by epidermal leaf cells of pea, *Pisum sativum* L., *J. Exp. Bot.*, 48, 2055–2061, doi:10.1093/jxb/48.12.2055, 1997.

Fall, R. and Benson, A. A.: Leaf methanol – the simplest natural product from plants, *Trends Plant Sci.*, 1, 296–301, 1996.

Folkers, A., Hüve, K., Ammann, C., Dindorf, T., Kesselmeier, J., Kleist, E., Kuhn, U., Uerlings, R., and Wildt, J.: Methanol emissions from deciduous tree species: dependence on temperature and light intensity, *Plant Biol.*, 10, 65–75, 10.1111/j.1438-8677.2007.00012.x, 2008.

Frenkel, C., Peters, J. S., Tieman, D. M., Tiznado, M. E., and Handa, A. K.: Pectin methylesterase regulates methanol and ethanol accumulation in ripening tomato (*Lycopersicon esculentum*) fruit, *J. Biol. Chem.*, 273, 4293–4295, 1998.

Fuentes, J. D. and Wang, D.: On the seasonality of isoprene emissions from a mixed temperate forest, *Ecol. Appl.*, 9, 1118–1131, 1999.

Galbally, I. E. and Kirstine, W.: The production of methanol by flowering plants and the global cycle of methanol, *J. Atmos. Chem.*, 43, 195–229, 2002.

Geron, C., Guenther, A., Sharkey, T., and Arnts, R. R.: Temporal variability in basal isoprene emission factor, *Tree Physiol.*, 20, 799–805, 2000.

Goldberg, R., Morvan, C., Jauneau, A., and Jarvis, M. C.: Methyl-esterification, de-esterification and gelation of pectins in the primary cell wall, in: *Pectins and Pectinases*, edited by: Visser, J. and Voragen, A. G. J., Elsevier Science, Amsterdam, 151–172, 1996.

Gray, D. W., Goldstein, A. H., and Lerdau, M. T.: The influence of light environment on photosynthesis and basal methylbutenol emission from *Pinus ponderosa*, *Plant Cell Environ.*, 28, 1463–1474, 2005.

Gray, D. W., Goldstein, A. H., and Lerdau, M. T.: Thermal history regulates methylbutenol basal emission rate in *Pinus ponderosa*, *Plant Cell Environ.*, 29, 1298–1308, 2006.

BGD

8, 415–434, 2011

Short term changes in methanol emission

P. Y. Oikawa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Short term changes in methanol emission

P. Y. Oikawa et al.

Hanson, D. T. and Sharkey, T. D.: Effect of growth conditions on isoprene emission and other thermotolerance-enhancing compounds, *Plant Cell Environ.*, 24, 929–936, 2001.

5 Harley, P., Guenther, A., and Zimmerman, P.: Effects of light, temperature and canopy position on net photosynthesis and isoprene emission from sweetgum (*Liquidambar styraciflua*) leaves, *Tree Physiol.*, 16, 25–32, 1996.

Harley, P., Guenther, A., and Zimmerman, P.: Environmental controls over isoprene emission in 10 deciduous oak canopies, *Tree Physiol.*, 17, 705–714, 1997.

10 Harley, P., Greenberg, J., Niinemets, Ü., and Guenther, A.: Environmental controls over methanol emission from leaves, *Biogeosciences*, 4, 1083–1099, doi:10.5194/bg-4-1083-2007, 2007.

Harriman, R. W., Tieman, D. M., and Handa, A. K.: Molecular cloning of tomato pectin methylesterase gene and its expression in Rutgers, ripening inhibitor, nonripening, and never ripe tomato fruits, *Plant Physiol.*, 97, 80–87, 1991.

15 Heikes, B. G., Chang, W. N., Pilson, M. E. Q., Swift, E., Singh, H. B., Guenther, A., Jacob, D. J., Field, B. D., Fall, R., Riemer, D., and Brand, L.: Atmospheric methanol budget and ocean implication, *Global Biogeochem. Cy.*, 16, 1133, doi:10.1029/2002GB001895, 2002.

Hüve, K., Christ, M. M., Kleist, E., Uerlings, R., Niinemets, U., Walter, A., and Wildt, J.: Simultaneous growth and emission measurements demonstrate an interactive control of methanol release by leaf expansion and stomata, *J. Exp. Bot.*, 58, 1783–1793, 2007.

20 Jacob, D. J., Field, B. D., Li, Q. B., Blake, D. R., de Gouw, J., Warneke, C., Hansel, A., Wisthaler, A., Singh, H. B., and Guenther, A.: Global budget of methanol: constraints from atmospheric observations, *J. Geophys. Res.-Atmos.*, 110, D08303, doi:10.1029/2004JD005172, 2005.

Karl, T., Guenther, A., Spirig, C., Hansel, A., and Fall, R.: Seasonal variation of biogenic VOC 25 emissions above a mixed hardwood forest in Northern Michigan, *Geophys. Res. Lett.*, 30, 2186, doi:10.1029/2003GL018432, 2003.

Keppler, F., Kalin, R. M., Harper, D. B., McRoberts, W. C., and Hamilton, J. T. G.: Carbon isotope anomaly in the major plant C₁ pool and its global biogeochemical implications, *Biogeosciences*, 1, 123–131, doi:10.5194/bg-1-123-2004, 2004.

30 Lehning, A., Zimmer, I., Steinbrecher, R., Bruggemann, N., and Schnitzler, J. P.: Isoprene synthase activity and its relation to isoprene emission in *Quercus robur* L-leaves, *Plant Cell Environ.*, 22, 495–504, 1999.

Lehning, A., Zimmer, W., Zimmer, I., and Schnitzler, J. P.: Modeling of annual variations of oak

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 Lindinger, W., Hansel, A., and Jordan, A.: Proton-transfer-reaction mass spectrometry (PTR-MS): on-line monitoring of volatile organic compounds at pptv levels, *Chem. Soc. Rev.*, 27, 347–354, 1998.

10 Loreto, F., Ciccioli, P., Cecinato, A., Brancaleoni, E., Frattoni, M., Fabozzi, C., and Tricoli, D.: Evidence of the photosynthetic origin of monoterpenes emitted by *Quercus ilex* L. leaves by ^{13}C labeling, *Plant Physiol.*, 110, 1317–1322, doi:10.1104/pp.110.4.1317, 1996a.

15 Loreto, F., Ciccioli, P., Cecinato, A., Brancaleoni, E., Frattoni, M., and Tricoli, D.: Influence of environmental factors and air composition on the emission of [alpha]-pinene from *Quercus ilex* leaves, *Plant Physiol.*, 110, 267–275, doi:10.1104/pp.110.1.267, 1996b.

20 Macdonald, R. C. and Fall, R.: Detection of substantial emissions of methanol from plants to the atmosphere, *Atmos. Environ. A-Gen.*, 27, 1709–1713, 1993.

25 Meissner, R., Jacobson, Y., Melamed, S., Levyatuv, S., Shalev, G., Ashri, A., Elkind, Y., and Levy, A.: A new model system for tomato genetics, *Plant J.*, 12, 1465–1472, 1997.

Nemecek-Marshall, M., Macdonald, R. C., Franzen, F. J., Wojciechowski, C. L., and Fall, R.: Methanol emission from leaves – enzymatic detection of gas-phase methanol and relation of methanol fluxes to stomatal conductance and leaf development, *Plant Physiol.*, 108, 1359–1368, 1995.

30 Niinemets, U. and Reichstein, M.: Controls on the emission of plant volatiles through stomata: differential sensitivity of emission rates to stomatal closure explained, *J. Geophys. Res.-Atmos.*, 108(17), 4208, doi:10.1029/2002JD002620, 2003a.

Niinemets, U. and Reichstein, M.: Controls on the emission of plant volatiles through stomata: a sensitivity analysis, *J. Geophys. Res.-Atmos.*, 108, 4211, doi:10.1029/2002JD002626, 2003b.

Pelloux, J., Rusterucci, C., and Mellerowicz, E. J.: New insights into pectin methylesterase structure and function, *Trends Plant Sci.*, 12, 267–277, doi:10.1016/j.tplants.2007.04.001, 2007.

35 Petron, G., Harley, P., Greenberg, J., and Guenther, A.: Seasonal temperature variations influence isoprene emission, *Geophys. Res. Lett.*, 28, 1707–1710, 2001.

Refregier, G., Pelletier, S., Jaillard, D., and Hofte, H.: Interaction between wall deposition and cell elongation in dark-grown hypocotyl cells in *Arabidopsis*, *Plant Physiol.*, 135, 959–968, doi:10.1104/pp.104.038711, 2004.

Short term changes in methanol emission

P. Y. Oikawa et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Short term changes in methanol emission

P. Y. Oikawa et al.

Rose, J. K. C. and Bennett, A. B.: Cooperative disassembly of the cellulose-xyloglucan network of plant cell walls: parallels between cell expansion and fruit ripening, *Trends Plant Sci.*, 4, 176–183, 1999.

Sharkey, T. D., Singsaas, E. L., Lerdau, M. T., and Geron, C. D.: Weather effects on isoprene emission capacity and applications in emissions algorithms, *Ecol. Appl.*, 9, 1132–1137, 1999.

Singh, H., Chen, Y., Tabazadeh, A., Fukui, Y., Bey, I., Yantosca, R., Jacob, D., Arnold, F., Wohlfstrom, K., Atlas, E., Flocke, F., Blake, D., Blake, N., Heikes, B., Snow, J., Talbot, R., Gregory, G., Sachse, G., Vay, S., and Kondo, Y.: Distribution and fate of selected oxygenated organic species in the troposphere and lower stratosphere over the Atlantic, *J. Geophys. Res.-Atmos.*, 105, 3795–3805, 2000.

Singh, H., Chen, Y., Staudt, A., Jacob, D., Blake, D., Heikes, B., and Snow, J.: Evidence from the Pacific troposphere for large global sources of oxygenated organic compounds, *Nature*, 410, 1078–1081, 2001.

Somerville, C., Bauer, S., Brininstool, G., Facette, M., Hamann, T., Milne, J., Osborne, E., Paredez, A., Persson, S., Raab, T., Vorwerk, S., and Youngs, H.: Toward a systems approach to understanding plant cell walls, *Science*, 306, 2206–2211, doi:10.1126/science.1102765, 2004.

Stahlberg, R. and Van Volkenburgh, E.: The effect of light on membrane potential, apoplastic pH and cell expansion in leaves of *Pisum sativum* L. var. Argenteum., *Planta*, 208, 188–195, 1999.

Tie, X., Guenther, A., and Holland, E.: Biogenic methanol and its impacts on tropospheric oxidants, *Geophys. Res. Lett.*, 30, 1881, 2003.

Van Volkenburgh, E.: Leaf expansion – an integrating plant behaviour, *Plant Cell Environ.*, 22, 1463–1473, 1999.

von Kuhlmann, R., Lawrence, M. G., Crutzen, P. J., and Rasch, P. J.: A model for studies of tropospheric ozone and nonmethane hydrocarbons: model description and ozone results, *J. Geophys. Res.-Atmos.*, 108(22), 4294, doi:10.1029/2002jd002893, 2003a.

von Kuhlmann, R., Lawrence, M. G., Crutzen, P. J., and Rasch, P. J.: A model for studies of tropospheric ozone and nonmethane hydrocarbons: model evaluation of ozone-related species, *J. Geophys. Res.-Atmos.*, 108(26), 4729, doi:10.1029/2002jd003348, 2003b.

Willats, W. G. T., McCartney, L., Mackie, W., and Knox, J. P.: Pectin: cell biology and prospects for functional analysis, *Plant Mol. Biol.*, 47, 9–27, 2001.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 1. **(a)** MeOH emission, **(b)** stomatal conductance (g_s), and **(c)** MeOH emission normalized for g_s from mature ($n = 4$; black circles) and immature ($n = 4$; open circles) *L. esculentum* over five light levels at constant temperature (29 ± 1 °C). MeOH emission (slope = 0.0015, $t = 4.09$, $P = 0.001$ for mature; slope = 0.0044, $t = 10.52$, $P < 0.0001$ for immature) and g_s ($F = 112.94$ $P < 0.0001$ $R^2 = 0.86$ for mature; $F = 19.06$ $P = 0.004$ $R^2 = 0.96$ for immature) significantly increased with PPFD. Normalized MeOH emissions were negatively associated with PPFD (regression coefficient = -0.0006 , $z = -5.53$, $P < 0.0001$ for mature; regression coefficient = -0.0008 , $z = -50.20$, $P < 0.0001$ for immature).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Short term changes
in methanol emission

P. Y. Oikawa et al.




Fig. 2. Average MeOH flux values normalized for stomatal conductance (g_s) over five light levels for $n = 5$ mature (black circles) and immature (white circles) *L. esculentum* (each plant was measured 5x; standard error bars shown). PPFD had no significant effect on MeOH flux for either leaf type ($F = 2.96$; $df = 1, 19$; $P = 0.10$ for mature; $F = 0.6$; $df = 1, 19$; $P = 0.45$ for immature).

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)



Short term changes
in methanol emission

P. Y. Oikawa et al.




Fig. 3. Average MeOH flux values normalized for stomatal conductance (g_s) over five light levels for $n = 5$ mature (black circles) and immature (white circles) *L. esculentum* (standard error bars shown). PPFD had no significant effect on MeOH flux ($F = 0.81$ $df = 4$ $P = 0.53$). Plant type had a significant effect on MeOH emission overall ($F = 79.43$ $df = 1$ $P < 0.0001$).

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)




Fig. 4. Average PME activity rates over five light levels for $n = 5$ mature (black circles) and immature (white circles) *L. esculentum* (standard error bars shown). PPFD had no significant effect on enzyme activity ($F = 0.40$ $df = 4$ $P = 0.81$). Plant type had a significant effect on PME activity overall ($F = 14.46$ $df = 1$ $P = 0.0005$).