



This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga¹, S. Chatani^{1,2}, S. Nakatsuka¹, D. Kusumoto³, K. Kubota⁴,
Y. Utsumi⁴, T. Enoki⁴, A. Tani⁵, and T. Hiura⁶

¹Auto Oil and New Fuels Department, Japan Petroleum Energy Center, 4-3-9 Toranomon, Minato-Ku, Tokyo 105-0001, Japan

²Toyota Central R&D Laboratories Inc., 41-1 Aza Yokomichi, Oaza Nagakute, Nagakute-cho, Aichi-gun, Aichi 480-1192, Japan

³Experimental Station at Tanashi, University Forests, Graduate school of agricultural and life science, the University of Tokyo, 1-1-8 Midorimachi, Nishi-Tokyo, Tokyo 188-0002, Japan

⁴Shiiba Research Forest, Kyushu University, 949 Okawachi, Shiiba, Higashi-Usuki, Miyazaki 883-0402, Japan

⁵Institute for Environmental Science, University of Shizuoka, 52-1 Yada, Suruga-Ku, Shizuoka 422-8526, Japan

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

⁶Tomakomai Research Station, Field Science Center for Northern Biosphere, Hokkaido University, Takaoka, Tomakomai, Hokkaido 053-0035, Japan

Received: 26 May 2011 – Accepted: 4 July 2011 – Published: 8 July 2011

Correspondence to: S. N. Matsunaga (s-matsunaga@pecj.or.jp)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Abstract

Reactive volatile organic compounds (VOCs) are known to affect atmospheric chemistry, especially biogenic VOCs (BVOCs) which have a significant impact on regional air quality due to their large emission rates and high reactivities. Diterpenes (Kaur-16-ene and a diterpene likely to be Hibaene) were detected in all of the 205 enclosure air samples collected over multiple seasons at two different sites from *Cryptomeria japonica* and *Chamaecyparis obtusa* trees, the dominant coniferous trees in Japan. The emission rate of Kaur-16-ene, was determined to be from 0.01 to 7.1 $\mu\text{g dwg}^{-1} \text{h}^{-1}$ (average: 0.61 $\mu\text{g dwg}^{-1} \text{h}^{-1}$) by branch enclosure measurements and solid phase – liquid extraction techniques. The emission rate was an order of magnitude higher than that of monoterpenes which are known to comprise a major portion of total BVOC emissions. In addition, total emission of Kaur-16-ene at 30 °C was estimated to be comparable to that of total anthropogenic VOC emissions. To our knowledge, this is the first report of volatile diterpene emissions from trees.

15 1 Introduction

Reactive volatile organic compounds (VOCs) have multiple important impacts on atmospheric chemistry and physics (Guenther et al., 2006). For example, VOCs are a precursor to tropospheric ozone, an effective greenhouse gas known to be toxic at sufficiently high concentrations (Mustafa, 1990). Reactive VOCs at high concentrations 20 may also decrease atmospheric concentrations of the hydroxyl radical (OH), a strong oxidant and major sink of less reactive greenhouse gases such as methane in the atmosphere (Kulumala et al., 2004). VOCs also form secondary organic aerosols which may produce direct and indirect effects on the earth's radiation budget (Kavouras et al., 1998).

25 VOCs are known to have both anthropogenic and biogenic sources. Biogenic VOCs (BVOCs) are mainly emitted from terrestrial higher plants such as trees, and are

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



5 rates of the heavier BVOCs. Kaur-16-ene and other compounds likely to be diterpenes were detected in all of the emission samples collected from *C. japonica* and *C. obtusa* over multiple seasons at two different sites. These two endemic tree species are the most dominant trees in Japan accounting for approximately 60 % of total leaf mass in Japanese forests (Matsunaga, 2011). Here, we present the first observations of biogenic emissions of diterpenes in remarkable concentrations.

10 Although the analytical technique cannot distinguish stereo-isomers compounds produced in leaves (Railton et al., 1984; Perry and Weavers, 1985; Cool et al., 1998), the Kaur-16-ene detected in this study is most likely to be ent-Kaurene. The emission rate of Kaur-16-ene was generally more than an order of magnitude higher than that of total monoterpane emissions. Assuming a temperature dependence of Kaur-16-ene emission used in the G93 model (Guenther et al., 1993), the emission was estimated to be comparable to total monoterpane emission, a group of commonly emitted BVOCs, and also to total anthropogenic VOC emission in Japan in summer.

15 2 Experiment

2.1 Site description and field campaign

Seven field campaigns have been conducted at two sites, Tanashi Experimental Station of the Tokyo University Forests ($35^{\circ}44'21''$ N, $139^{\circ}32'18''$ E) in Tokyo and Miyazaki Research Forest of Kyushu University ($32^{\circ}21'23''$ N, $131^{\circ}6'2''$ E) in Shiiba, Miyazaki, 20 Japan. Both of the experimental forests have a planted area of mature *C. japonica* and *C. obtusa* trees. The campaigns were conducted in January, August, and November of 2009 and in April of 2010 at Tanashi and in May, August, and November of 2010 at Shiiba. Each sampling period was 5 days.

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

2.2 Branch enclosure system

Leaves were sampled at branch ends about 5–15 nodes from the trunk and were located at the top of the canopy approximately 10–15 m above the ground level. Tree ages of *C. japonica* at Tanashi and Shiiba were 54 and 56 yr, respectively and for *C. obtusa*, at Tanashi and Shiiba were 33 and 56 yr, respectively.

The branches were enclosed with a 15 l Teflon[®] bag (transparent to more than 95 % of PAR). The enclosure samples were collected from the branch of *C. japonica* at canopy top and 5 m (3 m for *C. obtusa*) below the top within same individual. The bags were installed at least 24 h prior to the first sample collection to avoid disturbance due to the installation of the bag. The air inside the bag was ventilated with a VOC free air at a flow rate of 4–5 l min⁻¹ to avoid water condensation and excessive temperature increase in the bag (see Fig. 1 for an overview). The bags were installed onto the branches on the first day of the campaign and kept on the branches until the sampling period ended 5 days later. The air was supplied from a commercial compressed air cylinder and cleaned by an activated charcoal trap. The flow rate was controlled by a mass flow controller. Air entering the bag was supplied through a Teflon[®] tube ring roughly 1.5 mm in diameter with approximately 20 holes so that the inside air could be circulated and homogenized. The compressed air was taken from the atmosphere and, thus, the CO₂ concentrations are assumed to remain at average ambient level. The maximum relative humidity in the bag has been calculated to be 70–75 % in the daytime. Considering that water condensation was observed when the air flow rate was reduced to below 4 l min⁻¹, relative humidity in the bag was not too dry during sampling. Air temperature inside of the bag was recorded every 2 min by a data logger equipped with a Teflon[®] coated thermo sensor (RS-12, Especmic, Ohguchi, Niwa, Aichi, Japan). Branches were harvested after the completion of each field campaign and dried at 60 °C for 48 h to determine dry weight of the leaves, therefore, samples were collected from different branches of same individuals with other sampling periods in every season. For *C. japonica*, the dry weight was determined for whole branches

BGD

8, 6681–6700, 2011

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



harvested because stems also have green needle leaves. Therefore, the dry weight for *C. japonica* includes stem, bark and leaves.

2.3 Sample collection

Diterpene samples were collected from the enclosure bag described in Sect. 2.2.

5 Monoterpene and sesquiterpenes, which are C_{10} and C_{15} BVOCs, respectively, were also collected from the same bag and analyzed to confirm any disturbance (i.e., extraordinarily high concentrations from damage to storage reservoirs of the BVOCs).

The diterpene samples were collected into a 6.35 mm (1/4 inches) o.d. glass tube filled with approximately 60 mg of HayeSep Q (Hayes Separations, Inc., Bandera TX, USA). Sample flow rate was controlled at 200 ml min^{-1} by a mass flow controller connected to an electronic pump. The collection time ranged from 0.5 to 2 h. Diterpenes collected on the adsorbent were extracted with approximately 2 ml of hexane (Special grade, Wako chemical, Chuo-Ku, Osaka, Japan) at the site. The extracts were dropped into 2 ml glass vials (Supelco, Bellefonte PA, USA) directly from the sampling tube and stored in a freezer, which was kept at approximately -15°C , at the site. The samples were transported to the laboratory by a commercial transport service, which kept the packages at -18°C ("Cool Takkyu-bin", Yamato Transport Co., Ltd., Chuo-Ku, Tokyo, Japan) before further procedures were performed.

2.4 GC-MS analysis

20 A one micro liter of internal standard solution ($20\text{--}100 \text{ ng } \mu\text{l}^{-1}$ of cyclopentadecane in hexane) was added into the extracts prior to the concentration. Then, the extracts were evaporated to approximately $5\text{--}10 \mu\text{l}$ by a gentle argon flow at the laboratory. The internal standard solution was prepared for each field campaign. One μl of the concentrated extract containing the internal standard was injected into the split-splitless injector of the gas chromatograph (GC; Agilent 7890A GC, Agilent Technologies Inc., Santa Clara CA, USA) connected to a mass spectrometer (MS; Agilent 5975C MSD). Diterpenes

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



were separated using a fused silica capillary column (HP-5ms, 30 m × 0.25 mm i.d., film thickness of 0.25 µm; Agilent Technologies Inc.) and helium carrier gas. The injector (300 °C) was equipped with a deactivated quartz insert tube and operated in the pulsed splitless mode. GC oven temperatures were programmed to start at 60 °C and increase to 150 °C at 2.5 °C min⁻¹, then raised to 300 °C at 30 °C min⁻¹ and held for 10 min. The identification of Kaur-16-ene was made comparing its GC retention time and mass spectra with those of the authentic standard (Figs. 2 and 3). The authentic standard was purchased from OIChemIm Ltd., Olomouc, Czech Republic. Only Kaur-16-ene and an unknown diterpene, likely to be Hibaene, were detected as two abundant peaks of diterpenes (Fig. 2).

Kaur-16-ene was quantified using the MS in the selected ion monitoring mode. The amount of the Kaur-16-ene was determined by comparing response intensity (peak area on a chromatogram for the selected ions) between the internal standard and the target compounds. The response of the Kaur-16-ene was adjusted based on the ratio of the response for internal standard to Kaur-16-ene at the same amount because the selected ions are different between Kaur-16-ene and the standard. The ratio was determined by analyzing authentic standard for Kaur-16-ene and internal standard (cyclopentadecane) purchased from Wako and Sigma-Aldrich (Co., St. Louis MO, USA). The amount of the standard was gravimetrically determined. The precision of the whole analytical procedure was around 5 %. Detection limit was approximately 0.1 ng g⁻¹ h⁻¹ (depending on the leaf mass enclosed in the bag and the sampling time).

2.5 Calculation for the emission rate of Kaur-16-ene

The emission rate was determined based on the Eq. 1 below.

$$E = \frac{F_{\text{air}} \times C - C_{\text{blank}}}{f_{\text{samp}} \times t_{\text{samp}} \times W_{\text{leaf}}} \quad (1)$$

where, E and C are the emission rate in µg g⁻¹ h⁻¹ and determined amount of Kaur-16-ene by the GC-MS analysis in µg, respectively. F_{air} , C_{blank} , F_{samp} , t_{samp} and W_{leaf} are 6688

flow rate of the purified air supplied to the bag in ml min^{-1} , amount of Kaur-16-ene in the blank sample in μg adjusted for the actual sampling time, sample flow rate, sampling time in hour and dry weight of the leaf in g, respectively. C_{blank} for Kaur-16-ene was below the detection limit through all field campaigns.

5 3 Results and discussion

3.1 Basal emission rate of Kaur-16-ene

Only Kaur-16-ene was identified and quantified in the diterpenes using an authentic standard. To estimate the regional emission rate, the basal emission rate of Kaur-16-ene was calculated. Basal emission rate is defined to be an emission rate of BVOC, which is emitted into the canopy, at the standard condition (30°C for temperature dependence) (Guenther et al., 1993, Guenther et al., 2006). Therefore, it is regarded to be a constant in most BVOC emission models. Figure 4a and b presents observed relationships between the inside air temperature of the enclosure bag and measured emission rate of Kaur-16-ene from *C. japonica* and *C. obtusa* observed at Tanashi and Shiiba. Plot for *C. obtusa* was only for data obtained at Shiiba due to insufficient number of samples at Tanashi. Assuming temperature dependence of the emission of Kaur-16-ene, the emission rate can be estimated according to the G93 algorithm (Eq. 2).

$$E = E_s \exp\{\beta(T - T_s)\} \quad (2)$$

$$20 \quad \ln E = \beta(T - T_s) + \ln E_s \quad (3)$$

where, E presents measured emission rate. T and T_s presents measured temperature and the standard temperature, respectively. An empirical coefficient β can be calculated as the slope of the liner regression for the plot and quantifies the intensity of temperature dependence (Eq. 3). Natural logarithm of the basal emission rate E_s is

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the intercept at y-axis (Guenther et al., 1993). The calculated E_s and β over the seven campaigns are presented in Table 1. Although there may be another factor which controls the emission (e.g. light intensity), we concluded that temperature is the most effective controlling factor of the emission of Kaur-16-ene based on comparisons of the emission rate between branch at the canopy top and the light attenuated branch. The comparisons have been conducted for both *C. japonica* and *C. obtusa* trees. The basal emission rate did not show any significant difference with light intensity.

The basal emission rate of Kaur-16-ene did not show significant seasonal variation. As mentioned above, basal emission rate is regarded to be a constant in most emission models because it is a normalized emission rate at a set of standard conditions. However, recent studies report that the basal emission often has a seasonal variation (Geron and Arnts, 2010). Matsunaga et al. (2011) also reports that the basal emission rates of monoterpenes and sesquiterpenes emitted from *C. japonica* change with season. However, the basal emission rate of Kaur-16-ene emitted from *C. japonica* and *C. obtusa* did not clearly change with seasons. This may suggest that diterpenes are produced for other purposes or are produced through different processes than mono- and sesquiterpenes.

3.2 Estimate on the Kaur-16-ene emission over Japan

The total emission rate of Kaur-16-ene over Japan at 30 °C and 5 °C was based on leaf mass calculated from trunk volumes for *C. japonica* and *C. obtusa*. The trunk volumes are recorded in a statistical table of timber resources by each prefectoral administration in Japan (Matsunaga, 2011). As shown in Fig. 5, the emission rate of Kaur-16-ene was estimated to be 630 and 3.3 ton h⁻¹, respectively, while the emission rate of total monoterpenes (molecular weight; MW: 136) in Japan were estimated to be 250 tons h⁻¹ at 30 °C and 26 tons h⁻¹ at 5 °C by the emission model MEGAN (Model of Emissions of Gases and Aerosols from Nature) (Guenther et al., 2006). On the other hand, the emission rate based on MEGAN employing observed basal emissions for Kaur-16-ene and the coefficient β was also estimated at 30 °C and 5 °C as 460 and

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

2.7 ton h⁻¹, respectively. It appears that the emission rate of Kaur-16-ene estimated at 30 °C exceeds the total emission rate of anthropogenic VOCs in Japan (220 ton h⁻¹) (Kannari et al., 2008). Considering the result, Kaur-16-ene can be regarded as a new and important atmospheric constituent that has a remarkable influence on the air quality in Japan.

4 Summary

4.1 Diterpene emission

A significant emission of diterpenes, which has not been recognized to be emitted into the atmosphere, has been determined employing both enclosure and ambient air measurements. Their reactivity is expected to be high due to their unsaturated structure and high molecular weight of molecules which contain many hydrogen atoms. In addition, the aerosol formation yield by the oxidation of diterpenes is also expected to be high due to their high molecular weight and reactivity. Considering the comparable emission rate to other classes of BVOCs such as monoterpenes and sesquiterpenes, which are known to be major BVOC constituents, impacts of the diterpene emissions can be important in atmospheric chemistry and physics.

4.2 Unknown diterpene and future study

Samples also contained other diterpene-like compounds, which have not been identified by the authentic standards. One abundant unknown compound in the samples seemed to be Hibaene which is a diterpene also known as a constituent of coniferous leaves (Yatagai et al., 1985; Nagahama and Tajima, 1996). The mass spectrum was similar to that in NIST library. The Hibaene like compound was detected in the samples at similar intensities with Kaur-16-ene. Therefore, total diterpene emission can be larger than presented in this study. Also, to confirm the importance of diterpene

emissions at larger scales, other conifers such as *Abies*, *Larix*, *Picea* and *Pinus* spp., which are common at the global scale, should be investigated for the diterpene emissions in the near future.

BGD

8, 6681–6700, 2011

Acknowledgements. This research was performed commissioned by the Ministry of Economy, Trade and Industry, and as part of a collaborative research program between auto and oil industries called Japan Auto–Oil Program (JATOP) run by Japan Petroleum Energy Center. We also thank staff at the Tanashi Experimental Station of the University of Tokyo and Shiiba Research Forest of Kyushu University for a lot of help provided to the field campaigns.

References

10 Appleton, R. A., McCrindle, R., and Overton, K. H.: The alternative occurrence of (-)-Kaurene or (+)-Phyllocladene in the leaves of *Cryptomeria japonica* D. Don, *Phytochemistry*, 7, 135–137, 1968.

15 Appleton, R. A., McGrindle, R., and Overton, K. H.: The diterpenes from the leaves of *Cryptomeria japonica*, *Phytochemistry*, 9, 581–583, 1970.

20 Beerling, D. J., Hewitt, C. N., Pyle, J. A., and Raven, J. A.: Critical issues in trace gas biogeochemistry and global change, *Phil. Trans. R. Soc. A*, 365, 1629–1642, 2011.

25 Cool, L. G., Hu, Z., and Zavarin, E.: Foliage terpenoids of Chinese *Cupressus* species, *Biochem. Sys. Ecol.*, 26, 899–913, 1998.

30 Di Carlo, P., Brune, W. H., Martinez, M., Harder, H., Lesher, R., Ren, X., Thornberry, T., Carroll, M., Valerie Young, Shepson, P. B., Riemer, D., Apel, E., and Campbell, C.: Missing OH Reactivity in a Forest: Evidence for Unknown Reactive Biogenic VOCs, *Science*, 304, 722–725, 2004.

35 Geron, C. D. and Arnts, R. R.: Seasonal monoterpane and sesquiterpene emissions from *Pinus taeda* and *Pinus virginiana*, *Atmos. Environ.*, 44, 4240–4251, 2010.

40 Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K., and Fall, R.: Isoprene and monoterpane emission rate variability: model evaluations and sensitivity analyses, *J. Geophys. Res.*, 98, 12609–12617, 1993.

45 Guenther, A. B., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.

Kannari, A., Streetsb, D. G., Tonooka, Y., Murano, K., and Baba, T.: MICS-Asia II: An inter-comparison study of emission inventories for the Japan region, *Atmos. Environ.*, 42, 3584–3591, 2008.

Kato, K.: Factors enabling *Epinotia granitalis* (Lepidoptera: Tortricidae) overwintered larvae to escape from oleoresin mortality in *Cryptomeria japonica* trees in comparison with *Semanotus japonicus* (Coleoptera: Cerambycidae), *J. For. Res.*, 10, 205–210, doi:10.1007/s10310-004-0132-9, 2005.

Kavouras, I. G., Mihalopoulos, N., and Stephanou, E. G.: Formation of atmospheric particles from organic acids produced by forests, *Nature*, 395, 683–686, 1998.

Keeling, C. I. and Bohlmann, J.: Diterpene resin acids in conifers, *Phytochemistry*, 67, 2415–2423, 2006.

Kim, S., Guenther, A., Karl, T., and Greenberg, J.: Branch-level measurement of total OH reactivity for constraining unknown BVOC emission during the CABINEX (Community Atmosphere-Biosphere INteractions Experiments)-09 Field Campaign, *Atmos. Chem. Phys. Discuss.*, 11, 7781–7809, doi:10.5194/acpd-11-7781-2011, 2011.

Kulmala, M., Suni, T., Lehtinen, K. E. J., Dal Maso, M., Boy, M., Reissell, A., Rannik, Ü., Aalto, P., Keronen, P., Hakola, H., Bäck, J., Hoffmann, T., Vesala, T., and Hari, P.: A new feedback mechanism linking forests, aerosols, and climate, *Atmos. Chem. Phys.*, 4, 557–562, doi:10.5194/acp-4-557-2004, 2004.

Lee, J.-H., Lee, B.-K., Lee, S. H., and Hong, S.-K.: Comparison of chemical compositions and antimicrobial activities of essential oils from three conifer trees; *Pinus densiflora*, *Cryptomeria japonica* and *Chamaecyparis obtusa*, *J. Microbiol. Biotechnol.*, 19(4), 391–396, doi:10.4014/jmb.0803.191, 2009.

Matsunaga, S. N.: Biogenic VOC emission inventory, JATOP Res. Rep. FY2010, Section 1.4, Japan Petro. Energy Center, Toranomon, Tokyo, 2011 (in Japanese).

Matsunaga, S. N., Mochizuki, T., Ohno, T., Endo, Y., Kusumoto, D., and Tani, A.: Monoterpene and sesquiterpene emissions from Sugi (*Cryptomeria japonica*) based on a branch enclosure measurements, *Atmos. Pollu. Res.*, 2, 16–23, doi:10.5094/APR.2011.003, 2011.

Mustafa, M. G.: Biochemical basis of ozone toxicity, *Free Rad. Biol. Med.*, 9, 245–265, doi:10.1016/0891-5849(90)90035-H, 1990.

Nagahama, S. and Tajima, M.: Diterpene Hydrocarbons from the leaf oil of *Thujopsis dolabrata*

var. dolabrata, *Biochem. Systematics Ecol.*, 24(1), 49–52, 1996.

Nakatsuka, T., Hirose, Y., and Matsuda, M.: Terpenoids. II Composition of the essential oil from leaves of “Sugi” (*Cryptomeria japonica* D. Don), *Mokuzai Gakkaishi*, 3, 206–208, 1957.

Perry, N. B. and Weavers, R. T.: Foliage diterpenes of *Dacrydium Intermedium*: identification, variation and biosynthesis, *Phytochemistry*, 24, 2899–2904, 1985.

Railton, I. D., Fellows B., and West C. A.: Ent-kaurene synthesis in chloroplasts from higher plants, *Phytochemistry*, 23, 1261–1267, 1984.

Sinha, V., Williams, J., Lelieveld, J., Ruuskanen, T. M., Kajos, M. K., Patokoski, J., Hellen, H., Hakola, H., Mogensen, D., Boy, M., Rinne, J., and Kulmala, M.: OH reactivity measurement within a boreal forest: Evidence for unknown reactive emissions, *Environ. Sci. Technol.*, 44, 6614–6620, 2010.

von Rudloff, E.: Volatile Leaf Oil Analysis in Chemosystematic Studies of North American Conifers, *Biochem. Systematics Ecol.*, 2, 131–167, 1975.

Yasue, M., Ogiyama, K., Suto, S., Tsukahara, H., Miyahara, F., and Ohba, K.: Geographical differentiation of natural *Cryptomeria* stands analyzed by diterpene hydrocarbon constituents of individual trees, *J. Japanese Forest Soc.*, 69, 152–156, 1987.

Yatagai, M., Sato, T., and Takahashi, T.: Terpenes of leaf oils from Cupressaceae, *Biochem. Systematics Ecol.*, 13(4), 377–385, 1985.

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 1. Calculated basal emission E_s of Kaurene and total monoterpenes in $\mu\text{g dwg}^{-1} \text{h}^{-1}$, coefficient β and correlation coefficient obtained from seven field measurements conducted at two sites.

	Tree species	Number of individuals	Number of samples	E_s	β	r^2
Tanashi, Tokyo, 2009–2010 ¹						
Kaur-16-ene	<i>C. japonica</i>	3	72	1.05	0.15	0.61
	<i>C. japonica</i> ²			2.79	0.21	
	<i>C. obtusa</i>	1	17	2.20	0.21	0.50
Total monoterpenes	<i>C. japonica</i>	2	90	0.27	0.10	0.64
Shiiba, Miyazaki, 2010 ³						
Kaur-16-ene	<i>C. japonica</i>	2	57	8.74	0.22	0.49
	<i>C. obtusa</i>	2	59	0.99	0.20	0.53
Total monoterpenes ⁴	<i>C. japonica</i>	2	14	0.48	0.10	
	<i>C. obtusa</i>	2	12	0.26	0.10	

¹ The measurements were conducted on January, August, November of 2009 and April 2010.

² Averaged E_s calculated assuming β of 0.21 by the equation of $E_s = E/\exp(\beta(T - T_s))$.

³ The measurements were conducted on May, August and November of 2010.

⁴ Averaged E_s calculated assuming β of 0.10 by the equation of $E_s = E/\exp(\beta(T - T_s))$ measured on May 2010.

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

Title Page

Abstract

Introduction

sions

References

es

Figures

1

1

Ba

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion




Fig. 1. An overview for the branch enclosure sampling system.

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.




Fig. 2. Typical gas chromatogram of samples collected from *C. japonica*.

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.

Scan 1790 (39.760 min): SYCS101104C_01.D\data.ms

Mass spectrum for peak of Kaur-16-ene in sample collected from *C. japonica*

Scan 1788 (39.752 min): KSTD100928B_01.D\data.ms

Authentic standard of Kaur-16-ene

NIST library

(main) Kaurene

Chemical structure of Kaur-16-ene: A tricyclic diterpene with a cyclohexenyl group at C16.

Fig. 3. Structure and typical sample mass spectrum of Kaur-16-ene, analysis of the authentic standard, and NIST library.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

18

1

Bac

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion




Fig. 4. A liner relationship between the temperature and natural logarithm for the measured emission rate of Kaur-16-ene collected from (a) *C. japonica* and (b) *C. obtusa* (only for Shiiba).

Title Page

Abstract

Introduction

Conclusion

References

Tables

Figures

1

Back

Close

Full Screen / Esc

Volatile diterpene emission from dominant conifers in Japan

S. N. Matsunaga et al.




Fig. 5. Estimated emission rate of Kaur-16-ene at 30 °C over Japan.