

Interactive
Comment

Interactive comment on “Stand age and tree species affect N₂O and CH₄ exchange from afforested soils” by J. R. Christiansen and P. Gundersen

Anonymous Referee #1

Received and published: 25 July 2011

General comments The paper by Christiansen and Gundersen presents interesting results on the effect of tree species and stand age on the exchange of nitrous oxide and methane between forest soil and the atmosphere. The study is well planned and in most parts the interpretation of results is sound. However, I have concerns about some of the conducted statistical analyses that have to be addressed and I would like to suggest refocusing parts of the paper (see below).

One of the objectives of the study was to “identify the environmental factors responsible for the differences in GHG exchange”, which is approached by regression analyses. However, from my point of view several of the regression analyses are not appropriate.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Firstly, the regression between the annual N₂O emissions and the %NO₃ (percentage of NO₃- of total mineral N) is inappropriate as the %NO₃ was only measured once at the end of the two year flux measurement period. Given that the concentrations of mineral N as well as the ratio between NO₃- and NH₄⁺ can vary during the course of a year as well as between years, the representativeness of the measured %NO₃ can be questioned. Secondly, to support the relation between N₂O fluxes and %NO₃ the authors further conducted regressions between the annual N₂O fluxes and NO₃- leaching as well as sub-root NO₃- concentrations measured over a 3-yr period by Hansen et al (2007). However, these data came from the period 2000-2002, while N₂O measurements were conducted 2008-2010, questioning the appropriateness of the regression analysis. It should be noted that the authors acknowledge the weakness of these regressions. However, from my point of view all the mentioned regressions should be removed from the paper.

Regarding the result section I would like to suggest an alternative structure of the subsections as follows: 3.1 Abiotic soil properties: This could include the text presented p. 5739, line 22 – p. 5740, line 7 and p. 5741, line 8 – p. 5742, line 14, with potentially shortening of some parts. 3.2 N₂O exchange: containing p. 5738, line 24 – p. 5739, line 10 and p. 5740, line 9 – 18. 3.3 CH₄ exchange: containing p. 5739, line 11 – 21 and p. 5740, line 19 – p. 5741, line 3. 3.4 Relation between GHG exchange and soil properties: containing p. 5742, line 15 – p. 5743, line 3 (but excluding the above mentioned regressions!) as well as p. 5740, line 15-16 and p. 5741, line 4-7.

Please check the paper carefully for phrasings that indicate that the N₂O emissions or CH₄ uptake are by the trees (e.g. CH₄ oxidation in oak), as the exchange is between soil and atmosphere.

The GHG fluxes were measured in each stand in three plots, that each consists of three chambers. It is not clear if for statistics an n=3 (for plots) or an n=9 (for chambers) was used. It is, moreover, somewhat confusing that results were partly presented as stand averages, but partly as chamber fluxes, which also resulted in some repetitions. I

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

suggest focusing on the stand average fluxes and discuss individual chamber fluxes only in respect to spatial variability.

Specific comments p. 5730, l. 17: could you specify which physico-chemical properties? p. 5731, l. 4-6: suggest moving this sentence to after the next paragraph p. 5732, l. 1ff: I think the cited study from Höglwald (Papen and Butterbach-Bahl, 1999) is specific as the spruce forest exhibit about four-time higher NO than N₂O fluxes (see Gasche and Papen, 1999). Consequently, the total NO+N₂O emission is higher from the spruce compared to the beech site. This may deserve mentioning here. p. 5732, l. 10-12: needs rephrasing p. 5732, l. 12: write “deciduous” instead of “hardwood” p. 5732, l. 23: replace “so” with “in order” p. 5732, l. 25: it is not clear if you mean that N₂O emissions increase or decrease. Please rephrase. p. 5736, l. 10: Unclear what is meant by “due to limitation of concentration data points during enclosure”. p. 5736, l. 20: How many soil cores were taken? p. 5737, l. 18: How did you deal with negative fluxes when log-transforming the data? p. 5739, l. 5-7: repetition from p. 5738, l. 27ff. p. 5739, l. 11: should read “ $\mu\text{g CH}_4\text{-C}$ ”? p. 5739, l. 12: “... only above zero during winter and early spring” seems to contrast with line 16-17 “Emission of CH₄ was observed ... predominantly in spring and autumn” p. 5739, l. 18: as maximum emission a value of $66 \mu\text{g CH}_4\text{-C m}^{-2} \text{ h}^{-1}$ is given, but line 11 reads that “CH₄ fluxes ranged between -30 and 9”. And why was not maximum CH₄ uptake discussed, as more often uptake then emission was observed? p. 5739, l. 24-25: Suggest: “Soil water content ranged from 44 to 6 vol%”. I don’t think that much information is provided by mentioning in which plot the extremes were measured, as it seems that it is just by chance that highest moisture was in O-70 and not O-93, as well as lowest moisture in S-69 and not S-97. p. 5739, l. 26-29: Suggest deleting whole sentence from “Only ...” p. 5740, l. 11: Suggest “average values” for consistency. p. 5741, l. 2: Delete “and” after significantly p. 5741, l. 17: The statement “even though soil temperature increased with stand age in oak and decreased in Norway Spruce” appears to be incorrect, as Fig. 3b indicate that both these changes were non significant! p. 5741, l. 19: Delete “marginal” p. 5744, l. 7: “nitrous oxide reductase” p. 5743, l. 9: Suggest to only pro-

C2150

BGD

8, C2148–C2153, 2011

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



vide mean and SE. p. 5744, l. 11ff: One additional aspect of the lower diffusivity in the younger stands is that the retention time of N₂O in the soil is prolonged, which in turn increases the likelihood for a reduction to N₂ and hence can reduce the N₂O emissions. p. 5745, l. 5ff: The difference in C/N ratio between the stands at Vestskoven is rather small (10.6 – 14.6) compared to the range for European forest (13.4 – 37.7; Pilegaard et al. 2006) as well as organic forest soils (13 – 90; Klemetsson et al. 2005). Moreover, the C/N ratio at Vestskoven is for all four stands in the lower range of the European compilations, indicating the potential for significant N₂O emissions. As the C/N ratio is only a general proxy for N₂O emissions, one can question its applicability for comparing the plot N₂O fluxes. p. 5745, l. 8: Suggest “higher net nitrification” p. 5745, l. 13: It is not clear if you mean net or gross nitrification rates. Moreover, I do not see the functional relationship between C/N ratio and nitrification (particularly not for autotrophic nitrification). Why should nitrification be higher at lower C/N ratios? I also think that gross nitrification may actually be lower in the younger stands, as the competition for NH₄⁺ may be higher. p. 5745, l. 17ff: I think that the paragraph on changing plant N demand after afforestation should be more highlighted, as from my point of view this may be at least as important as the changes in physico-chemical soil properties. p. 5748, l. 18-19: Suggest: “CH₄ uptake increased in oak but remained constant in Norway Spruce stands” (see statistics). p. 5748, l. 24: Suggest adding that low N availability is due to high plant N demand. p. 5749, l. 5: You have not talked earlier about the N cycling, hence it should not be included in the conclusions. However, it may be helpful to discuss earlier your results in the light of differences in N transformations between coniferous and broad-leaf forest soils, which was the subject to several studies (e.g. Brüggemann et al., 2005; Christenson et al., 2009; Staelens et al., 2011; Ste-Marie and Houle, 2006; Zeller et al., 2007).

Fig. 5: Irregardless of my general doubts to perform this regression (see above) it appears that the correlation between N₂O flux and soil water NO₃⁻ is almost exclusively due to one of the twelve plots. Therefore, such a regression has to be interpreted with care. Table S-1: Plots are here labelled A – C, while in Table 1 by numbers (1 – 3).

C2151

BGD

8, C2148–C2153, 2011

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Please be consistent.

Cited references: - Brüggemann, N., Rosenkranz, P., Papen, H., Pilegaard, K., and Butterbach-Bahl, K.: Pure stands of temperate forest tree species modify soil respiration and N turnover, *Biogeosciences Discussions*, 2, 303-331, 2005. - Christenson, L. M., Lovett, G. M., Weathers, K. C., and Arthur, M. A.: The influence of tree species, nitrogen fertilization, and soil C to N ratio on gross soil nitrogen transformations, *Soil Science Society of America Journal*, 73, 638-646, 2009. - Gasche, R., and Papen, H.: A 3-year continuous record of nitrogen trace gas fluxes from untreated and limed soil of a N-saturated spruce and beech forest ecosystem in Germany 2. NO and NO₂ fluxes, *Journal of Geophysical Research*, 104, 18505-18520, 1999. - Hansen, K., Rosenqvist, L., Vesterdal, L., and Gundersen, P.: Nitrate leaching from three afforestation chronosequences on former arable land in Denmark, *Glob. Change Biol.*, 13, 1250-1264, 2007. - Klemetsson, L., von Arnold, K., Weslien, P., and Gundersen, P.: Soil CN ratio as a scalar parameter to predict nitrous oxide emissions, *Global Change Biology*, 11, 1142-1147, 2005. - Papen, H., and Butterbach-Bahl, K.: A 3-year continuous record of nitrogen trace gas fluxes from untreated and limed soil of a N-saturated spruce and beech forest ecosystem in Germany 1. N₂O emissions, *Journal of Geophysical Research*, 104, 18487-18503, 1999. - Pilegaard, K., Skiba, U., Ambus, P., Beier, C., Brüggemann, N., Butterbach-Bahl, K., Dick, J., Dorsey, J., Duyzer, J., Gallagher, M., Gasche, R., Horvath, L., Kitzler, B., Leip, A., Pihlatie, M. K., Rosenkranz, P., Seufert, G., Vesala, T., Westrate, H., and Zechmeister-Boltenstern, S.: Factors controlling regional differences in forest soil emission of nitrogen oxides (NO and N₂O), *Biogeosciences*, 3, 651–661, 2006. - Staelens, J., Rütting, T., Huygens, D., De Schrijver, A., Müller, C., Verheyen, K., and Boeckx, P.: In situ gross nitrogen transformations differ between temperate deciduous and coniferous forest soils, *Biogeochemistry*, DOI: 10.1007/s10533-10011-19598-10537, 2011. - Ste-Marie, C., and Houle, D.: Forest floor gross and net nitrogen mineralization in three forest types in Quebec, Canada, *Soil Biology & Biochemistry*, 38, 2135-2143, 2006. - Zeller, B., Recous, S., Kunze, M., Moukouri, J., Colin-Belgrand, M., Bienaimé, S., Ranger, J., and Dambrine, E.: Infl-

C2152

BGD

8, C2148–C2153, 2011

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



ence of tree species on gross and net N transformations in forest soil, *Annals of Forest Science*, 64, 151-158, 2007.

Interactive comment on *Biogeosciences Discuss.*, 8, 5729, 2011.

BGD

8, C2148–C2153, 2011

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C2153

