

Interactive
Comment

Interactive comment on “Tracing the transport of colored dissolved organic matter in water masses of the Southern Beaufort Sea: relationship with hydrographic characteristics” by A. Matsuoka et al.

Anonymous Referee #2

Received and published: 6 January 2012

Tracing the transport of colored dissolved organic matter in water masses of the Southern Beaufort Sea: relationship with hydrographic characteristics A Matsuoka et al.

General Comments: The paper investigates light absorption properties of colored dissolved organic matter (CDOM) in the Southern Beaufort Sea. The paper adds to a growing body of CDOM and DOC studies in the Arctic Ocean and the impact of Rivers such as the Mackenzie discharging into the Arctic This is an excellent study of CDOM optical properties (absorption and spectral slope) and DOC in relationship to hydrography and linkages to the various Arctic water masses; the study also supports the

C5192

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



potential use of CDOM as a tracer of water masses in the Arctic. Further, observed CDOM-salinity and CDOM-DOC relationships in surface waters could be used to study carbon budgets using ocean color remote data in the study region. This is a well written manuscript; my concerns with the manuscript are listed below by a few comments or questions that can be addressed in the revised version:

Specific comments: i) A surface layer of river plume water with a conservative CDOM-salinity relationship is observed to be dispersed over a large area of the Southern Beaufort Sea. However, this large riverine CDOM footprint in the shallow surface layer (0 – 20m) of the Southern Beaufort Sea would likely be impacted by variable river discharge, wind field and circulation features in the region. Summer variability in the intensity and frequency of the northeasterly (upwelling-favorable) winds that generally prevail in the region (MacDonald et al., 1999) would most likely impact the horizontal extent of the dispersal, surface mixing and the sea-ice melt conditions that would likely change the dynamics of the relationship between CDOM, DOC and salinity (Figures 1, 8). With a large fraction of the data points obtained at stations along transects with greatest influence of the Mackenzie River plume (Figure 1), a conservative CDOM-salinity relationship in the UPML is more likely (Figure 5). However, it is possible that the fraction of freshwater due to ice-melt in surface waters could change under variable physical influences thus biasing ocean color derived estimates of CDOM or DOC using the relationships derived in this study. It is pertinent to address this issue in the revised paper.

ii) Page 11012, line 25: “At salinity < 28 (in the UPML) aCDOM(440) values showed a strong and negative correlation with salinity except for some stations where waters were influenced by sea ice melt (dotted circle in Fig 5a).” It would helpful to know what % of data points were influenced by sea ice melt.

iii) Page 11020, line 16: “In this study, we found that DOC concentrations were highly correlated with aCDOM(440) in the UPML ($r^2=0.97$). This result suggests that when aCDOM(440) values are obtained using ocean color remote sensing, DOC concentra-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

tions can be also estimated from satellite.” Please provide the number stations used in obtaining the correlation shown in Figure 8. Do the data points used also include stations influenced by sea ice melt?

iv) An additional source of CDOM/DOC in the Arctic Ocean is likely due to biological processes associated with polar phytoplankton blooms within the Arctic Ocean. Figure 3 shows a substantial phytoplankton fluorescence peak. This bloom would likely be much stronger earlier in the spring in the near surface waters. It might be worth commenting on the potential for autochthonous CDOM contribution associated with the large phytoplankton production generally observed during the early stages of the polar blooms or the subsurface phytoplankton peaks observed in the water column during the field campaign.

Reference: Macdonald, R.W., E.C. Carmack, F.A. McLaughlin, K.K. Falkner and J.H. Swift. 1999. Connections among ice, runoff, and atmospheric forcing in the Beaufort Gyre. *Geophys. Res. Lett.* 26(15):2223–2226. doi:10.1029/1999GL900508.

Interactive comment on Biogeosciences Discuss., 8, 11003, 2011.

BGD

8, C5192–C5194, 2012

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

