



BGD

8, C5208–C5209, 2012

Interactive Comment

Interactive comment on "Carbon dynamics in the western Arctic Ocean: insights from full-depth carbon isotope profiles of DIC, DOC, and POC" by D. R. Griffith et al.

R. Benner (Referee)

benner@mailbox.sc.edu

Received and published: 9 January 2012

The authors present novel and interesting data on the isotopic compositions of the major forms of carbon in the Canada Basin. There are many potential uses for these data, and the authors highlight one particularly interesting observation - the potential for an isotopic signature of chemolithoautrophic carbon fixation at intermediate depths in the basin. Indications of chemolithoautotrophy have been observed at intermediate depths in other ocean basins, so the authors' observations are consistent with the growing recognition of this process of carbon fixation in the ocean interior.

The low DO14C values in the deep Canada Basin are an interesting feature that the





authors do not discuss. The DO14C in the deep Canada Basin is depleted (-450‰) in comparison to values for the deep North Atlantic (-390‰ Bauer 2002). It appears this could be due to aging in the Canada Basin, which is consistent with the increase in age of DIC in basin deep waters. However, the DO13C in the deep Canada Basin is also depleted (-23‰ in comparison to the deep North Atlantic (-21‰ Bauer 2002). This suggests a different source of DOC could also be impacting its DO14C content. It would be useful for the authors to expand on this and make a more quantitative comparison between the depletion of radiocarbon in DOC and DIC in the Canada Basin.

Specific comment: Pg 10687, line 9 - the rate of carbon fixation should include the unit of time (3-16 μmol C m-3 y-1)

BGD

8, C5208–C5209, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive comment on Biogeosciences Discuss., 8, 10677, 2011.