

Regional differences in modelled net production and shallow remineralization in the North Atlantic subtropical Gyre

B. Fernández-Castro¹, L. Anderson², E. Marañón¹, S. Neuer³, B. Ausín¹,
M. González-Dávila⁴, J. M. Santana-Casiano⁴, A. Cianca⁴, R. Santana⁴,
O. Llinás⁵, M. J. Rueda⁶, and B. Mouriño-Carballido¹

¹Departamento de Ecología e Bioloxía Animal, Universidade de Vigo, Vigo, Pontevedra, 36200, Spain.

²Applied Ocean Physics and Engineering Department, Woods Hole Oceanographic Institution, Woods Hole, MA 02543-1541, USA.

³School of Life Sciences, Arizona State University, Tempe, AZ 85287-4501, United States.

⁴Universidad de Las Palmas de Gran Canaria (ULPGC), Campus Universitario de Tafira s/n. 35017 Las Palmas de Gran Canaria, Spain.

⁵Plataforma Oceánica de Canarias (PLOCAN), Crta. de Taliarte s/n, P.O. Box 413, 35200 Telde, Gran Canaria, Spain

⁶Instituto Canario de Ciencias Marinas, Gobierno de Canarias, Telde, Gran Canaria, Spain.

Correspondence to: B. Fernández-Castro
(bieito.fernandez@uvigo.es)

Abstract. We used 5-year concomitant data of tracer distribution from the BATS (Bermuda Time-series Study) and ESTOC (European Station for Time-Series in the Ocean, Canary Islands) sites to build a 1-D tracer model conservation including horizontal advection and compute net production and shallow remineralization rates at both sites. Our main goal was to verify if differences in these rates are consistent with the lower export rates of particulate organic carbon observed at ESTOC. Net production rates computed below the mixed layer to 110 m from April to December for oxygen, dissolved inorganic carbon and nitrate at BATS ($1.34 \pm 0.79 \text{ mol O}_2 \text{ m}^{-2}$, $-1.73 \pm 0.52 \text{ mol C m}^{-2}$ and $-125 \pm 36 \text{ mmol N m}^{-2}$) showed no statistically significant differences compared to ESTOC ($1.03 \pm 0.62 \text{ mol O}_2 \text{ m}^{-2}$, $-1.42 \pm 0.30 \text{ mol C m}^{-2}$ and $-213 \pm 56 \text{ mmol N m}^{-2}$). Shallow remineralization rates between 110 and 250 m computed at ESTOC ($-3.9 \pm 1.0 \text{ mol O}_2 \text{ m}^{-2}$, $1.53 \pm 0.43 \text{ mol C m}^{-2}$ and $38 \pm 155 \text{ mmol N m}^{-2}$) were statistically higher for oxygen compared to BATS ($-1.81 \pm 0.37 \text{ mol O}_2 \text{ m}^{-2}$, $1.52 \pm 0.30 \text{ mol C m}^{-2}$ and $147 \pm 43 \text{ mmol N m}^{-2}$). The divergence in the lateral transport of tracers, which was more significant at ESTOC, was responsible for the differences in estimated oxygen remineralization rates between both stations. According to these results, the differences in net production and shallow remineralization are not consistent with the differences in the flux of sinking organic matter observed between both stations, suggesting a convergence in the horizontal advection of slowly sinking, suspended or dissolved organic matter at ESTOC.

1 Introduction

20 The ocean is responsible for an annual photosynthetic fixation of ~ 50 Pg of carbon which represents
around half of the global primary production (Field et al., 1998). Carbon fixation by marine phyto-
plankton and its transport to the deep ocean, known as the biological carbon pump, plays a key role
in the ocean-atmosphere CO_2 exchange, and hence affects climate on time-scales from decades to
25 thousands of years. The carbon pump efficiency is not constant and depends on several factors such
as the input of nutrients into the euphotic layer, the balance between synthesis and remineralization
of organic matter or the planktonic community composition.

Subtropical gyres represent the central part of the ocean and are characterized by a strong strat-
ification of the water column. This translates into a weak input of nutrients into the euphotic layer
and, as a consequence, low phytoplankton biomass and productivity. Because of their oligotrophic
30 characteristics they have been traditionally considered oceanic deserts. However, from the point of
view of carbon export from the euphotic zone, subtropical gyres were found to be surprisingly pro-
ductive (Emerson et al., 1997). This fact, combined with their vast extension ($\sim 60\%$ of the total
ocean surface), results in their contribution of at least 30% of total marine new production (Najjar
and Keeling, 2000).

35 The North Atlantic subtropical gyre (NASG) is one of the best studied open ocean regions, and
in the past decades, it has been a major contributor to the development of our understanding of
biogeochemical cycles in subtropical regions. Traditionally subtropical gyres have been considered
relatively constant ecosystems in time and space. However, the comparative study of time-series
data revealed that spatial heterogeneity is important in understanding the biogeochemistry of this
40 biome (Neuer et al., 2002a; Mouriño-Carballido and Neuer, 2008). Two time-series stations, BATS
(Bermuda Atlantic Time-series Study, 31.7°N - 64.2°W) and ESTOC (European Station for Time
series in the Ocean, Canary Islands, 29.16°N - 15.5°W), are located at about the same latitude in
the western (NASW) and eastern (NASE) portions of NASG. The monthly sampling program was
initiated in October 1988 at BATS and in February 1994 at ESTOC, with the aim of tracking sea-
45 sonal, inter-annual and inter-decadal changes in several biogeochemical variables (Steinberg et al.,
2001; Neuer et al., 2007). Although both stations exhibit typical oligotrophic characteristics, they
are characterized by different hydrographic dynamics. BATS is located in the recirculation of the
Gulf Stream and it is strongly affected by mesoscale activity (Cianca et al., 2007). ESTOC is en-
trained by the Canary Current and is indirectly influenced by the coastal African upwelling, which
50 export nutrients and organic matter towards the center of the gyre by means of upwelling filaments
and Ekman transport (Pelegrí et al., 2005; Álvarez-Salgado et al., 2007). It is important to note that
upwelling filaments do not influence ESTOC directly (Davenport et al., 2002), and this station is
considered oligotrophic based on nutrient scarcity and phytoplankton biomass and production rates
(Neuer et al., 2007).

55 Although both stations are characterized by similar phytoplankton biomass and primary produc-

tion rates, annual mean carbon export measured by using sediment traps at 150, 200 and 300 m is significantly lower at ESTOC, by a factor of 3–5, than at BATS (Neuer et al., 2002a; Helmke et al., 2010). Assuming that the euphotic zone is in steady state, higher export production should be sustained by larger inputs of nutrients and/or organic matter into the euphotic zone. It has been proposed that differences in the nutrients input could be due to the more intense mesoscale activity observed at BATS (Siegel et al., 1999; Mouriño et al., 2003). A comparative study using 10-year BATS and ESTOC data combined with satellite altimetry data indicated that the higher physical forcing dominant at BATS, deeper mixed layers and more intense mesoscale dynamics, is partly compensated by the shallower nutricline observed at ESTOC (Cianca et al., 2007). According to these authors, NASE receives ~75% of the nutrients available for new production at NASW, although this difference is not statistically significant. Another source of new nitrogen that could explain the observed differences in carbon export between the two stations is biological nitrogen fixation. Recent studies indicate that this process has a significant contribution to total new nitrogen inputs in oligotrophic waters (Capone et al., 2005; Mouriño-Carballido et al., 2011; Bonnet et al., 2011). Biogeochemical estimates suggest a lower contribution of nitrogen fixation at ESTOC compared with BATS (Neuer et al., 2002a).

The differences observed between the two stations in export rates of particulate organic carbon could also be a consequence of differences in the remineralization rates of the sinking organic matter. The comparative analysis of respiration rates determined from in vitro oxygen evolution experiments conducted in the NASE and NASW indicated higher oxygen consumption rates in the eastern part (Mouriño-Carballido and Neuer, 2008). It has been shown that atmospheric dust deposition can strongly stimulate bacterial respiration (Pulido-Villena et al., 2008). A recent study demonstrated that the most frequent and intense response of the microbial plankton to atmospheric dust deposition in the Tropical Atlantic is the stimulation of the bacterial activity rather than phytoplankton primary production (Marañón et al., 2010). This is consistent with the results found at ESTOC, which is strongly influenced by the natural atmospheric deposition of Saharan dust and, where phytoplanktonic production seems not to be affected by aerosol inputs (Neuer et al., 2004).

For the first time, concomitant data of tracer distribution from the BATS and ESTOC time-series sites were analysed and used to build up a 1D tracer conservation diagnostic model. The model was used to compute net production and shallow remineralization rates at both sites. The main goal of this study was to verify if differences in the synthesis and consumption of organic matter could explain the lower export rates of particulate organic carbon observed at ESTOC.

2 Comparison of the seasonal cycles at BATS and ESTOC

Monthly climatologies were calculated for temperature, salinity, density anomaly (σ_T), oxygen, dissolved inorganic carbon (DIC), nitrate (actually nitrate+nitrite) and chlorophyll *a*, using the BATS

and ESTOC data collected for the period 1996-2001. The BATS data were obtained from the BATS Web site (<http://bats.bios.edu>). BATS and ESTOC measurements were made monthly except during the spring bloom period at BATS (February-April) when biweekly samplings were conducted. Methodologies and results have been reported earlier for BATS (Michaels and Knap, 1996; Steinberg et al., 2001) and ESTOC (Neuer et al., 2007). Discrete data were first interpolated to a grid with the following depths: 5, 10, 20, 40, 60, 80, 100, 110, 120, 140, 160, 200 and 250 m. The mean annual cycle was then constructed by averaging data from each depth onto a temporal grid of 15 days using a weighted moving average with a normal weighting factor, f_i^n , and a window of 50 days:

$$\langle C^n \rangle = \frac{\sum_i f_i^n C_i}{\sum_i f_i^n} \quad (1)$$

$$f_i^n = \exp \left[- \left(\frac{t^n - t_i}{\sigma} \right)^2 \right]$$

where C is the averaged variable, t is the time of the year in days (0-365), i is the index of the discrete data point, n is the index for the temporal grid and $\sigma = 35$ days is a constant time scale. The mixed layer depth was calculated as the depth where temperature differs 0.1°C from the 10 m value.

BATS and ESTOC exhibit a similar seasonality in the hydrographic conditions characterized by a period of winter mixing from November to March followed by a period of summer stratification from May to September (Figure 1). The maximum winter mixing occurs later (mid-March) at BATS compared to ESTOC (February). The transition between both periods is controlled by the seasonal cycle of solar irradiation and changes in the wind speed. Seasonality is more intense at BATS where surface temperature increases from a minimum of $\sim 20^\circ\text{C}$ in mid-March to a maximum of $\sim 28^\circ\text{C}$ in August. At ESTOC temperature reaches a minimum of $18 - 19^\circ\text{C}$ in February and a maximum of $\sim 24^\circ\text{C}$ in September. This maximum appears later than at BATS because the intensification of the Trade Winds affects the ESTOC site during the summer period. Maxima mixed layers are deeper at BATS (193 ± 26 m) than at ESTOC (146 ± 32 m). Summer stratification is also more intense at BATS, where mixed layers during this period reach ~ 20 m. Mixed layers at ESTOC extend to a depth of ~ 50 m due to the Trade Winds intensification that enhances mixing during July-August. The main difference between both stations in salinity appears during the summer, when mixed layer values are fresher at BATS due to the rain intensification.

Chlorophyll concentration is strongly influenced by the physical forcing at both stations. Phytoplankton blooms occur in late winter/spring, after the mixed layer extends below the nutricline, with greatest monthly values of chlorophyll concentration similar for both stations (ca. 0.3 mg m^{-3}). As water column stratification increases, surface chlorophyll decreases and a deep chlorophyll maximum establishes at ca. 100 m at BATS and ESTOC. This biological activity as well as the physical forcing determine the concentration of the chemical tracers considered in this study.

The oxygen seasonal cycle in the mixed layer at both stations is characterized by maximum values during the winter period due to the increase in solubility. The temperature increase during the summer months causes a drop in solubility and hence in surface oxygen concentration. As stratification increases during summer, oxygen accumulates below the mixed layer (~ 40 m at BATS and ~ 60 m

at ESTOC). This accumulation is clearly pictured by the oxygen anomaly distribution that represents the excess of oxygen concentration above the solubility equilibrium. The oxygen anomaly distribution shows that whereas the photic layer (ca. 100 m at both sites, Cianca et al. (2007)) at ESTOC is oversaturated during the whole year, at BATS it is undersaturated during the winter period. This difference is probably due to the different intensity in the mixing and ventilation of deep waters that characterizes both stations. The seasonal variability in the oxygen cycle is also weaker at ESTOC, where the maximum amplitude of the accumulation in oxygen anomaly below the mixed layer was $\sim 9 \text{ mmol m}^{-3}$ (from February to August), versus $\sim 17 \text{ mmol m}^{-3}$ (from February to September) at BATS. The oxygen anomaly distribution in the euphotic zone has a mirror-like image below this layer. Oxygen concentration increases during winter-spring as a result of the mixing with oxygen-rich surface waters and it is maximum in April-May. Then, during summer stratification oxygen concentration decreases, with the minimum values observed in January (BATS) and November (ESTOC) at 200 m. Despite the differences observed in the euphotic zone, the maximum amplitude of the oxygen signal at 200 m ($\sim 7 \text{ mmol m}^{-3}$) is similar in both stations.

DIC concentrations in the mixed layer are greatest in April (BATS) and March (ESTOC), due to mixing with deeper waters and, to a lesser extent, to air-sea exchange (Marchal et al., 1996; Gruber et al., 1998; González-Dávila et al., 2003). Surface DIC concentrations decline to minimum values in October, this decline is more pronounced at BATS ($\sim 44 \text{ mmol m}^{-3}$) compared to ESTOC ($\sim 15 \text{ mmol m}^{-3}$). Below the euphotic zone DIC accumulation occurs in concert with the observed oxygen decline. The DIC accumulation at 200 m from April to January (BATS) and from April to October (ESTOC) is 6 mmol m^{-3} and 8.7 mmol m^{-3} , respectively.

Nitrate concentration is very low in the euphotic zone at both stations. The nitracline ($> 0.5 \text{ mmol m}^{-3}$) is generally shallower at ESTOC ($90.3 \pm 5.0 \text{ m}$) compared to BATS ($104 \pm 14 \text{ m}$) but with a greater vertical variability at the western station. As a consequence of the influence of the nutrient poor 18°C Subtropical Mode Water at BATS, nitrate concentration below the nitracline is higher at ESTOC. Below the euphotic zone nitrate accumulates from June to January at BATS and from May to December at ESTOC, the amplitude of the accumulation at 200 m is very similar in both stations (0.7 mmol m^{-3}).

The seasonal variability of the tracers considered in this study, in agreement with previous reports for BATS (Menzel and Ryther, 1959; Jenkins and Goldman, 1985; Marchal et al., 1996) and ESTOC (González-Dávila et al., 2003; Neuer et al., 2007; Santana-Casiano et al., 2007), is consistent with the fact that synthesis of organic matter occurs in the euphotic layer and that this matter is remineralized, at least partially, in the shallow aphotic zone between 100-250 m. In order to quantify the contribution of the biological processes and the physical forcing to the observed seasonal variability of the chemical tracers, a 1-D diagnostic model was implemented.

3 Model implementation

3.1 Model general description

We adapted the tracer conservation model based on Ono et al. (2001), who estimated the shallow
 165 remineralization at BATS for the period 1992-1998. This approach includes the main physical pro-
 cesses that occur below the mixed layer when convective winter mixing is not the dominant process.
 These processes are vertical diffusion, vertical advection (Ekman transport) and lateral advection.
 The used tracer conservation equation was

$$\frac{\partial C}{\partial t} = -u \frac{\partial C}{\partial x} - v \frac{\partial C}{\partial y} - w \frac{\partial C}{\partial z} + K \frac{\partial^2 C}{\partial z^2} + J_C \quad (2)$$

170 where $C = C(t, z)$ represents temperature (T) or the tracer (oxygen, DIC, nitrate) concentration for
 each depth, z , and time t ; $u(t, z)$ and $v(t, z)$ are the longitudinal and latitudinal geostrophical veloc-
 ities, respectively; $\partial C / \partial x$ and $\partial C / \partial y$ are the longitudinal and latitudinal gradients of temperature
 and tracer concentration; $K(t)$ is the vertical diffusivity and $J_C(t, z)$ represents the sources minus
 sinks term. For temperature J_C represents the effect of the solar shortwave radiation that penetrates
 175 below the mixed layer depth, whereas for oxygen, DIC and nitrate it represents the net effect of
 photosynthesis and respiration, therefore net production and shallow remineralization.

The model was executed 23 times for each variable in periods of 15 days during the seasonal
 cycle, with a time step $dt = 0.005$ days and an uniform vertical grid ($dz = 1$ m) from the mixed
 layer depth to 250 m. In each of these runs the model was initialized with the profile of the
 180 tracer for the initial time t , $C^{obs}(t, z)$, and produces the modelled profile for the final time $t + \Delta t$,
 $C^{model}(t + \Delta t, z)$, where $\Delta t = 15$ days. For the biological tracers J_C was computed diagnostically,
 as described by Mourino-Carballido and Anderson (2009), so that the output of the model for each
 run fits the profile for the corresponding final time. An initial guess for the $J_C(t, z)$ term was made
 as $(C_{obs}(t + \Delta t, z) - C_{obs}(t, z)) / \Delta t$. The model was run forward from time t to time $t + \Delta t$ and a
 185 mean squared misfit was computed as:

$$Cost = \left(\int_{MLD}^{250m} (C_{obs}(t + \Delta t, z) - C_{mod}(t + \Delta t, z))^2 dz \right)^{1/2}. \quad (3)$$

When the difference of $Cost$ with the previous estimate was less than $0.002 \text{ mmol m}^{-2}$ the simula-
 tion was ended and the simulation of the next 15 day period was initiated. Otherwise, $J_C(t, z)$ was
 corrected as:

$$190 \quad J_C^{new}(t, z) = J_C(t, z) + 0.75(C_{obs}(t + \Delta t, z) - C_{mod}(t + \Delta t, z)) / \Delta t, \quad (4)$$

and a new run was conducted. In this way $J_C(t, z)$ was optimized so that $C^{mod}(t + \Delta t, z)$ fits
 $C^{obs}(t + \Delta t, z)$.

In order to compute the biological source term (J_C), tracer data, horizontal and vertical velocities,
 were calculated as described below and linearly interpolated to the model grid.

3.2.1 Horizontal advection

Horizontal gradients of temperature, oxygen and nitrate were calculated using the four grid points surrounding BATS and ESTOC at the World Ocean Atlas 2009 monthly climatology (WOA09) (Locarnini et al., 2010; Garcia et al., 2010b,a). Longitudinal and latitudinal gradients were computed using averaged values for each x and y components. DIC data for the same locations were obtained from the Global Distribution of Total Inorganic Carbon and Total Alkalinity Below the Deepest Winter Mixed Layer Depths climatology (Goyet et al., 2000), which includes lower temporal resolution trimonthly averaged data.

Horizontal velocities, u and v , were assumed to be geostrophic and computed from the monthly temperature and salinity WOA09 data (Locarnini et al., 2010; Antonov et al., 2010) and the thermal wind equations. Water density was calculated using the Millero and Poisson (1981) parametrization. The thermal wind equations were integrated down to 3000 m, which was assumed as the level of no motion following Siegel and Deuser (1997). The calculated geostrophic flow for BATS ($0.02 - 0.04 \text{ m s}^{-1}$) was directed southwest during the whole year (see Figure 2), consistent with the fact that BATS is influenced by the Gulf Stream recirculation (Siegel and Deuser, 1997; Ono et al., 2001). The flow intensity was more variable near ESTOC ($0.02 - 0.07 \text{ m s}^{-1}$), where the current was directed southwest during most of the year except between March and April, when u was eastward, and during the transition from spring to summer (May-July) when v was northward. This pattern is consistent with the results described by Neuer et al. (2007) and Pelegrí et al. (2005).

Due to the implementation imposed to ensure volume conservation (see next section), lateral transport included a non-geostrophic component. Further on we will use geostrophic horizontal advection to refer to the geostrophic component of the lateral transport and horizontal advection to refer to the total corrected lateral transport.

Additionally, model runs where the geostrophic transport term was set to zero ($u = v = 0$) were used to determine the influence of the geostrophic transport in the computed biological rates.

3.2.2 Vertical advection

Ekman downwelling/upwelling velocity, w , was computed for both stations from the wind stress monthly climatological data included in the International Comprehensive Ocean-Atmosphere Data Set, with a spatial resolution of $2^\circ \times 2^\circ$ (Leetmaa and Bunker, 1978). The computed Ekman velocity at BATS was negative during most of the year and characterized by a clear seasonal cycle (see Figure 3a). Downwelling was maximum in February (-96 m yr^{-1}) and minimum in September, when a weak upwelling was computed (10 m yr^{-1}). These results are in agreement with the harmonic function used by Musgrave et al. (1988) and Ono et al. (2001) at BATS. The Ekman velocity was lower at ESTOC where the seasonal cycle was characterized by relatively weak upwelling during

230 the summer (ca. 20 m yr⁻¹) and relatively weak downwelling during the rest of the year.

The Ekman velocity was set to zero at the surface and increased linearly to the Ekman depth, which was considered as the minimum value of 30 m and the mixed layer depth, and decreased linearly to zero down to 250 m (Ono et al., 2001). As the depth-dependent w requires horizontal convergence or divergence for volume conservation, horizontal advection included a correction term.

235 This was accomplished numerically by implicitly evaluating $w\partial C/\partial z$ at the grid box interfaces.

3.2.3 Shortwave solar radiation

The effect of the solar shortwave radiation that penetrates below the mixed layer (J_C term for the temperature model) was computed as:

$$J_C^T(t, z) = \frac{1}{\rho(t, z)C_p(t, z)} \frac{\partial I(t, z)}{\partial z}, \quad (5)$$

240 where ρ is the water density computed from temperature and salinity seasonal cycles using the Millero and Poisson (1981) formulation, C_p is the specific heat (Fofonoff and Millard, 1983), and $I(t, z)$ is the shortwave radiation flux computed by using the attenuation model of Paulson and Simpson (1977) for Type I water and the surface shortwave radiation values (Figure 3b). These values were obtained by fitting to an harmonic function monthly data for the period 1996-2001 obtained
245 from the CORE.2 Global Air-Sea flux dataset close to both sites. Annual means and amplitudes of the harmonic function were 182 ± 5 W m⁻² and 87 ± 6 W m⁻² at BATS, and 191 ± 5 W m⁻² and 68 ± 8 W m⁻² at ESTOC, respectively. Surface I was minimum in January for both stations.

3.3 Temperature model and K optimization

Similar to the approach followed by Musgrave et al. (1988), Ono et al. (2001) and Mouriño-Carballido
250 and Anderson (2009) we computed the vertical diffusivity (K) from the optimization of the simulated temperature (T) seasonal cycle (Figure 4). An optimized value of K was obtained every 15 days by minimizing the following function for each run:

$$Cost(t) = \left(\int_{MLD(t)}^{250m} (T_{obs}(t, z) - T_{mod}(t, z))^2 dz \right)^{1/2}. \quad (6)$$

The annual mean K was 2.0 ± 1.5 cm² s⁻¹ for BATS and 1.5 ± 1.6 cm² s⁻¹ for ESTOC. The
255 maximum K at BATS was set to 5 cm² s⁻¹ on account of numerical limitations in the biological source term calculations due to the high values computed during the winter mixing. At ESTOC higher K values were computed in July and August, possibly related to Trade Wind intensification during the summer months.

The comparison of the temperature rate of change computed from the observed and simulated
260 seasonal cycles showed a good agreement for both stations (Figure 5). The errors in the simulation

of the temperature seasonal cycle were computed as:

$$Err(t, z) = abs \left(\frac{T_{mod}(t, z) - T_{obs}(t, z)}{T_{obs}(t, z)} \right). \quad (7)$$

Model errors were related to a deficient parametrization of the constant vertical diffusivity especially during strong mixing events. Higher errors were computed during the periods of strong winter mixing for both stations, just below the mixed layer in June-July (BATS) and August-September (ESTOC), and between 100 and 250 m in September at BATS and June-September at ESTOC. Ono et al. (2001) used a single K optimized for the whole year, however we observed that optimizing K for each 15-day period reduced the total integrated error in the simulation of the temperature seasonal cycle by 13% at BATS and 20% at ESTOC (data not shown). Figure 5d shows the contribution of the geostrophic horizontal advection to the temperature rate of change. This contribution was more important at ESTOC, where an input of warmer water was computed in June-July and colder water in September.

3.4 Integrated budgets and errors estimation

Net production rates below the mixed layer were computed by integrating the biological source term (J_C) from this depth to the estimated compensation depth. This depth was set to 110 m based on the J_C distribution and the examination of rates sensitivity to this limit (see below). To compute shallow remineralization, J_C was integrated from the compensation depth down to 250 m. Both rates were integrated between April and December in order to avoid the period of intense and intermittent winter mixing, which is not accurately simulated by the model. The physical model terms were also integrated for the same period and depth intervals to evaluate their contribution to the change in the tracer inventories.

Uncertainties associated with the integrated budgets were estimated by using Monte Carlo simulations. A high number ($N > 100$) of model runs for all the tracers (including temperature for K optimization) were performed with each element of the model inputs being randomly generated from a Gaussian distribution. The standard deviation of the Gaussian distribution was taken from the estimated error of the model inputs. The weighted standard error was used for the seasonal cycles of temperature and the chemical tracers, the standard error provided by the WOA09 data base was used for the variables obtained from this climatology, a 75% error was assigned to the DIC lateral gradients consistent with the magnitude of computed errors for the lateral gradients of oxygen and nitrate, for the Ekman velocity (w) the error considered was 25% according to Ono et al. (2001). The uncertainties associated with the integrated budgets were the standard deviation of the values of the integrated budgets produced by each model realization.

4 Results and discussion

4.1 Distribution of biological sources and sinks

295 Figure 6 shows the biological source term (J_C) computed for oxygen, DIC and nitrate by using the diagnostic model at BATS and ESTOC and the geostrophic horizontal flux divergence (i.e. the contribution of the geostrophic horizontal advection to the total rate of tracer change).

Despite the signal being noisier, probably due to using a shorter data set and a time-dependent K optimization, the distribution of J_C computed for all the tracers at BATS was in general in agreement
300 with the results reported by Ono et al. (2001). The biological oxygen (DIC and nitrate) production (consumption) computed below the mixed layer was consistent with the synthesis of organic matter, whereas oxygen (DIC and nitrate) consumption (production) computed in the shallow aphotic zone indicated remineralization of organic matter. Oxygen production occurred below the mixed layer during summer stratification with the maximum at ca. 40 m. The compensation depth ($J_C = 0$) was
305 between 80-100 m. Below the compensation depth, maximum oxygen consumption was computed at ca. 120 m during late spring and early summer. The high oxygen consumption computed just below the mixed layer coincided with large errors in the temperature simulation (see Figure 5c), and they are probably related to limitations in the diffusivity optimization. DIC consumption occurred just below the mixed layer during summer stratification. For this tracer the compensation depth
310 was slightly shallower (ca. 60-80 m), and maximum production was computed at 120 m during the summer. The distribution of J_C for nitrate was very similar to DIC but the compensation depth was located deeper (ca. 110-115 m). Maximum nitrate consumption was computed at 80 m during May-June. Maximum production occurred at 120 m coinciding with the maximum rates of DIC (oxygen) production (consumption), and slightly shallower than the maximum described by Ono et al. (2001)
315 at 140 m. A second maximum in nitrate remineralization was observed at 250 m close to the model border.

The distribution of J_C at ESTOC was in general very similar to the patterns described for BATS although a few differences were observed. The oxygen compensation depth was located deeper at ca. 115 m. Highest rates of oxygen production were also located deeper (ca. 100 m), coinciding
320 with the deep chlorophyll maximum. The vertical structure of the spring-summer remineralization was also different compared to BATS, as maxima rates instead of being located at specific depths occurred through out the whole water column. Another important difference with BATS was observed in September-November when high oxygen (DIC, nitrate) consumption (production) was computed through the whole water column. This feature coincided with significant inputs of oxy-
325 gen and nitrate, and also cold water (Figure 5d), through horizontal advection. The role of the horizontal transport in the seasonality of the synthesis and remineralization of organic matter has been previously reported in the Canary region, which is under the influence of the coastal African upwelling (Neuer et al., 2002b; Arístegui et al., 2003; Pelegrí et al., 2005; Alonso-González et al.,

2009). Upwelling filaments and Ekman transport export particulate and dissolved organic matter
330 from the coastal upwelling to the open ocean. During this transport organic matter is remineralized,
at least partially, providing an external source for nutrients (Pelegrí et al., 2005). According to these
authors the relevance of this process is higher during the fall. These facts indicate that at ESTOC,
even though removed from direct influence of the upwelling zone, allochthonous inputs of organic
and inorganic matter could represent sources and sinks for oxygen, DIC and nutrients that must be
335 considered when interpreting our model results.

4.2 Net production rates

We set 110 m as the mean compensation depth because it is the approximate depth of maxima in the
integrated net production and remineralization rates for nitrate and for oxygen at ESTOC (see Figure
7). DIC and oxygen at BATS actually have maxima at 80-90 m, but they all have inflection points
340 near 110 m, and the 110 m integrals are not greatly different (20% lower for DIC at ESTOC and
O₂ at BATS; 2% lower for DIC at BATS). Net production rates and all the integrated model terms,
including the associated errors computed below the mixed layer to 110 m from April to December
for both stations, are shown in Figure 8.

Net production rates computed for oxygen ($1.34 \pm 0.79 \text{ mol O}_2 \text{ m}^{-2}$), dissolved inorganic carbon
345 ($-1.73 \pm 0.52 \text{ mol C m}^{-2}$) and nitrate ($-125 \pm 36 \text{ mmol N m}^{-2}$) at BATS were slightly higher
for oxygen and carbon compared to ESTOC ($1.03 \pm 0.62 \text{ mol O}_2 \text{ m}^{-2}$, $-1.42 \pm 0.30 \text{ mol C m}^{-2}$
and $-213 \pm 56 \text{ mmol N m}^{-2}$) but showed no statistically significant differences. In agreement with
previous geochemical estimates (Riser and Johnson, 2008) our results indicate that in these regions,
at least for the depths (mixed layer base to 110 m) and time period (April-December) we considered,
350 photosynthesis exceeds remineralization of organic matter.

The comparison of the observed changes in the tracer inventories and the modelled net production
evidences the relevance of the physical fluxes, as the change in the tracers, in general, underesti-
mated biological net production. The change in the tracer inventories underestimated oxygen net
production rates by 163% at BATS and 136% at ESTOC. For DIC the underestimation was 63% at
355 ESTOC, whereas it was overestimated by 17% at BATS. For nitrate the underestimation was 97% at
BATS and 102% at ESTOC. The contribution of diffusion was larger than total advection (horizon-
tal+vertical) for all the tracers at both stations except DIC at BATS.

Computed net production rates compared with a summary of previous values reported for BATS
and ESTOC are presented in Table 1, where different integration periods must be interpreted care-
fully. Gruber et al. (1998) and Brix et al. (2006) estimated that net production at BATS during the
360 spring-summer period represents 60 – 80% of the total annual net production. Similar results were
obtained at ESTOC by González-Dávila et al. (2003). As the integration interval (April-December)
used for our estimates misses part of the winter-spring bloom (Brix et al., 2006; González-Dávila
et al., 2007, 2010), our rates are probably lower than annual estimates. According to Marchal et al.

365 (1996), net production in the mixed layer in the Sargasso Sea represented 60% of the photic layer
net production. Assuming this is true also for ESTOC, our rates would represent about 40% of the
total net production, i.e. $3.4 \text{ mol O}_2 \text{ m}^{-2}$ and $-4.3 \text{ mol C m}^{-2}$ at BATS and $2.6 \text{ mol O}_2 \text{ m}^{-2}$ and
 $-3.6 \text{ mol C m}^{-2}$ at ESTOC. As nitrate consumption is only observed in the lower photic layer, this
extrapolation is not suitable for this tracer. Using an oxygen tracer model, Musgrave et al. (1988)
370 estimated the rate of net production in the photic layer at BATS to be $3-4 \text{ mol O}_2 \text{ m}^{-2} \text{ yr}^{-1}$, very
similar to our estimate ($3.4 \text{ mol O}_2 \text{ m}^{-2}$). Our rate for net production of carbon at BATS ($1.73-4.3$
 mol C m^{-2}) is in the range of previous estimates using mixed layer model approaches ($1.8-4.0 \text{ mol}$
 C m^{-2} , Marchal et al. (1996); Gruber et al. (1998); Brix et al. (2006)). A compilation of nitrogen
inputs carried out in the Sargasso Sea including wet and dry deposition, convective, diapycnal and
375 isopycnal mixing, Ekman transport, mesoscale eddies and nitrogen fixation quantified annual pro-
duction in $0.37-1.39 \text{ mol N m}^{-2} \text{ yr}^{-1}$ (Lipschultz et al., 2002). Our estimate for net production
of nitrogen at BATS ($0.125 \pm 0.036 \text{ mol N m}^{-2}$), which neglects important sources of nitrogen as
nitrogen fixation, convective mixing and mesoscale structures, does not reach the lower end of this
compilation. Our estimate is also lower than the value obtained by Siegel et al. (1999) (0.48 ± 0.12
380 $\text{mol N m}^{-2} \text{ yr}^{-1}$), who reported a compilation of new nitrogen inputs where nitrogen fixation was
not included.

González-Dávila et al. (2003), by using a mixed-layer model for ESTOC that did not include
lateral advection, estimated net production of carbon in the mixed layer between April and October,
and the period 1996-2001, to be $25.5 \pm 5.7 \text{ mmol C m}^{-3}$. This value corresponds to a mixed layer net
385 production rate of $1.55 \pm 0.75 \text{ mol C m}^{-2}$, computed considering the averaged depth of the mixed
layer during this period ($61 \pm 26 \text{ m}$). Following the same approach but using a longer data set (1995-
2004), González-Dávila et al. (2007) estimated net production in the mixed layer to be 3.3 ± 0.8
 mol C m^{-2} . Both rates are in close agreement with our estimate ($1.42-3.6 \text{ mol C m}^{-2}$). Our rate
of net production based on nitrate ($0.213 \pm 0.053 \text{ mol N m}^{-2}$) is only slightly lower than the total
390 new nitrogen inputs reported by Cianca et al. (2007) at ESTOC ($0.28 \pm 0.09 \text{ mol N m}^{-2} \text{ yr}^{-1}$), who
considered eddy pumping, convective, diapycnal and isopycnal mixing, and Ekman transport.

The $-\text{O}_2:\text{C}$ ratios computed from the net production estimates at BATS (0.77 ± 0.51) and ESTOC
(0.72 ± 0.46) are lower than the ratio for NO_3 -based production (1.4, Gruber (2008)) and closer to
the ratio for NH_4 -based production (1.1). The lower $-\text{O}_2:\text{C}$ ratio at ESTOC could be related to the
395 remineralization peak observed during the fall (September-November), which was more intense for
oxygen than for DIC, probably due to the larger lateral inputs (see Figure 6). When the biological
source term at ESTOC was integrated between April and September, excluding the remineralization
pulse observed in autumn, the $-\text{O}_2:\text{C}$ ratio (1.30 ± 0.62) was closer to the standard values. Net pro-
duction rates computed between April and December with the geostrophic model term set to zero
400 (see section 3.2.1) gave a $-\text{O}_2:\text{C}$ ratio of ~ 1.0 , also in better agreement with the canonic values.
At BATS, we attributed the low $-\text{O}_2:\text{C}$ ratio to the diffusion simulation that causes a strong oxygen

consumption between the mixed-layer depth and 35 m. If this region was not included in the integration, oxygen production rate would be increased ($2.14 \pm 0.37 \text{ mol O}_2 \text{ m}^{-2}$) and DIC production rate lowered ($-1.20 \pm 0.21 \text{ mol C m}^{-2}$), resulting in higher than Redfield ratio (1.78 ± 0.86). The
405 C:N value at BATS (13.9 ± 5.8) was higher than the Redfield ratio (6.6) and very similar at ESTOC (6.7 ± 2.3). However, this is slightly deceiving in that most of the oxygen production and DIC uptake occurs above 80 m, while most of the NO_3 uptake occurs between 80-110 m, which is not constant Redfield C:N coupling.

4.3 Remineralization rates

410 All the model terms were integrated between April and December and 110-250 m to obtain shallow remineralization rates (see Figure 9). Oxygen ($-1.81 \pm 0.37 \text{ mol O}_2 \text{ m}^{-2}$) and DIC ($1.52 \pm 0.30 \text{ mol C m}^{-2}$) remineralization rates computed for BATS were in good agreement with those reported by Ono et al. (2001) ($-2.08 \pm 0.38 \text{ mol O}_2 \text{ m}^{-2}$ and $-1.53 \pm 0.35 \text{ mol C m}^{-2}$), whereas nitrate
415 remineralization rate ($147 \pm 43 \text{ mmol N m}^{-2}$) was twice the rate reported by these authors ($80 \pm 46 \text{ mmol N m}^{-2}$). This mismatch was due to the differences in the geostrophic lateral advection term ($-51 \pm 44 \text{ mmol N m}^{-2}$ versus $39 \pm 39 \text{ mmol N m}^{-2}$ reported by Ono et al.) that resulted from the use of different climatologies. Changes in the tracer inventories also underestimated shallow remineralization, as they represented 51%, 45% and 21% of remineralization rates computed for oxygen, DIC and nitrate, respectively.

420 Shallow remineralization rates computed for ESTOC were significantly higher than at BATS for oxygen ($-3.9 \pm 1.0 \text{ mol O}_2 \text{ m}^{-2}$) but showed no significant differences for DIC ($1.53 \pm 0.43 \text{ mol C m}^{-2}$) and nitrate ($38 \pm 155 \text{ mmol N m}^{-2}$). The latter was associated with large errors. The change in the tracer inventories underestimated remineralization rates for oxygen and DIC as represented
425 17 and 36% of these rates. However, nitrate remineralization rate was overestimated (340%) by the change in the inventory.

Diffusion was a source for oxygen and a sink for DIC and nitrate in both stations because oxygen (DIC/nitrate) gradient was negative (positive) downwards and greater at 110 m than 250 m. Vertical advection was a source for all the tracers at BATS because w was negative at 110 m and zero at 250 m, whereas it was a sink at ESTOC because upwelling was dominant during the investigation period.
430 Lateral transport (geostrophic horizontal advection) was a source (sink) for oxygen (nitrate/DIC) at BATS and a source (sink) for oxygen and nitrate (DIC) at ESTOC. The divergence of the lateral transport of oxygen caused the differences in the remineralization rates computed for both stations, as the 1D model without geostrophic advection resulted in oxygen consumption rates reduced by
435 61% at ESTOC ($-1.51 \pm 0.29 \text{ mol O}_2 \text{ m}^{-2}$) and only by 27% at BATS ($-1.31 \pm 0.80 \text{ mol O}_2 \text{ m}^{-2}$). The input of nitrate through lateral advection was also responsible for the low remineralization rate and the large error computed for nitrate at ESTOC, as the 1D model computed a higher rate and lower associated error ($135 \pm 23 \text{ mmol N m}^{-2}$). The fact that differences in remineralization

rates between both stations were only statistically significant for oxygen could be related to the different temporal resolution of the databases used to compute the lateral gradients of the tracers.

440 Oxygen and nitrate horizontal gradients were calculated from the World Ocean Atlas 2009 monthly climatology, whereas DIC gradients were obtained from the Global Distribution of Total Inorganic Carbon and Total Alkalinity Below the Deepest Winter Mixed Layer Depths climatology, which includes lower temporal resolution tri-monthly averaged data (see section 3.2.1). Model results show that inconsistency in remineralization rates differences between the tracers decreased if oxygen and
445 nitrate variability was reduced through smoothing (data not shown). Large errors associated to nitrate remineralization were probably due to the lower number of observations used to compute the nitrate, compared to oxygen, climatology.

Remineralization rates computed at BATS were comparable with previous estimates reported in the region (see Table 2). Jenkins and Goldman (1985) calculated the shallow (100-250 m) oxygen
450 consumption rate between April and November at station S in the Sargasso Sea to be $3 \text{ mol O}_2 \text{ m}^{-2}$. Sarmiento et al. (1990) summarized remineralization rates determined from a variety of tracer based techniques in the Sargasso Sea to be $0.6 - 3.3 \text{ mol O}_2 \text{ m}^{-2} \text{ yr}^{-1}$. Carlson et al. (1994) estimated the remineralization of dissolved organic matter from spring to fall as $0.99 - 1.21 \text{ mol C m}^{-2}$, lower than our estimate as it did not include the particulate organic fraction.

455 To our knowledge, geochemical estimates of shallow remineralization based on tracer distributions have not been conducted at ESTOC so far. Alonso-González et al. (2009) used a box model approach to calculate the lateral transport and consumption of suspended particulate organic matter (POC) in the Canary Current region. They found that lateral POC fluxes, which were up to 3 orders of magnitude higher than vertical fluxes, accounted for $1.42 \text{ mmol C m}^{-2} \text{ d}^{-1}$, what represents
460 $28 - 59\%$ of the total mesopelagic (100-700 m) respiration. The total carbon consumption estimated for mesopelagic waters in the fall was $2.4 - 5.1 \text{ mmol C m}^{-2} \text{ d}^{-1}$, slightly lower than our estimate ($6.7 \pm 1.9 \text{ mmol C m}^{-2} \text{ d}^{-1}$) and one order of magnitude lower than respiration rates obtained during the summer by using combined in vitro oxygen consumption and enzymatic activity techniques ($68 \text{ mmol C m}^{-2} \text{ d}^{-1}$) in the subtropical Northeast Atlantic (Arístegui et al., 2005). Alonso-González
465 et al. hypothesized that this mismatch could be due to the different times of the year when the experiments were carried out. However, recent studies indicate that long incubations may lead to overestimations of community respiration as a result of an increase in the activity of heterotrophic bacteria (Marañón et al., 2007; Calvo-Díaz et al., 2011). Due to the differences in integration depths, comparison with these studies must be done carefully.

470 Shallow remineralization rates computed by our model can also be compared to particulate organic matter attenuation flux computed from sediment traps deployed at BATS and ESTOC. Comparative information for sediment traps for both stations corresponds to 150, 200 and 300 m at BATS and 200, 300 and 500 m at ESTOC (Helmke et al., 2010). In order to estimate remineralization rates between 110 and 250 m the particle flux at 110 and 250 m needs to be inferred. The former was

475 calculated using the Martin's attenuation equation (Martin et al., 1987), considering the seasonal b
exponents calculated by Helmke et al. (2010) ($b = 0.6 - 1.19$) for BATS, and the value reported by
Neuer et al. (2007) ($b = 0.81$) for ESTOC. The particulate flux at 250 m was calculated by linearly
interpolating the flux measured at 200 and 300 m. Shallow remineralization computed from spring-
time to fall was $0.38 \text{ mol C m}^{-2}$ at BATS and $0.08 \text{ mol C m}^{-2}$ at ESTOC. Both values were lower
480 than our estimates, which included the dissolved and particulate organic fractions, and represented
25 and 5% of the carbon remineralization computed for BATS and ESTOC, respectively. The sum
of the remineralization rates of particulate and dissolved organic matter at BATS ($1.37 - 1.59 \text{ mol C}$
 m^{-2}) matches our estimation of carbon remineralization ($1.52 \pm 0.30 \text{ mol C m}^{-2}$). Unfortunately,
no information about the seasonal variability of the dissolved organic carbon cycle is available at
485 ESTOC.

Our results indicated that lateral transport is significant to the seasonality of the hydrodynamic
conditions and the tracer distribution at ESTOC. Previous studies indicated that higher remineraliza-
tion in the subtropical Northeast Atlantic could be supported by dissolved (Álvarez-Salgado et al.,
2007) and slow sinking particulate organic matter (Alonso-González et al., 2009) exported from
490 the high productive African coastal upwelling, and also by dark dissolved inorganic carbon fixation
(Baltar et al., 2010). In this respect, Arístegui et al. (2003) estimated that the contribution of dis-
solved organic carbon to the mesopelagic oxygen consumption in the Canary Current region was
about 30%. Alonso-González et al. (2009) indicated that the reported differences in the carbon ex-
port rates between both stations could be, at least partially, explained by lower sedimentation rates
495 of the particulate organic matter in the eastern part of the gyre, as slow-sinking or suspended POC
does not accumulate in the sediment traps and can be laterally advected (Arístegui et al., 2009).
Although organic matter was not explicitly taken into account in our model, our results support this
hypothesis as they show stronger lateral transport and higher remineralization rates for oxygen at
ESTOC, which can not be sustained by the flux of sinking organic matter.

500 According to our results shallow remineralization represents at BATS $54 \pm 33\%$ and $51 \pm 36\%$
of the net production computed in the photic zone for oxygen and DIC, respectively, whereas at
ESTOC it represents $151 \pm 99\%$ and $43 \pm 15\%$ respectively. The high value for oxygen at ESTOC is
related to the strong lateral inputs computed in the fall (see oxygen geostrophic horizontal advection
in Figure 6), that also resulted in negative net production computed during these months (see oxygen
505 biological source in Figure 6), suggesting that high remineralization rates could partially be sustained
by non-locally produced organic matter.

The $-\text{O}_2:\text{C}$ ratio for the shallow remineralization computed at BATS (1.19 ± 0.34) was in good
agreement with the Redfield ratio (1.43), whereas at ESTOC (2.52 ± 0.98) was slightly higher. C:N
ratios were in excess of the Redfield ratio at both stations, and were higher at ESTOC (40 ± 162)
510 than at BATS (10.2 ± 3.6), pointing to the export of carbon enriched organic matter. The C:N ratio
reported by Ono et al. (2001) for BATS (19 ± 12) was twice our estimate, as a consequence of the

differences in nitrate remineralization estimates. The higher C:N ratio computed for ESTOC was consistent with the carbon rich organic matter observed in the mixed layer (Koeve, 2006) and the sinking organic matter (Neuer et al., 2007) in the region, and with close to Redfield ratios observed in the sinking organic matter at BATS (Schnitzer and Steinberg, 2002). Moreover, ecological models employed to study the sensitivity of carbon export to atmospheric nitrogen inputs in NASW and NASE showed higher C:N values in the sinking organic matter in NASE (Mouriño-Carballido et al., in revision).

5 Conclusions

Subtropical gyres play a crucial role in the global carbon cycle due to the large extension they occupy which is expected to get bigger as a consequence of global warming (Polovina et al., 2008). Understanding the heterogeneity of these biomes is critical to comprehend the role of the oceans in the future carbon cycle. By using a 1D diagnostic tracer model we investigated if potential differences in net production and shallow remineralization are consistent with the lower export of particulate organic carbon observed at the eastern (ESTOC) compared to the western side (BATS), of the North Atlantic Subtropical gyre. Our results showed that net production was not statistically different at both stations, whereas oxygen shallow remineralization was higher at ESTOC. As our approach does not estimate mixed-layer net production, particularly during the winter-spring bloom period, we can not discard that it underestimates the differences in net production between the two stations. The comparison of our estimates with vertical fluxes of POC shows that whereas a significant fraction (25%) of the shallow remineralization is sustained by sinking organic matter at BATS, this fraction is much lower at ESTOC (5%). The importance of horizontal processes in the seasonal cycles at ESTOC indicates that oxygen consumption at this station could be also supported by allochthonous inputs of organic matter. There is still work to be done to fully understand the mechanisms behind the regional differences observed in the vertical export of carbon at the North Atlantic subtropical gyre. In the light of this and other studies it seems that crucial information must come from the study of horizontal fluxes at ESTOC.

Acknowledgements. We are grateful to all the technicians and crew members involved in the collection, preparation, and analysis of the large data sets used in this study, in particular those from the BATS and ESTOC programs. B. Mouriño was supported by the Ramón y Cajal program from the Spanish Minister of Science and Technology. Funding for this study was provided by the Xunta de Galicia under the research project VAR-ITROP (09MDS001312PR, PI B. Mouriño) and by the Ministerio de Ciencia e Innovation MOMAC project (CTM2008-05914/MAR).

References

- 545 Alonso-González, I. J., Arístegui, J., Vilas, J. C., and Hernández-Guerra, A.: Lateral POC transport and consumption in surface and deep waters of the Canary Current region: A box model study, *Global Biogeochem. Cy.*, 23, GB2007, doi:10.1029/2008GB003185, 2009.
- Álvarez-Salgado, X. A., Arístegui, J., Barton, E. D., and Hansell, D. A.: Contribution of upwelling filaments to offshore carbon export in the subtropical Northeast Atlantic Ocean, *Limnol. Oceanogr.*, 52, 1287–1292, 550 2007.
- Antonov, J. I., Seidov, D., Boyer, T. P., Locarni, R. A., Mishonov, A. V., Garcia, H. E., Baranova, O. K., Zweng, M. M., and Johnson, D. R.: *World Ocean Atlas 2009, Volume 2: Salinity.*, Ed. NOAA Atlas NESDIS 69, U.S. Government Printing Office, Washington, D.C., 2010.
- Arístegui, J., Barton, E., Montero, M., García-Muñoz, M., and Escáñez, J.: Organic carbon distribution and 555 water column respiration in the NW Africa-Canaries Coastal Transition Zone, *Aquat. Microb. Ecol.*, 33, 289–301, 2003.
- Arístegui, J., Duarte, C. M., Gasol, J. M., and Alonso-Sáez, L.: Active mesopelagic prokaryotes support high respiration in the subtropical northeast Atlantic Ocean, *Geophys. Res. Lett.*, 32, L03 608, doi: 10.1029/2004GL021863, 2005.
- 560 Arístegui, J., Gasol, J. M., Duarte, C. M., and Herndl, G. J.: Microbial oceanography of the dark ocean's pelagic realm, *Limnol. Oceanogr.*, 54, 1501–1529, doi:10.4319/lo.2009.54.5.1501, 2009.
- Baltar, F., Arístegui, J., Sintés, E., Gasol, J. M., Reinthaler, T., and Herndl, G. J.: Significance of non-sinking particulate organic carbon and dark CO₂ fixation to heterotrophic carbon demand in the mesopelagic north-east Atlantic, *Geophys. Res. Lett.*, 37, L09 602, doi:10.1029/2010GL043105, 2010.
- 565 Bonnet, S., Grosso, O., and Moutin, T.: Planktonic dinitrogen fixation along a longitudinal gradient across the Mediterranean Sea during the stratified period (BOUM cruise), *Biogeosciences*, 8, 2257–2267, doi: 10.5194/bg-8-2257-2011, 2011.
- Brix, H., Gruber, N., Karl, D. M., and Bates, N. R.: On the relationships between primary, net community, and export production in subtropical gyres, *Deep Sea. Res. Pt. II*, 53, 698–717, 2006.
- 570 Calvo-Díaz, A., Díaz-Pérez, L., G Morán, X. A., Teira, E., and Marañón, E.: Bottle enclosure causes a decrease in the autotrophic to heterotrophic biomass ratio of picoplankton in oligotrophic marine waters, *Appl. Environ. Microb.*, 2011.
- Capone, D. G., Burns, J. A., Montoya, J. P., Subramaniam, A., Mahaffey, C., Gunderson, T., Michaels, A. F., and Carpenter, E. J.: Nitrogen fixation by *Trichodesmium* spp.: An important source of new nitrogen to the tropical and subtropical North Atlantic Ocean, *Global Biogeochem. Cy.*, 19, GB2024, doi: 575 10.1029/2004GB002331, 2005.
- Carlson, C. A., Ducklow, H. W., and Michaels, A. F.: Annual flux of dissolved organic carbon from the euphotic zone in the northwestern Sargasso Sea, *Nature*, 371, 405–408, 10.1038/371405a0, 1994.
- Cianca, A., Helmke, P., Mouriño, B., Rueda, M. J., Llinás, O., and Neuer, S.: Decadal analysis of hydrography and in situ nutrient budgets in the western and eastern North Atlantic subtropical gyre, *J. Geophys. Res. - Oceans*, 112, C07 025, doi:10.1029/2006JC003788, 2007.
- 580 Davenport, R., Neuer, S., Helmke, P., Perez-Marrero, J., and Llinás, O.: Primary productivity in the northern Canary Islands region as inferred from SeaWiFS imagery, *Deep-Sea Res. Pt. II*, 49, 3481 – 3496, doi:10.

1016/S0967-0645(02)00095-4, 2002.

- 585 Emerson, S., Quay, P., Karl, D., Winn, C., Tupas, L., and Landry, M.: Experimental determination of the organic carbon flux from open-ocean surface waters, *Nature*, 389, 951–954, 1997.
- Field, C. B., Behrenfeld, M. J., Randerson, J. T., and Falkowski, P.: Primary Production of the Biosphere: Integrating Terrestrial and Oceanic Components, *Science*, 281, 237–240, doi:10.1126/science.281.5374.237, 1998.
- 590 Fofonoff, N. P. and Millard, R. C.: Algorithms for computation of fundamental properties of seawater, *Unesco Technical Papers in Marine Science*, 44, 53, 1983.
- Garcia, H. E., Locarnini, R. A., Boyer, T., Antonov, J. I., Zweng, M., and Johnson, D. R.: *World Ocean Atlas 2009, Volume 4: Nutrients (phosphate, nitrate, silicate)*, NOAA Atlas NESDIS 71, U.S. Government Printing Office, Washington, D.C., 2010a.
- 595 Garcia, H. E., Locarnini, R. A., Boyer, T., Antonov, J. I., Zweng, M., and Johnson, D. R.: *World Ocean Atlas 2009, Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Oxygen Saturation.*, NOAA Atlas NESDIS 70, U.S. Government Printing Office, Washington, D.C., 2010b.
- González-Dávila, M., Santana-Casiano, J. M., Rueda, M.-J., Llinás, O., and González-Dávila, E.-F.: Seasonal and interannual variability of sea-surface carbon dioxide species at the European Station for Time Series
600 in the Ocean at the Canary Islands (ESTOC) between 1996 and 2000, *Global Biogeochem. Cy.*, 17, 1076, doi:10.1029/2002GB001993, 2003.
- González-Dávila, M., Santana-Casiano, J. M., and González-Dávila, E. F.: Interannual variability of the upper ocean carbon cycle in the northeast Atlantic Ocean, *Geophys. Res. Lett.*, 34, L07 608, doi:10.1029/2006GL028145, 2007.
- 605 González-Dávila, M., Santana-Casiano, J. M., Rueda, M. J., and Llinás, O.: The water column distribution of carbonate system variables at the ESTOC site from 1995 to 2004, *Biogeosciences*, 7, 3067–3081, 2010.
- Goyet, C., Healy, R., and Ryan, J.: *Global distribution of total inorganic carbon and total alkalinity below the deepest winter mixed layer depths*, Environmental Sciences Division, U.S. Department of Energy, 2000.
- Gruber, N.: *Nitrogen in the Marine Environment*, chap. The marine nitrogen cycle: overview and challenges,
610 pp. 1–50, Academic Press, Elsevier Inc., Amsterdam., 2008.
- Gruber, N., Keeling, C. D., and Stocker, T. F.: Carbon-13 constraints on the seasonal inorganic carbon budget at the BATS site in the northwestern Sargasso Sea, *Deep-Sea Res. Pt. I*, 45, 673–717, 1998.
- Helmke, P., Neuer, S., Lomas, M. W., Conte, M., and Freudenthal, T.: Cross-basin differences in particulate organic carbon export and flux attenuation in the subtropical North Atlantic gyre, *Deep-Sea Res. Pt. I*, 57,
615 213–227, doi:10.1016/j.dsr.2009.11.001, 2010.
- Jenkins, W. and Goldman, J.: Seasonal oxygen cycling and primary production in the Sargasso Sea, *J. Mar. Res.*, 43, 465–491, 1985.
- Koeve, W.: C:N stoichiometry of the biological pump in the North Atlantic: Constraints from climatological data, *Global Biogeochem. Cy.*, 20, GB3018, doi:10.1029/2004GB002407, 2006.
- 620 Leetmaa, A. and Bunker, A. F.: Updated charts of the mean annual wind stress, convergences in Ekman Layers, and Sverdrup transport in the North Atlantic, *J. Mar. Res.*, 36, 311–322, 1978.
- Lipschultz, F., Bates, N., Carlson, C., and Hansell, D.: New production in the Sargasso Sea: History and current status, *Global Biogeochem. Cy.*, 16, 1001, doi:10.1029/2000GB001319, 2002.

- Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T., Garcia, H., Baranova, O. K., Zweng, M. M.,
625 and Johnson, D. R.: World Ocean Atlas 2009, Volume 1: Temperature, NOAA Atlas NESDIS 68, U.S.
Government Printing Office, Washington, D.C., 2010.
- Marañón, E., Pérez, V., Fernández, E., Anadón, R., Bode, A., González, N., Huskin, I., Isla, A., Morán, X.
A. G., Mouriño, B., Quevedo, M., Robinson, C., Serret, P., Teira, E., Varela, M. M., Woodward, E. M. S.,
630 and Zubkov, M. V.: Planktonic carbon budget in the eastern subtropical North Atlantic, *Aquat. Microb. Ecol.*,
48, 261–275, 2007.
- Marañón, E., Fernández, A., Mouriño-Carballido, B., Martínez-García, S., Teira, E., Cermeño, P., Chouciño,
P., Huete-Ortega, M., Fernández, E., Calvo-Díaz, A., Morán, X. A. G., Bode, A., Moreno-Ostos, E., Varela,
M. M., Patey, M. D., and Achterberg, E. P.: Degree of oligotrophy controls the response of microbial plankton
to Saharan dust, *Limnol. Oceanogr.*, 55, 2339–2352, 2010.
- 635 Marchal, O., Monfray, P., and Bates, N. R.: Spring summer imbalance of dissolved inorganic carbon in the
mixed layer of the northwestern Sargasso Sea, *Tellus B-Chem. Phys. Meteorol.*, 48, 115–134, 1996.
- Martin, J. H., Knauer, G. A., Karl, D. M., and Broenkow, W. W.: VERTEX: carbon cycling in the northeast
Pacific, *Deep-Sea Res. Pt. A.*, 34, 267–285, 1987.
- Menzel, D. and Ryther, J.: The annual cycle of primary production in the Sargasso Sea off Bermuda, *Deep-Sea*
640 *Res. (1953)*, 6, 351–367, 1959.
- Michaels, A. F. and Knap, A. H.: Overview of the U.S. JGOFS Bermuda Atlantic Time-series Study and the
Hydrostation S program, *Deep-Sea Res. Pt. II*, 43, 157–198, 1996.
- Millero, F. J. and Poisson, A.: International one-atmosphere equation of state of seawater, *Deep-Sea Res. Pt. A.*,
28, 625–629, 1981.
- 645 Mouriño, B., Fernández, E., Etienne, H., Hernández, F., and Giraud, S.: Significance of cyclonic SubTropical
Oceanic Rings of Magnitude (STORM) eddies for the carbon budget of the euphotic layer in the subtropical
northeast Atlantic, *J. Geophys. Res.*, 108, 3383, doi:10.1029/2003JC001884, 2003.
- Mouriño-Carballido, B. and Anderson, L. A.: Net community production of oxygen derived from in vitro
and in situ 1-D modeling techniques in a cyclonic mesoscale eddy in the Sargasso Sea, *Biogeosciences*, 6,
650 1799–1810, 2009.
- Mouriño-Carballido, B. and Neuer, S.: Regional Differences in the Role of Eddy Pumping in the North Atlantic
Subtropical Gyre: Historical conundrums revisited, *Oceanography*, 21, 52–61, 2008.
- Mouriño-Carballido, B., Marañón, E., Fernández, A., Graña, R., Bode, A., Varela, V., Domínguez, F., Escáñez,
J., and de Armas, D.: Importance of N₂ fixation versus nitrate eddy diffusion across the Atlantic Ocean,
655 *Limnol. and Oceanogr.*, 56(3), 999–1007, 2011.
- Mouriño-Carballido, B., Pahlow, M., and Oschlies, A.: High sensitivity of ultra-oligotrophic marine ecosystems
to atmospheric nitrogen deposition, *Geophys. Res. Lett.*, in revision, 2011.
- Musgrave, D. L., Chou, J., and Jenkins, W. J.: Application of a Model of Upper-Ocean Physics for Studying
Seasonal Cycles of Oxygen, *J. Geophys. Res.*, 93, 15 679–15 700, 1988.
- 660 Najjar, R. G. and Keeling, R. F.: Mean annual cycle of the air-sea oxygen flux: A global view, *Global Bio-
geochem. Cy.*, 14, 573–584, 2000.
- Neuer, S., Davenport, R., Freudenthal, T., Wefer, G., Llinás, O., Rueda, M., Steinberg, D., and Karl, D.:
Differences in the biological carbon pump at three subtropical ocean sites, *Geophys. Res. Lett.*, 29, 32,

doi:10.1029/2002GL015393, 2002a.

- 665 Neuer, S., Freudenthal, T., Davenport, R., Llinás, O., and Rueda, M.-J.: Seasonality of surface water properties and particle flux along a productivity gradient off NW Africa, *Deep-Sea Res. Pt. II: Topical Studies in Oceanography*, 49, 3561–3576, 2002b.
- Neuer, S., Torres-Padron, M., Gelado-Caballero, M., Rueda, M., Hernández-Brito, J., Davenport, R., and Wefer, G.: Dust deposition pulses to the eastern subtropical North Atlantic gyre: Does ocean's biogeochemistry respond?, *Global Biogeochem. Cy.*, 18, GB4020, doi:10.1029/2004GB002228, 2004.
- 670 Neuer, S., Cianca, A., Helmke, P., Freudenthal, T., Davenport, R., Meggers, H., Knoll, M., Santana-Casiano, J. M., González-Dávila, M., Rueda, M.-J., and Llinás, O.: Biogeochemistry and hydrography in the eastern subtropical North Atlantic gyre. Results from the European time-series station ESTOC, *Progr. Oceanogr.*, 72, 1–29, doi:10.1016/j.pocean.2006.08.001, 2007.
- 675 Ono, S., Ennyu, A., Najjar, R. G., and Bates, N. R.: Shallow remineralization in the Sargasso Sea estimated from seasonal variations in oxygen, dissolved inorganic carbon and nitrate, *Deep-Sea Res. Pt. II*, 2001, 48, 1567–1582, 2001.
- Pätsch, J., Kühn, W., Radach, G., Santana Casiano, J. M., Gonzalez Davila, M., Neuer, S., Freudenthal, T., and Llinás, O.: Interannual variability of carbon fluxes at the North Atlantic Station ESTOC, *Deep-Sea Res. Pt. II*, 49, 253–288, 2002.
- 680 Paulson, C. A. and Simpson, J. J.: Irradiance Measurements in the Upper Ocean, *J. Phys. Oceanogr.*, 7, 952–956, 1977.
- Pelegrí, J., Arístegui, J., Cana, L., González-Dávila, M., Hernández-Guerra, A., Hernández-León, S., Marrero-Díaz, A., Montero, M., Sangrà, P., and Santana-Casiano, M.: Coupling between the open ocean and the coastal upwelling region off northwest Africa: water recirculation and offshore pumping of organic matter, *J. Mar. Syst.*, 54, 3–37, 2005.
- Polovina, J. J., Howell, E. A., and Abecassis, M.: Ocean's least productive waters are expanding, *Geophys. Res. Lett.*, 35, L03 618, doi:10.1029/2007GL031745, 2008.
- Pulido-Villena, E., Wagener, T., and Guieu, C.: Bacterial response to dust pulses in the western Mediterranean: Implications for carbon cycling in the oligotrophic ocean, *Global Biochem. Cy.*, 22, GB1020, doi:10.1029/2007GB003091, 2008.
- 690 Riser, S. C. and Johnson, K. S.: Net production of oxygen in the subtropical ocean, *Nature*, 451, 323–U5, doi:10.1038/nature06441, 2008.
- Santana-Casiano, J. M., González-Dávila, M., Rueda, M.-J., Llinás, O., and González-Dávila, E.-F.: The interannual variability of oceanic CO₂ parameters in the northeast Atlantic subtropical gyre at the ESTOC site, *Global Biogeochem. Cy.*, 21, GB1015, doi:10.1029/2006GB002788, 2007.
- Sarmiento, J., Thiele, G., Key, R., and Moore, W.: Oxygen and nitrate new production and remineralization in the North-Atlantic Subtropical Gyre, *J. Geophys. Res. - Oceans*, 95, 18 303–18 315, 1990.
- Schnetzer, A. and Steinberg, D. K.: Active transport of particulate organic carbon and nitrogen by vertically migrating zooplankton in the Sargasso Sea, *Mar. Ecol. - Prog. Ser.*, 234, 71–84, 10.3354/meps234071, 2002.
- 700 Siegel, D. A. and Deuser, W. G.: Trajectories of sinking particles in the Sargasso Sea: modeling of statistical funnels above deep-ocean sediment traps, *Deep-Sea Res. Pt. I*, 44, 1519–1541, 1997.
- Siegel, D. A., McGillicuddy, D. J., and Fields, E. A.: Mesoscale eddies, satellite altimetry, and new production

in the Sargasso Sea, *J. Geophys. Res.*, 104, 13 359–13 379, 1999.

- 705 Steinberg, D. K., Carlson, C. A., Bates, N. R., Johnson, R. J., Michaels, A. F., and Knap, A. H.: Overview of the US JGOFS Bermuda Atlantic Time-series Study (BATS): a decade-scale look at ocean biology and biogeochemistry, *Deep-Sea Res. Pt. II*, 48, 1405–1447, 2001.




Fig. 1. Seasonal cycles of the vertical distribution of temperature ($^{\circ}\text{C}$), salinity (psu), density anomaly σ_T (kg m^{-3}), chlorophyll a (mg m^{-3}), nitrate + nitrite (mmol m^{-3}), oxygen (mmol m^{-3}), oxygen anomaly (mmol m^{-3}) and dissolved inorganic carbon (DIC) (mmol m^{-3}). Black discontinuous line represents the mixed layer depth. Black thick line represents the isoline of zero oxygen anomaly and the nitracline ($[\text{NO}_3]=0.5 \text{ mmol m}^{-3}$).




Fig. 2. Computed geostrophic horizontal velocities (u, v) for BATS and ESTOC. The thick black line represents the null velocity component.




Fig. 3. (a) Vertical Ekman velocity (w) at the Ekman depth (w_E) and (b) surface shortwave solar radiation computed for the period 1996-2001 near BATS and ESTOC. The continuous lines represent the fit to a single harmonic function, whereas the dotted line correspond to the 95% confidence intervals. Fitting of w_E to the harmonic function was not used in the model.




Fig. 4. Optimized diffusivity (K) computed for BATS and ESTOC. Values higher than $5 \text{ cm}^2\text{s}^{-1}$ where set to $5 \text{ cm}^2\text{s}^{-1}$ on account of numerical limitations in the biological source term calculations.




Fig. 5. (a) Observed and (b) simulated temperature rate of change, (c) errors in the simulated temperature seasonal cycle, and (d) geostrophic horizontal advection flux divergence at BATS and ESTOC. The black discontinuous line represents the mixed layer depth and the black thick line represents the zero rate of change isoline.




Fig. 6. Biological source term (J_C) and geostrophic horizontal advection flux divergence computed for oxygen, DIC and nitrate at BATS and ESTOC. The discontinuous line represents the mixed-layer depth and the black thick line the zero iseline. Isoline separations are $0.1 \text{ mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$, $0.1 \text{ mmol C m}^{-2} \text{ d}^{-1}$ and $0.01 \text{ mmol N m}^{-2} \text{ d}^{-1}$, for oxygen, DIC and nitrate, respectively.




Fig. 7. Time and depth-integrated net production rates as a function of the lower limit used for the integration at BATS and ESTOC for oxygen (solid), DIC (dashed) and nitrate (dot-dashed).




Fig. 8. Integrated model terms computed from the mixed layer depth to 110 m, and from April to December. NP is net production. Integrated geostrophic horizontal advection term is shown in brackets. The white (black) arrows and boxes represent net gain (lost). Rates are expressed in mol m⁻² for oxygen and DIC and in mmol m⁻² for nitrate.




Fig. 9. Integrated model terms computed from 110 m down to 250 m, and from April to December. REM is remineralization. Integrated geostrophic horizontal advection term is shown in brackets. White (black) arrows and boxed represent net gain (lost). Rates are expressed in mol m^{-2} for oxygen and DIC and in mmol m^{-2} for nitrate.

Table 1. Summary of net production estimates reported for BATS and ESTOC. MLD is the mixed-layer depth.

Region	Source	Technique	Period	Depths	Estimate mol m ⁻²
Sargasso Sea	Musgrave et al. (1988)	Tracer model	Annual	Photic Layer (PL)	3 – 4 (O ₂)
BATS	Marchal et al. (1996)	Mixed-layer model	March-Oct	Extrapolation PL	1.8 – 2.9 (C)
BATS	Gruber et al. (1998)	Mixed-layer model	Annual	Mixed Layer	2.3 ± 0.9 (C)
BATS	Brix et al. (2006)	Mixed-layer model	Annual	PL	4.9 ± 0.3 (C)
BATS	Siegel et al. (1999)	Nitrogen budget	Annual	PL	0.48 ± 0.12 (N)
Sargasso Sea	Lipschultz et al. (2002)	N budget	Annual	PL	0.37 – 1.39 (N)
BATS	Cianca et al. (2007)	N budget	Annual	PL	0.38 ± 0.08 (N)
BATS	This study	Tracer Model	Apr-Dec	MLD to 110 m	1.34 ± 0.79 (O ₂) 1.73 ± 0.52 (C) 0.125 ± 0.036 (N)
ESTOC	González-Dávila et al. (2003)	Mixed-layer model	Mar-Oct	Mixed Layer	1.55 ± 0.75 (C)
ESTOC	González-Dávila et al. (2007)	Mixed-layer model	Annual	Mixed Layer	3.3 ± 0.8 (C)
ESTOC	Cianca et al. (2007)	N budget	Annual	PL	0.28 ± 0.09 (N)
ESTOC	This study	Tracer Model	Apr-Dec	MLD to 110 m	1.03 ± 0.62 (O ₂) 1.42 ± 0.30 (C) 0.213 ± 0.053 (N)

Table 2. Summary for remineralization estimates at BATS and ESTOC. AOU is apparent oxygen utilization; DOC, dissolved organic carbon; POC, particulate organic carbon; ETS, electron transport system and OC is organic carbon.

Region	Source	Technique	Period	Depths m	Estimate mol m ⁻²
Sargasso Sea	Jenkins and Goldman (1985)	AOU seasonal cycle	Apr-Nov	100 – 250	3.0 (O ₂)
Sargasso Sea	Sarmiento et al. (1990)	Tracer Distribution	Annual	100 – 250	0.6 – 3.3 (O ₂)
BATS	Carlson et al. (1994)	DOC changes	Apr-Nov	100 – 250	0.99 – 1.21 (C)
BATS	from Helmke et al. (2010)	POC Fluxes	Spring-Fall	110 – 250	0.38 (C)
BATS	Ono et al. (2001)	Tracer Model	Apr-Dec	100 – 250	2.08 ± 0.38 (O ₂)
				100 – 250	1.53 ± 0.35 (C)
				100 – 250	0.080 ± 0.046 (N)
BATS	This study	Tracer Model	Apr-Dec	110 – 250	1.81 ± 0.37 (O ₂)
					1.52 ± 0.30 (C)
					0.147 ± 0.043 (N)
ESTOC	from Helmke et al. (2010)	POC fluxes	Spring-Fall	110 – 250	0.08 (C)
ESTOC	This Study	Tracer Model	Apr-Dec	110 – 250	3.9 ± 1.0 (O ₂)
					1.53 ± 0.43 (C)
					0.04 ± 0.16 (N)
					mmol m ⁻² d ⁻¹
Canary Current	Arístegui et al. (2003)	ETS Activity	August	200 – 1000	7 (C)
Canary Current	Arístegui et al. (2005)	ETS/O ₂ <i>in vitro</i>	Summer	200 – 1000	68 (C)
Canary Current	Alonso-González et al. (2009)	OC balance	Fall	100 – 700	2.4 – 5.1 (C)