



## Abstract

Legumes can be considered as pioneer plants during ecosystem development, as they form a symbiosis with different nitrogen fixing rhizobia species, which enable the plants to grow on soils with low available nitrogen content. In this study we compared the abundance and diversity of nitrogen fixing microbes based on the functional marker gene *nifH*, which codes for a subunit of the Fe-protein of the dinitrogenase reductase, in nodules of different size classes of *Trifolium arvense* (L.). Additionally, carbon and nitrogen contents of the bulk soil and plant material were measured. Plants were harvested from different sites, reflecting 2 (2a) and 5 (5a) yr of ecosystem development, of an opencast lignite mining area in the south of Cottbus, Lower Lusatia (Germany) where the artificial catchment “Chicken Creek” was constructed to study the development of terrestrial ecosystems.

Plants from the 5a site revealed higher amounts of carbon and nitrogen, although *nifH* gene abundances in the nodules and carbon and nitrogen contents between the two soils did not differ significantly. Analysis of the *nifH* clone libraries showed a significant effect of the nodule size on the community composition of nitrogen fixing microbes. Medium sized nodules (2–5 mm) contained a uniform community composed of *Rhizobium leguminosarum* bv. *trifolij*, whereas the small nodules (< 2 mm) consisted of a diverse community including clones with non-*Rhizobium nifH* gene sequences. Regarding the impact of the soil age on the community composition a clear distinction between the small and the medium nodules can be made. While clone libraries from the medium nodules were pretty similar at both soil ages, soil age had a significant effect on the community compositions of the small nodules, where the proportion of *R. leguminosarum* bv. *trifolij* increased with soil age.

**BGD**

9, 13135–13160, 2012

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 1 Introduction

As a result of global change, in the last century an increase in initial ecosystems can be observed worldwide. Initial ecosystems can evolve naturally for example during the retreat of glaciers or in response to human activities mainly in areas with opencast mining activities. Therefore questions addressing food web dynamics, plant establishment and soil formation in initial ecosystems are not only of basic ecological interest but also of high commercial concern.

To address these questions an experimental catchment called “Chicken Creek” was constructed in the opencast lignite mining area in the south of Cottbus, Lower Lusatia (Germany) (Kendzia et al., 2008), which provides a unique opportunity to study initial ecosystem development. Surprisingly already after five years of soil development increasing complexity of food webs and increasing density of woody plants like birch, pine, and black locust was monitored (Elmer et al., 2011; Schaaf et al., 2010), which indicates a very fast progress in ecosystem development.

In this respect, monitoring of the vegetation revealed that the legume *Trifolium arvense* (L.) appeared as one of the pioneering plants already after two years of ecosystem development. Its abundance steadily increased in the following years and peaked after four years (Elmer et al., 2011). Already after 5 yr of ecosystem development significant concentrations of carbon in the soil developed, whereas the nitrogen pools remained low. These results underline the advantage of legumes in nitrogen poor ecosystems.

As legumes form symbiotic interactions with rhizobia, they are able to survive in soils with low amounts of available nitrogen as 75 and 90 % of the nitrogen needed for growth is provided by the symbiotic partner (Jacot et al., 2000). Boswell et al. (2007) demonstrated that up to  $11 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  can enter the soil by symbiotic nitrogen fixation from *T. arvense*. However, the successful establishment of the symbiosis strongly depends on (i) the survival of the bacteria in bulk soil mainly during autumn and winter, (ii) the colonization of the rhizosphere of the host plant, which in the case of clover, is followed

**BGD**

9, 13135–13160, 2012

### Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)**

S. Schulz et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

by root hair curling and the formation of an infection thread, and (iii) the intracellular infection and nodule development (Crespi and Gálvez, 2000; Sessitsch et al., 2002; Markmann and Parniske, 2009). These high numbers of different interim stages make the process prone to disturbances or environmental stresses. In principal, plants can only form a symbiosis with a restricted number of rhizobia for example *Trifolium* species preferably form a symbiosis with *Rhizobium leguminosarum* bv. *trifolii*, while for *Phaseolus* it is known that they can be infected by different *R. leguminosarum* species and even by *Burkholderia* (Doyle, 1998; Gyaneshwar et al., 2011). Furthermore it has been often postulated that the inter- and intra-species diversity of rhizobia, which are able to form nodules with selected legumes strongly depend on environmental conditions like soil pH and water content or biogeography. For example, it was shown that *R. etli* as microsymbiont of beans can be replaced by *R. tropici* under acidic conditions (Anyango et al., 1995; Sadowsky and Graham, 2006) and that diversity of rhizobia infecting chickpea (Alexandre et al., 2009) or *Lotus corniculatus* (Sotelo et al., 2011) depends on the investigated biogeographic region.

In this study we compared the diversity of nitrogen fixing microbes associated with *T. arvense* (L.) at the “Chicken Creek” catchment collected from nodules from a site 2 yr (2a) after initial ecosystem development started, where *T. arvense* recently appeared, and a site 5 yr (5a) after initial ecosystem development started, where the clover already exceeded highest density. We characterized the abundance and diversity of nitrogen fixing microbial communities in the nodules by targeting the *nifH* gene, which codes for the Fe-protein of the dinitrogenase reductase. Additionally, we analyzed nodule numbers, different soil parameter and carbon and nitrogen content of the plants. Due to that and the higher environmental stress level at the very beginning of ecosystem development, we assumed that the level of diversity of rhizobia nodulating clover will change with ongoing succession, being more diverse at the beginning of ecosystem development.

## 2 Material and methods

### 2.1 Sampling site

Samples were taken from two sites directly adjacent to the artificial catchment “Chicken Creek”, which is located in an opencast lignite mining area in the south of Cottbus, Lower Lusatia (N 51.360408, E 14.160988). This region is characterized by a sub-continental climate with a mean annual temperature of 8.9 °C and relatively low annual precipitation with 563 mm (Gerwin et al., 2009).

The catchment has a size of 6.5 ha and was composed of a sandy substrate (84.8 % sand, 9.1 % silt, 6.1 % clay) from Pleistocene sediments, which were deposited as terminal moraine during the Saale-glacial period. Below the sandy surface a base layer of Tertiary clay was constructed as a barrier for seepage water. Further details about the technical construction and the initial conditions are given by Gerwin et al. (2009) and Kendzia et al. (2008). The construction of the catchment was finished in September 2005, which was therefore defined as “point zero” for ecosystem development.

The sampling site in the east of the Chicken Creek was constructed as part of the catchment and has thus the same development stage. The second site in the west of the Chicken Creek was restored to “point zero” in 2008, thus the progress of soil development is three years behind the artificial catchment. At both sites three plots (3 × 3 m) were selected. Each subplot was 50 m apart from each other.

### 2.2 Sampling procedure and sample preparation

Sampling took place in July 2010. Thus, the eastern site had an age of 5 yr (5a) and the western one of 2 yr (2a). From each plot three *Trifolium arvense* (L.) plants at the flowering stage and bulk soil (0–5 cm) samples were taken. Plants were separated in green biomass and roots. The aboveground biomass was used to determine the carbon and nitrogen content. The roots were directly rinsed with sterile water and nodules from all plants were counted and classified in three groups of different size (< 2 mm, 2–5 mm,

**BGD**

9, 13135–13160, 2012

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and > 5 mm). Afterwards all nodules were surface sterilized by incubating the nodules for 5 min with 1 % Chloramin T (AppliChem, Germany) and three times rinsing with sterile water for 5 min (Singh, 1992). Sterility was tested by dipping the nodules on LB agar plates followed by an incubation of the plates for 48 h at 30 °C. Afterwards nodules were directly frozen at –20 °C. Bulk soil samples were stored at 4 °C for subsequent soil chemical analysis.

### 2.3 Chemical analysis of soil and plant samples

Each individual plant and bulk soil material was dried at 65 °C for 2 days. Afterwards, it was ball-milled (Retsch MM2, Retsch GmbH, Germany) and 1.5 mg plant material and 80 mg soil were weighted into 3.3 × 5 mm and 5 × 9 mm tin capsules (IVA Analy-sentechnik, Germany), respectively. The total carbon and nitrogen contents were measured with the Elemental-Analysator Euro-EA (Eurovector, Italy).

Nitrate (NO<sub>3</sub><sup>-</sup>-N), ammonium (NH<sub>4</sub><sup>+</sup>-N), dissolved organic carbon (DOC) and nitrogen (DON) of all soil samples were determined in CaCl<sub>2</sub> extracts. For the extraction, 20 g of fresh soil was overhead shaken for 45 min with 20 ml 0.01 M CaCl<sub>2</sub>. Afterwards, the extracts were filtered through a Millex HV Millipore filter (pore size, 0.45 μm). Nitrate, ammonium and DON were determined by continuous flow analysis with a photometric autoanalyzer (CFA-SAN Plus, Skalar Analytik, Germany). DOC was determined with the DIMA-TOC 100 (Dima Tec, Germany).

### 2.4 Nucleic acid extraction from nodules

For nucleic acid extraction, from each plant a small (< 2 mm) and a medium (2–5 mm) sized nodule was used. The small respectively medium nodules from the three plants per plot sampled were pooled and DNA was extracted by thermolysis. Plants from the three different plots were treated as true replicates. Nodules were crushed under sterile conditions with a pestle in 300 μl lysis buffer (Krasova-Wade and Neyra, 2007). The lysis buffer consisted of 0.2 mg ml<sup>-1</sup> ProteinaseK (Biozym, Germany), 0.05 % Tween20

**BGD**

9, 13135–13160, 2012

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Sigma, Germany) and 10 % Taq Polymerase PCR-Buffer (Invitrogen, Germany). Afterwards, samples were incubated for 1 h at 55 °C and 10 min at 95 °C. Cell debris were centrifuged for 5 min at 16 000 g (modified after Sambrook and Russell, 2001). The quantity of DNA in the supernatant was checked with a spectrophotometer (Nanodrop, PeqLab, Germany). Afterwards, samples were frozen at –20 °C.

## 2.5 Quantitative real-time PCR of the *nifH* gene

Quantitative real-time PCR was conducted on a 7300 Real-Time PCR System (Applied Biosystems, Germany) using SybrGreen as fluorescent dye. For the quantitative real-time PCR (qPCR) *nifH* primers described by Rösch et al. (2002) were applied, which produced a 458 bp PCR fragment. As standard curve a serial dilution ( $10^2$  to  $10^7$  copies  $\mu\text{l}^{-1}$ ) of a plasmid containing the *nifH* gene fragment of *Azospirillum irakense* DSM 11568 was used. The reaction mixture consisted of 12.5  $\mu\text{l}$  Power SybrGreen Master Mix (Applied Biosystems, Germany), 0.12  $\mu\text{M}$  of each primer (Metabion, Germany), 0.5  $\mu\text{l}$  of 3 % bovine serum albumin (Sigma, Germany), and 2  $\mu\text{l}$  DNA. The reaction volume was adjusted to 25  $\mu\text{l}$  with nuclease-free water. The thermal profile was as follows: a 10 min hot start at 95 °C, followed by 40 repetitions of 45 s at 95 °C, 45 s at 55 °C and 45 s at 72 °C. (Töwe et al., 2010). For confirming specificity of the amplicons after each PCR run a melting curve and a 2 % agarose gel were conducted. The amplification efficiency was calculated with the equation  $\text{Eff} = [10^{(-1/-\text{slope})} - 1]$  and resulted in 83 %.

## 2.6 Preparation of *nifH* gene clone libraries and sequencing

In total 12 clone libraries were prepared (2 sites, 2 nodule sizes, and 3 plots). For the amplification of *nifH* the same primers as for the qPCR were used. The reaction mixture consisted of 2.5  $\mu\text{l}$  TopTaq buffer, 5  $\mu\text{l}$  Q-Solution, 2.5  $\mu\text{l}$  of 2 mM dNTPs (Fermentas, Germany), 0.2  $\mu\text{M}$  of each primer (Metabion, Germany), 0.5  $\mu\text{l}$  BSA (3 %), 1  $\mu\text{l}$  TopTaq Polymerase (Qiagen, Germany) and 1  $\mu\text{l}$  DNA. The reaction volume was adjusted to

**BGD**

9, 13135–13160, 2012

### Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





25 µl with nuclease-free water. The thermal profile was as follows: a 10 min hot start at 94 °C, followed by 35 repetitions of 45 s at 94 °C, 45 s at 57 °C, 45 s at 72 °C and a final prolongation of 10 min at 72 °C. The PCR products were then checked for the right size of 458 bp and cleaned up with the Qiagen PCR Purification Kit.

For cloning and transformation the Zero Blunt® Topo® Cloning Kit (Invitrogen, Germany) was used as recommended by the manufacturer. After the transformation, 30 clones per sample were cultivated overnight in 5 ml LB containing 50 µg ml<sup>-1</sup> kanamycin. Plasmids were isolated with the NucleoSpin® Plasmid Kit (Macherey Nagel, Germany) and inserts were checked for the right size by EcoRI digestion (Fermentas, Germany). Finally, clones with the right size were used for sequencing.

Inserts were sequenced on the ABI PRISM® 3730 DNA Sequencer (Applied Biosystems, USA) using the BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems). The sequencing reaction consisted of 2 µl Buffer Descriptor Table Buffer, 2 µl Big Dye Terminator v3.1 Mix, 1 µl M13 forward sequencing primer (10 pmol), 100 ng DNA and was adjusted with nuclease free water to a final volume of 10 µl. The thermal profile started with 1 min 96 °C and was followed by 35 repetitions of 15 s at 96 °C, 15 s at 50 °C and 4 min at 60 °C.

## 2.7 Phylogenetic analysis

Sequences were verified using the BioEdit Sequence Alignment Editor v. 7.1.3. (Hall, 1999). Nucleotide sequences were translated in protein sequences with the ExpASY Translate Tool (<http://web.expasy.org/translate/>) and BlastX was used to confirm that the sequence codes for *nifH*. All sequences including stop codons or deletions were omitted from further analyses. In total, 233 sequences were used for further analyses, wherefrom 114 originated from the 2a site and 119 from the 5a site. Sequence alignments, construction of the phylogenetic tree and calculation of a distance matrix was done with the ARB software package ([www.arb-home.de](http://www.arb-home.de)) (Ludwig et al., 2004). For the phylogenetic analyses of the *nifH* transcripts, we used the updated *nifH* database from

**BGD**

9, 13135–13160, 2012

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Zehr et al. (2003) (<http://www.es.ucsc.edu/~wwwzehr/research/database/>). Sequences were imported to the databases, and the alignments were checked and manually corrected. The phylogenetic tree was calculated by applying the neighbor-joining algorithm (Saitou and Nei, 1987) (1000 bootstrap replications). All phylogenetic analyses were done on DNA level as strongly related *nifH* sequences can be expected in clone libraries from nodules (Opperdoes and Lemey, 2009).

Calculations of operational taxonomic units (OTU) were done with mothur v. 1.26.0. (Schloss et al., 2009). As estimators for diversity and evenness the Invsimpson and the Shannon index were applied, respectively. Retrieved unique *nifH* nucleic acid sequences are available at the GenBank database under the Accession No JX501529-JX501657. For details about the clones comprising identical sequences see Table S1.

## 2.8 Statistic

Statistical analyses were carried out with the R environment (<http://www.r-project.org>). Prior to analysis, data were tested for normal distribution by Q-Q plots and the Kolmogorov-Smirnoff test. If necessary data were log transformed prior to further analysis. Soil and plant parameter were tested for significant differences by one-factor ANOVA. Gene abundance data and the amount of the differently sized nodules per plant were tested by two-factor ANOVA. Pairwise comparisons between the different nodule size classes were done with t-tests, which were corrected for multiple testing with the Holm method. The influence of soil age and nodule size on the *nifH* gene diversity was tested on the basis of the distance matrix with Adonis, implementing 999 permutations.

**BGD**

9, 13135–13160, 2012

### Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3 Results

#### 3.1 Chemical analysis of soil and plant samples

Total carbon (TC) and nitrogen (TN) concentrations were higher in soil samples from the 5a site compared to the 2a site. However, whereas TN content was 1.36 times higher in the soil samples from the 5a site ( $5.91 \mu\text{g g}^{-1}$  respectively  $8.01 \mu\text{g g}^{-1}$ ), the total carbon content increased only about a factor of 1.1 from 136.91 (2a) to  $149 \mu\text{g g}^{-1}$  (5a), resulting in a decreased C/N ratio in soil samples from the 5a site. The nitrate concentrations were below the detection limit for both soils ( $< 0.3 \mu\text{g g}^{-1}$ ). Also ammonium concentrations were low, however higher values were measured in the 2a soil, reaching  $0.04 \mu\text{g g}^{-1}$  compared to  $0.02 \mu\text{g g}^{-1}$  in soil samples from the 5a site. In contrast dissolved organic nitrogen (DON) was higher in the 5a soil reaching  $0.46 \mu\text{g g}^{-1}$ . Dissolved organic carbon (DOC) concentrations did not differ significantly between soil samples from the two sites and were in the range of  $3.04 \mu\text{g g}^{-1}$  to  $3.53 \mu\text{g g}^{-1}$ .

Parallel to the higher DON and DOC values in the 5a soil, plants from the 5a site comprised significantly higher carbon and nitrogen contents, 41.4% carbon and 2.16% nitrogen compared to 37.1 and 1.68%. However, as both, carbon and nitrogen, increased, the C/N ratio of the plants did not differ significantly between both sites. Data are summarized in Table 1.

#### 3.2 Distribution of nodules

In total 297 nodules were collected from 18 plants, 154 from the 2a site and 143 from the 5a site. Nodules were classified into three classes: (i) small nodules ( $< 2 \text{ mm}$ ), (ii) medium nodules ( $2\text{--}5 \text{ mm}$ ), and (iii) large nodules ( $> 5 \text{ mm}$ ). The distribution of the nodules is shown in Fig. 1. Overall, the amount of nodules significantly decreased with their size ( $p < 0.001$ ). Thus, small nodules, made up 67% (107 nodules) in plants derived from the 2a soil and 80% (116) in plants derived from the 5a soil. Consequently, more medium and large nodules were collected from plants from the 2a site comprising

**BGD**

9, 13135–13160, 2012

### Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



26 and 7 %, respectively, while from plants sampled from the 5a site only 17 and 3 % belonged to the medium and large sized nodule class.

### 3.3 Abundance of *nifH* genes in the nodules

The amount of *nifH* gene copies per nodule is shown in Fig. 2. Overall, the *nifH* gene abundance ranged around  $5 \times 10^5$  copies per nodule in the small nodules independent from the site the plants were sampled. Regarding the medium nodules, similar values were measured for plants grown on the 5a soil, whereas *nifH* gene copy numbers per nodule were 2.5 times higher in the nodules from the 2a site. However, as the values strongly fluctuated no significant influence of soil age or nodule size was detected.

### 3.4 Phylogenetic diversity of nodule associated diazotrophs

The analysis of rarefaction curves on the 97 % DNA homology level revealed no differences in the diversity of *nifH* harboring microbes in medium sized nodules from plants grown on the two different sites (Fig. 3). The maximum number of OTUs per nodule was 3. If the rarefaction curves were analyzed on a 99 % similarity level for the medium sized nodules however, up to 6 OTUs for nodules from plants grown on the 2a site were detected, whereas the number of OTUs from nodules grown on the 5a site were significantly lower (3 OTUs), indicating a higher diversity in nodules of this size class in plants grown on the younger site. Overall the diversity of *nifH* harboring microbes was bigger in the small nodules compared to the medium sized nodules, where up to 10 OTUs (on a 99 % similarity level) could be detected. In contrast to the data described for the medium sized nodules, the number of OTUs from nodules that were assigned to the small size class was higher in plants grown on the 5a site. As the number of analyzed clones was too low for nodules of the small class from plants grown on the 5a site as well as for nodules of the medium size class grown on the 2a site to reach full coverage, which is needed for a quantitative diversity assessment, the following analytic steps were performed on a 97 % homology level.

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Based on this 97% homology calculation, all obtained *nifH* sequences were distributed among 9 OTUs (Table 2). In the clone libraries from the small and medium sized nodules from plants grown on the 2a site as well as the medium sized nodules grown on the 5a site, sequences clustered in three OTUs each, while calculations resulted in six OTUs for the sequences of the small nodules from plants grown on the 5a site.

Adonis calculations revealed a significant influence of the nodule size ( $p = 0.001$ ) on the *nifH* gene diversity. Regarding the small nodules, highest diversity was estimated for nodules from plants sampled on the 5a site with an Invsimpson value of 3.59, whereas the distribution of the sequences among the OTUs was not as even as for the small nodules from the 2a site, which were divided almost equally between OTU1, 5 and 7. As a result, clone libraries of the small nodules from the two sites were significantly different ( $p = 0.048$ ). In contrast to the small nodules, clone libraries from the medium nodules were very similar ( $p = 0.745$ ) and diversity and evenness were very low, as more than 95% of the sequences clustered in OTU5.

The phylogenetic affiliation of the *nifH* gene sequences, as depicted in Fig. 4, revealed that two main clusters were formed. From all sequenced clones 24 fell in Cluster I, while 209 could be assigned to Cluster II, which were interspersed throughout OTU5-8. Generally, Cluster II was composed of *Rhizobium* and *Sinorhizobium*, but all sequences exhibited strong homology with *Rhizobium leguminosarum* bv. *trifolii*. Despite the high similarity to *R. leguminosarum* bv. *trifolii*, sequence variations among the OTUs ranged from 100 to 92.4% similarity. Surprisingly besides sequences which were closely related to rhizobial *nifH* sequences, the OTUs, which could be assigned to cluster I, were closely related to *Bradyrhizobia*, *Burkholderia*, *Leptothrix* and *Azospirillum*. Based on the phylogenetic affiliation it is very likely that OTU9 is homolog to *Azospirillum*. OTU1-4 were assigned to a very diverse subcluster composed of *Azospira*, *Stenotrophomas* and *Burkholderia*, *Azoarcus* and *Leptothrix*, which made a clear affiliation almost impossible.

## 4 Discussion

A lot of research has been done on legume-*Rhizobium* symbiosis in the last century mainly in the context of agriculture to improve efficiency of plant microbe interactions with the aim to enhance yields (Fischer et al., 2012; Perrineau et al., 2011; Naeem et al., 2004). However, little is known about that interaction of rhizobia and legumes in initial ecosystems, where nutrients and microbial biomass are low and inoculation with *Rhizobium* strains does not occur. Consequently in these ecosystems the success of a symbiosis depends on the indigenous rhizobial community being already present in the soil. Therefore it was the aim of this study to compare the diversity of *nifH* carrying diazotrophs in the nodules of the pioneering clover *Trifolium arvense* (L.), taken from a 2 yr (2a) and a 5 yr (5a) old soil. Both sites developed under similar climatic conditions and from the same starting material, thus these effects as drivers for the observed differences in diversity pattern could be excluded.

As we were especially interested in rhizobia, which have the potential to fix atmospheric N<sub>2</sub>, this study focused on the analysis of the *nifH* gene instead of other house-keeping genes or genes involved in nodulation (Sotelo et al., 2011; Rogel et al., 2011; Talbi et al., 2010). Thus, we can exclude that *nod*<sup>+</sup> but *nifH*<sup>-</sup> rhizobia were investigated, which are not advantageous for the nitrogen budget of the host plant. Moreover, analyses were done on the level of DNA to distinguish sequences of closely related biovars. For example NifH of *R. leguminosarum* bv. *trifolii* and bv. *viciae* share 95% of the amino acids, while on DNA level similarity of *nifH* is below 85%.

Measurements of the nitrogen content of the plants showed that the clover from the 5a site exhibited significantly higher nitrogen concentrations, although DON and NH<sub>4</sub><sup>+</sup> did not differ significantly between the two soils. Therefore, higher nitrogen contents can be related to higher nitrogen fixation rates, which might be a fact of higher number of nodules per plant or a generally higher activity of nitrogen fixing microbes. Voisin et al. (2010) could prove that nodule formation is induced under nitrate limitation and as no nitrate was detectable at both sites, it is not surprising that plants developed a

**BGD**

9, 13135–13160, 2012

### Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



relatively high number of nodules. In average, 16 indeterminate nodules per plant were detected, which is comparable or even higher compared to other studies where systems without inoculation of rhizobial strains were studied (Naeem et al., 2004; Wielbo et al., 2010). The evaluation of the amount and distribution of small, medium and large nodules resulted in no significant difference between plants from the two different sites. Therefore, higher nitrogen contents of the plants from the 5a site might be attributed to more efficient strains which comprise higher nitrogen fixing activities compared to strains that are associated with nodules of plants derived from the 2a site. To address that question, *nifH* clone libraries and the influence of soil age on the nodule community were compared.

Clone libraries of the medium sized nodules showed similar diversity pattern for both sites. The community was mainly composed of *Rhizobium leguminosarum* bv. *trifolii*, which exclusively formed one OTU (Table 2). *R. leguminosarum* bv. *trifolii* is the common and most effective strain infecting *Trifolium* species (Doyle, 1998). It is known that the plants are able to control nodule size and therefore promote growth of nodules formed by effective rhizobia (Voisin et al., 2010; Kouchi and Yoneyama, 1984). Hence it is very likely that the medium sized nodules represent the most active nodules, which is further underlined by the tendency of higher *nifH* copy numbers in the nodules of the medium size class mainly in plants from the 2a site and by findings from Tajima et al. (2007), who stated that at the flowering stage of peanuts nitrogen fixation activity and nodule size are positively correlated.

However, obviously the medium sized nodules did not drive the difference between the sites. In contrast to the medium sized nodules, Adonis permutation tests revealed a significant influence of soil age on the *nifH* harboring microbial community of the small sized nodules, which is reflected by a high *nifH* gene diversity and evenness in the nodules from plants derived from the 2a site. The differences are mainly based on an increase of clones coding for *R. leguminosarum* bv. *trifolii* and in addition to a decrease of species from cluster I. The increasing amount of clones which harbor the insert of *nifH* related to *R. leguminosarum* bv. *trifolii*, especially from OTU5, where also

**BGD**

9, 13135–13160, 2012

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the clones from the medium nodules clustered, might be attributed to an enrichment effect of that bacteria in the 5a soil, where *T. arvense* was already present in the previous years. As *T. arvense* is an annual plant species nodule inhabiting bacteria are frequently released at the end of the vegetation period (Sadowsky and Graham, 2006).

5 In contrast to inoculated species, which are often replaced by the indigenous species until the next vegetation period, naturally occurring rhizobia, which were enriched in the nodules, survive (Sessitsch et al., 2002). Consequently, the probability of a successful development of a legume-*Rhizobium* symbiosis increased year by year. However, one might ask why these nodules did not develop to a medium nodule. Due to low nutrient contents in the soil and the fact that the plant does not spend more energy than needed in nodule production, it is obvious that there might be not enough energy available to promote the maturation of more medium or large nodules. It is known that nodules formed in response to the presence of indigenous rhizobia in soil often exhibit a higher diversity (Liu et al., 2007; Zakhia et al., 2006) compared to systems where high numbers of selected rhizobial strains have been inoculated. In the small nodules from the 5a site highest rhizobial diversity was observed. While most sequences belonged to OTU5, which is the dominant OTU from the medium nodules, some sequences clustered in OTU6, 7 and 8. These organisms might play an important role to setup optimal conditions for nitrogen fixation in the nodules and improve fitness of effective *R. leguminosarum* bv. *trifolii* strains. Moreover, the presence of other than *Rhizobium* sequences (OTU1-4, 9) might influence nodule establishment. As shown for *Pseudomonas*, *Bacillus*, *Sphingomonas* and others this might be due to co-invasion of rhizosphere bacteria (Zakhia et al., 2006; Muresu et al., 2008). Therefore it is not surprising that similarities with typical species that colonize the rhizosphere of annual plants were found in cluster I. However, the role of non-rhizobial species in nodules is discussed controversially. For example Plazinksi et al. (1985) showed that *Azospirillum* inhibited the symbiosis of *Rhizobium* with *Trifolium repens*. In contrast, the presence of *Pseudomonas* species can improve nodulation (Bolton et al., 1990). As the cluster I was very diverse, it is difficult to figure out the specific role of the sequences found.



## 5 Conclusion

In summary the data indicate that the plants from the 5a site were more effective in nitrogen fixation, as their nitrogen content was significantly higher, although the ammonium concentrations in soil did not differ significantly. This development might be rather attributed to a change in the legume-*Rhizobium* symbiosis than an overall increase of nodules. In this regard, the hypothesis, that higher *nifH* diversity can be expected in the nodules from the 2a soil, was partly disproven. While no difference was observed for the medium nodules Adonis calculations revealed a significant influence of soil age on the diversity of the small nodules. Furthermore the proportion of *R. leguminosarum* bv. *trifolii* clones was higher at plants from the 5a site, what might be mainly attributed to an enrichment of this bacteria due to the longer *T. arvense* history of this site and might be the reason for the higher nitrogen content in those plants. In this regard, it would be interesting for future studies to address the question if similar observations can be made for perennial legumes like *Lotus corniculatus* (L.), which is also a pioneering plant at the Chicken Creek (Schaaf et al., 2010), where it is known that nitrogen is accumulated in the rhizosphere (Boldt et al., 2012) and how the development continues during the next phases of ecosystem development.

**Supplementary material related to this article is available online at:**

**<http://www.biogeosciences-discuss.net/9/13135/2012/>**

**[bgd-9-13135-2012-supplement.pdf](#)**

*Acknowledgements.* This study is part of the Transregional Collaborative Research Centre 38 (SFB/TRR38) which is financially supported by the Deutsche Forschungsgemeinschaft (DFG, Bonn) and the Brandenburg Ministry of Science, Research and Culture (MWFK, Potsdam). We thank Vattenfall Europe Mining AG for providing the research site Chicken Creek.

**BGD**

9, 13135–13160, 2012

### Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## References

- Alexandre, A., Brígido, C., Laranjo, M., Rodrigues, S., and Oliveira, S.: Survey of chickpea rhizobia diversity in Portugal reveals the predominance of species distinct from *Mesorhizobium ciceri* and *Mesorhizobium mediterraneum*, *Microb. Ecol.*, 58, 930–941, 2009.
- 5 Anyango, B., Wilson, K. J., Beynon, J. L., and Giller, K. E.: Diversity of rhizobia nodulating *Phaseolus vulgaris* L. in two Kenyan soils with contrasting pHs, *Appl. Environ. Microbiol.*, 61, 4016–4021, 1995.
- Boldt, K., Uwe Schneider, B., Fritsch, S., and Hüttl, R. F.: Influence of root growth of two pioneering plant species on soil development during the initial stage of ecosystem genesis in the Lusatian post mining landscape, *Can. J. Soil. Sci.*, 92, 67–76, 2012.
- 10 Bolton, H., Elliott, L., Turco, R., and Kennedy, A.: Rhizoplane colonization of pea seedlings by *Rhizobium leguminosarum* and a deleterious root colonizing *Pseudomonas* sp. and effects on plant growth, *Plant Soil*, 123, 121–124, 1990.
- Boswell, C. C., Lowther, W. L., and Rutherford, A. J.: Symbiotic nitrogen fixation by *Trifolium arvense* in semi-arid short tussock grasslands, *New Zeal. J. Agr. Res.*, 50, 511–521, 2007.
- 15 Crespi, M. and Gálvez, S.: Molecular mechanisms in root nodule development, *J. Plant Growth Regul.*, 19, 155–166, 2000.
- Doyle, J. J.: Phylogenetic perspectives on nodulation: evolving views of plants and symbiotic bacteria, *Trends Plant Sci.*, 3, 473–478, 1998.
- 20 Elmer, M., Schaaf, W., Biemelt, D., Gerwin, W., and Hüttl, R. F.: The artificial catchment “Chicken Creek” – initial ecosystem development 2005–2010, *Research Centre Landscape Development and Mining Landscapes (FZLB)*, Cottbus, 168, 2011.
- Fischer, D., Pfitzner, B., Schmid, M., Simoes-Araujo, J. L., Reis, V. M., Pereira, W., Ormeno-Orrillo, E., Hai, B., Hofmann, A., Schloter, M., Martinez-Romero, E., Baldani, J. I., and Hartmann, A.: Molecular characterisation of the diazotrophic bacterial community in uninoculated and inoculated field-grown sugarcane (*Saccharum* sp.), *Plant Soil*, 356, 83–99, 2012.
- 25 Gerwin, W., Schaaf, W., Biemelt, D., Fischer, A., Winter, S., and Hüttl, R. F.: The artificial catchment “Chicken Creek” (Lusatia, Germany) – a landscape laboratory for interdisciplinary studies of initial ecosystem development, *Ecol. Eng.*, 35, 1786–1796, 2009.
- 30 Gyaneshwar, P., Hirsch, A. M., Moulin, L., Chen, W.-M., Elliott, G. N., Bontemps, C., Estrada-de los Santos, P., Gross, E., dos Reis, F. B., Sprent, J. I., Young, J. P. W., and James, E. K.:

---

### Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Legume-nodulating *Betaproteobacteria*: diversity, host range, and future prospects, Mol. Plant Microbe In., 24, 1276–1288, 2011.

Hall, T. A.: BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT, Nucl. Acid S., 41, 95–98, 1999.

Jacot, K. A., Lüscher, A., Nösberger, J., and Hartwig, U. A.: Symbiotic N<sub>2</sub> fixation of various legume species along an altitudinal gradient in the Swiss Alps, Soil Biol. Biochem., 32, 1043–1052, 2000.

Kendzia, G., Reißmann, R., and Neumann, T.: Targeted development of wetland habitats for nature conservation fed by natural inflow in the post-mining landscape of Lusatia, World Min., 2, 88–95, 2008.

Kouchi, H. and Yoneyama, T.: Dynamics of carbon photosynthetically assimilated in nodulated soya bean plants under steady-state conditions 1. Development and application of <sup>13</sup>C<sub>2</sub> assimilation system at a constant <sup>13</sup>C abundance, Ann. Bot., 53, 875–882, 1984.

Krasova-Wade, T. and Neyra, M.: Optimization of DNA isolation from legume nodules, Lett. Appl. Microbiol., 45, 95–99, 2007.

Liu, X., Wang, E., Li, Y., and Chen, W.: Diverse bacteria isolated from root nodules of *Trifolium*, *Crotalaria* and *Mimosa* grown in the subtropical regions of China, Arch. Microbiol., 188, 1–14, 2007.

Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadhukumar, Buchner, A., Lai, T., Steppi, S., Jobb, G., Förster, W., Brettske, I., Gerber, S., Ginhart, A. W., Gross, O., Grumann, S., Hermann, S., Jost, R., König, A., Liss, T., Lüßmann, R., May, M., Nonhoff, B., Reichel, B., Strehlow, R., Stamatakis, A., Stuckmann, N., Vilbig, A., Lenke, M., Ludwig, T., Bode, A., and Schleifer, K. H.: ARB: a software environment for sequence data, Nucleic Acids Res., 32, 1363–1371, 2004.

Markmann, K. and Parniske, M.: Evolution of root endosymbiosis with bacteria: how novel are nodules?, Trends Plant. Sci., 14, 77–86, 2009.

Muresu, R., Polone, E., Sulas, L., Baldan, B., Tondello, A., Delogu, G., Cappuccinelli, P., Alberghini, S., Benhizia, Y., Benhizia, H., Benguedouar, A., Mori, B., Calamassi, R., Dazzo, F. B., and Squartini, A.: Coexistence of predominantly nonculturable rhizobia with diverse, endophytic bacterial taxa within nodules of wild legumes, FEMS Microbiol. Ecol., 63, 383–400, 2008.

Naeem, F. I., Ashraf, M. M., Malik, K. A., and Hafeez, F. Y.: Competitiveness of introduced *Rhizobium* strains for nodulation in fodder legumes, Pak. J. Bot., 36, 159–166, 2004.

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Opperdoes, F. R. and Lemey, P.: Phylogenetic analysis using protein sequences, in: The Phylogenetic Handbook, 2 edn., edited by: Lemey, P., Salemi, M., and Vandamme, A.-M., Cambridge University Press, Cambridge, 313–342, 2009.
- Perrineau, M. M., Le Roux, C., de Faria, S. M., de Carvalho Balieiro, F., Galiana, A., Prin, Y., and Béna, G.: Genetic diversity of symbiotic *Bradyrhizobium elkanii* populations recovered from inoculated and non-inoculated *Acacia mangium* field trials in Brazil, *Syst. Appl. Microbiol.*, 34, 376–384, 2011.
- Plazinski, J. and Rolfe, B. G.: Interaction of azospirillum and rhizobium strains leading to inhibition of nodulation, *Appl. Environ. Microbiol.*, 49, 990–993, 1985.
- Rogel, M. A., Ormeño-Orrillo, E., and Martínez Romero, E.: Symbiovars in rhizobia reflect bacterial adaptation to legumes, *Syst. Appl. Microbiol.*, 34, 96–104, 2011.
- Rösch, C., Mergel, A., and Bothe, H.: Biodiversity of denitrifying and dinitrogen-fixing bacteria in an acid forest soil, *Appl. Environ. Microbiol.*, 68, 3818–3829, 2002.
- Sadowsky, M. and Graham, P.: Root and stem nodule bacteria of legumes in: The Prokaryotes, edited by: Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.-H., and Stackebrandt, E., Springer Science + Business Media, New York, 818–841, 2006.
- Saitou, N. and Nei, M.: The neighbor-joining method: a new method for reconstructing phylogenetic trees, *Mol. Biol. Evol.*, 4, 406–425, 1987.
- Sambrook, J. and Russell, D. W.: *Molecular Cloning – A Laboratory Manual*, 3 edn., Cold Spring Harbor Laboratory Press, New York, 2001.
- Schaaf, W., Biemelt, D., and Hüttl, R. F.: Initial development of the artificial catchment “Chicken Creek” – monitoring program and survey 2005–2008, *Research Centre Landscape Development and Mining Landscapes (FZLB)*, Cottbus, 2010.
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski, R. A., Oakley, B. B., Parks, D. H., Robinson, C. J., Sahl, J. W., Stres, B., Thallinger, G. G., Van Horn, D. J., and Weber, C. F.: Introducing mothur: open-source, platform-independent, vommunity-supported software for describing and comparing microbial communities, *Appl. Environ. Microbiol.*, 75, 7537–7541, 2009.
- Sessitsch, A., Howieson, J. G., Perret, X., Antoun, H., and Romero, M. I.: Advances in *Rhizobium* research, *Crit. Rev. Plant Sci.*, 21, 323–378, 2002.
- Singh, C. S.: Prevalence of *Azospirillum* within the stem nodules of *Aeschynomene* spp. and *Neptunia* sp., *Zbl. Mikrobiol.*, 147, 455–458, 1992.

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Sotelo, M., Irisarri, P., Lorite, M. J., Casaretto, E., Rebuffo, M. N., Sanjuán, J., and Monza, J.: Diversity of rhizobia nodulating *Lotus corniculatus* grown in northern and southern regions of Uruguay, *Appl. Soil Ecol.*, 49, 197–207, 2011.
- Tajima, R., Lee, O. N., Abe, J., Lux, A., and Morita, S.: Nitrogen-fixing activity of root nodules in relation to their size in peanut (*Arachis hypogaea* L.), *Plant Prod. Sci.*, 10, 423–429, 2007.
- Talbi, C., Delgado, M. J., Girard, L., Ramirez-Trujillo, A., Caballero-Mellado, J., and Bedmar, E. J.: *Burkholderia phymatum* strains capable of nodulating *Phaseolus vulgaris* are present in Moroccan soils, *Appl. Environ. Microbiol.*, 76, 4587–4591, 2010.
- Töwe, S., Albert, A., Kleineidam, K., Brankatschk, R., Dümig, A., Welzl, G., Munch, J., Zeyer, J., and Schloter, M.: Abundance of microbes involved in nitrogen transformation in the rhizosphere of *Leucantheropsis alpina* (L.) heywood grown in soils from different sites of the Damma glacier forefield, *Microb. Ecol.*, 60, 762–770, 2010.
- Voisin, A.-S., Munier-Jolain, N., and Salon, C.: The nodulation process is tightly adjusted to plant growth. An analysis using environmentally and genetically induced variation of nodule number and biomass in pea, *Plant Soil*, 337, 399–412, 2010.
- Wielbo, J., Marek-Kozaczuk, M., Mazur, A., Kubik-Komar, A., and Skorupska, A.: Genetic and metabolic divergence within a *Rhizobium leguminosarum* bv. *trifolii* population recovered from clover nodules, *Appl. Environ. Microbiol.*, 76, 4593–4600, 2010.
- Zakhia, F., Jeder, H., Willems, A., Gillis, M., Dreyfus, B., and de Lajudie, P.: Diverse bacteria associated with root nodules of spontaneous legumes in Tunisia and first report for *nifH*-like gene within the genera *Microbacterium* and *Starkeya*, *Microb. Ecol.*, 51, 375–393, 2006.
- Zehr, J. P., Jenkins, B. D., Short, S. M., and Steward, G. F.: Nitrogenase gene diversity and microbial community structure: a cross-system comparison, *Environ. Microbiol.*, 5, 539–554, 2003.

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)

S. Schulz et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 1.** Chemical parameters of bulk soil and plants. Carbon and nitrogen contents were measured in bulk soil samples from the 2a and the 5a site ( $n = 3$ , standard deviations in parenthesis). Plant carbon and nitrogen contents were measured in *Trifolium arvense* (L.) from the 2a and the 5a site ( $n = 9$ , standard deviations in parenthesis). Nitrate concentration was below the detection limit for all samples ( $< 0.3 \mu\text{g g}^{-1}$ ). A significant impact of soil age on the measured parameters was tested with one-factor ANOVA ( $p < 0.05$ ) and indicated by an asterisk.

	soil ( $\mu\text{g g}^{-1}$ )					plant (%)		
	$\text{NH}_4^+$	DON	TN	DOC	TC	C*	N*	C/N
2a	0.04 (0.02)	0.33 (0.03)	5.91 (2.54)	3.04 (0.64)	136.91 (8.44)	37.1 (4.7)	1.68 (0.3)	22.32 (2.9)
5a	0.02 (0.003)	0.46 (0.12)	8.01 (0.93)	3.53 (0.38)	149 (3.09)	41.4 (1.0)	2.16 (0.46)	19.93 (3.9)

## Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)


S. Schulz et al.

**Table 2.** Distribution of sequences among different operational taxonomic units (OTU). In total sequences were consolidated in 4 groups: 2a small, 2a medium, 5a small and 5a medium. The subsequent classification of sequences in different OTUs was done with mothur v. 1.26.0. on a 97 % similarity level for each sample individually. As indices for diversity and evenness the Invsimpson index and the Shannon Evenness index were applied, respectively.

OTU	number of sequences			
	2a small	2a medium	5a small	5a medium
1	13	2	–	–
2	–	–	6	–
3	–	–	1	–
4	–	–	–	1
5	15	74	19	77
6	–	–	6	–
7	9	1	6	–
8	–	–	2	–
9	–	–	–	1
Invsimpson	3.08	1.08	3.59	1.05
Shannon Evenness	0.98	0.17	0.72	0.12

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



**Fig. 1.** Distribution of the different nodule size classes. Nodules from each site were grouped after their size in three classes: < 2 mm (small), 2–5 mm (medium) and > 5 mm (large) ( $n = 9$ , error bars represent standard deviations). Significant differences between the three nodule size classes are indicated by different capital letters for the 2a site and by small letters for the 5a site. Pairwise comparisons were done with a  $t$ -test where the Holm method was applied for multiple testing.

**Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)**

S. Schulz et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 2.** Copy numbers per nodule of the functional gene *nifH*. Gene abundances were measured in small and medium sized nodules from the 2a and the 5a site ( $n = 3$ , error bars represent standard deviations).

**Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)**

S. Schulz et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)**

S. Schulz et al.




**Fig. 3.** Rarefaction curves of the different clone libraries. In total sequences were consolidated in 4 groups: 2a small, 2a medium, 5a small and 5a medium. Calculations were made on 99 and 97% similarity level for the small (left) and the medium (right) sized nodules from plants grown on the 2a and the 5a site. Calculations were done with mothur v. 1.26.0.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Diversity pattern of N fixing microbes in nodules of *Trifolium arvense* (L.)**

S. Schulz et al.



**Fig. 4.** Phylogenetic tree of the obtained *nifH* gene sequences in comparison to previously published sequences (accession numbers in brackets), analyses were conducted with the ARB software package. The tree was calculated using the neighbor-joining algorithm. Bootstrap values are given in percent next to the branches (1000 replicates). The outgroup is composed of *Methanopyrus kandleri* AV19 (AE009439), *Methanosarcina acetivorans* C2A (AE010299) and *Clostridium kluyveri* DSM 555 (CP000673).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

