Biogeosciences Discuss., 9, 17519–17538, 2012 www.biogeosciences-discuss.net/9/17519/2012/ doi:10.5194/bgd-9-17519-2012 © Author(s) 2012. CC Attribution 3.0 License.



This discussion paper is/has been under review for the journal Biogeosciences (BG). Please refer to the corresponding final paper in BG if available.

Phosphorus sorption and buffering mechanisms in suspended sediments from the Yangtze Estuary and Hangzhou Bay, China

M. Li^1 , M. J. Whelan², G. Wang³, and S. M. White²

¹College of Environmental Science and Engineering, Beijing Forestry University, Beijing 100083, China

²School of Applied Sciences, Cranfield University, Bedfordshire, MK43 0AL, UK ³State Key Laboratory of Hydroscience and Engineering, Tsinghua University,

Beijing 100084, China

Received: 19 October 2012 - Accepted: 19 November 2012 - Published: 10 December 2012

Correspondence to: M. Li (liminbjfu@126.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.





Abstract

The adsorption isotherm and the mechanism of the buffering effect are important controls on phosphorus behaviors in estuaries and are important for estimating phosphate concentrations in aquatic environments. In this paper, we derive phosphate adsorption isotherms in order to investigate sediment adsorption and buffering capacity for phosphorus discharged from sewage outfalls in the Yangtze Estuary and Hangzhou Bay near Shanghai, China. Experiments were also carried out at different temperatures in order to explore the buffering effects for phosphate. The results show that P sorption in sediments with low fine particle fractions was best described using exponential equations. Some P interactions between water and sediment may be caused by the precipitation of CaHPO₄ from Ca²⁺ and HPO₄²⁻ when the phosphate concentration in the liquid phase is high. Results from the buffering experiments suggest that the Zero Equilibrium Phosphate Concentrations (EPC₀) vary from 0.014 mgI⁻¹ to 0.061 mgI⁻¹, which are consistent with measured phosphate concentrations in water samples col-

¹⁵ lected at the same time as sediment sampling. Values of EPC_0 and linear sorption coefficients (*K*) in sediments with high fine particle and organic matter contents are relatively high, which implies that they have high buffering capacity. Both EPC_0 and *K* increase with increasing temperature, indicating a higher P buffering capacity at high temperatures.

20 1 Introduction

25

Phosphorus (P) is a major limiting nutrient in many surface waters and is often responsible for triggering eutrophication and toxic algal blooms (Kawasaki et al., 2010; Pratt and Shilton, 2009). In estuarine and coastal waters, P exchange at seawater-sediment interfaces can exert significant influence on P availability and it is, therefore, important to understand and describe this process well. Three equations (Langmuir, Freundlich





and Temkin) are commonly used to describe sorption equilibria in water-sediment systems.

P can rapidly react with particle surfaces in natural systems (Froelich, 1988; Borggaard et al., 2004). The physico-chemical reactions of P between overlying wa ter and sediment in natural systems include adsorption/desorption and precipitation/dissolution reactions, which together, regulate soluble P concentrations in natural rivers, streams and estuaries (Wauchope and McDowell, 1984; Price et al., 2010). This phenomenon is known as the "phosphate buffer mechanism" (Richardson, 1985), which assumes that sediment, whether benthic or suspended, maintain dissolved re active phosphate concentrations at some near-constant value regardless of biological removal and exogenic inputs (Froelich, 1988). P is sorbed onto particles in rivers, and

- removal and exogenic inputs (Froelich, 1988). P is sorbed onto particles in rivers, and then carried into estuaries. In estuaries, where freshwater and seawater mix, phosphorus is desorbed from particles to augment dissolved phosphate, especially in anoxic, saline sediment (Conley et al., 1995). This buffering effect is partly attributed to the adsorption of P onto metal oxide surfaces. Particulate phosphorus associated with
- iron and aluminium oxides often decreases with increasing salinity, suggesting that dissolved phosphate concentrations might be partly buffered by desorption from aluminium and iron oxide phases (Conley et al., 1995; Schippers et al., 2006).

The Yangtze Estuary in Eastern China frequently experiences eutrophication prob lems which are often manifested as red tides and fish kills (Xiang, 2011). Phosphate, primarily from treated wastewater outfalls is believed to be the principal driver for these phenomena (Li et al., 2004). High sediment loads are transported to the estuary by the Yangtze River every year (Liu et al., 2008). This causes the aggregation of sediment particles in the Yangtze Estuary and Hangzhou Bay, which in turn maintains the availability of dissolved-phase P via adsorption and/or desorption reactions.

In this paper, P sorption behavior in sediments collected close to wastewater outfalls in the Yangtze Estuary and Hangzhou Bay was investigated. The suitability of three commonly used sorption isotherm models was evaluated, along with the potential for predicting the dissolved reactive phosphate concentration by buffering experiments.





2 Materials and methods

2.1 Sampling

Surface sediment samples (0–2 cm) were collected from effluent outfall areas in the Yangtze Estuary and Hangzhou Bay using a CN-11 sediment sampler (Beijing
Zhuochuan Electronic Technology Co., Ltd.) (Li et al., 2004). The locations of the three outfalls and the five sediment sampling sites are shown in Fig. 1. Two sampling sites (Z4 and W4) are located around the Zhuyuan and Bailonggang outfall areas in the Yangtze Estuary. The distances between sampling sites and the outfalls are all approximately 0.5 km. These two outfalls discharge effluent from municipal wastewater treatment plants with chemical enhanced primary treatment processes. The other three sampling sites (A, B and F) are located in the area around the Xinghuo outfall, which discharges effluent from an industrial park wastewater treatment plant, in Hangzhou Bay. The distance between sampling site B and the outfall is also 0.5 km. Three sedi-

¹⁵ ments were obtained at each sampling site and stored together in one air-sealed plastic bag at 4 °C before being transported back to the laboratory (within 12 h). All sediments were subsequently air-dried, homogenized and passed through a 0.15-mm sieve and then stored in wide-mouthed amber glass jars in a dry, ventilated place.

A description and characterization of all samples is shown in Table 1. The organic ²⁰ matter (OM) content in the sediment was determined by the potassium bichromatedilution heat colorimetric method (Bao and Jiang, 1999). The particle size distributions of the sediment samples on a weight basis were determined by the pipette method (Beuselinck et al., 1998). Water samples were also collected around each sediment sampling site at the same time, which were analysed for dissolved inorganic phos-²⁵ phate (DIP) concentration (Table 1). Triplicate samples were used for all determinations and results are presented as means of three replicate analyses. Standard errors (not shown) were always less than ±5% of the mean value.





2.2 Isothermal adsorption experiments

Phosphate sorption isotherms were determined experimentally. Aliquots of dry, sieved and ground sediment (0.5 g ± 0.0001 g) were placed in 250-ml glass conical flasks. Then, artificial sea water (100 ml, salinity 5), prepared according to the method proposed by Liu et al. (1982), with P were added to each flask. This solution was spiked with KH₂PO₄ to give one of seven nominal phosphorus concentrations (0.5, 2, 4, 8, 12, 16 and 20 mg Pl⁻¹). Blank flasks containing no sediment were also prepared. Initial P concentrations in each incubation solution were determined by molybdate colorimetry (Kara et al., 1997). pH, including the blank experiments, was adjusted to a value of 8 using diluted KOH or HNO₃ in order to minimize pH variations due to the different amounts of KH₂PO₄ added. The conical flasks were sealed and shaken continuously at 200 rpm and 25 ± 1 °C for 24 h and then centrifuged at 3000 rpm (maximum relative centrifugal force: 1207 g) for 10 min. Phosphate concentrations in the supernatant after centrifugation were determined and assumed to be the final P concentration. The difference between initial and final P concentrations was used to calculate phosphate adcording to a phosphate.

adsorption capacity. Triplicate experiments were conducted for all samples and results were presented as means of three parallel determinations. Standard errors are not shown here since they were always less than $\pm 5\%$ of the mean value.

2.3 Phosphate buffer experiments

Phosphate buffering experiments, similar to the adsorption experiments described above, were carried out, except using very low initial P concentrations of 0, 0.04, 0.08, 0.12, 0.2, 0.3, 0.4 and 0.5 mg Pl⁻¹, which represent the range of concentrations commonly encountered under natural conditions. The experiments were conducted at three constant temperatures of 12 °C, 22 °C and 32 °C, respectively, to investigate the influees ence of temperature on buffering capacity.

 $[\mathsf{P}_D]_{initial}$ represents the initial liquid-phase phosphate concentration before shaking, while $[\mathsf{P}_D]_{final}$ represents the equilibrium liquid-phase phosphate concentration. The





difference between $[P_D]_{initial}$ and $[P_D]_{final}$ is expressed as ΔP_D . P is absorbed by sediments when ΔP_D is positive and desorbed from sediments when ΔP_D is negative. ΔP_S represents the P adsorbed or desorbed per unit mass of sediment.

3 Results and discussion

20

25

5 3.1 Adsorption isotherm models

The adsorption isotherms of five sediment samples are shown in Fig. 2a. Langmuir, Freundlich and Temkin adsorption isotherm equations were well known and used in many previous studies (Borggaard et al., 2005). These three isotherm equations were obtained by least-squares fitting to experimental sorption data. The isotherm equations, derived constants (Γ∞, *b*, *K* and *n*), correlation coefficients (*r*) and standard deviation (*S*) are shown in Table 2. The theoretical Langmuir and exponential isotherms are also plotted in Fig. 2a. Samples A, Z4 and W4 were all well described using the Langmuir adsorption isotherm with similar parameters. These curves are clearly distinct from the best-fit sorption isotherms obtained for samples B and F which were best described
using exponential isotherms.

It is clear from Table 2 that the P-adsorption data for samples A, Z4 and W4 can be well described using all three isotherm models ($r \ge 0.968$). The best suitable isotherm model for these three sediments was selected on the basis of the highest mean correlation coefficients (r) and lowest mean standard deviation (S) values for the three samples. Overall, the Langmuir equation was considered to give the best fit for these samples.

Sorption data from samples B and F were not well described by any of the isotherm equations (0.463 < r < 0.801). Visual inspection of the sorption data for these samples (Fig. 2a) suggests that an exponential model may be more appropriate for describing sorption in these samples. Least squares fitting of exponential equations to data from





samples B and F yielded the following results:

 $B: \Gamma = 16.464 \exp(0.1125c) \quad r = 0.930$

 $F: \Gamma = 30.65 \exp(0.0588c)$ r = 0.844

- ⁵ Equilibrium P-adsorption capacity versus initial liquid-phase phosphate concentrations for samples B and F are shown in Fig. 2b. When the initial P concentrations of the solution are lower than 12mgl⁻¹, the P-adsorption capacities of both samples are relatively low and remain approximately constant with increased phosphate concentration. When the initial phosphate concentrations are higher than 12mgl⁻¹, on the other hand, the adsorption capacities increase rapidly. This suggests that precipitation reactions may occur during the adsorption process at high initial phosphate concentrations (Li and Stanforth, 2000). Phosphate exists as HPO₄²⁻ during the adsorption process because
- the pH of this experimental system is 8. The Ca²⁺ concentration of artificial seawater with salinity 5 used in the experiment is 0.00143M and the solubility product constant K_{sp} of CaHPO₄ is 1 × 10⁻⁷ at a temperature of 25 °C. Therefore, when the initial phos-

phate concentration of the solution is 12 mgI^{-1} , the ion product of HPO_4^{2-} and Ca^{2+} can be calculated as follows:

 $K = [Ca^{2+}][HPO_4^{2-}] = 0.00143 \times 12 \times 0.001/31 = 5.5 \times 10^{-7} > K_{sp} = 1 \times 10^{-7}$

The result shows that the condition for precipitation of CaHPO₄ is satisfied, which means that CaHPO₄ can precipitate on the surfaces of sediments B and F and results in an increase in apparent P-adsorption capacity in these samples. This may also explain why the P-adsorption data of these samples are well described by the exponential equations. Note that the adsorption isotherms of samples A, Z4 and W4 do not show the same distinctive increase of P-adsorption capacity when the initial liquidphase phosphate concentration is higher than 12 mg I⁻¹ (Fig. 2a). The following factors help to explain these different behaviors. First, as shown in Table 1, the percentage of the particles smaller than 0.005 mm in samples B and F is only 5–6%, which is





much lower than that in samples A, Z4 and W4 (20–28 %). This means that particles in samples A, Z4 and W4 are likely to have higher specific surface areas than those in samples B and F. Second, from Table 2, the P binding energy constants (*b*) calculated using the Langmuir equation for samples A, Z4 and W4 (0.346–0.428) are much higher than those for samples B and F (0.064–0.181). The constant *b* represents the binding capacity of the sediment surface for phosphate. This means that when the ion product of HPO_4^{2-} and Ca^{2+} in the experimental system is saturated, the adsorption reaction continues to dominate in samples A, Z4 and W4 because of the high specific surface areas and strong P-binding capacities of these sediments (Özacar, 2003).

3.2 Buffering effects

5

The results of buffering adsorption experiments are shown in Fig. 3. At low initial P concentrations, the adsorption curve crosses the x-axis, where ΔP_S is 0. This indicates the point at which there is no net P exchange between sediment and the incubation solution. The concentration of P in the equilibrium solution at $\Delta P_S = 0$ is termed the "Zero Equilibrium Phosphate Concentration (EPC₀)" and represents the concentration at which the sediments typically display their maximum buffering capacities to changes in solution P (Bridgham et al., 2001). The EPC₀ value is traditionally considered as the soluble P concentration for natural systems in long-term equilibrium.

The slope of the adsorption curve at $\Delta P_S = 0$ is the linear adsorption coefficient (*K*), which indicates the sediment's buffering intensity at EPC₀ (and also the maximum buffering intensity for certain sediments). The P adsorbed to sediment at EPC₀ is called the equilibrium-adsorbed phosphate (Ps-EPC₀) and is a measure of the mass of reactive phosphate carried on per gram of sediment that is available for release to a solution devoid of phosphate. Ps-EPC₀ can be calculated by

²⁵ Ps - EPC₀ = $\mathcal{K} \cdot \text{EPC}_0$ (µg Pg⁻¹)

The values of EPC_0 , K and Ps- EPC_0 (Table 3) for four sediment samples used in this study were obtained from the buffering adsorption data. The EPC_0 values ranged from





- 0.014 to 0.061 mgl⁻¹. The maximum EPC₀ was found in sample A (0.061 mgl⁻¹), although the EPC₀ value for sample Z4 was similar (0.057 mgl⁻¹). The EPC₀ values for samples B and F were relatively low (0.014 and 0.029 mgl⁻¹, respectively). Similar patterns were observed for *K* values, which ranged between 0.177 and 0.562 lg⁻¹. The highest *K* value (0.562 lg⁻¹) was observed in sample Z4, which also had the highest Ps-EPC₀ value (32.164 µgPg⁻¹). Sediment A also had a relatively high Ps-EPC₀ value (19.185 µgPg⁻¹). One possible explanation for the higher *K* and Ps-EPC₀ values derived for samples A and Z4 compared to those of sediments B and F is the higher content of fine particles (< 0.005 mm) and organic matter in these sediments (fine particle fraction 27 % and 28 %, OM 0.66 % and 0.82 %, for A and Z4, respectively). Sediments with higher fine particle contents will tend to absorb more phosphorus and show bet-
- ter phosphorus buffering capacities due to their higher surface areas. Organic matter is very important for maintaining soil moisture (Krol and Bronstert, 2007) contributing to sediment phosphorus concentrations and for phosphorus cycling (de Assis et al.,
- ¹⁵ 2010). Ohno et al. (2005) also reported a linear relationship between dissolved organic carbon and water-soluble P concentrations in soil extracts, implying organic matter may increase soil P availability. It is, therefore, likely that high sediment organic matter contents will contribute to the exchange of P between the particulate and dissolved phases.
- Water quality monitoring conducted at the same time as sediment sampling showed that the concentrations of soluble reactive phosphate in seawater close to the Zhuyuan outfall areas ranged between 0.004 and 0.076 mgl⁻¹ (Table 1), coinciding with the value of EPC₀ (0.057 mgl⁻¹) determined by the buffering experiment using sample Z4 (collected from the same area). There was similar agreement between experimental and water quality data for the Bailonggang and Xinghuo outfalls, suggesting that the con-
- ²⁵ water quality data for the Bailonggang and Xinghuo outfalls, suggesting that the concentration of phosphate in seawater is related to phosphate sorption to sediment.

	BGD 9, 17519–17538, 2012								
	Phosphorus sorption and buffering mechanisms in sediments M. Li et al.								
D 2		—							
5	Title Page	Title Page							
-	Abstract Introduction	n							
	Conclusions References	S							
))))	Tables Figures								
	I∢ ►I								
5	۰ (۲								
-	Back Close								
	Full Screen / Esc								
2	Printer-friendly Version								
Interactive Discussion									



3.3 Influence of temperature on sediment buffer capacity to phosphorus

The results of buffering experiments at different temperatures are shown in Fig. 4a for sample Z4. Values of EPC_0 and K for this sediment at different temperatures are shown in Fig. 4b, c, respectively.

- As the temperature (*T*) increases, values of EPC_0 and *K* also increase (due to a shift in the sorption isotherm). However, the slope of the regression line for EPC_0 is not significant. Increasing the temperature from 12 °C to 32 °C resulted in an increase in EPC_0 for sediment Z4 from 0.04 to 0.06 mg l⁻¹, which was similar to the observed concentration of soluble reactive phosphate in seawater near the Zhuyuan outfall. The value of
- 10 K also increases with temperature, suggesting that buffering capability of sediment for phosphate is enhanced as the temperature increases. The rate of both adsorption and desorption will increase with temperature. Thus, as well as shifting the equilibrium ratio between sorbed and dissolved phase P concentrations, the response time to changes in P availability will reduce at higher temperatures.
- ¹⁵ Most red tide algae tend to grow quickly at high temperatures (Dae-II et al., 2004). For example, Yamaguchi et al. (2010) reported that *Chattonella ovata* could grow rapidly at temperatures over 20°C and that maximal growth rates were between 25 and 30°C. Our finding of increased P buffering capacity at high temperatures might be an explanation for this phenomenon.

20 4 Conclusions

- P sorption in sediments with a high fraction of fine particles was best described using the Langmuir equation. Sorption of P in sediments with a low fraction of fine particles was best described using an exponential sorption equation.
- The precipitation reaction of CaHPO₄ from Ca²⁺ and HPO₄²⁻ occurs in parallel with the sorption process in some samples when the dissolved phosphate concentration is higher than 12 mgI^{-1} . The increase in apparent P-sorption capacities 17528





25

at high initial P-concentrations in these samples may be caused by this precipitation reaction and may explain why the sorption data were best described using an exponential model.

- The EPC₀ values derived from the buffer experiments ranged between 0.014 and 0.061 mg l⁻¹. These concentrations are consistent with measured phosphate concentrations in water samples collected from the Yangtze Estuary and Hangzhou Bay. They reveal a potential for predicting the dissolved P concentration using a sediment buffer experiment.
- The K values vary from 0.177 to $0.562 \, \text{Ig}^{-1}$ and increase with increasing fine particle and organic matter content. It is suggested that sediments which are rich in fine particles and organic matter have stronger buffering capacities for phosphate.
- Values of sediment EPC₀ and *K* all increase with increasing temperature. This
 may provide one explanation for why "red tides" tend to occur in warm weather,
 due to higher P buffering capacity at high temperatures.
- Acknowledgements. This work was supported by "the Fundamental Research Funds for the Central Universities" (No. YX 2010-28), the National Natural Science Foundation of China (No. 50709001) and the Natural Science Foundation of Beijing, China (No. 8083029). Li was also supported by the State Scholarship Fund of China (No. 2009651510) to pursue her research at Cranfield University as a visiting scholar from August 2010 to August 2011.

20 References

5

10

Bao, S. D. and Jiang, R. F.: Measurement of organic matter content in soil, in: Analysis of Agri-Chemistry in Soil, 3rd edn., Chinese Agricultural Science and Technology Press, Beijing, China, 1999.

Beuselinck, L., Govers, G., Poesen, J., Degraer, G., and Froyen, L.: Grain-size analysis by

²⁵ laser diffractometry: comparison with the sieve-pipette method, CATENA, 32, 193–208, doi:10.1016/s0341-8162(98)00051-4, 1998.





Borggaard, O. K., Szilas, C., Gimsing, A. L., and Rasmussen, L. H.: Estimation of soil phosphate adsorption capacity by means of a pedotransfer function, Geoderma, 118, 55-61, doi:10.1016/s0016-7061(03)00183-6, 2004.

Bridgham, S. D., Johnston, C. A., Schubauer-Berigan, J. P., and Weishampel, P.: Phosphorus sorption dynamics in soils and coupling with surface and pore water in riverine wetlands, Soil 5 Sci. Soc. Am. J., 65, 577-588, 2001.

Conley, D. J., Smith, W. M., Cornwell, J. C., and Fisher, T. R.: Transformation of particle-bound phosphorus at the land-sea interface, Estuar. Coast. Shelf Sci., 40, 161–176, 1995.

Dae-II, K., Yukihiko, M., Sou, N., Mineo, Y., Yang-Ho, Y., Yuji, O., Nobuyoshi, I., and Tsuneo, H.:

Effects of temperature, salinity and irradiance on the growth of the harmful red tide dinoflagel-10 late cochlodinium polykrikoides margalef (Dinophyceae), J. Plankton Res., 26, 61-66, 2004. de Assis, C. P., de Oliveira, T. S., da Nóbrega Dantas, J. D. A., and de Sá Mendonça, E.: Organic matter and phosphorus fractions in irrigated agroecosystems in a semi-arid region of Northeastern Brazil, Agr. Ecosyst. Environ., 138, 74-82, doi:10.1016/j.agee.2010.04.002, 2010.

15

20

30

- Froelich, P. N.: Kinetic control of dissolved phosphate in natural rivers and estuaries: a primer on the phosphate buffer mechanism, Limnol. Oceanogr., 33, 649–668, 1988.
- Kara, D., Özsavasçi, C., and Alkan, M.: Investigation of suitable digestion methods for the determination of total phosphorus in soils, Talanta, 44, 2027-2032, doi:10.1016/s0039-9140(97)00014-3, 1997.
- Kawasaki, N., Ogata, F., and Tominaga, H.: Selective adsorption behavior of phosphate onto aluminum hydroxide gel, J. Hazard. Mater., 181, 574-579, doi:10.1016/j.jhazmat.2010.05.051, 2010.

Krol, M. S. and Bronstert, A.: Regional integrated modelling of climate change impacts on natural resources and resource usage in semi-arid Northeast Brazil, Environ. Modell. Softw.,

- 25 22, 259-268, 2007.
 - Laly, J. P. and Sgluo, G.: Chemistry Oceanography, 2nd edn., Ocean Press, Beijing, China, 1982.
 - Li, L. and Stanforth, R.: Distinguishing adsorption and surface precipitation of phosphate on goethite (α -FeOOH), J. Colloid Interf. Sci., 230, 12–21, 2000.
 - Li, M., Wei, H. P., Wang, G. Q., and Ni, J. R.: Study on adsorption behavior of phosphate by sediments from Changjiang estuary and Hangzhou bay, Acta Oceanol. Sin., 26, 132-136, 2004.





Ohno, T., Griffin, T. S., Liebman, M., and Porter, G. A.: Chemical characterization of soil phosphorus and organic matter in different cropping systems in Maine, USA, Agr. Ecosyst. Environ., 105, 625–634, doi:10.1016/j.agee.2004.08.001, 2005.

Özacar, M.: Adsorption of phosphate from aqueous solution onto alunite, Chemosphere, 51, 321–327, doi:10.1016/s0045-6535(02)00847-0, 2003.

Pratt, C. and Shilton, A.: Suitability of adsorption isotherms for predicting the retention capacity of active slag filters removing phosphorus from wastewater, Water Sci. Technol., 59, 1673–1678, 2009.

5

Price, R. M., Savabi, M. R., Jolicoeur, J. L., and Roy, S.: Adsorption and desorption of phos-

¹⁰ phate on limestone in experiments simulating seawater intrusion, Appl. Geochem., 25, 1085– 1091, doi:10.1016/j.apgeochem.2010.04.013, 2010.

Richardson, C. J.: Mechanisms controlling phosphorus retention capacity in freshwater wetlands, Science, 228, 1424–1427, 1985.

Schippers, P., van de Weerd, H., de Klein, J., de Jong, B., and Scheffer, M.: Impacts of agricul-

- tural phosphorus use in catchments on shallow lake water quality: about buffers, time delays and equilibria, Sci. Total Environ., 369, 280–294, doi:10.1016/j.scitotenv.2006.04.028, 2006.
 Wauchope, R. D. and McDowell, L. L.: Adsorption of phosphate, arsenate, methanearsonate, and cacodylate by lake and stream sediments: comparisons with soils, J. Environ. Qual., 13, 499–504, 1984.
- Yamaguchi, H., Mizushima, K., Sakamoto, S., and Yamaguchi, M.: Effects of temperature, salinity and irradiance on growth of the novel red tide flagellate *Chattonella ovata* (Raphi-dophyceae), Harmful Algae, 9, 398–401, doi:10.1016/j.hal.2010.02.001, 2010.





Site DIP ^a (mgl ⁻		I ⁻¹) OM ^b (%)		Particle size composition (% total volume)			
		Range	Mean		0.25–0.074 mm	0.074–0.005 mm	< 0.005 mm
	А			0.66	0	73	27
Xinghuo	В	0.040-0.077	0.048	0.02	0	95	5
	F			0.25	58	36	6
Zhuyuan	Z4	0.004–0.076	0.042	0.82	0	72	28
Bailonggang	W4	0.028–0.116	0.049	0.77	0	80	20

 Table 1. Description of different sediment and water samples.

^aDIP: dissolved inorganic phosphate of the water samples.

^bOM: organic matter content of the sediment samples.



Table 2. Langmuir, Freundlich, Temkin isotherm equations and constants for adsorption of phosphate by different sediments.

	Langmuir equation				
Sample	$\frac{c}{\Gamma} = \frac{c}{\Gamma_{m}} + \frac{1}{\Gamma_{m}b}$	Γ_{∞} (µg P g ⁻¹)	<i>b</i> (Img ⁻¹)	r	S
А	$\frac{c}{\Gamma} = 0.0042c + 0.0098$	238.09	0.428	0.996	12.632
В	$\frac{c}{\Gamma} = 0.0062c + 0.0969$	161.29	0.064	0.463	63.425
F	$\frac{c}{\Gamma} = 0.0101c + 0.0557$	99.01	0.181	0.711	35.791
Z4	$\frac{c}{\Gamma} = 0.0027c + 0.0066$	370.37	0.409	0.992	12.885
W4	$\frac{c}{\Gamma} = 0.0028c + 0.0081$	357.14	0.346	0.997	13.461
	Freundlich equation				
	$\lg \Gamma = \frac{1}{n} \lg c + \lg k$	K	п	r	S
А	$\lg \Gamma = 0.3896 \lg c + 1.8721$	74.49	2.567	0.995	11.075
В	$\lg \Gamma = 0.5067 \lg c + 1.2752$	18.845	1.974	0.801	64.842
F	$\lg \Gamma = 0.2282 \lg c + 1.5456$	35.124	4.382	0.655	35.171
Z4	lgΓ = 0.7297lg <i>c</i> + 1.78	60.256	1.37	0.968	71.921
W4	$\lg \Gamma = 0.4974 \lg c + 1.9392$	86.936	2.01	0.974	30.954
	Temkin equation				
	$\Gamma = \frac{\Gamma_{\infty}RT}{\beta Q_0} \cdot \ln(C_0 c)$	r	S		
А	$\Gamma = 43.38 \ln c + 84.588$	0.987	11.433		
В	Γ = 32.103 ln <i>c</i> + 10.449	0.585	64.829		
F	Γ = 15.316 ln <i>c</i> + 33.008	0.556	35.094		
Z4	Γ = 56.739 ln <i>c</i> + 147.12	0.988	22.037		
W4	Γ = 68.553 ln <i>c</i> + 113.41	0.994	11.648		



CC D

 Γ : equilibrium P adsorption amount; *c*: equilibrium P concentration in liquid phase; Γ_{∞} : maximum adsorption amount; *b*: adsorption equilibrium constant; *r*: correlation coefficient; *S*: standard deviation; *K*: empirical constant; *n*: empirical constant.

Sediment samples	$EPC_0 \ (mgl^{-1})$	K (lg ⁻¹)	$Ps-EPC_0 (\mu g P g^{-1})$
A	0.061	0.314	19.185
В	0.014	0.177	2.490
F	0.029	0.220	6.304
Z4	0.057	0.562	32.164

 Table 3. Comparison of the buffering capacity of different sediments to phosphate.

EPC₀: zero equilibrium phosphate concentration.

K: linear adsorption coefficient.

Ps-EPC₀: equilibrium-adsorbed phosphate.

























c ()

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper







CC ① BY