

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# Occurrence, sources and transport pathways of natural and anthropogenic hydrocarbons in deep-sea sediments of the Eastern Mediterranean Sea

**C. Parinos<sup>1</sup>, A. Gogou<sup>1</sup>, I. Bouloubassi<sup>2</sup>, R. Pedrosa-Pàmies<sup>3</sup>, I. Hatzianestis<sup>1</sup>,  
A. Sànchez-Vidal<sup>3</sup>, G. Rousakis<sup>1</sup>, D. Velaoras<sup>1</sup>, G. Krokos<sup>1</sup>, and V. Lykousis<sup>1</sup>**

<sup>1</sup>Institute of Oceanography, Hellenic Centre for Marine Research (HCMR), 46.7 km Athens-Sounion Av., 19013 Anavyssos, Attiki, Greece

<sup>2</sup>Laboratoire d'Océanographie et du Climat: Expérimentation et Approches Numériques (LOCEAN/IPSL), Université Pierre et Marie Curie-CNRS-IRD-MNHN, UMR 7159, 4 Place Jussieu, 75252 Paris CEDEX 05, France

<sup>3</sup>GRC Geociències Marines, Departament d'Estratigrafia, Paleontologia i Geociències Marines, Facultat de Geologia, Universitat de Barcelona, Barcelona, Spain

Received: 27 November 2012 – Accepted: 29 November 2012

– Published: 13 December 2012

Correspondence to: C. Parinos (ksparinos@hcmr.gr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Surface sediments collected from deep basins (22 stations, 1018–4087 m depth) of the Eastern Mediterranean Sea (EMS) were analyzed for aliphatic, triterpenoid and polycyclic aromatic hydrocarbons (PAHs) as tracers of natural and anthropogenic inputs. Concentrations of total aliphatic hydrocarbons (TAHC), *n*-alkanes (NA) and the Unresolved Complex Mixture (UCM) of aliphatic hydrocarbons ranged from 1.34 to 49.2  $\mu\text{g g}^{-1}$ , 145 to 4810  $\text{ng g}^{-1}$  and 0.73 to 36.7  $\mu\text{g g}^{-1}$ , respectively, while total PAHs (TPAH<sub>25</sub>) concentrations ranged from 11.6 to 223  $\text{ng g}^{-1}$ . Molecular profiles of aliphatic hydrocarbons and PAHs reflect the contribution of both natural (epicuticular plant waxes) and anthropogenic (degraded petroleum products, unburned fossil fuels and combustion of petroleum, grass, wood and coal) compounds in deep EMS sediments, with hydrocarbon mixtures displaying significant regional variability. Hydrocarbon concentrations correlated significantly with the Total Organic Carbon (TOC) content of sediments, indicating that organic carbon exerts an important control on their transport and fate in the study area, while strong sub-basin and mesoscale variability of water masses also impact their regional characteristics. Major findings of this study support that deep basins/canyons of the EMS could act as traps of both natural and anthropogenic hydrocarbons.

## 1 Introduction

Hydrocarbons are abundant components of both natural and anthropogenic organic material introduced in coastal and open sites of the world's oceans. They enter the marine environment through both atmospheric (dry/wet deposition, gas exchange across the air–water interface) and aquatic pathways (continental run-offs, direct discharges, off-shelf export), or are produced autochthonously from marine organisms. Due to their hydrophobic nature hydrocarbons tend to associate with particles resulting in their transport and final accumulation in sediments (Bouloubassi et al., 2006, 2012;

**BGD**

9, 17999–18038, 2012

### Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Castro-Jiménez et al., 2012; Dachs et al., 2002; Lipiatou et al., 1997; Parinos et al., 2012; Prahla and Carpenter, 1979).

Polycyclic aromatic hydrocarbons (PAHs) are included in lists of priority chemical pollutants by international environmental agencies (EEA, EPA), since certain homologues are highly carcinogenic and mutagenic, exhibiting tendency to bioaccumulate in aquatic organisms (Samanta et al., 2002 and references therein). Similarly, the Unresolved Complex Mixture (UCM), consisting of branched alicyclic hydrocarbons (Gough and Rowland, 1990; Killops and Al-Juboori, 1990), which is a commonly observed contaminant of marine sediments in which it can persist for decades (Reddy et al., 2002), has been proven as toxic to sediment-dwelling organisms (Scarlett et al., 2007).

Natural sources of hydrocarbons in the marine environment include terrestrial plant waxes, diagenetic transformation of biogenic precursors, marine phytoplankton and bacteria (Gogou et al., 2000). Major anthropogenic sources are industrial outfalls, petroleum processing/seepage and accidental oil spills, since hydrocarbons are main constituents of petroleum, pyrolysis/combustion of organic material (e.g. biomass burning, incomplete combustion of fossil fuels) and several industrial processes (Countway et al., 2003; Dachs et al., 1997; Laflamme and Hites, 1978; Leister and Baker, 1994; Neff, 1979; Simoneit, 1984; Tolosa et al., 1996; Yunker et al., 2002).

The Eastern Mediterranean Sea (EMS), and especially its coastal area, is a region under intense anthropogenic pressure from increasing industrial and urban activities, resulting in pollutant discharges (EEA, 2006). Being an important geographical region for merchant shipping and oil transportation, the EMS receives substantial amounts of petroleum discharges, mainly along shipping routes (UNEP, 2010). Atmospheric deposition is also an important route for the introduction of hydrocarbons across the EMS (Castro-Jiménez et al., 2012; Gogou et al., 1996; Grimalt et al., 1988; Tsapakis 2003; Tsapakis and Stefanou, 2005), which also receives significant Saharan dust inputs (Jickells et al., 2005), during dust storm events, that contain both natural and anthropogenic hydrocarbons as recently reported (Ladji et al., 2010).

**BGD**

9, 17999–18038, 2012

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Aliphatic hydrocarbons and PAH burden of surficial sediments in the EMS has been studied over the past twenty five years in coastal or shallow basin sites, namely the Aegean Sea, Cretan Sea (Southern Aegean sea), northeastern Mediterranean coast, southern Italy, Egypt coast and Adriatic Sea (Aboul-Kassim and Simoneit, 1995; Alebic-Juretic, 2011; Annicchiarico et al., 2011; Barakat et al., 2011; Botsou and Hatzianestis, 2012; Cardellicchio et al., 2007; Gogou et al., 2000; Gonul and Kucuksezgin, 2012; Guzzella and De Paolis, 1994; Hatzianestis et al., 2001; Kucuksezgin et al., 2012; Marcomini et al., 1986; Nemr et al., 2007; Sklivagou et al., 2008; Yilmaz et al., 1998; Youssif, 2002). However, there is a lack of data regarding hydrocarbons occurrence in deep basins of the EMS. Unlike coastal settings, deep-sea sediments reflect averaged deposition patterns and concentrations of natural and anthropogenic material of wider timeframes, due to the low sediment accumulation rates. In this study we focus on aliphatic, triterpenoid and polycyclic aromatic compounds as tracers of natural and anthropogenic inputs, reporting the first data set being addressed for surface sediments in deep EMS basins (22 stations, 1018–4087 m depth). Our main goal was to infer conclusions regarding their occurrence, major sources and transport pathways and to examine the role of deep EMS basins as repositories of hydrocarbons.

## 2 Oceanographic setting

The EMS is a semi-enclosed basin that connects with the Western Mediterranean Sea through the straits of Sicily. It includes four major sub-basins, the Ionian and Levantine basins, the Adriatic Sea and the Aegean Sea (Malanotte – Rizzoli et al., 1988; Robinson et al., 1992, 2001). It has a complex circulation pattern with water masses distribution being influenced by both large-scale and mesoscale variability (Malanotte-Rizzoli et al., 1997; Millot, 2005). As a concentration basin, it is characterized by an anticyclonic circulation that transforms surface Atlantic Waters (AW), entering through the strait of Sicily at the upper 100–150 m of the water column, into Levantine Intermediate Waters (LIW) in the eastern part of the region (Rhodes gyre). LIW leave the basin as

# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a westward current through the Sicily strait at depths between 200 m and 500 m. The Aegean Sea occasionally contributes saline intermediate water masses to the EMS (Cretan Intermediate Water – CIW) with characteristics very close to those of LIW. Strong mesoscale variability, creating permanent and transient eddies and gyres, enhances exchanges between the continental shelf waters and slope waters (Danovaro et al., 2010).

Deep water layers of the EMS originate mainly in the Adriatic Sea and are exported by means of bottom-arrested currents toward the abyssal layers of the Ionian basin, flowing in an eastward path all the way towards the eastern Levantine basin. The Aegean Sea can also contribute in deep water formation and may even become more effective than the Adriatic as a deep water source for the EMS, as in the case of the Eastern Mediterranean Transient (EMT) during the early '90s (Civitaresse et al., 2010; Roether et al., 1996; Theoharis et al., 1999).

Mean sedimentation rates in the EMS basins are low and generally do not exceed 5 cm kyr<sup>-1</sup> mainly due to the oligotrophic character of the area (Garcia-Orellana et al., 2009).

## 3 Materials and methods

### 3.1 Sampling

Surface sediments were collected with a multicorer at twenty two stations in deep basins (1018–4087 m depth) of the Ionian Sea, Cretan Sea (southern Aegean Sea) and Levantine Sea, during four oceanographic cruises conducted between January 2007 and June 2012 (Fig. 1; Table 1). January 2007 samples were collected during the M71 Leg. 3 cruise of the R/V *Meteor* (University of Hamburg, Germany). May 2010, 2011 and June 2012 samples were collected during cruises of the R/V *Aegaeo* (HCMR, Greece). The undisturbed top centimeter (1 cm) of each core was recovered, wrapped in aluminum foil and stored at –20 °C till analysis.

## 3.2 Methods

Freeze-dried sediments were spiked with a mixture of internal standards ( $[^2\text{H}_{50}]$ tetracosane,  $[^2\text{H}_{10}]$ phenanthrene,  $[^2\text{H}_{10}]$ pyrene,  $[^2\text{H}_{12}]$ chrysene,  $[^2\text{H}_{12}]$ perylene and  $[^2\text{H}_{12}]$ benzo[ghi]perylene) and solvent extracted three times by sonification with a dichloromethane : methanol mixture (4 : 1 v/v). Combined extracts were fractionated on silica column, applying a modified protocol after Gogou et al. (1998). Aliphatic hydrocarbons were eluted with 6 mL *n*-hexane and PAHs with 10 mL *n*-hexane/toluene (9 : 1, v/v). Both fractions were concentrated by vacuum rotary evaporation, transferred to a 1.5 mL amber vial and finally evaporated under a gentle nitrogen stream.

Instrumental analysis was carried out by Gas Chromatography–Mass Spectrometry (GC–MS) on an Agilent 7890 GC, equipped with an HP-5MS capillary column (30 m × 0.25 mm i.d. × 0.25 μm phase film), coupled to an Agilent 5975C MSD. For the analysis of aliphatic hydrocarbons the MSD operated in full scan mode and the GC oven temperature was initially held at 60 °C for 2 min, brought to 80 °C at a rate of 25 °C min<sup>-1</sup>, then to 300 °C at a rate of 5 °C min<sup>-1</sup> and finally held at 300 °C for 35 min. PAHs were analyzed using a selected ion monitoring (SIM) acquisition program. The oven temperature program was the same as in the case of aliphatic hydrocarbons but with a 300 °C final isothermal of 6 min. Helium was used as carrier gas at a flow of 1.1 mL min<sup>-1</sup>.

Standard solutions of the targeted compounds, purchased from Dr. Ehrenstorfer GmbH, were spiked with the internal standards and run on each injection set in order to derive relative response factors (RRFs) of the analytes. The precision of the analytical method used for PAHs was evaluated by analyzing the National Institute of Standards (NIST) standard reference sediment SRM 1941a (Organics in Marine Sediment). The determined values ranged between 93 and 106 % of the certified values while in terms of repeatability the relative standard deviation was below 5 %. Procedural blanks processed were found to be free of contamination.

**BGD**

9, 17999–18038, 2012

### Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Spatial distributions of concentrations and characteristic parameters of aliphatic and polycyclic aromatic hydrocarbons were visualized using Ocean Data View – ODV (Schlitzer, 2011).

## 4 Results

### 4.1 Molecular profile of hydrocarbons

$C_{12}$  to  $C_{42}$  *n*-alkanes (NA) were the main constituents of the resolved aliphatic compounds, representing on average 56 % of their total sum. Their molecular profile (Fig. 2a) was dominated by long chain homologues ( $C_n \geq 24$ ) maximizing at  $C_{31}$  with an elevated odd/even carbon preference ( $CPI_{24-35} > 1.9$ , 4.3 on average; Table 2). NA with  $C_n > 35$  constituted a large fraction ( $> 30\%$ ) at some stations, while NA with  $C_n \leq 23$  were mostly minor constituents and did not exhibit any odd/even carbon preference ( $CPI_{14-23} \sim 1$ ). A UCM, present as a unimodal hump centered at  $C_{30}$ , was the major component of total aliphatic hydrocarbons (69 % on average).

25 PAHs comprising parent (unsubstituted) compounds with 2–6 aromatic rings and alkyl-substituted homologues were determined. A typical molecular profile of PAHs is presented in Fig. 2b. Phenanthrene and its methyl, di- and tri- methyl homologues dominated the low-MW PAH ( $\leq 3$  aromatic rings). Their sum, referred to hereafter as  $\sum Phe$ , accounted for  $23 \pm 6\%$  of total PAHs. The high MW parent compounds ( $\geq 4$  aromatic rings), which are of pyrolytic origin (Neff, 1979), were dominated by benzo[fluoranthenes, indeno[1,2,3-*cd*]pyrene and to a lesser degree chrysene. Their sum (referred to hereafter as  $\sum COMB$ ), excluding perylene which may have natural sources (Venkatesan, 1988), represented  $55 \pm 8\%$  of total PAHs. Retene (methyl-1-isopropyl-7 phenanthrene) was the major naturally-derived PAH determined in the study area (Ramdahl, 1983).

**BGD**

9, 17999–18038, 2012

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 4.2 Concentrations and spatial distribution of hydrocarbons

Concentrations of total aliphatic hydrocarbons (TAHC: sum of total resolved compounds and UCM), NA and UCM are presented in Table 2. TAHC and UCM concentrations ranged from 1.34 to 49.2  $\mu\text{g g}^{-1}$  and 0.73 to 36.7  $\mu\text{g g}^{-1}$ , respectively, while NA concentrations ranged between 145 and 4810  $\text{ng g}^{-1}$ . Their spatial distribution shows a generally increasing trend from Cretan Sea and Levantine Sea stations towards the deep stations of Ionian Sea (Fig. 3a–c). Maximum concentrations are found at station H03 while minimum at station H12, both located in the Ionian Sea.

Table 3 summarizes concentrations of PAHs in surface sediments of the study area. TPAH<sub>25</sub> refers to the total sum of compounds monitored while  $\sum\text{PAH}_{13}$  refers to the sum of 13 parent compounds mainly considered in marine pollution studies, i.e. Fl, Phe, Anth, Flth, Pyr, BaA, Chry, BFlths, BeP, BaP, IndP, BgP and DBA (for abbreviations of PAH compounds see Fig. 2b). TPAH<sub>25</sub> and  $\sum\text{PAH}_{13}$  concentrations ranged from 11.6 to 223  $\text{ng g}^{-1}$  and from 5.90 to 130  $\text{ng g}^{-1}$ , respectively. The concentrations of  $\sum\text{COMB}$  ranged from 5.08 to 118  $\text{ng g}^{-1}$  while  $\sum\text{Phe}$  ranged between 2.74 and 38.3  $\text{ng g}^{-1}$ . The spatial distribution of TPAH<sub>25</sub>,  $\sum\text{Phe}$  and  $\sum\text{COMB}$  concentrations is presented in Fig. 3d–f. Highest TPAH<sub>25</sub>,  $\sum\text{COMB}$  and  $\sum\text{Phe}$  concentrations were recorded at station H05, while lowest ones were found at station H12, both located in the Ionian Sea. The spatial distribution of TPAH<sub>25</sub> (Fig. 3d) is similar to the one observed for TAHC (Fig. 3a).

## 5 Discussion

### 5.1 Hydrocarbons levels

Table 4 summarizes concentrations of aliphatic and polycyclic aromatic hydrocarbons in the study area and data reported in deep-sea sediments of the Mediterranean Sea and other open sites of the world ocean. TAHC concentrations in the study area are

**BGD**

9, 17999–18038, 2012

### Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





slightly higher than those reported in the Cretan Sea (Gogou et al., 2000) and comparable to those reported in the Black Sea (Wakeham et al., 1996). As for UCM and NA, concentrations reported in this study are higher than those reported for the open Western Mediterranean Sea (Tolosa et al., 1996) and Cretan Sea (Gogou et al., 2000), but lower than those reported in the Arctic Ocean (Yunker et al., 2011). Total PAH concentrations are comparable to those reported for deep-sea settings such as Cretan Sea (Gogou et al., 2000), central Pacific Ocean (Ohkouchi et al., 1999) and South China Sea (Liu et al., 2012; Yang et al., 2000), while are considerably lower from those recorded in other deep oceanic sites: the Black Sea (Wakeham et al., 1996), the Western Mediterranean Sea (Lipiatou and Saliot, 1991; Tolosa et al., 1996), the Atlantic Ocean (OSPAR QRS, 2000) and the Arctic Ocean (Yunker et al., 2011).

## 5.2 Sources of natural and anthropogenic hydrocarbons

### 5.2.1 Natural sources

The dominance of  $C_{25}$  to  $C_{35}$  *n*-alkanes (Fig. 2a) with an elevated odd/even carbon preference shows the importance of natural terrestrial inputs, derived from epicuticular plant waxes (Eglinton and Hamilton, 1967), throughout the study area. The sum of major terrestrial *n*-alkanes ( $C_{27}$ ,  $C_{29}$ ,  $C_{31}$  and  $C_{33}$ ), referred to hereafter as Ter.NA, averaged 50 % of total NA (Table 2).

The low abundance of  $C_{15}$ ,  $C_{17}$  and  $C_{19}$  *n*-alkanes (< 5 % of NA) reflects a minor contribution from marine sources. This is consistent with both the ultra-oligotrophic character of the Eastern Mediterranean Sea (Krom et al., 2003) and the preferential preservation of terrestrial over planktonic hydrocarbons in the marine environment (Prah and Carpenter, 1984). Also, bacterial sources, evidenced by the presence of hopenes (hop-17(21)-ene, hop-13(18)-ene and 17 $\beta$ (H)-hop-22(29)-ene; Fig. 4) are also of minor importance.

Among PAHs, the presence of retene reflects inputs from terrestrial plants, mainly conifers (Wakeham et al., 1980) or inputs related to pinewood combustion (Ramdahl,

**BGD**

9, 17999–18038, 2012

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1983). Retene represented less than 4 % of TPAH<sub>25</sub> in all stations, except station H05 in which it accounted for ~ 18 % of total PAHs (see below Sect. 5.3).

### 5.2.2 Anthropogenic sources

The elevated relative abundance of UCM compared to the total resolved aliphatic hydrocarbons (TRes) (UCM/TRes = 1.6–5.7; Table 2) is a positive indication of contribution from degraded petroleum products (Brassell and Eglinton, 1980; Farrington and Quinn, 1973). Consistently, a series of C<sub>29</sub>–C<sub>35</sub> 17 $\alpha$ (H),21 $\beta$ (H)-hopanes were identified in all samples (Fig. 4). The extended, C<sub>31</sub>–C<sub>35</sub>, homologues were present as pairs of the C<sub>22</sub> diastereoisomers (22S and 22R), with a 22S/22S + 22R ratio close to 0.6, typical for petrogenic hydrocarbons (Mackenzie, 1984). At some stations, low-MW *n*-alkanes (C<sub>n</sub> ≤ 23) without odd/even carbon preference likely derive from fossil inputs (light diesel), although they may also represent reworked algal material (Salot, 1981; Wang et al., 1999), while large amounts of *n*-alkanes with C<sub>n</sub> > 35 point to enhanced contribution of heavy fuel oil residuals (Brooks et al., 1954; Hsieh et al., 2000).

Among PAHs, the dominance of alkylated phenanthrenes in the low-MW PAH and the high abundance of parent PAH with ≥ 4 aromatic rings (Fig. 2b), reflect an admixture of unburned fossil sources (petroleum) and combustion/pyrolysis of fossil fuels (LaFlamme and Hites, 1978; Sporstol et al., 1983; Wakeham et al., 1980). Source specific diagnostic ratios have been used in order to further assess PAH sources (Yunker et al., 2002). The Flth/(Flth + Pyr) and IndP/(IndP + BgP) ratios exhibit values > 0.50, averaging 0.60 ± 0.04 and 0.56 ± 0.02 respectively, indicating grass, wood and coal combustion as primary sources for these PAHs. The BaA/(BaA + Chry) ratio ranged between 0.15–0.35 (average 0.26 ± 0.04), indicating an admixture of pyrolytic and fossil contributions. As for low-MW PAH compounds, the C<sub>0</sub>/(C<sub>0</sub> + C<sub>1</sub>) Phe ratio ranged between 0.33–0.70 (average 0.51 ± 0.10) indicating a varying contribution of pyrolytic (ratio > 0.5) and petroleum (ratio < 0.5) phenanthrenes sources in the study area.

The ratio of total phenanthrenes ( $\Sigma$ Phe), mostly derived from petrogenic inputs, and  $\Sigma$ COMB, representing pyrolytic PAH, ( $\Sigma$ Phe/ $\Sigma$ COMB, 0.22 to 0.87, average

18008

BGD

9, 17999–18038, 2012

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



0.45 ± 0.19; Table 3), reflects dominant pyrolytic sources for PAH mixtures in the study area (Bouloubassi and Salot, 1993; Bouloubassi et al., 2001; Gogou et al., 2000; Parinos et al., 2012).

### 5.3 Regional characteristics of hydrocarbons

Cluster analysis (joining clustering method) was used to group stations with similar *n*-alkanes or PAH profiles. Ward's method was used for amalgamation of clusters while distances between objects were measured as Euclidean distances.

Figure 5 presents the hierarchical diagram obtained from the cluster analysis of stations with regard to their *n*-alkanes profile, using the relative homologues concentrations for the total sum of NA as starting data, clustered station sites and characteristic GC chromatographs for each returning cluster. Stations are clustered into three main groups. Cluster I stations (RED2.1, RED4, RED5, H04 and H07) are characterized by high abundances of Ter.NA along with low-MW homologues ( $C_n \leq 23$ ). Cluster II stations (RED8, RED1.1, IER01, H12, RED15.1, RED9, H01, RED13 and Rho02) are characterized by high abundances of Ter.NA and low concentrations of both low and higher ( $C_n > 35$ ) molecular weight compounds. Stations in cluster III (RED3, RED3.1, RED7, H02, Her01, Her03, H05 and H03) are characterized by high abundances of Ter.NA along with significant presence of long chain homologues ( $C_n > 35$ ).

Stations were also grouped using cluster analysis with regard to PAH profile characteristics (relative compounds concentrations for TPAH<sub>25</sub>). Figure 6 presents the hierarchical diagram obtained from cluster analysis, clustered station sites and molecular profile for each returning cluster. Stations are also clustered into three main groups. Cluster I stations (RED2.1, RED3.1, RED3, RED4, RED5, RED7, H03, H07, H04, H01, IER01 and H12) are characterized by the elevated abundance of all alkylated homologues monitored within the phenanthrene, dibenzothiophene, pyrene and chrysene series, indicating enhanced contribution of petrogenic PAHs. Cluster II stations (RED8, RED1.1, RED13, RED9, RED15.1, H02, Her03, Her01 and Rho02) on the other hand displays maximum concentrations for parent phenanthrene, pyrene and chrysene and

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



less than 0.1 % of TPAH<sub>25</sub> for dibenzothiophene and its alkylated homologues, together with maximum values for high-MW parent compounds, indicating dominant pyrolytic PAH contribution. Cluster III corresponding only to station H05 is characterized by the highest concentration of retene, while the distribution of other compounds, e.g. low abundance of alkylated compounds and  $\sum\text{COMB}$  being > 50 % of TPAH<sub>25</sub>, point to dominant pyrolytic sources.

### 5.3.1 Ionian Sea

The  $\sum\text{Phe}/\sum\text{COMB}$  ratio shows an increasing trend from Aegean Sea to Sicily straits (stations H02, H03 and H04; Fig. 7a) in agreement with decreasing trend of  $\text{C}_0/(\text{C}_0 + \text{C}_1)$  Phe ratio values for the same stations (Fig. 7b), indicating an increase in the relative contribution of fossil fuels from east to west. Consistently, stations H02 and H03 display relatively elevated contribution of heavy oil residuals as inferred by  $\text{C}_n > 35$  *n*-alkanes (Fig. 5c), station H03 presents maximum UCM concentrations (Fig. 3b; Table 2) and station H04 is characterized by  $\text{C}_n \leq 23$  *n*-alkanes (Fig. 5c). The Ionian sea stations H02, H03 and H04 are located close to the main shipping route from Greece to Italy, displaying high oil spill density, with maximum in the area around station H04 (Ferraro et al., 2009). Additionally, atmospheric inputs should also be considered as a major source of fossil compounds at stations located in the open Ionian Sea, since the abundance of long chain *n*-alkanes ( $\text{C}_n > 35$ ) has been reported in rural aerosol molecular profiles of the EMS (Gogou et al., 1996), while Castro-Jiménez et al. (2012) also reported high diffusive deposition fluxes for Phe, DBT,  $\text{C}_1$ -Phe and  $\text{C}_1$ -DBT in the area.

Station H07, off the Gulf of Taranto in southern Italy, lies in an area subjected to strong anthropogenic pressure (Annicchiarico et al., 2011; Cardellicchio et al., 2007). Our hydrocarbon data show indeed strong fossil inputs, reflected in the abundance of short chain *n*-alkanes ( $\text{C}_n \leq 23$ ), PAHs profile and in the high  $\sum\text{Phe}/\sum\text{COMB}$  ratio values (Figs. 5c, 6c, 7a; Table 3). Station H12 in the eastern Ionian Sea is located along a main shipping route from Aegean Sea to Adriatic Sea characterized by high

**BGD**

9, 17999–18038, 2012

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



density of likely oil spills (Ferraro et al., 2009). However, this station displays minimum values for TAHC, UCM and TPAH<sub>25</sub> concentrations (Tables 2 and 3; Fig. 3) and low  $\sum\text{Phe}/\sum\text{COMB}$  ratio value (Table 3), indicating low pollutant load and pyrolytic PAH predominance. The low hydrocarbon content of this station is probably related to strong near bottom currents (Poulos et al., 1999) that favour resuspension and dispersal of sedimentary material.

The renewal time of the Ionian deep and bottom waters is approximately 58 yr (Schlitzer et al., 1991). Thus, the deep Ionian basin may act as a long term depository of hydrocarbons exported from the Adriatic Sea, since the latter is the main deep water source (see Sect. 2). Supportive to this higher concentrations of particle-associated PAHs are reported for near bottom rather than surface waters of the Otranto Straits (Parinos, unpublished data) while near bottom concentrations also display a decreasing trend moving from the Otranto Straits to the deep basins of the central Ionian Sea (stations H05 to H03 of this study).

### 5.3.2 Cretan Sea (southern Aegean) and western Cretan-Antikythera straits

For station RED7, maximum values for  $\sum\text{Phe}/\sum\text{COMB}$  ratio along with low  $C_0/(C_0 + C_1)$  Phe ratio values (Table 3), abundance of long chain *n*-alkanes (Fig. 5c) and finally, PAHs profile (Fig. 6c), indicate enhanced fossil inputs. Stations RED3 and RED3.1 located in the western Cretan-Antikythera straits also display enhanced fossil inputs, in contrast with stations in neighboring regions (Ionian Sea and Levantine Sea) where  $\sum\text{Phe}/\sum\text{COMB}$  and  $C_0/(C_0 + C_1)$  Phe ratio values indicate dominant pyrolytic sources (Fig. 7a, b).

Elevated concentrations for petrogenic PAHs ( $\sum\text{Phe}$ ) have been reported in central Aegean Sea and Cretan Sea sediments (Hatzianestis and Sklivagou, 2001; Hatzianestis et al., personal communication). These data combined with our results (Fig. 8) evidence a decreasing southward trend for % $\sum\text{Phe}$  from central Aegean Sea to Cretan Sea and western Cretan-Antikythera straits. This trend could be attributed to

**BGD**

9, 17999–18038, 2012

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the circulation patterns of the Aegean Sea and the chemical characteristics of low-MW petrogenic PAHs.

The latter are less hydrophobic and occur mainly in the dissolved phase, thus they display longer residence time in the water column and could be transported at long distances. The general cyclonic circulation of the Aegean Sea and the net outflow, in the upper 400 m of depth, towards the EMS observed in the western Cretan straits (Kontoyiannis et al., 1999, 2005) likely contribute to the transfer of petrogenic hydrocarbons from central Aegean Sea towards the western Cretan straits. This transfer may be enhanced during periods of deep water formation in the Aegean Sea, when dense water outflows through the eastern and especially the western Cretan strait canyons (Lykousis et al., 2011). The western Cretan strait canyons (stations RED3.1 and RED7) may thus represent a significant sink of hydrocarbons (Tables 2 and 3), in agreement with earlier observations regarding the transfer and distributions of organic pollutants in deep canyon settings (Bouloubassi et al., 2012; Dachs et al., 1997; Fang et al., 2009; Liu et al., 2009; Salvadó et al., 2012).

### 5.3.3 Levantine Sea

The presence of long chain ( $C_n > 35$ ) *n*-alkanes (Fig. 5c) at stations located in Herodotus basin (Her01 and Her03) supports the importance of atmospheric deposition in this area. Her01 and Her03 also display a dominant pyrolytic PAH burden, as shown by the low  $\sum \text{Phe} / \sum \text{COMB}$  values (Table 3) and PAHs profile (Fig. 6c). In contrast, at stations around Ierapetra deep (IER01, RED1.1 and RED15.1) the values of  $\sum \text{Phe} / \sum \text{COMB}$  and  $C_0 / (C_0 + C_1)$  Phe and PAHs profile (Table 3; Fig. 6c) reflect an elevated petrogenic PAH contribution. The presence of the well documented Ierapetra anticyclone (Larnicol et al., 2002; Taupier-Letage, 2008), a semi-permanent eddy, further acts to entrap water masses exiting the Cretan sea, increasing the residence time of hydrocarbons and enhancing their downward transfer.

**BGD**

9, 17999–18038, 2012

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 5.4 Drivers of hydrocarbon's distribution in deep-sea sediments of the Eastern Mediterranean Sea

In order to assess the main processes driving hydrocarbon distributions in the study area, correlation analysis was performed for aliphatic and polycyclic aromatic hydrocarbons concentrations and Total Organic Carbon (TOC) content of sediments.

TAHC, UCM, NA, Ter.NA, TPAH<sub>25</sub>,  $\sum$ Phe and  $\sum$ COMB concentrations show significant correlation with TOC (Table 5), indicating that organic carbon exerts an important control on the distribution of both natural and anthropogenic hydrocarbons in the study area. This is in agreement with earlier observations in coastal and open sea marine sediments (Bouloubassi et al., 2012; Charlesworth et al., 2002; De Luca et al., 2004; Gogou et al., 2000; Mayer, 1993; Oros and Ross, 2004; Readman et al., 2002; Tsapakis et al., 2003; Witt et al., 1995; Yang et al., 2000) and is attributed to the high affinity of hydrophobic aliphatic and polycyclic aromatic hydrocarbons to organic matter.

Overall, the concentrations of aliphatic hydrocarbons correlate with the water column depth indicating that sediments in the deep EMS basins act as their repository (Table 5). The significant correlation of UCM/TRes ratio to depth ( $r = 0.556$ ,  $p = 0.007$ ), imply that the accumulation is stronger for chronic anthropogenic (petrogenic) pollutant burden rather than for natural, biogenic hydrocarbons.

Although TPAH<sub>25</sub>,  $\sum$ Phe and  $\sum$ COMB concentrations correlate to organic carbon, they do not correlate with water column depth (Table 5). Also, as discussed in Sect. 4.3, aliphatic and polycyclic aromatic hydrocarbons display different maximum concentration points, i.e. station H03 for aliphatic hydrocarbons and H05 for PAHs. The above imply that although organic carbon exerts the main control on the distribution of both TAHC and TPAH<sub>25</sub> in the study area, an additional factor affects the dispersal of PAHs. Elemental (black) carbon and especially the partitioning between natural organic carbon and combustion derived black carbon has been reported to exert the main control on PAHs regional distribution (Accardi-Dey and Gschwend, 2002, 2003). It is well documented that high-MW pyrolytic PAH show greater affinity to soot (black carbon)

**BGD**

9, 17999–18038, 2012

### Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rather than organic carbon (Boehm and Farrington, 1984; Dachs and Eisenreich, 2000; Gustaffson et al., 1997), while low-MW PAH (mostly of fossil origin) as more soluble are more efficiently scavenged by organic rich particles e.g phytoplankton and fecal pellets (Dachs et al., 1996, 1997).

## 6 Conclusions

Aliphatic hydrocarbons and PAHs in surface deep-sea sediments of the Eastern Mediterranean Sea (EMS) occur at levels in the same order of magnitude as those found in deep-sea settings in the Western Mediterranean Sea and worldwide. Their distribution is characterised by an increasing trend moving from Cretan Sea (Southern Aegean Sea) and Levantine Sea towards the deep basins of the Ionian Sea. Their molecular profiles evidence contributions of both natural and anthropogenic inputs in the study area. The former derive from terrestrial vegetation while the latter are related to both intense maritime traffic and atmospheric inputs. The composition of hydrocarbon mixtures displays significant regional variability reflecting the relative importance of sources.

Concentrations of both aliphatic and polycyclic aromatic hydrocarbons exhibit significant correlations with organic carbon contents, indicating that the latter exerts a main control on the transport, fate and ultimate accumulation of both natural and anthropogenic chemical species in the deep basins of the EMS. Aliphatic hydrocarbon concentrations correlate with water column depth, emerging the role of deep EMS basins as repositories for these compounds, while the lack of correlation between PAH concentrations and depth imply that PAHs distribution is not controlled by organic carbon alone, but probably from the partitioning between natural organic carbon and combustion derived elemental (black or soot) carbon.

Distribution of hydrocarbons in the study area is also driven by strong sub-basin and mesoscale variability of water masses that impact their regional characteristics. Our results imply a transport of more soluble petrogenic PAHs from the central Aegean Sea

BGD

9, 17999–18038, 2012

### Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





through the western Cretan-Antikythera straits to the western Cretan straits canyons, the latter acting as a sink of hydrocarbons. Further investigation is needed on the impact of organic pollutants to deep-sea biological communities.

*Acknowledgements.* This work has been supported by the HERMIONE IP (EU/FP7), MEDE-COS and REDECO (Marin-ERA, EU/FP6) Projects and a grant by Generalitat de Catalunya to University of Barcelona (2009 SGR 1305). We sincerely thank chief scientists K.-C. Emeis, N. Lampadariou and the officers and crews of R/V *Aegaeo* (Hellenic Centre for Marine Research, Greece) and R/V *Meteor* (University of Hamburg, Germany) for their precious help during cruises, and Chara Kyriakidou (HCMR, Athens) for preparation of the study area map.

## References

- Aboul-Kassim, T. A. T. and Simoneit, B. R. T.: Petroleum hydrocarbon fingerprinting and sediment transport assessed by molecular biomarker and multivariate statistical analyses in the Eastern Harbour of Alexandria, Egypt, *Mar. Pollut. Bull.*, 30, 63–73, doi:10.1016/0025-326x(94)00102-f, 1995.
- Accardi-Dey, A. and Gschwend, P. M.: Assessing the combined roles of natural organic matter and black carbon as sorbents in sediments, *Environ. Sci. Technol.*, 36, 21–29, doi:10.1021/es010953c, 2002.
- Accardi-Dey, A. and Gschwend, P. M.: Reinterpreting literature sorption data considering both absorption into organic carbon and adsorption onto black carbon, *Environ. Sci. Technol.*, 37, 99–106, doi:10.1021/es020569v, 2003.
- Alebic-Juretic, A.: Polycyclic aromatic hydrocarbons in marine sediments from the Rijeka Bay area, Northern Adriatic, Croatia, 1998–2006, *Mar. Pollut. Bull.*, 62, 863–869, doi:10.1016/j.marpolbul.2011.01.035, 2011.
- Annicchiarico, C., Buonocore, M., Cardellicchio, N., Di Leo, A., Giandomenico, S., and Spada, L.: PCBs, PAHs and metal contamination and quality index in marine sediments of the Taranto Gulf, *Chem. Ecol.*, 27, 21–32, doi:10.1080/02757540.2010.536156, 2011.
- Barakat, A. O., Mostafa, A., Wade, T. L., Sweet, S. T., and El Sayed, N. B.: Distribution and characteristics of PAHs in sediments from the Mediterranean coastal environment of Egypt, *Mar. Pollut. Bull.*, 62, 1969–1978, doi:10.1016/j.marpolbul.2011.06.024, 2011.

**BGD**

9, 17999–18038, 2012

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Boehm, P. D. and Farrington, J. W.: Aspects of the polycyclic aromatic hydrocarbon geochemistry of recent sediments in the Georges Bank Region, *Environ. Sci. Technol.*, 18, 840–845, 1984.
- Botsou, F. and Hatzianestis, I.: Polycyclic aromatic hydrocarbons (PAHs) in marine sediments of the Hellenic coastal zone, eastern Mediterranean: levels, sources and toxicological significance, *J. Soil. Sediment.*, 12, 265–277, doi:10.1007/s11368-011-0453-1, 2012.
- Bouloubassi, I. and Saliot, A.: Investigation of anthropogenic and natural organic inputs in estuarine sediments using hydrocarbon markers (NAH, LAB, PAH), *Oceanol. Acta*, 16, 145–161, 1993.
- Bouloubassi, I., Lipiatou, E., Saliot, A., Tolosa, I., Bayona, J. M., and Albaiges, J.: Carbon sources and cycle in the western Mediterranean – the use of molecular markers to determine the origin of organic matter, *Deep-Sea Res. Pt II*, 44, 781–799, doi:10.1016/s0967-0645(96)00094-x, 1997.
- Bouloubassi, I., Fillaux, J., and Saliot, A.: Hydrocarbons in surface sediments from there Changjiang (Yangtze River) Estuary, East China Sea, *Mar. Pollut. Bull.*, 42, 1335–1346, doi:10.1016/s0025-326x(01)00149-7, 2001.
- Bouloubassi, I., Méjanelle, L., Pete, R., Fillaux, J., Lorre, A., and Point, V.: PAH transport by sinking particles in the open Mediterranean Sea: a 1 year sediment trap study, *Mar. Pollut. Bull.*, 52, 560–571, doi:10.1016/j.marpolbul.2005.10.003, 2006.
- Bouloubassi, I., Roussiez, V., Azzoug, M., and Lorre, A.: Sources, dispersal pathways and mass budget of sedimentary polycyclic aromatic hydrocarbons (PAH) in the NW Mediterranean margin, Gulf of Lions, *Mar. Chem.*, 142, 18–28, doi:10.1016/j.marchem.2012.07.003, 2012.
- Brassell, S. C. and Eglinton, G.: Environmental chemistry – an interdisciplinary subject. Natural and pollutant organic compounds in contemporary aquatic environments, in: *Analytical Techniques in Environmental Chemistry*, edited by: Albaiges, J., Pergamon, Oxford, 1980.
- Brooks, B. T., Kurtz, S. S., Boord, C. E., and Schmerling, L. (Eds.): *The Chemistry of Petroleum Hydrocarbons*, Vol. 1, Chapt. 3., Reinhold Publishing Corp., New York, 1954.
- Cardellicchio, N., Buccolieri, A., Giandomenico, S., Lopez, L., Pizzulli, F., and Spada, L.: Organic pollutants (PAHs, PCBs) in sediments from the Mar Piccolo in Taranto (Ionian Sea, Southern Italy), *Mar. Pollut. Bull.*, 55, 451–458, doi:10.1016/j.marpolbul.2007.09.007, 2007.
- Castro-Jiménez, J., Berrojalbiz, N., Wollgast, J., and Dachs, J.: Polycyclic aromatic hydrocarbons (PAHs) in the Mediterranean Sea: atmospheric occurrence, deposition and

decoupling with settling fluxes in the water column, *Environ. Pollut.*, 166, 40–47, doi:10.1016/j.envpol.2012.03.003, 2012.

Countway, R. E., Dickhut, R. M., and Canuel, E. A.: Polycyclic aromatic hydrocarbon (PAH) distributions and associations with organic matter in surface waters of the York River, VA Estuary, *Org. Geochem.*, 34, 209–224, doi:10.1016/s0146-6380(02)00162-6, 2003.

Civitaresse, G., Gačić, M., Lipizer, M., and Eusebi Borzelli, G. L.: On the impact of the Bi-modal Oscillating System (BiOS) on the biogeochemistry and biology of the Adriatic and Ionian Seas (Eastern Mediterranean), *Biogeosciences*, 7, 3987–3997, doi:10.5194/bg-7-3987-2010, 2010.

Dachs, J., Bayona, J. M., Fowler, S. W., Miquel, J. C., and Albaigés, J.: Vertical fluxes of polycyclic aromatic hydrocarbons and organochlorine compounds in the western Alboran Sea (southwestern Mediterranean), *Mar. Chem.*, 52, 75–86, 1996.

Dachs, J., Bayona, J. M., Raoux, C., and Albaiges, J.: Spatial, vertical distribution and budget of polycyclic aromatic hydrocarbons in the Western Mediterranean seawater, *Environ. Sci. Technol.*, 31, 682–688, 1997.

Dachs, J. and Eisenreich, S. J.: Adsorption onto aerosol soot carbon dominates gas-particle partitioning of polycyclic aromatic hydrocarbons, *Environ. Sci. Technol.*, 34, 3690–3697, doi:10.1021/es991201, 2000.

Dachs, J., Lohmann, R., Ockenden, W. A., Méjanelle, L., Eisenreich, S. J., and Jones, K. C.: Oceanic biogeochemical controls on global dynamics of persistent organic pollutants, *Environ. Sci. Technol.*, 36, 4229–4237, doi:10.1021/es025724k, 2002.

Danovaro, R., Company, J. B., Corinaldesi, C., D’Onghia, G., Galil, B., Gambi, C., Goaday, A. J., Lampadariou, N., Luna, G. M., Morigi, C., Olu, K., Polymenakou, P., Ramirez-Llodra, E., Sabbatini, A., Sardá, F., Sibuet, M., and Tselepidis, A.: Deep-sea biodiversity in the Mediterranean Sea: the known, the unknown, and the unknowable, *PLoS ONE*, 5, e11832, doi:10.1371/journal.pone.0011832, 2010.

De Luca, G., Furesi, A., Leardi, R., Micera, G., Panzanelli, A., Costantina Piu, P., and Sanna, G.: Polycyclic aromatic hydrocarbons assessment in the sediments of the Porto Torres Harbor (Northern Sardinia, Italy), *Mar. Chem.*, 86, 15–32, 2004.

Eglinton, G. and Hamilton, R. J.: Leaf epicuticular waxes, *Science*, 156, 1322–1335, 1967.

European Environment Agency: Priority Issues in the Mediterranean Environment, EEA Report no. 4, EEA, Copenhagen, 2006.

**BGD**

9, 17999–18038, 2012

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fang, M. D., Chang, W. K., Lee, C. L., and Liu, J. T.: The use of polycyclic aromatic hydrocarbons as a particulate tracer in the water column of Gaoping (Kaoping) Submarine Canyon, J. Marine Syst., 76, 457–467, 2009.

Farrington J. W. and Quinn, J. G.: Petroleum hydrocarbons in Narragansett Bay: I. Survey of hydrocarbons in sediments and clams (*Mercenaria mercenaria*), Estuar. Coast. Mar. Sci., 1, 71–79, 1973.

Ferraro, G., Meyer-Roux, S., Muellenhoff, O., Pavliha, M., Svetak, J., Tarchi, D., and Topouzelis, K.: Long term monitoring of oil spills in European seas, Int. J. Remote Sens., 30, 627–645, doi:10.1080/01431160802339464, 2009.

Garcia-Orellana, J., Pates, J. M., Masqué, P., Bruach, J. M., and Sanchez-Cabeza, J. A.: Distribution of artificial radionuclides in deep sediments of the Mediterranean Sea, Sci. Total Environ., 407, 887–898, doi:10.1016/j.scitotenv.2008.09.018, 2009.

Gogou, A., Stratigakis, N., Kanakidou, M., and Stephanou, E. G.: Organic aerosols in Eastern Mediterranean: components source reconciliation by using molecular markers and atmospheric back trajectories, Org. Geochem., 25, 79–96, doi:10.1016/s0146-6380(96)00105-2, 1996.

Gogou, A. I., Apostolaki, M., and Stephanou, E. G.: Determination of organic molecular markers in marine aerosols and sediments: one-step flash chromatography compound class fractionation and capillary gas chromatographic analysis, J. Chromatogr. A, 799, 215–231, doi:10.1016/s0021-9673(97)01106-0, 1998.

Gogou, A., Bouloubassi, I., and Stephanou, E. G.: Marine organic geochemistry of the Eastern Mediterranean: 1. Aliphatic and polyaromatic hydrocarbons in Cretan Sea surficial sediments, Mar. Chem., 68, 265–282, doi:10.1016/s0304-4203(99)00082-1, 2000.

Gonul, L. T. and Kucuksezgin, F.: Aliphatic and polycyclic aromatic hydrocarbons in the surface sediments from the Eastern Aegean: assessment and source recognition of petroleum hydrocarbons, Environ. Sci. Pollut. R., 19, 31–41, doi:10.1007/s11356-011-0524-2, 2012.

Gough, M. A. and Rowland, S. J.: Characterization of unresolved complex mixtures of hydrocarbons in petroleum, Nature, 344, 648–650, 1990.

Grimalt, J., Albaigés, J., Sicre, M. A., Marty, J. C., and Salot, A.: Aerosol transport of polynuclear aromatic hydrocarbons over the Mediterranean Sea, Naturwissenschaften, 75, 39–42, doi:10.1007/bf00367439, 1988.

- Gustafsson, Ö., Haghseta, F., Chan, C., Macfarlane, J., and Gschwend, P. M.: Quantification of the dilute sedimentary soot phase: implications for PAH speciation and bioavailability, *Environ. Sci. Technol.*, 31, 203–209, doi:10.1021/es960317s, 1997.
- Guzzella, L. and De Paolis, A.: Polycyclic aromatic hydrocarbons in sediments of the Adriatic Sea, *Mar. Pollut. Bull.*, 28, 159–165, doi:10.1016/0025-326x(94)90392-1, 1994.
- Hatzianestis, I. and Sklivagou, E.: Distribution of Polycyclic Aromatic Hydrocarbons in Surface Sediments of the Aegean Sea (Eastern Mediterranean), presented at the 7th Greece-Cyprus chemistry conference, Nikosia, 55–59, 2001.
- Hatzianestis, J., Sklivagou, E., and Georgakopoulou, E.: Hydrocarbons, pesticides and PCBs in sediments from Thermaikos Gulf, Greece, *Fresen. Environ. Bull.*, 10, 63–68, 2001.
- Hsieh, M., Philp, R. P., and Del Rio, J. C.: Characterization of high molecular weight biomarkers in crude oils, *Org. Geochem.*, 31, 1581–1588, doi:10.1016/s0146-6380(00)00085-1, 2000.
- IOC, IHO, and BODC: Centenary Edition of the GEBCO Digital Atlas, published on CD-ROM on behalf of the Intergovernmental Oceanographic Commission and the International Hydrographic Organization as part of the General Bathymetric Chart of the Oceans, British Oceanographic Data Centre, Liverpool, UK, 2003.
- Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, C., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., LaRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global iron connections between desert dust, ocean biogeochemistry, and climate, *Science*, 308, 67–71, doi:10.1126/science.1105959, 2005.
- Killops, S. D. and Al-Juboori, M. A. H. A.: Characterisation of the unresolved complex mixture (UCM) in the gas chromatograms of biodegraded petroleum, *Org. Geochem.*, 15, 147–160, 1990.
- Kontoyiannis, H., Theocharis, A., Balopoulos, E., Kioroglou, S., Papadopoulos, V., Collins, M., Velegarakis, A. F., and Iona, A.: Water fluxes through the Cretan Arc Straits, Eastern Mediterranean Sea: March 1994 to June 1995, *Prog. Oceanogr.*, 44, 511–529, doi:10.1016/s0079-6611(99)00044-0, 1999.
- Kontoyiannis, H., Balopoulos, E., Gotsis-Skretas, O., Pavlidou, A., Assimakopoulou, G., and Papageorgiou, E.: The hydrology and biochemistry of the Cretan Straits (Antikithira and Kassos Straits) revisited in the period June 1997–May 1998, *J. Marine Syst.*, 53, 37–57, doi:10.1016/j.jmarsys.2004.06.007, 2005.

# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Krom, M. D., Groom, S., and Zohary, T.: The Eastern Mediterranean, in: The Biogeochemistry of Marine Systems, edited by: Black, K. D. and Shimmield, G. B., Blackwell Publishing, Oxford, 91–122, 2003.
- Kucuksezgin, F., Pazi, I., and Gonul, L. T.: Marine organic pollutants of the Eastern Aegean: aliphatic and polycyclic aromatic hydrocarbons in Candarli Gulf surficial sediments, *Mar. Pollut. Bull.*, 64, 2569–2575, doi:10.1016/j.marpolbul.2012.07.019, 2012.
- Ladji, R., Yassaa, N., Balducci, C., and Cecinato, A.: Organic components of Algerian desert dusts, *Chemosphere*, 81, 925–931, doi:10.1016/j.chemosphere.2010.07.059, 2010.
- Laflamme, R. E. and Hites, R. A.: The global distribution of polycyclic aromatic hydrocarbons in recent sediments, *Geochim. Cosmochim. Ac.*, 42, 289–303, 1978.
- Larnicol, G., Ayoub, N., and Le Traon, P. Y.: Major changes in Mediterranean Sea level variability from 7 years of TOPEX/Poseidon and ERS-1/2 data, *J. Marine Syst.*, 33–34, 63–89, doi:10.1016/s0924-7963(02)00053-2, 2002.
- Leister, D. L. and Baker, J. E.: Atmospheric deposition of organic contaminants to the Chesapeake Bay, *Atmos. Environ.*, 28, 1499–1520, doi:10.1016/1352-2310(94)90210-0, 1994.
- Lipiatou, E. and Saliot, A.: Fluxes and transport of anthropogenic and natural polycyclic aromatic hydrocarbons in the western Mediterranean Sea, *Mar. Chem.*, 32, 51–71, 1991.
- Lipiatou, E., Tolosa, I., Simó, R., Bouloubassi, I., Dachs, J., Marti, S., Sicre, M. A., Bayona, J. M., Grimalt, J. O., Saliot, A., and Albaigés, J.: Mass budget and dynamics of polycyclic aromatic hydrocarbons in the Mediterranean Sea, *Deep-Sea Res. Pt II*, 44, 881–905, doi:10.1016/s0967-0645(96)00093-8, 1997.
- Liu, J. T., Hung, J.-J., Lin, H.-L., Huh, C.-A., Lee, C.-L., Hsu, R. T., Huang, Y.-W., and Chu, J. C.: From suspended particles to strata: the fate of terrestrial substances in the Gaoping (Kaoping) submarine canyon, *J. Marine Syst.*, 76, 417–432, doi:10.1016/j.jmarsys.2008.01.010, 2009.
- Liu, L.-Y., Wang, J.-Z., Wei, G.-L., Guan, Y.-F., and Zeng, E. Y.: Polycyclic aromatic hydrocarbons (PAHs) in continental shelf sediment of China: implications for anthropogenic influences on coastal marine environment, *Environ. Pollut.*, 167, 155–162, doi:10.1016/j.envpol.2012.03.038, 2012.
- Lykousis, V., Smith, C., Salomidi, M., Stavrakakis, S., and Kontogiannis, H.: Hydro-sedimentary processes, deep corals and anthropogenic impacts in the canyons of the W. Cretan straits: new findings from HERMIONE ROV dives and fixed moorings, HERMIONE workshop, Malaga, 2011.

- Mackenzie, A. S.: Applications of biological biomarkers in petroleum geochemistry, in: *Advances in Petroleum Geochemistry*, edited by: Brooks, J. and Welte, D., Academic Press, London, 115–214, 1984.
- 5 Malanotte-Rizzoli, P. and Hecht, A.: Large scale properties of the Eastern Mediterranean: a review, *Oceanol. Acta*, 11, 323–335, 1988.
- Malanotte-Rizzoli, P., Manca, B. B., D'Alcalà, M. R., Theocharis, A., Bergamasco, A., Bregant, D., Budillon, G., Clvitarese, G., Georgopoulos, D., Michelato, A., Sansone, E., Scarazato, P., and Souvermezoglou, E.: A synthesis of the Ionian Sea hydrography, circulation and water mass pathways during POEM-phase I, *Prog. Oceanogr.*, 39, 153–204, doi:10.1016/s0079-6611(97)00013-x, 1997.
- 10 Marcomini, A., Pavoni, B., Donazzolo, R., and Orio, A. A.: Combined preparative and analytical use of normal-phase and reversed-phase high-performance liquid chromatography for the determination of aliphatic and polycyclic aromatic hydrocarbons in sediments of the Adriatic Sea, *Mar. Chem.*, 18, 71–84, 1986.
- 15 Mayer, L. M.: Organic matter at the sediment–water interface, in: *Org. Geochem., Principles and Applications*, edited by: Engel, M. N. and Macko, S. A., Plenum Press, New York, 171–184, 1993.
- Millot, C. and Taupier-Letage, I.: Circulation in the Mediterranean Sea, *Handbook Environ. Chem.*, 5K, 29–66, 2005.
- 20 Neff, J.: *Polycyclic Aromatic Hydrocarbons. Sources, Fates and Biological Effects*, Applied Science Publ., London, 262 pp., 1979.
- Nemr, A. E., Said, T. O., Khaled, A., El-Sikaily, A., and Abd-Allah, A. M. A.: The distribution and sources of polycyclic aromatic hydrocarbons in surface sediments along the Egyptian Mediterranean coast, *Environ. Monit. Assess.*, 124, 343–359, doi:10.1007/s10661-006-9231-8, 2007.
- 25 Ohkouchi, N., Kawamura, K., and Wahata, H.: Distributions of three- to seven-ring polynuclear aromatic hydrocarbons on the deep sea floor in the Central Pacific, *Environ. Sci. Technol.*, 33, 3086–3090, 1999.
- Oros, D. R. and Ross, J. R. M.: Polycyclic aromatic hydrocarbons in San Francisco Estuary sediments, *Mar. Chem.*, 86, 169–184, 2004.
- 30 OSPAR: Quality Status Report 2000: Region IV – Bay of Biscay and Iberian Coast, OSPAR Commission, London, 2000.

# **Natural and anthropogenic hydrocarbons in deep EMS sediments**

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Parinos, C., Gogou, A., Bouloubassi, I., Stavrakakis, S., Plakidi, E., and Hatzianestis, I.: Sources and downward fluxes of polycyclic aromatic hydrocarbons in the open southwestern Black Sea, *Org. Geochem.*, submitted, 2012.

Poulos, S. E., Lykousis, V., Collins, M. B., Rohling, E. J., and Pattiaratchi, C. B.: Sedimentation processes in a tectonically active environment: the Kerkyra-Kefalonia submarine valley system (NE Ionian Sea), *Mar. Geol.*, 160, 25–44, doi:10.1016/s0025-3227(99)00016-x, 1999.

Prahl, F. G. and Carpenter, R.: The role of zooplankton fecal pellets in the sedimentation of polycyclic aromatic hydrocarbons in Dabob Bay, Washington, *Geochim. Cosmochim. Ac.*, 43, 1959–1972, 1979.

Prahl, F. G. and Carpenter, R.: Hydrocarbons in Washington coastal sediments, *Estuar. Coast. Shelf Sci.*, 18, 703–720, 1984.

Ramdahl, T.: Retene – a molecular marker of wood combustion in ambient air, *Nature*, 306, 580–582, 1983.

Readman, J. W., Fillmann, G., Tolosa, I., Bartocci, J., Villeneuve, J.-P., Catinni, C., and Mee, L. D.: Petroleum and PAH contamination of the Black Sea, *Mar. Pollut. Bull.*, 44, 48–62, 2002.

Reddy, C. M., Eglinton, T. I., Hounshell, A., White, H. K., Xu, L., Gaines, R. B., and Frysinger, G. S.: The West Falmouth oil spill after thirty years: the persistence of petroleum hydrocarbons in marsh sediments, *Environ. Sci. Technol.*, 36, 4754–4760, doi:10.1021/es020656n, 2002.

Robinson, A. R., Malanotte-Rizzoli, P., Hecht, A., Michelato, A., Roether, W., Theocharis, A., Ünlüata, Ü., Pinardi, N., Artegiani, A., Bergamasco, A., Bishop, J., Brenner, S., Christianidis, S., Gacic, M., Georgopoulos, D., Golnaraghi, M., Hausmann, M., Junghaus, H. G., Lascaratos, A., Latif, M. A., Leslie, W. G., Lozano, C. J., Oğuz, T., Özsoy, E., Papageorgiou, E., Paschini, E., Rozentroub, Z., Sansone, E., Scarazzato, P., Schlitzer, R., Spezie, G. C., Tziperman, E., Zodiatis, G., Athanassiadou, L., Gerges, M., and Osman, M.: General circulation of the Eastern Mediterranean, *Earth-Sci. Rev.*, 32, 285–309, doi:10.1016/0012-8252(92)90002-b, 1992.

Robinson, A. R., Leslie, W. G., Theocharis A., and Lascaratos A.: Mediterranean Sea circulation, in: *Encyclopedia of Ocean Sciences*, Steelr, J., Turekian, K., Thorpe, S., Academic Press, London, 1689–1706, 2001.

# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Roether, W., Manca, B. B., Klein, B., Bregant, D., Georgopoulos, D., Beitzel, V., Kovačević, V., and Luchetta, A.: Recent changes in eastern Mediterranean deep waters, *Science*, 271, 333–335, 1996.

Salot, A.: Natural Hydrocarbons in Sea Water, Chapt. 11, in: Elsevier Oceanography Series, Vol. 31, edited by: Duursma, E. K. and Dawson, R., Elsevier, 327–374, 1981.

Salvadó, J. A., Grimalt, J. O., López, J. F., Palanques, A., Heussner, S., Pasqual, C., Sanchez-Vidal, A., and Canals, M.: Role of dense shelf water cascading in the transfer of organochlorine compounds to open marine waters, *Environ. Sci. Technol.*, 46, 2624–2632, doi:10.1021/es2038189, 2012.

Samanta, S. K., Singh, O. V., and Jain, R. K.: Polycyclic aromatic hydrocarbons: environmental pollution and bioremediation, *Trends Biotechnol.*, 20, 243–248, doi:10.1016/s0167-7799(02)01943-1, 2002.

Scarlett, A., Galloway, T. S., and Rowland, S. J.: Chronic toxicity of unresolved complex mixtures (UCM) of hydrocarbons in marine sediments, *J. Soil. Sediment.*, 7, 200–206, doi:10.1065/jss2007.06.232, 2007.

Schlitzer, R., Roether, W., Oster, H., Junghans, H. G., Hausmann, M., Johannsen, H., and Michelato, A.: Chlorofluoromethane and oxygen in the Eastern Mediterranean, *Deep-Sea Res. A*, 38, 1531–1551, 1991.

Schlitzer, R.: Ocean Data View, <http://odv.awi.de>, 2011.

Simoneit, B. R. T.: Organic matter of the trophosphere – III. Characterization and sources of petroleum and pyrogenic residues in aerosols over the western United States, *Atmos. Environ.*, 18, 51–67, 1984.

Sklivagou, E., Varnavas, S. P., Hatzianestis, I., and Kaniyas, G.: Assessment of aliphatic and polycyclic aromatic hydrocarbons and trace elements in coastal sediments of the Saronikos Gulf, Greece (Eastern Mediterranean), *Mar. Georesour. Geotechnol.*, 26, 372–393, doi:10.1080/10641190802474554, 2008.

Sporstøl, S., Gjøs, N., Lichtenthaler, R. G., Gustavsen, K. O., Urdal, K., Orelid, F., and Skei, J.: Source identification of aromatic hydrocarbons in sediments using GC/MS, *Environ. Sci. Technol.*, 17, 282–286, 1983.

Taupier-Letage, I.: On the use of thermal images for circulation studies: application to the eastern Mediterranean basin, in: Remote Sensing of the European Sea, edited by: Barale, V. and Gade, M., Springer Verlag, 153–164, 2008.

# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Theocharis, A., Nittis, K., Kontoyiannis, H., Papageorgiou, E., and Balopoulos, E.: Climatic changes in the Aegean Sea influence the Eastern Mediterranean thermohaline circulation (1986–1997), *Geophys. Res. Lett.*, 26, 1617–1620, doi:10.1029/1999gl900320, 1999.
- Tolosa, I., Bayona, J. M., and Albaiges, J.: Aliphatic and polycyclic aromatic hydrocarbons and sulfur/oxygen derivatives in Northwestern Mediterranean sediments: spatial and temporal variability, fluxes, and budgets, *Environ. Sci. Technol.*, 30, 2495–2503, 1996.
- Tsapakis, M., Stephanou, E. G., and Karakassis, I.: Evaluation of atmospheric transport as a nonpoint source of polycyclic aromatic hydrocarbons in marine sediments of the Eastern Mediterranean, *Mar. Chem.*, 80, 283–298, doi:10.1016/s0304-4203(02)00132-9, 2003.
- Tsapakis, M. and Stephanou, E. G.: Polycyclic aromatic hydrocarbons in the atmosphere of the Eastern Mediterranean, *Environ. Sci. Technol.*, 39, 6584–6590, doi:10.1021/es050532l, 2005.
- UNEP/MAP/MEDPOL: Sub-Regional Assessment of the Status of Marine and Coastal Ecosystems and of Pressures to the Marine and Coastal Environment Eastern Mediterranean Sea, UNEP(DEPI)/MED WG.350/Inf.4, Barcelona, Spain, 37–38, 2010.
- Venkatesan, M. I.: Occurrence and possible sources of perylene in marine sediments – a review, *Mar. Chem.*, 25, 1–27, 1988.
- Wakeham, S. G., Schaffner, C., and Giger, W.: Polycyclic aromatic hydrocarbons in recent lake sediments. I. Compounds having anthropogenic origins, *Geochim. Cosmochim. Ac.*, 44, 403–413, 1980.
- Wakeham, S. G.: Aliphatic and polycyclic aromatic hydrocarbons in Black Sea sediments, *Mar. Chem.*, 53, 187–205, doi:10.1016/0304-4203(96)00003-5, 1996.
- Wang, Z., Fingas, M., and Page, D. S.: Oil spill identification, *J. Chromatogr. A*, 843, 369–411, doi:10.1016/s0021-9673(99)00120-x, 1999.
- Witt, G.: Polycyclic aromatic hydrocarbons in water and sediment of the Baltic Sea, *Mar. Pollut. Bull.*, 31, 237–248, 1995.
- Yang, G. P.: Polycyclic aromatic hydrocarbons in the sediments of the South China Sea, *Environ. Pollut.*, 108, 163–171, doi:10.1016/s0269-7491(99)00245-6, 2000.
- Yilmaz, K., Yilmaz, A., Yemenicioglu, S., Sur, M., Salihoglu, I., Karabulut, Z., Karakoç, F. T., Hatipoglu, E., Gaines, A. F., Phillips, D., and Hewer, A.: Polynuclear aromatic hydrocarbons (PAHs) in the Eastern Mediterranean Sea, *Mar. Pollut. Bull.*, 36, 922–925, doi:10.1016/s0025-326x(98)00094-0, 1998.

Youssif, M. M. S.: Identification of petroleum hydrocarbon contamination in the Egyptian Mediterranean seashore, Water Stud. Ser., 11, 105–114, 2002.

Yunker, M. B., Macdonald, R. W., Vingarzan, R., Mitchell, R. H., Goyette, D., and Sylvestre, S.: PAHs in the Fraser River Basin: a critical appraisal of PAH ratios as indicators of PAH source and composition, Org. Geochem., 33, 489–515, doi:10.1016/s0146-6380(02)00002-5, 2002.

Yunker, M. B., Macdonald, R. W., Snowdon, L. R., and Fowler, B. R.: Alkane and PAH biomarkers as tracers of terrigenous organic carbon in Arctic Ocean sediments, Org. Geochem., 42, 1109–1146, 2011.

**BGD**

9, 17999–18038, 2012

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Detailed location and characteristics of sampling sites.

| Station | Location<br>Lat. N | Long. E    | Water depth (m) | Sampling date | TOC (%) <sup>a</sup> |
|---------|--------------------|------------|-----------------|---------------|----------------------|
| RED2.1  | 33° 42.81′         | 26° 20.45′ | 2720            | May 2010      | 0.37                 |
| RED3    | 35° 24.14′         | 23° 24.10′ | 2976            | May 2010      | 0.37                 |
| RED3.1  | 35° 18.17′         | 23° 18.92′ | 3314            | May 2010      | 0.58                 |
| RED4    | 35° 45.71′         | 25° 06.00′ | 1620            | May 2010      | 0.42                 |
| RED5    | 35° 40.57′         | 25° 06.12′ | 1018            | May 2010      | 0.39                 |
| RED7    | 34° 36.11′         | 24° 08.73′ | 3589            | May 2011      | 0.51                 |
| RED8    | 36° 04.47′         | 25° 17.07′ | 1715            | May 2011      | 0.33                 |
| RED9    | 36° 00.03′         | 23° 53.55′ | 1200            | May 2011      | 0.40                 |
| RED1.1  | 34° 24.13′         | 26° 14.67′ | 3568            | Jun 2012      | 0.54                 |
| RED15.1 | 34° 36.55′         | 25° 55.49′ | 2428            | Jun 2012      | 0.62                 |
| RED13   | 34° 57.01′         | 25° 54.90′ | 1101            | Jun 2012      | 0.46                 |
| H01     | 35° 45.00′         | 23° 00.00′ | 2117            | Jan 2007      | 0.29                 |
| H02     | 35° 45.00′         | 21° 00.00′ | 3008            | Jan 2007      | 0.45                 |
| H03     | 35° 45.00′         | 18° 30.00′ | 4087            | Jan 2007      | 0.63                 |
| H04     | 35° 55.00′         | 16° 00.00′ | 3750            | Jan 2007      | 0.65                 |
| H05     | 37° 30.00′         | 18° 30.00′ | 3154            | Jan 2007      | 0.57                 |
| H07     | 39° 10.00′         | 17° 45.00′ | 1866            | Jan 2007      | 1.15                 |
| H12     | 38° 50.00′         | 19° 45.00′ | 1450            | Jan 2007      | 0.15                 |
| Her01   | 33° 55.44′         | 27° 44.45′ | 2680            | Jan 2007      | 0.31                 |
| Her03   | 33° 40.00′         | 29° 00.00′ | 3090            | Jan 2007      | 0.49                 |
| IER01   | 34° 26.54′         | 26° 11.51′ | 3626            | Jan 2007      | 0.52                 |
| Rho02   | 35° 37.12′         | 27° 42.03′ | 1305            | Jan 2007      | 0.47                 |

<sup>a</sup> TOC values from Pedrosa-Pàmies et al. (2012).

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 2.** Concentrations and characteristic parameters of aliphatic hydrocarbons determined in deep-sea sediments of the Eastern Mediterranean Sea.

| Station | TAHC<br>( $\mu\text{g g}^{-1}$ ) | UCM<br>( $\mu\text{g g}^{-1}$ ) | NA<br>( $\text{ng g}^{-1}$ ) | Ter. NA<br>( $\text{ng g}^{-1}$ ) | CPI <sub>24–35</sub> | UCM/TRes |
|---------|----------------------------------|---------------------------------|------------------------------|-----------------------------------|----------------------|----------|
| RED2.1  | 37.5                             | 24.8                            | 4810                         | 1800                              | 1.78                 | 3.2      |
| RED3    | 16.2                             | 10.8                            | 1680                         | 442                               | 2.37                 | 2.9      |
| RED3.1  | 43.2                             | 31.7                            | 3820                         | 1260                              | 2.78                 | 4.1      |
| RED4    | 13.0                             | 7.75                            | 1790                         | 819                               | 4.25                 | 2.3      |
| RED5    | 11.6                             | 6.41                            | 1660                         | 805                               | 4.07                 | 1.8      |
| RED7    | 28.6                             | 21.7                            | 2410                         | 938                               | 3.42                 | 4.9      |
| RED8    | 7.32                             | 5.32                            | 681                          | 414                               | 5.84                 | 4.0      |
| RED9    | 6.84                             | 5.11                            | 586                          | 400                               | 6.67                 | 4.5      |
| RED1.1  | 15.5                             | 11.3                            | 1500                         | 892                               | 5.51                 | 4.2      |
| RED15.1 | 15.1                             | 10.8                            | 1530                         | 772                               | 4.71                 | 4.0      |
| RED13   | 8.47                             | 5.66                            | 1050                         | 720                               | 7.28                 | 3.2      |
| H01     | 3.23                             | 1.71                            | 550                          | 374                               | 6.18                 | 1.8      |
| H02     | 20.1                             | 14.5                            | 2160                         | 800                               | 3.07                 | 4.2      |
| H03     | 49.2                             | 36.7                            | 4670                         | 1350                              | 2.26                 | 4.7      |
| H04     | 24.2                             | 18.6                            | 2090                         | 1100                              | 3.64                 | 5.3      |
| H05     | 32.0                             | 23.9                            | 2930                         | 1210                              | 3.51                 | 4.5      |
| H07     | 15.7                             | 12.5                            | 997                          | 467                               | 3.64                 | 5.7      |
| H12     | 1.34                             | 0.73                            | 145                          | 80.7                              | 3.66                 | 1.6      |
| Her01   | 12.4                             | 8.64                            | 1460                         | 603                               | 3.86                 | 3.7      |
| Her03   | 20.0                             | 14.5                            | 2100                         | 767                               | 3.13                 | 4.2      |
| IER01   | 15.1                             | 10.6                            | 1800                         | 1070                              | 5.75                 | 3.9      |
| Rho02   | 8.06                             | 5.00                            | 1160                         | 774                               | 6.26                 | 2.6      |

## Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

**Table 4.** Concentrations of aliphatic and polycyclic aromatic hydrocarbons in the study area and data reported for deep-sea sediments of the Mediterranean Sea and other open sites of the world ocean.

| Location                           | Depth (m) | Sediment Layer | TPAH ( $\text{ng g}^{-1}$ )         | TAHC ( $\mu\text{g g}^{-1}$ ) | UCM ( $\mu\text{g g}^{-1}$ ) | NA ( $\text{ng g}^{-1}$ ) | Reference                  |
|------------------------------------|-----------|----------------|-------------------------------------|-------------------------------|------------------------------|---------------------------|----------------------------|
| Eastern Mediterranean Sea          | 1101–4087 | 1 cm           | 11.6–213 ( $\sum\text{PAH}_{25}$ )  | 1.34–49.2                     | 0.73–36.7                    | 145–4810                  | This study                 |
| Black Sea                          | 1500–2210 | 1 or 2 cm      | 320–1250 ( $\sum\text{PAH}_{24}$ )  | < 22 <sup>a</sup>             |                              |                           | Wakeham et al. (1996)      |
| Cretan Sea (Southern Aegean Sea)   | 700–1570  | 1 cm           | 14.7–128 ( $\sum\text{PAH}_{24}$ )  | 0.59–3.02                     | 0.34–2.56                    | 80–497                    | Gogou et al. (2000)        |
| Central Pacific ocean              | 2505–5906 | 2 cm           | 0.81–60.6 ( $\sum\text{PAH}_{14}$ ) |                               |                              |                           | Ohkouchi et al. (1999)     |
| South China Sea                    | 1045–2432 | 2 cm           | 125–199 ( $\sum\text{PAH}_{10}$ )   |                               |                              |                           | Yang et al. (2000)         |
| South China Sea                    | 1050–2456 | 5 cm           | 40–109 ( $\sum\text{PAH}_{18}$ )    |                               |                              |                           | Liu et al. (2012)          |
| Western Mediterranean Sea          | 750–2200  | 1 cm           | 325–1437 ( $\sum\text{PAH}_{31}$ )  |                               |                              |                           | Lipiatou and Saliot (1991) |
| Northwestern Mediterranean Sea     | 1790–2700 | 1 cm           | 100–500 ( $\sum\text{PAH}_{22}$ )   |                               | 1–1.7                        | 1000–1700                 | Tolosa et al. (1996)       |
| Cap Ferret Canyon (Atlantic ocean) | 615–1040  | n.a.           | 615–1040 (n.a.)                     |                               |                              |                           | OSPAR, QRS (2000)          |
| Arctic Ocean                       | 2265–4230 | 1 cm           | 113–2504 ( $\sum\text{PAH}_{29}$ )  |                               |                              | 1760–12400                | Yunker et al. (2011)       |

n.a.: not available.

<sup>a</sup> all stations except Danube estuary (station 4T).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

**Table 5.** Statistically significant correlations for aliphatic and polycyclic aromatic hydrocarbon concentrations, TOC content and water column depth in the study area.

|                    |                         | TAHC     | UCM      | NA       | Ter.NA   | TPAH <sub>25</sub> | $\Sigma$ COMB | $\Sigma$ Phe |
|--------------------|-------------------------|----------|----------|----------|----------|--------------------|---------------|--------------|
| (n = 22)           |                         |          |          |          |          |                    |               |              |
| TOC                | Pearson Correlation     | 0.625**  | 0.652**  | 0.525*   | 0.634**  | 0.544*             | 0.548*        | 0.514*       |
|                    | Significance (2-tailed) | 0.002    | 0.001    | 0.015    | 0.002    | 0.011              | 0.010         | 0.017        |
| Depth              | Pearson Correlation     | 0.706**  | 0.730**  | 0.622**  | 0.565**  |                    |               |              |
|                    | Significance (2-tailed) | < 0.0005 | < 0.0005 | 0.002    | 0.006    |                    |               |              |
| TAHC               | Pearson Correlation     | 1        | 0.995**  | 0.955**  | 0.834**  | 0.617**            | 0.587**       | 0.654**      |
|                    | Significance (2-tailed) |          | < 0.0005 | < 0.0005 | < 0.0005 | 0.002              | 0.004         | 0.001        |
| TPAH <sub>25</sub> | Pearson Correlation     | 0.617**  | 0.645**  | 0.478*   | 0.476*   | 1                  | 0.981**       | 0.909**      |
|                    | Significance (2-tailed) | 0.002    | 0.001    | 0.024    | 0.025    |                    | < 0.0005      | < 0.0005     |

\*\* Correlation is significant at the 0.01 level (2-tailed).

\* Correlation is significant at the 0.05 level (2-tailed).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 1.** Study area and location of sampling sites across the Eastern Mediterranean Sea. Sites presented as circles represent sediments collected during January 2007 cruise, while sites presented as triangles represent sediments collected during May 2010, 2011 and July 2012 cruises (Table 1). The map was produced using GEBCO Digital Atlas (IOC, IHO and BODC, 2003).

# **Natural and anthropogenic hydrocarbons in deep EMS sediments**

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 2.** Molecular profile for *n*-alkanes (**A**) and PAHs (**B**) in deep-sea sediments collected across the Eastern Mediterranean Sea. Individual compound abundances are normalized as percentage of the total sum of NA or PAHs (TPAH<sub>25</sub>) respectively and presented along with min-max deviation lines. NA homologues are assigned with their carbon atoms number. PAH abbreviations: Fluorene (Fl); dibenzothiophene (DBT); methyl-dibenzothiophenes (C<sub>1</sub>-DBT); dimethyl-dibenzothiophenes (C<sub>2</sub>-DBT); phenanthrene (Phe); methyl-phenanthrenes (C<sub>1</sub>-Phe); dimethyl-phenanthrenes (C<sub>2</sub>-Phe); trimethyl-phenanthrenes (C<sub>3</sub>-Phe); anthracene (Anth); fluo-ranthrene (Flth); pyrene (Pyr); methyl-pyrenes (C<sub>1</sub>-Pyr); dimethyl-pyrenes (C<sub>2</sub>-Pyr); retene (Ret); benzo[*a*]anthracene (BaA); chrysene/triphenylene (Chry/Tri); methyl-chrysenes (C<sub>1</sub>-Chry); dimethyl-chrysenes (C<sub>2</sub>-Chry); benzo[*b/j/k*]fluoranthene (BFlths); benzo[*e*]pyrene (BeP); benzo[*a*]pyrene (BaP); perylene (Per); indeno[1,2,3-*cd*]pyrene (IndP); benzo[*ghi*]perylene (BgP) and dibenzo[*a,h*]anthracene (DBA).



**Fig. 3.** Spatial distribution of aliphatic and polycyclic aromatic hydrocarbons concentrations in deep-sea sediments across the Eastern Mediterranean Sea.

# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 4.** Characteristic ion fragmentogram ( $m/z$  191). Hopanes: Ts:  $18\alpha(\text{H})$ -22, 29,30-trisnorneohopane, Tm:  $17\alpha(\text{H})$ -22,29,30-trisnorhopane.  $29\alpha\beta$ :  $17\alpha(\text{H})$ , $21\beta(\text{H})$ -30-norhopane.  $30\alpha\beta$ :  $17\alpha(\text{H})$ ,  $21\beta(\text{H})$ -hopane.  $31\alpha\beta$ :  $17\alpha(\text{H})$ ,  $21\beta(\text{H})$ -homohopanes (22S and 22R).  $32\alpha\beta$ :  $17\alpha(\text{H})$ , $21\beta(\text{H})$ -bishomohopane (22S and 22R).  $33\alpha\beta$ :  $17\alpha(\text{H})$ , $21\beta(\text{H})$ -trishomohopane (22S and 22R).  $34\alpha\beta$ :  $17\alpha(\text{H})$ , $21\beta(\text{H})$ -tetrakishomohopane (22S and 22R).  $35\alpha\beta$ :  $17\alpha(\text{H})$ , $21\beta(\text{H})$ -pentakishomohopane (22S and 22R). Hopenes: A: hop-17(21)-ene. B: hop-13(18)-ene. C:  $17\beta(\text{H})$ -hop-22(29)-ene (Diploptene).

Natural and  
anthropogenic  
hydrocarbons in  
deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.




**Fig. 5.** Hierarchical diagram obtained from cluster analysis of all stations with regard to *n*-alkanes profile **(A)**, clustered station sites **(B)** and characteristic GC-MS chromatographs for each returning cluster **(C)**. Cluster I: Station H03, Cluster II: Station RED9 and Cluster III: Station Her03. I.S: Internal Standard.



# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.




**Fig. 6.** Hierarchical diagram obtained from cluster analysis of all stations with regard to PAH profile characteristics (A), clustered station sites (B) and molecular profile for each returning cluster (C). Individual compound abundances are normalized as percentage of TPAH<sub>25</sub> and presented along with min-max deviation lines.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.



**Fig. 7.** Spatial distribution of  $\Sigma\text{Phe}/\Sigma\text{COMB}$  (**A**) and  $C_0/(C_0 + C_1) \text{ Phe}$  (**B**) ratios in the study area.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

# Natural and anthropogenic hydrocarbons in deep EMS sediments

C. Parinos et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 8.** Spatial distribution of  $\% \Sigma \text{Phe}$  in Cretan Sea (southern Aegean Sea) and western Cretan-Antikythera straits surface sediments. Data for Cretan Sea sediments are a combination of data reported in this study and data reported (stations not shown) by Hatzianestis and Sklivagou (2001).