

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# Global-scale pattern of peatland *Sphagnum* growth driven by photosynthetically active radiation and growing season length

J. Loisel<sup>1</sup>, A. V. Gallego-Sala<sup>2,3,4</sup>, and Z. Yu<sup>1</sup>

<sup>1</sup>Department of Earth and Environmental Sciences, Lehigh University, 1 West Packer Avenue, Bethlehem, PA 18015-3001, USA

<sup>2</sup>Department of Earth and Ecosystem Science, Lunds Universitet, Sölvegatan 12, 223 62 Lund, Sweden

<sup>3</sup>Geography, College of Life and Environmental Sciences, University of Exeter, Amory Building, Rennes Drive, Exeter, EX4 4RJ, UK

<sup>4</sup>School of Geographical Sciences, University of Bristol, University Road, Bristol BS8 1SS, UK

Received: 18 January 2012 – Accepted: 8 February 2012 – Published: 27 February 2012

Correspondence to: J. Loisel (jul208@lehigh.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**BGD**

9, 2169–2196, 2012

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

High-latitude peatlands contain about one third of the world's soil organic carbon, most of which is derived from partly decomposed *Sphagnum* (peat moss) plants. We conducted a meta-analysis based on a global dataset of *Sphagnum* growth measurements collected from published literature to investigate the effects of bioclimatic variables on *Sphagnum* growth. Analysis of variance and general linear models were used to relate *Sphagnum magellanicum* and *S. fuscum* growth rates to photosynthetically active radiation integrated over the growing season ( $PAR_0$ ) and a moisture index. We found that  $PAR_0$  was the main predictor of *Sphagnum* growth for the global dataset, and effective moisture was only correlated with moss growth at continental sites. The strong correlation between *Sphagnum* growth and  $PAR_0$  suggests the existence of a global pattern of growth, with slow rates under cool climate and short growing seasons, highlighting the important role of temperature and growing season length in explaining peatland biomass production. Large-scale patterns of cloudiness during the growing season might also limit moss growth. Although considerable uncertainty remains over the carbon balance of peatlands under a changing climate, our results suggest that increasing  $PAR_0$  as a result of global warming and lengthening growing seasons could promote *Sphagnum* growth. Assuming that production and decomposition have the same sensitivity to temperature, this enhanced growth could lead to greater peat-carbon sequestration, inducing a negative feedback to climate change.

## 1 Introduction

Peatlands are wetland ecosystems that are primarily distributed across the boreal and subarctic regions of the world. They cover about 3% of the global land area, and represent at least half of the world's wetlands. In these ecosystems, net primary production (NPP) exceeds decomposition, resulting in the formation and accumulation of carbon-rich peat deposits that can be up to tens of meters thick (Rydin and

**BGD**

9, 2169–2196, 2012

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jeglum, 2006). Since the last deglaciation, peatlands have accumulated over 600 gigatons of organic carbon (Gt C) globally (Yu et al., 2010), which constitutes up to about a third of the global soil organic C. Many paleoecological and modeling studies have shown that inter-annual, centennial, and millennial changes in peat-C accumulation rates are mainly controlled by the regional climate (e.g., Mauquoy et al., 2002; Malmer and Wallén, 2004; Frohking et al., 2010), and that warming might promote peat-C sequestration in temperature-sensitive regions by increasing NPP more than decomposition (Frohking et al., 2003; Beilman et al., 2009; Jones and Yu, 2010; Charman et al., 2012; Loisel and Yu, 2012). These results point to a potential increase in the peat C-sink capacity under the ongoing and projected climate warming, which would result in a negative feedback to climate change. However, this finding remains a matter of considerable debate because warming is also expected to promote peat decomposition via direct enhanced microbial decomposition or indirect drying of the peatland surface (Ise et al., 2008; Dorrepaal et al., 2009).

*Sphagnum* (peat moss) forms vast, nearly continuous carpets that dominate the groundcover of most high-latitude peatlands. Although these bryophytes usually do not contribute the largest proportion of peatlands NPP (Blodau, 2002), they possess recalcitrant tissues that resist microbial breakdown (van Breemen, 1995) and release phenolic compounds that inhibit microbial decomposition (Freeman et al., 2001), making them more decay-resistant than most other peatland plants. As a result, partly decomposed *Sphagnum* remnants account for about 50 % of the peatland organic matter (Rydin and Jeglum, 2006). Over time, the imbalance between production and decomposition leads to the formation of *Sphagnum*-rich peat deposits. For this reason, *Sphagnum* is often referred to as the main “peat builder”, and peat mosses might store more organic C than any other plant genus worldwide (Clymo and Hayward, 1982; Wieder, 2006).

Each *Sphagnum* species possesses a well-defined ecological niche that is primarily based on peatland surface-moisture conditions, with dry-adapted species forming hummocks, wet-adapted species growing in hollows, and intermediate species colonizing

**BGD**

9, 2169–2196, 2012

## **Bioclimatic drivers of peatland *Sphagnum* growth**

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lawns. In general, *Sphagnum* NPP increases along this dry-to-wet moisture gradient, with highest productivity under wet conditions (van Breemen, 1995). However, the rate of decomposition is also highest in wet microhabitats, as wet-adapted species preferentially allocate their resources into labile metabolic carbohydrates rather than in recalcitrant structural carbohydrates (Turetsky et al., 2008). Mineral nutrient richness and abundance also influence *Sphagnum* distribution in peatlands with, in general, a higher diversity in groundwater-fed systems (fens) and less variety in acidic, precipitation-fed peatlands (bogs). Overall, species that inhabit wet fens (e.g., *Sphagnum cuspidatum*) are generally characterized by the highest growth values, whereas dry bog hummocks (e.g., *Sphagnum fuscum*) have the lowest productivity (Gunnarsson, 2005).

*Sphagnum* growth form is sensitive to local environmental conditions as mosses are unable to actively regulate carbon uptake and water loss because they lack stomata. For example, mosses growing in wet hollows reside in a less dense carpet than mosses growing on drier hummocks (Titus and Wagner, 1984). These wet-adapted species also possess less-developed capillary water transport systems and have a smaller water-holding capacity than dry-adapted species. As a result, mosses growing in wet hollows are typically responsible for the largest variability in growth rate at the local scale: as soon as the water table gets closer to the peatland surface, they grow quickly because of efficient resource allocation strategies, but they also desiccate quickly following water table drawdown (Titus and Wagner, 1984). Dry-adapted species are more resilient to local moisture changes, mostly because of their ability to retain moisture (Luken, 1985). For example, *Sphagnum fuscum* and *S. magellanicum* are generally considered dry-adapted species, as they typically grow on hummocks and lawns, where they form densely packed communities to avoid desiccation (Rydin and Jeglum, 2006). This growth form is a consequence of the preferential resource allocation to structural carbohydrates (Turetsky et al., 2008), and results in a strongly vertical growth component.

*Sphagnum* productivity has been the object of a series of observational and modeling studies over the past 30 yr (see Wieder (2006) for a review). Many studies have

**BGD**

9, 2169–2196, 2012

## **Bioclimatic drivers of peatland *Sphagnum* growth**

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



found a positive relationship between *Sphagnum* NPP and annual or summer temperature (Moore, 1989; Dorrepaal et al., 2003; Gunnarsson, 2005; Breeuwer et al., 2008), or higher *Sphagnum* productivity at lower latitudes (Wieder and Lang, 1983). However, several studies have found no temperature effect or even a reduction in NPP with increasing temperature, possibly because of moisture stress and dehydration (Weltzin et al., 2001; Gerdol et al., 2007). It was also suggested that higher evapotranspiration as a result of higher temperature might lead to reduced *Sphagnum* NPP (Skre and Oechel, 1981).

In his review on *Sphagnum* productivity, Gunnarsson (2005) found that moss growth was strongly correlated to taxonomic sections (*Acutifolia* vs. *Cuspidata*), site-specific microhabitat characteristics (hummocks vs. hollows), and broad climatic gradients. He shows that mean annual temperature and precipitation together explained about 40 % of the variance in moss growth, with temperature having the most important effect. The study by Gunnarsson (2005) constituted the first and only compilation effort that aimed at understanding both local- and large-scale controls on peat moss growth. In terms of identifying the effects of climate on *Sphagnum* growth, however, the work by Gunnarsson (2005) has one major drawback: the database included 31 *Sphagnum* species, from which many were (i) very poorly represented (22 out of the 31 species have less than 10 growth measurements), (ii) limited in their geographic range (some species are solely found in maritime, subarctic, or nutrient-rich peatlands), and (iii) characterized by very different productivity values due to phylogeny and microhabitat preferences (e.g., oligotrophic hummock species vs. eutrophic hollow species). This collection of a large number of species might have hindered the detection of fundamental effects of climatic gradients on *Sphagnum* productivity. In contrast, the present study focuses on a synthesis of the climatic factors that influence peatland *Sphagnum* growth using the two most well-studied and abundant species: *S. fuscum* and *S. magellanicum*. This approach should improve our ability to establish large-scale geographic and bioclimatic effects on moss growth by reducing the amount of noise induced by species-specific features.

**Bioclimatic drivers of peatland *Sphagnum* growth**

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 1.1 Research question and rationale

In general, warming is expected to increase plant productivity in cold regions (Nemani et al., 2003; Hudson and Henry, 2009). In the boreal and subarctic ecoregions, warming-induced increases in the growing season length (early spring warming) and earlier snowmelt (water availability) are also believed to change the phenology of plants and promote C sequestration in many ecosystems (e.g., Aurela et al., 2004; Menzel et al., 2006; Schwartz et al., 2006; Stelzer and Post, 2009). These effects should be especially beneficial for *Sphagnum* species because spring photosynthesis starts as soon as (i) the snow cover disappears (or even before complete disappearance), and (ii) daily temperatures reach 0 °C, since these non-vascular plants do not root into the colder soil (Moore et al., 2006).

The objective of the present study is to investigate whether the estimated annual growth of two *Sphagnum* species (*S. magellanicum* and *S. fuscum*) can be related to broad environmental controls, namely light, temperature, growing season length, and available moisture. Here we examine large-scale patterns of *Sphagnum* growth using published moss growth values as well as new thermal and moisture indexes. Given that a large amount of peatlands are characterized by extended *Sphagnum* carpets, this study intends to assess the impacts of past, present, and future change in growing season length, PAR, and available moisture on peat biomass production, one of the two most important C flux terms of peatland ecosystems (the other one being peat decomposition).

## 2 Methods

### 2.1 *Sphagnum* growth

We compiled a data set of 142 measurements of *Sphagnum* growth at 52 sites located in North America, Europe, and Southern South America. Site latitudes range from

**BGD**

9, 2169–2196, 2012

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



39° to 68° (Fig. 1 and Table 1). A detailed description of study sites and references can be found in Appendix S1. We examined each publication and only recorded data from studies that reported in situ, species-specific values of *Sphagnum* stem length growth (LG; cm yr<sup>-1</sup>). All these measurements were performed following the cranked wire method (Clymo, 1970), a simple but consistent approach that is widely used in peatland ecology. The accuracy of the cranked wire method is mostly limited by the starting and ending dates of the measurements: if the investigator misses out a portion of the early spring and/or late summer growth, the recorded growth increment will not be representative of the entire growing season, and will result in an underestimation of length growth. We assume that all moss growth values used in this study represent all or most of the growing season growth, but we acknowledge that these values most likely represent minimum growth estimates.

Although these LG terms do not take growth form into account, they were preferred over published data of *Sphagnum* biomass and density because the latter two types of data were derived using a wide array of techniques that could not be directly compared. For example, Bauer et al. (2007) used shoot density and shoot mass to determine moss bulk density, Berendse et al. (2001) used the dry weight of the upper 3 cm (including the capitula) of samples that were 7 or 8 cm in diameter, and Gerdol (1995) averaged the dry weight of 10 capitula. As directly comparing these biomass values would have introduced unquantifiable unknowns in our data set (Clymo, 1970), all the measurements were presented as LG values. This approach is justified since *Sphagnum* growth is predominantly apical, meaning that moss shoots elongate upwards from their apical meristem (Clymo, 1970). From a morphological standpoint, vertical shoot growth is inversely related to branch density so that slower stem growth typically leads to greater branch density. For example, *Sphagnum* length growth will be reduced under dry conditions, but the stem volumetric density will increase, allowing the moss to acquire and retain more moisture (Luken, 1985). This acclimation is partly attributable to the production and “delivery” mechanisms of branches, which are formed in the capitulum (plant apex) and transferred to the apical meristem as the stem elongates (Rydin

**Bioclimatic drivers of peatland *Sphagnum* growth**

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





and Jeglum, 2006). For these reasons, we assumed that environmental conditions exert a primary control on stem growth, which then influences branch density.

To minimize species-specific differences in length growth, only growth data for *Sphagnum fuscum* ( $n = 100$ ) and *S. magellanicum* ( $n = 42$ ) were compiled. These dry-adapted species were selected due to their ubiquity (Gunnarsson, 2005). We did not use data from fertilized and other experimental treatments, but *Sphagnum* measurements from control plots in experimental manipulation studies were included in the database. For each observation, the peatland type (bog, poor fen, or rich fen), the microform on which the *Sphagnum* was growing (hummock, lawn, or hollow), and site continentality (maritime or continental) were compiled (Table 1). Continental and maritime sites were classified using the temperature difference between June and January mean monthly temperatures. If this temperature difference was greater than 25 °C, they were classified as continental sites (96 sites); otherwise, they were classified as maritime (46 sites). Only one growth measurement per site, peatland type, microhabitat, year and species was used. If there were many measurements for one of these categories, the mean productivity value was compiled. Species were well distributed among the continentality classes, with each group composed of 60% *Sphagnum fuscum* and 40% *S. magellanicum*. Peatland types were also relatively well distributed among the continentality classes with, for example, 8.3% of continental peatlands being rich fens vs. 6.5% in the maritime group.

## 2.2 Bioclimatic variables

Effects of climate on *Sphagnum* length growth were examined using two compound bioclimatic variables. First, a measure of photosynthetically active radiation integrated over the growing season, i.e. over days with mean temperature above 0 °C, ( $PAR_0$ , in mol photons  $m^{-2}$  season $^{-1}$ ) was computed for each site based on a gridded long-term mean climatology (temperature, precipitation, and sunshine hours) with a spatial resolution of 0.5° for the period 1931–1960 (CLIMATE 2.2; data available at <http://www.bridge.bris.ac.uk/projects/PAIN>).  $PAR_0$  was calculated from latitude,

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





modern orbital parameters, daily mean temperature values and sunshine hours (Prentice et al., 1993). The instantaneous PAR flux density is first calculated from the incoming shortwave radiation. An integration of this value between sunrise and sunset yields the average daily PAR. This is averaged over the season taking into account latitude and cloudiness. As  $PAR_0$  controls plant  $CO_2$  uptake and exerts a direct control on NPP (Chapin et al., 2002), we argue that this variable can be used as a measure of *Sphagnum* carbon fixation potential. Second, the ratio between annual precipitation ( $P$ ) and the annually integrated equilibrium evapotranspiration ( $E_q$ ) was used as an index of moisture balance ( $P/E_q$ ). Both  $P$  and  $E_q$  were derived from the CLIMATE 2.2 dataset.  $E_q$  is a measure of annual potential evapotranspiration that is a function of daily net radiation and temperature (Prentice et al., 1993; Harrison et al., 2010). This moisture index provides a measure of annual plant water availability, and it was preferred to other moisture indexes based on empirical evidence that  $P/E_q$  is a superior predictor of the occurrence of peatlands and peatland types (Gallego-Sala et al., 2010; Gallego-Sala and Prentice, 2012). These bioclimatic variables were analyzed to avoid the multicollinearity that often emerges from studies in which predictor variables are closely related to one another (Zar, 2010). Indeed, preliminary analysis showed strong correlations (multicollinearity > 60%) between latitude, growing season temperature (GST) and growing degree days above 0 °C ( $GDD_0$ ) for our study sites (data not shown). We avoided such strong intercorrelations in our analysis.

### 2.3 Statistical analysis

*Sphagnum* length growth values ( $cm\ yr^{-1}$ ) were log-transformed to meet the assumption of homoscedasticity before statistical analysis. Growth differences between *Sphagnum* species, peatland types, microforms and continentality classes were tested using one-way analysis of variance (ANOVA). To examine the effect of  $PAR_0$  and  $P/E_q$  on *Sphagnum* growth, we first determined their respective correlation coefficient (Pearson correlation). Simple linear regressions were also performed to obtain the coefficient of determination between each independent variable and *Sphagnum* growth.

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



These results were then used to build a multiple regression model where the variable with the highest coefficient of correlation was incorporated first, followed by the second variable as well as by their combined effect. Values were centered prior to the multiple regression analyses. Simple and multiple regression analyses of *Sphagnum* growth values were performed on (i) all data altogether, and (ii) each continentality class separately. All analyses were performed using PASW Statistics version 18.0 (SPSS Inc., Chicago, IL, USA).

### 3 Results

#### 3.1 Effect of species, peatland type, microform, and continentality on *Sphagnum* productivity

*Sphagnum* growth rates ranged from 0.1 to 6.0 cm yr<sup>-1</sup>, with a mean value of 1.4 ± 0.8 (1 SD) cm yr<sup>-1</sup>. The ANOVA revealed a significant effect of species on productivity ( $F(1, 140) = 5.931, p = 0.016$ , Table 2), with *Sphagnum magellanicum* growing significantly more rapidly than *S. fuscum* (Fig. 2). Similarly, continentality had a strong effect on moss growth ( $F(1, 140) = 13.704, p < 0.0001$ , Table 2), with continental sites characterized by faster growing samples than maritime sites (Fig. 2). Peatland type also had a significant effect on *Sphagnum* growth ( $F(2, 139) = 3.798, p = 0.025$ , Table 2), with rich fens being significantly more productive than bogs (Tukey's LSD:  $p = 0.01$ , Fig. 2). Finally, the ANOVA performed on microform did not reveal significant moss growth differences between hummocks and lawns-hollows combined ( $p = 0.181$ , Table 2; Fig. 2). For all four ANOVAs, the assumption of equality of variance was met (Levene's test:  $p > 0.05$ ).

#### 3.2 Relationships between *Sphagnum* productivity and bioclimatic parameters

The Pearson correlation coefficient between PAR<sub>0</sub> and *Sphagnum* growth revealed a positive correlation between these variables ( $r = 0.48, p < 0.0001$ ). However,  $P/E_q$

**BGD**

9, 2169–2196, 2012

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and *Sphagnum* growth were not correlated ( $r = 0.01$ ,  $p = 0.456$ ). Similarly, simple linear regressions using all data indicated that  $PAR_0$  explained most of the variance in *Sphagnum* growth ( $R^2 = 0.23$ ,  $p < 0.0001$ , Table 3, Fig. 3), but that  $P/E_q$  did not have a significant effect ( $R^2 = 0.00$ ,  $p = 0.912$ , Table 3). These patterns were somewhat different when continental and maritime sites were examined separately. While  $PAR_0$  explained more of the growth variance among continental sites than when all sites were considered ( $R^2 = 0.31$ ,  $p < 0.0001$ , Table 3, Fig. 3), it was not significant among maritime sites ( $R^2 = 0.02$ ,  $p = 0.172$ , Table 3, Fig. 3). Interestingly,  $P/E_q$  came out as a significant (though poor) predictor of moss growth in continental settings ( $R^2 = 0.06$ ,  $p = 0.009$ , Table 3), but was not significant across the maritime sites ( $R^2 = 0.00$ ,  $p = 0.690$ , Table 3). A multiple regression analysis performed on  $PAR_0$  and  $P/E_q$  for continental sites indicated that the interaction effect was not significant ( $p = 0.444$ , Table 3), and that these two variables did not explain significantly more variability in *Sphagnum* growth ( $R^2 = 0.31$ , Table 3) than  $PAR_0$  alone ( $R^2 = 0.31$ ).

## 4 Discussion

### 4.1 Species, peatland type, and microform effects

At the peatland scale, a large range of *Sphagnum* growth values is generally found (e.g., Moore, 1989; Campbell et al., 2000). This variability has often been attributed to species-specific biochemical mechanisms (Turetsky et al., 2008), and to microhabitat characteristics such as depth to water table (Rydin and Jeglum, 2006) or nutrient input (Bubier et al., 2007). In line with these previous studies, our results indicate that *Sphagnum* growth was different among species (*S. fuscum* vs. *S. magellanicum*) and peatland types (bogs vs. poor fens vs. rich fens). However, growth was not correlated to microform (hummock vs. lawn-hollow), contrary to the idea that wetter microhabitats such as lawns and hollows favor *Sphagnum* growth. Since microform classification is arbitrary, it is possible that microforms with similar hydrological and nutrient conditions

**BGD**

9, 2169–2196, 2012

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were classified as low hummocks at a study site but as lawns in another peatland, limiting our ability to distinguish a microform effect on *Sphagnum* growth. Direct measurements of water table depths would be more useful in analyzing microform influence on moss productivity. A smaller size sample for the lawn-hollow group ( $n = 28$ ) than for the hummock group ( $n = 114$ ) might also have limited the potential for a microform effect to be detected (statistical Type II error). Alternatively, it is possible that a microform effect on growth can only be apparent at the peatland scale (i.e., relative differences within a single ecosystem), but cannot be detected at the global scale.

## 4.2 Climatic controls

PAR integrated over the growing season should exert a positive control on peat moss growth in high-latitude regions because plant growth increases with the length and temperature of the growing season. Supporting this hypothesis,  $PAR_0$  was the most important variable explaining *Sphagnum* growth in our analysis, with higher  $PAR_0$  values associated with greater growth rates. These results suggest that peat moss growth is primarily driven by PAR and the growing season length, and imply that broad-scale controls on *Sphagnum* growth are significant despite the effect of local or site-specific factors such as nutrient influx or depth to water table.

Adequate effective moisture is necessary for peatland development and peat moss growth. In our data synthesis, however, no relationship between moss growth and the moisture index ( $P/E_q$ ) was found when all sites were considered. These results suggest that, over broad spatial scales, moisture availability does not play a significant role on *Sphagnum* growth. An adequate moisture supply at the local scale is necessary for peat moss growth, but our results suggest it might only play a secondary role when compared to growing season length, temperature, and PAR.

*Sphagnum* growth from continental sites was mostly sensitive to  $PAR_0$  ( $p < 0.0001$ ) and showed an influence of  $P/E_q$  ( $p = 0.009$ ). These results suggest that peat moss growth becomes limited with decreasing  $PAR_0$  and decreasing moisture availability. In contrast, growth values from maritime sites were neither associated with  $PAR_0$  nor

## **Bioclimatic drivers of peatland *Sphagnum* growth**

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$P/E_q$ . These differences between the continentality classes were not due to a species effect since each group was composed of 60% *Sphagnum fuscum* and 40% *S. magellanicum*. Similarly, peatland types were also relatively well distributed among the continentality classes with, for example, 8.3% of continental peatlands being rich fens vs. 6.5% in the maritime group.

The climate space of  $PAR_0$  and  $P/E_q$  for the northern peatland domain (Yu et al., 2010) is presented to provide context for our results and future work on peatland-climate relationships (Fig. 4). Northern peatlands typically occur where  $PAR_0$  is between 2000 and 7000 mol photons  $m^{-2}$  season $^{-1}$  (mean = 4205), and  $P/E_q$  is between 0.3 and 3 (mean = 1.3). Maritime sites from our study were characterized by significantly lower  $PAR_0$  and higher  $P/E_q$  values than their continental counterpart (Fig. 4). The study sites cover the  $PAR_0$  and  $P/E_q$  climate space relatively well, except for a near-complete absence of data points at  $PAR_0$  values lower than 4000 mol photons  $m^{-2}$  season $^{-1}$ . Peatland sites where flux tower measurements provide contemporary net ecosystem carbon balance budgets are similarly found in the highest  $PAR_0$  and lowest  $P/E_q$  portion of the climate space (Fig. 4). These results imply that much research is needed in colder, higher-latitude regions (with low  $PAR_0$ ), as well as in areas with higher effective moisture (high  $P/E_q$ ).

Bioclimatic differences between continental and maritime sites could explain the patterns in *Sphagnum* growth. For example, continental sites were characterized by significantly higher  $PAR_0$  values than the maritime sites (Fig. 4b). In addition, growing season temperature (GST) and July temperature were significantly different between maritime and continental sites (data not shown), with maritime regions characterized by cooler GST and July T than their continental counterparts. Cloudiness may be an important limiting factor for moss growth in maritime settings (Blodau, 2002; Mauquoy et al., 2002). In addition, the short duration or thin snow cover that characterizes peatlands in maritime regions could negatively impact peat moss growth, as snow cover may play a beneficial role in peatlands by preventing freezing-induced damage to *Sphagnum capitula* (Dorrepaal et al., 2003). An extended snow cover during spring

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

time in continental regions might also create a “warm” thermal environment that would allow rapid moss growth during early spring (Tim Moore, personal communication). Finally, the effect of diurnal temperature on moisture availability could partly explain why continental sites exhibit significantly higher moss growth than their maritime counterparts, as a greater diurnal temperature difference in continental settings more likely leads to dew formation that provides additional moisture to mosses.

A sampling bias among the maritime sites could also explain the weak correlation between  $PAR_0$ ,  $P/E_q$  and moss growth, as well as the overall lower moss growth rates under maritime conditions. As growing season length is relatively long under maritime settings due to mild spring, fall and winter conditions, it is possible that part of the early spring or late fall growth were not measured in the field, resulting in “minimal” *Sphagnum* growth estimates that are not representative of the entire growing season length. In a study on oxygen isotope analysis ( $\delta^{18}O$ ) of modern *Sphagnum* cellulose in Northern England, Daley et al. (2010) showed significant differences between the oxygen isotopic signature of *Sphagnum* cellulose during “winter” (February, March, November) and “summer” (May to August), which they attributed to direct water uptake from precipitation year-round. In their study, *Sphagnum* tissues were collected every month of the year and the  $\delta^{18}O$  value of alpha-cellulose was measured for each of these monthly samples, as well as in corresponding rain water samples. The isotopic signature of moss samples tracked values obtained from the rain samples, suggesting that water uptake is occurring during “winter” months in some maritime regions, and that  $CO_2$  uptake and associated plant growth could also be taking place over winter time.

### 4.3 Role of *Sphagnum* growth in regulating peat-C accumulation in northern peatlands

The response of peatlands to warming conditions is tightly coupled to plant growth and peat decomposition. A recent synthesis of 33 northern peatland sites found highly variable peat-carbon accumulation rates over the past 12 000 yr, with warming periods

**BGD**

9, 2169–2196, 2012

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



coinciding with increases in C accumulation rates (Yu et al., 2009). Similarly, using 77 sites along a north-south transect across the West Siberian Lowlands, Beilman et al. (2009) found a significant correlation between C accumulation and modern mean annual temperature, with warmer sites accumulating more C than colder sites over the past 2000 yr. A clear relationship between degree-days and mean annual temperature was also found for long-term peat-C accumulation in Finland and Canada (Clymo et al., 1998), similar to a positive relation between photosynthetically active radiation and growing season length for peat-C accumulation rates over the last millennium throughout the Northern Hemisphere (Charman et al., 2012). Peatland models have also reported an overall positive effect of warmer temperatures on peat-C sequestration: under a scenario where production and decomposition had the same sensitivity to temperature, productivity effects dominated and enhanced peat accumulation (Frolking et al., 2003). To reverse this trend, a  $Q_{10}$  value of 3 for decomposition rate and a  $Q_{10}$  value of 2 for productivity were needed. These results clearly show the important role of NPP in controlling peat-C sequestration, and results from our meta-analysis lend support to these paleoecological evidences and modeling results by showing that moss growth primarily responds to photosynthetically active radiation above 0 °C over large spatial scales.

## 5 Conclusions and implication

We showed that *Sphagnum* growth is most strongly controlled by photosynthetically active radiation integrated over the growing season ( $PAR_0$ ), which reflects latitudinal, temperature, and cloudiness gradients. These results indicate the existence of a global pattern in *Sphagnum* growth in relation with broad-scale bioclimatic and geographic gradients. Since *Sphagnum* is the main peat builder in most high-latitude peatlands, these patterns have important implications for our understanding of the biogeographic scaling of *Sphagnum*'s carbon sink function. Our results suggest higher peat production under warming climates if there is no moisture stress, in agreement with the idea

**BGD**

9, 2169–2196, 2012

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





that warming could lead to a negative feedback from terrestrial ecosystems as a result of increased net primary production and subsequent increased carbon storage (see Field et al. (2007) for a review). This also supports the recent findings in paleoecological studies by Yu et al. (2009), Beilman et al. (2009), Charman et al. (2012), and Loisel and Yu (2012), which suggest that increasing net primary production under warmer conditions during the Holocene may enhance long-term peat-carbon sequestration, contrary to a hypothesized overriding effect of increased peat decomposition under warming conditions (Ise et al., 2008; Dorrepaal et al., 2009). Specifically, increasing *Sphagnum* growth under a warming climate may shift the balance of productivity and decomposition, favoring peat-carbon sequestration. However, the relative importance of change in production and decomposition under warm climates in the future will depend on the direction and magnitude of moisture changes, which can best be evaluated with peatland models that incorporate both temperature and moisture functions of production and decomposition processes. To this effect, the present meta-analysis is particularly useful for testing and validating the peatland ecosystem models that are used for future projections, as it clearly shows the important role of PAR<sub>0</sub> on peatland biomass production, a component that is neglected in peatland ecosystem models (e.g., Frolking et al., 2010). Furthermore, important data gaps have been identified from the climate space of PAR<sub>0</sub> and available moisture ( $P/E_q$ ) of the northern peatland domain. Future studies on modern *Sphagnum* growth and contemporary peatland flux measurements should focus on site in these regions to generate maximum knowledge in climate controls over plant production and carbon sequestration in peatlands.

**Supplementary material related to this article is available online at:**

**<http://www.biogeosciences-discuss.net/9/2169/2012/>**

**[bgd-9-2169-2012-supplement.pdf](http://www.biogeosciences-discuss.net/9/2169/2012/bgd-9-2169-2012-supplement.pdf).**

**BGD**

9, 2169–2196, 2012

## **Bioclimatic drivers of peatland *Sphagnum* growth**

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



*Acknowledgements.* Alex Ireland, Eric Klein, Michael Clifford and Steph Hunt are acknowledged for discussion on this topic; Dan Brosseau and Chris Burke provided technical support; Alex Ireland, Tim Moore and David Currie provided helpful comments on previous versions of this manuscript. The research was funded by a US NSF grant (ATM # 0628455) and by an NSERC fellowship (BESC-D3–362645-2008).

## References

- Aurela, M., Laurila, T., and Tuovinen, J.-P.: The timing of snow melt controls the annual CO<sub>2</sub> balance in a subarctic fen, *Geophys. Res. Lett.*, 31, L16119, doi:10.1029/2004GL020315, 2004.
- Bauer, I. E., Tirlea, D., Bhatti, J. S., and Errington, R. C.: Environmental and biotic controls on bryophyte productivity along forest to peatland ecotones, *Can. J. Botany*, 85, 463–475, 2007.
- Beilman, D. W., MacDonald, G. M., Smith, L. C., and Reimer, P. J.: Carbon accumulation in peatlands of West Siberia over the last 2000 years, *Global Biogeochem. Cy.*, 23, GB1012. doi:10.1029/2007GB003112, 2009.
- Berendse, F., van Breemen, N., Rydin, H., Buttler, A., Heijmans, M., Hoosbeek, M. R., Lee, J. A., Mitchell, E., Saarinen, T., Vasander, H., and Wallén, B.: Raised atmospheric CO<sub>2</sub> levels and increased N deposition cause shifts in plant species composition and production in *Sphagnum* bogs, *Global Change Biol.*, 7, 591–598, 2001.
- Blodau, C.: Carbon cycling in peatlands – a review of processes and controls, *Environ. Rev.*, 10, 111–134, 2002.
- Breeuwer, A., Heijmans, M. P. D., Robroek, B. J. M., and Berendse, F.: The effect of temperature on growth and competition between *Sphagnum* species, *Oecologia*, 156, 155–167, 2008.
- Bubier, J. L., Moore, T. R., and Bledzki, L. A.: Effects of nutrient addition on vegetation and carbon cycling in an ombrotrophic bog, *Global Change Biol.*, 13, 1–19, 2007.
- Campbell, C., Vitt, D. H., Halsey, L. A., Campbell, I. D., Thormann, M. N., and Bayley, S. E.: Net primary production and standing biomass in northern continental wetlands, *Natural Resources Canada, Report NOR-X-369*, Canadian Forest Service, Edmonton, 2000.
- Chapin III, F. S., Matson, P. A., and Mooney, H. A.: *Principles of Terrestrial Ecosystem Ecology*, Springer, New York, USA, 2002.

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Charman, D., Beilman, D., Blaauw, M., Booth, R. K., Brewer, S., Chambers, F., Christen, J. A., Gallego-Sala, A. V., Harrison, S. P., Hughes, P. D. M., Jackson, S., Korhola, A., Mauquoy, D., Mitchell, F., Prentice, I. C., van der Linden, M., De Vleeschouwer, F., Yu, Z., Alm, J., Bauer, I. E., McCorish, Y., Garneau, M., Hohl, V., Huang, Y., Karofeld, E., Le Roux, G., Loisel, J., Moschen, R., Nichols, J. E., Nieminen, T. M., MacDonald, G. M., Phadtare, N. R., Rausch, N., Sillasoo, Ü., Swindles, G. T., Tuittila, E.-S., Ukonmaanaho, L., Väliranta, M., van Bellen, S., van Geel, B., Vitt, D., and Zhao, Y.: Climate-driven changes in peatland carbon accumulation during the last millennium, *Proc. Natl. Acad. Sci. USA*, submitted, 2011.
- Clymo, R. S.: The growth of *Sphagnum*: methods of measurement, *J. Ecol.*, 58(1), 13–49, 1970.
- Clymo, R. S. and Hayward, P. M.: The Ecology of *Sphagnum*. *Bryophyte Ecology*, edited by: Smith, A. J. E., Chapman & Hall, London, 229–289, 1982.
- Clymo, R. S., Turunen, J., and Tolonen, K.: Carbon accumulation in peatlands, *Oikos*, 81(2), 368–388, 1998.
- Daley, T. J., Barber, K. E., Street-Perrott, F. A., Loader, N. J., Marshall, J. D., Crowley, S. F., and Fisher, E. H. Holocene climate variability revealed by oxygen isotope analysis of *Sphagnum* cellulose from Walton Moss, Northern England, *Quaternary Sci. Rev.*, 29, 1590–1601, 2010.
- Dorrepaal, E., Aerts, R., Cornelissen, J. H. C., Callaghan, T. V., and van Logtestijn, R. S. P.: Summer warming and increased winter snow cover affect *Sphagnum fuscum* growth, structure and production in a sub-arctic bog, *Global Change Biol.*, 10, 93–104, 2003.
- Dorrepaal, E., Toet, S., van Logtestijn, R. S. P., Swart, E., van de Weg, M. J., Callaghan, T. V., and Aerts, R.: Carbon respiration from subsurface peat accelerated by climate warming in the subarctic, *Nature*, 460, 616–620, 2009.
- Field, C. B., Lobell, D. B., Peters, H. A., and Chiariello, N. R.: Feedbacks of terrestrial ecosystems to climate change, *Ann. Rev. Environ. Resour.*, 32, 1–29, 2007.
- Freeman, C., Ostle, N., and Kang, H.: An enzymic “latch” on a global carbon store, *Nature*, 409, 149, 2001.
- Frolking, S., Roulet, N. T., Moore, T. R., Richard, P. J. H., Lavoie, M., and Muller, S. D.: Modeling northern peatland decomposition and peat accumulation, *Ecosystems*, 4, 479–498, 2003.
- Frolking, S., Roulet, N. T., Tuittila, E., Bubier, J. L., Quillet, A., Talbot, J., and Richard, P. J. H.: A new model of Holocene peatland net primary production, decomposition, water balance, and peat accumulation, *Earth Syst. Dynam.*, 1, 1–21, doi:10.5194/esd-1-1-2010, 2010.
- Gallego-Sala, A. V. and Prentice, I. C.: Blanket bog, an endangered biome, *Nature Climate*

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Change, submitted, 2012.

Gallego-Sala, A. V., Clark, J. M., House, J. I., Orr, H. G., Prentice, I. C., Smith, P., Farewell, T., and Chapman, S. J.: Bioclimatic envelope model of climate change impacts on blanket peatland distribution in Great Britain, *Clim. Res.*, 34, C911, doi:10.3354/cr00911, 2010.

5 Gerdol, R.: The growth dynamics of *Sphagnum* based on field measurements in a temperate bog and on laboratory cultures, *J. Ecol.*, 83(3), 431–437, 1995.

Gerdol, R., Petraglia, A., Bragazza, L., Iacumin, P., and Brancaleoni, L.: Nitrogen deposition interacts with climate in affecting production and decomposition rates in *Sphagnum* mosses, *Global Change Biol.*, 13, 1810–1821, 2007.

10 Gunnarsson, U.: Global patterns of *Sphagnum* productivity, *Bryologist*, 27, 269–279, 2005.

Harrison, S. P., Prentice, I. C., Barboni, D., Kohfeld, K. E., Ni, J., and Sutra, J.-P.: Ecophysiological and bioclimatic foundations for a global plant functional classification, *Journal of Vegetation Science*, 21(2), 300–317, 2010.

Hudson, J. M. G. and Henry, G. H. R.: Increased plant biomass in a High Arctic heath community from 1981 to 2008, *Ecology*, 90(10), 2657–2663, 2009.

15 Ise, T., Dunn, A. L., Wosfy, S. C., and Moorcroft, P. R.: High sensitivity of peat decomposition to climate change through water-table feedback, *Nat. Geosci.*, 1, 763–766, 2008.

Jones, M. C. and Yu, Z.: Rapid deglacial and early Holocene expansion of peatlands in Alaska, *Proc. Natl. Acad. Sci. USA*, 107(16), 7347–7352, 2010.

20 Koehler, A.-K., Sottocornola, M., and Kiely, G.: How strong is the current carbon sequestration of an Atlantic blanket bog?, *Global change Biol.*, 17, 309–319, 2011.

Loisel, J. and Yu, Z.: Recent acceleration of carbon accumulation in a boreal peatland, south-central Alaska, 2012-JG-001978, submitted to *J. Geophys. Res.-Biogeosci.*, 2012.

25 Lund, M., Lafleur, P. M., Roulet, N. T., Lindroth, A., Christensen, T. R., Aurela, M., Chojnicki, B. H., Flanagan, L. B., Humphreys, E. R., Laurila, T., Oechel, W. C., Olejnik, J., Rinne, J., Schubert, P., and Nilsson, M. B.: Variability in exchange of CO<sub>2</sub> across 12 northern peatland and tundra sites, *Global Change Biol.*, 16, 2436–2448, 2010.

Luken, J. O.: Zonation of *Sphagnum* mosses: interactions among shoot growth, growth form, and water balance, *Bryologist*, 88(4), 374–379, 1985.

30 Malmer, N. and Wallén, B.: Input rates, decay losses and accumulation rates of carbon in bogs during the last millenium: internal processes and environmental changes, *Holocene*, 14(1), 111–117, 2004.

Mauquoy, D., Engelkes, T., Groot, M. H. M., Markesteijn, F., Oudejans, M. G., van der Plicht, J.,

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and van Geel, B.: High-resolution records of late-Holocene climate change and carbon accumulation in two north-west European ombrotrophic peat bogs. *Paleogeogr. Paleoclimatol. Paleoecol.*, 186, 275–310, 2002.

5 Menzel, A., Sparks, T. H., Estrella, N., Koch, E., Aasa, A., Ahas, R., Alm-Kübler, K., Bissolli, P., Braslavska, O., Briede, A., Chmielewski, F. M., Crepinsek, Z., Curnel, Y., Dahl, A., De Fila, C., Donnelly, A., Filella, Y., Jatcza, K., Måge, F., Mestre, A., Nordli, O., Peñuelas, J., Pirinen, P., Remišová, V., Scheifinger, H., Striz, M., Susnik, A., van Vliet, A. J. H., Wielgolaski, F.-E., Zach, S., and Zust, A.: European phenological response to climate change matches the warming pattern, *Global Change Biol.*, 12, 1969–1976, 2006.

10 Moore, T. R.: Growth and net production of *Sphagnum* at five fen sites, subarctic Eastern Canada, *Can. J. Botany*, 67, 1203–1207, 1989.

Moore, T. R., Lafleur, P. M., Poon, D. M. I., Heumann, B. W., Seaquist, J. W., and Roulet, N. T.: Spring photosynthesis in a cool temperate bog, *Global Change Biol.*, 12, 2323–2335, 2006.

15 Nemani, R. R., Keeling, C. D., Hashimoto, H., Jolly, W. M., Piper, S. C., Tucker, C. J., Myneli, R. B., and Running, S. W.: Climate-driven increases in global terrestrial net primary production from 1982 to 1999, *Science*, 300, 1560–1563, 2003.

Prentice, I. C., Sykes, M. T., and Cramer, W.: A simulation model for the transient effects of climate change on forest landscapes, *Ecol. Model.*, 65, 51–70, 1993.

Rydin, H. and Jeglum, J.: *The Biology of Peatlands*, Oxford University Press, Oxford, 2006.

20 Schwartz, M. D., Ahas, R., and Aasa, A.: Onset of spring starting earlier across the Northern Hemisphere, *Global Change Biol.*, 12, 343–351, 2006.

Skre, O. and Oechel, W. C.: Moss functioning in different taiga ecosystems in interior Alaska, *Oecologia*, 48(1), 50–59, 1981.

Steltzer, H. and Post, E.: Seasons and life cycles, *Science*, 324, 886–887, 2009.

25 Titus, J. E. and Wagner, D. J.: Carbon balance for two *Sphagnum* mosses: water balance resolves a physiological paradox, *Ecology*, 65(6), 1765–1774, 1984.

Turetsky, M. R., Crow, S. E., Evans, R. J., Vitt, D. H., and Wieder, R. K.: Trade-offs in resource allocation among *Sphagnum* species control decomposition in boreal peatlands, *J. Ecol.*, 96, 1297–1305, 2008.

30 van Breemen, N.: How *Sphagnum* bogs down other plants, *Trends Ecol. Evol.*, 10(7), 270–275, 1995.

Weltzin, J. F., Harth, C., Bridgham, S. D., Pastor, J., and Vonderharr, M.: Production and microtopography of bog bryophytes: response to warming and water-table manipulations, *Oe-*

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cologia, 128(4), 557–565, 2001.

Wieder, R. K.: Primary production in boreal peatlands. in: Boreal Peatland Ecosystems, edited by: Wieder, R. K. and Vitt, D. H., Ecological Studies 188, Springer-Verlag, Berlin, Heidelberg, 145–163, 2006.

5 Wieder, R. K. and Lang, G. E.: Net primary production of the dominant bryophytes in a *Sphagnum*-dominated wetland in West Virginia, *Bryologist*, 86(3), 280–286, 1983.

Yu, Z., Beilman, D. W., and Jones, M. C.: Sensitivity of northern peatland carbon dynamics to Holocene climate change. *Carbon Cycling in Northern Peatlands*, edited by: Baird, A. J., Belyea, L. R., Comas, X., Reeve, A. and Slater L., Geophysical Monograph 184, American Geophysical Union, Washington, 55–69, 2009.

10 Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J.: Global peatland dynamics since the last glacial maximum, *Geophys. Res. Lett.*, 37, L13402, doi:10.1029/2010GL043584, 2010.

15 Zar, J. H.: *Biostatistical Analysis*, 5th edn., Pearson Prentice Hall, Upper Saddle River, NJ, 2010.

## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

**Table 1.** Distribution of *Sphagnum* growth measurements in terms of species, peatland type, microform, and continentality.

		Number of samples
Species	<i>Sphagnum fuscum</i>	100
	<i>Sphagnum magellanicum</i>	42
Peatland type	Bog	111
	Poor fen	20
	Rich fen	11
Microform	Hummock	114
	Lawn	26
	Hollow	2
Continentality	Continental	96
	Maritime	46

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







## Bioclimatic drivers of peatland *Sphagnum* growth

J. Loisel et al.

**Table 3.** Results of univariate (a) and multiple (b) regression models of *Sphagnum* growth using all values, continental sites, and maritime sites. df: degree of freedom (the comma separates the number of df for the numerator and denominator);  $p$ : probability;  $PAR_0$  and  $P/E_q$  are defined in the text.

Sites	Predictor	df	adjusted $R^2$	$p$
<b>(a)</b>	$PAR_0$	1, 140	0.23	< 0.0001
	$P/E_q$	1, 140	0.00	0.912
Continental	$PAR_0$	1, 94	0.31	< 0.0001
	$P/E_q$	1, 94	0.06	0.009
Maritime	$PAR_0$	1, 44	0.02	0.172
	$P/E_q$	1, 44	0.00	0.690
<b>(b)</b>	$PAR_0$	2, 93	0.31	< 0.0001
	$P/E_q$	2, 93	0.01	0.832
	$PAR_0 \times P/E_q$	3, 92	0.04	0.444

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 1.** Global distribution of study sites ( $n = 52$ ) with *Sphagnum* growth measurements ( $n = 142$ ). Black circles indicate continental sites and open circles show maritime sites. The dark shaded area represents the northern peatland extent from Yu et al. (2010).

**Bioclimatic drivers of peatland *Sphagnum* growth**

J. Loisel et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Bioclimatic drivers of peatland *Sphagnum* growth


J. Loisel et al.



**Fig. 2.** *Sphagnum* growth values (mean and standard error) for each species **(a)**, continentality class **(b)**, peatland type **(c)**, and microform **(d)**.

**Bioclimatic drivers of  
peatland *Sphagnum*  
growth**

J. Loisel et al.



**Fig. 3.** Photosynthetically active radiation integrated over the growing season (PAR<sub>0</sub>) in relation to *Sphagnum* growth.  $R^2$  values are presented next to the regression lines.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 4.** *Sphagnum* growth sites in climate space. **(a)** Climate space of photosynthetically active radiation integrated over the growing season ( $PAR_0$ ) and the ratio between annual precipitation and the annually integrated equilibrium evapotranspiration ( $P/E_q$ ) of the northern peatland domain (grey dots) based on  $0.5^\circ \times 0.5^\circ$  gridded instrumental climate data (CLIMATE 2.2 dataset) for the period 1960–1990. The northern peatland area is from Yu et al. (2010). Open triangles represent the location (in climate space) of 8 peatland sites where flux tower measurements are available (Lund et al., 2010; Koehler et al., 2011). WP1: Western peatland 1 (Canada); WP2: western peatland 2 (Canada); Kaa: Kaamanen (Finland); Wet: Polwet (Poland); Faj: Fajemyr (Sweden); Mer Bleue (Canada); Sii: Siikaneva (Finland); Glencar (Ireland). **(b)** Box plot of PAR and  $P/E_q$  values of *Sphagnum* growth sites for continental and maritime sites. The boxes indicate the 25th and 75th percentiles, the whiskers above and below the boxes represent the 10th and 90th percentile, and the circles represent the 5th and 95th percentiles.