

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# Inverse method for estimating respiration rates from decay time series

**D. C. Forney<sup>1,2</sup> and D. H. Rothman<sup>1</sup>**

<sup>1</sup>Lorenz Center and Department of Earth, Atmospheric, and Planetary Sciences,  
Massachusetts Institute of Technology, Cambridge MA, 02139, USA

<sup>2</sup>Department of Mechanical Engineering, Massachusetts Institute of Technology, Cambridge  
MA, 02139, USA

Received: 8 March 2012 – Accepted: 8 March 2012 – Published: 26 March 2012

Correspondence to: D. C. Forney (dforney@mit.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**BGD**

9, 3795–3828, 2012

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

Long-term organic matter decomposition experiments typically measure the mass lost from decaying organic matter as a function of time. These experiments can provide information about the dynamics of carbon dioxide input to the atmosphere and controls on natural respiration processes. Decay slows down with time, suggesting that organic matter is composed of components (pools) with varied lability. Yet it is unclear how the appropriate rates, sizes, and number of pools vary with organic matter type, climate, and ecosystem. To better understand these relations, it is necessary to properly extract the decay rates from decomposition data. Here we present a regularized inverse method to identify an optimally-fitting distribution of decay rates associated with a decay time series. We motivate our study by first evaluating a standard, direct inversion of the data. The direct inversion identifies a discrete distribution of decay rates, where mass is concentrated in just a small number of discrete pools. It is consistent with identifying the best fitting “multi-pool” model, without prior assumption of the number of pools. However we find these multi-pool solutions are not robust to noise and are over-parametrized. We therefore introduce a method of regularized inversion, which identifies the solution which best fits the data but not the noise. This method shows that the data are described by a continuous distribution of rates which we find is well approximated by a lognormal distribution, and consistent with the idea that decomposition results from a continuum of processes at different rates. The ubiquity of the lognormal distribution suggest that decay may be simply described by just two parameters; a mean and a variance of log rates. We conclude by describing a procedure that estimates these two lognormal parameters from decay data. Matlab codes for all numerical methods and procedures are provided.

### Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 1 Introduction

Over  $10^7$  different types of organic substances (Wackett, 2006) comprise the roughly 1800 Gt of carbon in soils (Denman et al., 2007) and biomass and 750 Gt of dissolved and particulate organic carbon in the oceans (Benner and Herndl, 2011) and marine sediments (Denman et al., 2007). Diverse decomposer communities (up to 6000 bacterial species per gram soil and up to 11 000 bacterial species per gram marine sediments, Horner-Devine et al., 2004) respire these compounds, converting 75 Gt of terrestrial carbon (Schlesinger and Andrews, 2000) and 50 Gt of oceanic carbon to  $\text{CO}_2$  every year (Denman et al., 2007). During degradation, organic tissues are broken down to particulate or dissolved organic matter (Eijsackers and Zehnder, 1990), which are then processed microbially to carbon dioxide or converted to microbial compounds and by-products (Madigan et al., 2005) that are subsequently transformed and broken down again (Agren and Bosatta, 1998), eventually resulting in a complete conversion of organic carbon to carbon dioxide. Quantitatively estimating the rates of these processes is difficult for many reasons: the variety of components initially found in tissues vary in lability (Tenney and Waksman, 1929; Burdige, 2006; Lützow et al., 2006; Minderman, 1968; Berg and McClaugherty, 2007); as decomposition proceeds, organic molecules interact chemically (Berg and Laskowski, 2006; Lee et al., 2004; Lützow et al., 2006) forming humus and other hard-to-degrade compounds (Berg and McClaugherty, 2007; Paul, 2007; Eijsackers and Zehnder, 1990); microbial processes produce compounds which vary in lability (Andrén and Paustian, 1987; Agren and Bosatta, 1998); and particulate and dissolved carbon bond and sorb to clays and minerals (Oades, 1988; Hedges and Oades, 1997; Mayer, 1994; Vetter et al., 1998; Nieder and Benbi, 2008) forming organo-mineral complexes that also affect decomposability.

This wide range of decomposition processes and mechanisms lead to heterogeneous kinetics, with rates of decay ranging from weeks to thousands of years (Janssen, 1984; Yang and Janssen, 2000; Trumbore, 2000) or greater (Middelburg, 1989). Knowledge of rate heterogeneity is important because it tells us how long carbon resides in

**BGD**

9, 3795–3828, 2012

### Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





---

**Inverse method for  
estimating  
respiration rates**D. C. Forney and  
D. H. Rothman

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

between these parameters are not well understood (Adair et al., 2008; Feng, 2009b). The second class of models are continuous parallel models, also called reactive continuum, or disordered kinetic models. In these models, organic matter is described as a continuum of qualities, and decomposition proceeds by a continuous distribution of exponential decay rates (Boudreau and Ruddick, 1991; Feng, 2009b). These have only more recently been applied (Manzoni et al., 2009; Rothman and Forney, 2007; Feng, 2009a; Forney and Rothman, 2012). The third and most detailed type of models are transformational models, which incorporate transformations to other types of soil organic matter, decomposer biomass, humus, etc. These models can be discrete, consisting of a network of pools exchanging carbon with one another (van Veen and Paul, 1981; Parton et al., 1987, 1993; Eijsackers and Zehnder, 1990; Beare et al., 1992), or continuous (Carpenter, 1981; Bosatta, 1985; Agren and Bosatta, 1998). Some transformational models also couple the rates of organic carbon decomposition with nutrient dynamics (Manzoni and Porporato, 2007; Agren and Bosatta, 1998) and may be nonlinear (Manzoni and Porporato, 2007, 2009). Transformations in organic matter can also be described by a first-order decay constant which decreases according to a function of time (Janssen, 1984; Yang and Janssen, 2000; Bosatta, 1995).

The three types of models are similar. The multi-pool model can be written in terms of the disordered kinetic model when the continuous rate distribution is comprised of delta functions (Forney and Rothman, 2012). The disordered kinetic model can be represented by a transformational model when quality is preserved during transformation (Bosatta, 1995). When linear, transformational network models can sometimes be mathematically represented by a multi-pool model (Bolker et al., 1998). This simplification also occurs when the timescale of transformation is short with respect to the decay timescale of slowly degrading products (Forney and Rothman, 2012).

Using the disordered kinetic approach, we have recently found that plant matter decay is well described by a lognormal distribution of rates (Forney and Rothman, 2012) and that the two parameters of the lognormal distribution are qualitatively related to climatic and compositional variables. A distribution of rates provides a unique way to

visualize and pinpoint key dynamic information regarding carbon in the system and reveal the timescales of the retention of organic carbon in decomposition systems.

In this paper, we elaborate on the formation and implementation of the regularized inversion method that we employed to identify the best fitting rate distribution (Forney and Rothman, 2012). To do so, we first consider a simpler direct inversion of the data which forms the basis for the regularization method. The direct inversion tends to find discrete rate distributions with mass contained at only a few rates. In this sense, the direct inversion is a new procedure for finding the best fitting multi-pool solution without needing to assume how many pools are present. We use the resulting multi-pool solutions to motivate the use of a regularized inverse. We then describe the regularization method in detail and provide an example of its implementation. Because the regularization method identifies a common lognormal pattern (Forney and Rothman, 2012), we also present a simple procedure for estimating the lognormal parameters from the decay data. This procedure provides a more precise estimate of the best fitting lognormal, resulting in slightly better fits to the data than the regularized solution.

The remainder of the paper proceeds as follows: In Sect. 2 we discretize the disordered kinetic model for numerical manipulation. In Sect. 3 we calculate the model inverse. This basic inversion easily and rapidly identifies the appropriate multi-pool solution without prior assumption regarding the number and type of pools. Later in that section we show that the best fitting multi-pool solution is sensitive to noise in the data as both the number and type of best fitting pools fluctuate with the noise levels found in the data. Thus, in Sect. 4 we address the noise sensitivity by employing a method of regularization to invert the data and provide details of its implementation. Finally, in Sect. 5, we analyze litter decay data by assuming rates are distributed lognormally and fit just the two lognormal parameters to the data. Matlab codes for all numerical procedures and approaches are provided online in the Supplement.

**BGD**

9, 3795–3828, 2012

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 2 The parallel model of decay

Because organic matter is composed of many different compounds, some are more resistant to degradation and break down more slowly than others (Tenney and Waksman, 1929; Burdige, 2006; Lützow et al., 2006; Minderman, 1968; Berg and McLaugherty, 2007). Substrate heterogeneity suggests that degradation proceeds at different rates in parallel. This heterogeneity can result in either a discrete or a continuous distribution of decay rates.

We identify the best fitting rate distribution by inverting a continuous parallel model of decay. In this model, the fraction  $g(t)$  of original mass remaining is described by a continuous superposition of exponential decays (Boudreau and Ruddick, 1991),

$$g(t) = \int_0^{\infty} \rho(k) e^{-kt} dk \quad (1)$$

where  $\rho(k)$  is the probability distribution of components with rate  $k$  and  $\int_0^{\infty} \rho(k) dk = 1$ . We have previously found that litter decay rate distributions are well characterized by a lognormal  $\rho(k)$ , although other possible forms have been hypothesized (Boudreau and Ruddick, 1991; Bolker et al., 1998; Rothman and Forney, 2007; Feng, 2009b).

Mathematically, Eq. (1) is the Laplace transform of  $\rho(k)$ . We obtain the distribution  $\rho(k)$  by taking the inverse Laplace transform of the data  $g(t)$  (Forney and Rothman, 2012). To do so, we transform the model to facilitate calculations. We first make a change of variables from  $k$  to  $\ln k$  as we expect to find a wide range of decay rates.

Because probability is conserved,  $\rho(k)dk = \rho(\ln k)d \ln k$ , and Eq. (1) becomes

$$g(t) = \int_{-\infty}^{\infty} \rho(\ln k) e^{-kt} d \ln k \quad (2)$$

where  $\rho(\ln k)$  is the probability distribution in  $\ln k$  space.

**BGD**

9, 3795–3828, 2012

### Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To numerically calculate  $\rho(\ln k)$  we discretize Eq. (2). Although many discretization methods can be used, such as quadrature (Lamanna, 2005; Hansen, 1994), we choose to discretize Eq. (2) at  $n$  nodes  $\lambda_j$  spaced uniformly along the  $\ln k$  axis between the limits  $\ln k_{\min} < \ln k < \ln k_{\max}$ . The nodes are therefore spaced at intervals of width

$$\Delta\lambda = \frac{\ln k_{\max} - \ln k_{\min}}{n}.$$

This discretization is chosen in order to provide resolution over the appropriate wide range of  $k$ .<sup>1</sup> Equation (3) can then be written in matrix form as

$$\mathbf{g} = \mathbf{A}\boldsymbol{\rho}. \quad (3)$$

$\boldsymbol{\rho}$  is a vector of length  $n$ , representing the average value of  $\rho(\ln k)$  over an interval  $\Delta\lambda$ , centered at  $\lambda_j$ ,

$$\rho_j = \frac{1}{\Delta\lambda} \int_{\lambda_j - \frac{1}{2}\Delta\lambda}^{\lambda_j + \frac{1}{2}\Delta\lambda} \rho(\ln k) d \ln k.$$

The  $m \times n$  matrix  $\mathbf{A}$  is the discrete Laplace transform operator with elements

$$A_{ij} = e^{e^{\lambda_j} t_i} \Delta\lambda,$$

and  $\mathbf{g}$  is a vector of  $m$  predicted time series data points  $g_j = g(t_j)$ .

We proceed to identify the underlying rate distribution  $\rho(\ln k)$  from an observed decay time series  $\hat{g}(t)$  by inverting the model (Eq. 3) and solving for  $\boldsymbol{\rho}$ .

<sup>1</sup>While the domain of the inverse Laplace transform is infinite, we find that for litter decay data, the tails of the distributions vanish numerically and that the distribution is typically contained between  $10^{-6}\text{yr}^{-1} < k < 10^4\text{yr}^{-1}$ . We therefore set the values  $k_{\max} = 10^4\text{yr}^{-1}$  and  $k_{\min} = 10^{-6}\text{yr}^{-1}$ . We choose the number of discretization steps,  $n$ , to provide a reasonable resolution of  $\rho(\ln k)$  without being computationally burdensome.  $n \approx 100$  or  $n = 16m$  was satisfactory.

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





### 3 Direct inversion with constraints

Our approach in this section directly calculates the inverse Laplace transform of data with non-negative constraints. This approach identifies a handful of separate and distinct pools which are present during decay, having values  $\rho_j > 0$ ; the remaining pools are not present and have values  $\rho_j = 0$ . This technique therefore provides a direct estimate of the best fitting multi-pool model, expressed as a discrete distribution  $\rho(\ln k)$ . We find however that the multi-pool solution is very sensitive to noise in the data and is overparametrized. We therefore proceed in Sect. 4 to refine this approach using a regularization technique to invert the data (Forney and Rothman, 2012), providing a continuous rate distribution  $\rho(\ln k)$  which is both simpler and less sensitive to noise.

#### 3.1 Calculating the constrained direct inverse

The distribution  $\rho$  can be directly calculated by computing the inverse of (Eq. 3) from the measured decay data  $\hat{\mathbf{g}}$  which includes the first data point  $g_1 = 1$  at  $t_1 = 0$ ,

$$\rho = \mathbf{A}^{-1} \hat{\mathbf{g}} \quad (4)$$

Because solutions to (Eq. 4) fit the noisy data exactly, negative values of  $\rho_j$  are possible. In order to find solutions with non-negative  $\rho_j$ , we instead solve the constrained least squares problem

$$\min_{\rho} \|\mathbf{A}\rho - \hat{\mathbf{g}}\|^2 \quad (5)$$

such that

$$\rho_j > 0 \quad (6)$$

$$\sum_{j=1}^n A_{1j} \rho_j = 1, \quad (7)$$

**BGD**

9, 3795–3828, 2012

### Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







### 3.2 Analysis of the multi-pool solutions

We use the same inversion technique to find the best multi-pool solution associated with 191 datasets chosen (Forney and Rothman, 2012) from the LIDET study. We then determine whether pools are active, inert, or rapid as described above. If any pool contains a mass fraction of less than 1 %, we neglect that pool and distribute its mass proportionately amongst the remaining pools, as these small pools require additional model parametrization but have a negligible effect on the residual error.<sup>2</sup> Table 1 shows the results of the inversion and indicates that eight different types of multi-pool models are needed to describe all 191 of these LIDET data sets. 17 % of the datasets are described by one pool, 56 % are described by two pools, and 27 % are described by three or more pools. The maximum number of pools found was four and all datasets are characterized by 5 or less parameters. The data from Fig. 1a is of type 4 in Table 1; 24 other LIDET datasets also exhibit the same behavior. This information can be used to determine some patterns underlying litter decomposition. For example, we find that 29 % of all datasets involving either needles or wood appear to decay exponentially while roots on the other hand decay exponentially only for 2 % of their datasets. Roots tend to have 3 or 4 pools for 48 % of their datasets, while leaves, needles, and wood each have 3 or 4 pools less than 25 % of the time. However, extracting further information from the inversion results in order to determine which model and parameters are appropriate for different combinations of tissue types, substrates, or environments is difficult if not impossible since there is too much variation in  $\rho$  and model type. Indeed this is one of the shortcomings of the multi-pool model. It is too overparametrized to be useful in identifying fundamental constitutive relations between composition and dynamics.

<sup>2</sup>This occurred for only 7/191 datasets, 5 of which were two pool models that became one pool models.

**BGD**

9, 3795–3828, 2012

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.3 Sensitivity to noise

In this section we test the direct inversion method and evaluate the appropriateness of multi-pool models for describing noisy decay data. The standard shortcoming of a multi-pool solution is its inherent sensitivity to noise in the data  $\hat{\mathbf{g}}$  (Kroeker and Henkelman, 1986; Kleinberg, 1996), as models with differing numbers of pools can well approximate one another to a high degree of accuracy (Yeramian and Claverie, 1987). We proceed to investigate the solution's sensitivity to noise levels present in litter decay data.

If we assume that the multi-pool model is correct, then the line in Fig. 1a is the true decay  $g(t)$ . We then estimate the variance of the data noise  $\sigma_e^2$  by calculating its maximum likelihood  $\sigma_e^2 = \sum_i \hat{g}_i^2 / m$  (Konishi and Kitagawa, 2010), assuming the noise is Gaussian with zero mean. We then generate numerical trials of decay data by adding Gaussian noise of the same strength, in this case  $\sigma_e = 0.024$ , to the predicted decay  $g(t)$ . Then, for each trial we recalculate the associated inverse solution  $\rho$ . The results of the inversion of 60 000 numerical trials is shown in Fig. 2a–g. There is wide variation in the active pools just due to the noise in the data. Figure 2b shows how the model type varies due to noise. The “true” model, type 4, is only identified 39% of the time. The other LIDET datasets show similar results. These results highlight why it is difficult to gain fundamental understanding of decomposition processes from fitting a multi-pool model to the data.

### 4 Regularized inversion method

Because the inverse Laplace transform Eq. (4) is sensitive to noise in the data, it is ill-posed (Hansen, 1987, 1994). In the previous section, we have demonstrated that the inverse Laplace transform with non-negative constraints is also ill-posed (Yeramian and Claverie, 1987) at the level of noise present in our decay functions  $g(t)$ . There are a number of approaches to solve the inverse Laplace transform robustly, without being sensitive to small changes in the data. These include Laplace-Padé

**BGD**

9, 3795–3828, 2012

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





roughness,  $\omega$  is typically varied many orders of magnitude from  $\omega \ll 1$ , which emphasizes the importance of residual error, to  $\omega \gg 1$ , which emphasizes the smoothest solution. The best  $\rho(\ln k)$  is determined by identifying the value of  $\omega$  that sets an optimal balance between the residual error and the roughness.

#### 4.1 The L-curve

An approach for finding the optimal  $\omega$  is to use the “L-curve” technique (Hansen, 1987, 1994). An L-curve is generated by parametrically varying  $\omega$  and solving Eq. (9) for  $\rho$ , obtaining values for the residual error norm  $\|\hat{\mathbf{g}} - \mathbf{A}\rho\|$  and roughness norm  $\|\mathbf{R}\rho\|$  for each value of  $\omega$ . The L-curve is made by plotting  $\|\mathbf{R}\rho\|$  vs.  $\|\hat{\mathbf{g}} - \mathbf{A}\rho\|$  on a log-scale, resulting in a characteristic “L” shape. Figure 3a shows the L-curve for the dataset shown in Fig. 1a.

While each point on the curve is associated with an optimal solution for a certain value of  $\omega$ , the corner of the L-curve is associated with the regularized solution. The corner represents the point where increasing the information contained in the solution no longer improves the residual error. Solutions at the upper right of the L-curve contain high information as  $n$  elements  $\rho_j$  are free and independent (Pierce, 1980) whereas fewer modes are active in the smooth solution at the corner. The corner is also the point where relatively little decrease in information results in large increases in the residual error. Often, the corner is associated with a change in the number of maxima in  $\rho(\ln k)$ ; solutions above the corner may have more modes than solutions below.

Generally, the L-curve method identifies a smooth solution which predicts a decay having a residual error approximately equal to the noise in the dataset. Solutions for three different values of  $\omega$  are shown in Fig. 3b–d with the unregularized solution shown in Fig. 3b. Decays predicted from these three solutions are shown in Fig. 3e.

When there are multiple corners, as shown here, there are a few approaches to choosing the appropriate one. We use the following method to identify the corner associated with the simplest solution:

**BGD**

9, 3795–3828, 2012

### Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1. For each value of  $\omega$ , calculate the rank correlation (Kendall and Gibbons, 1990) of the residual error and the corresponding  $P$  value.
2. Identify the region of the L-curve where the rank correlation of solutions have  $P > 0.1$ , suggesting no significant trend in the residual.
3. Choose the corner within the region of the L-curve having  $P > 0.1$ ;
4. If a corner seems too rounded or undefined, the L-curve can also be plotted on linear  $\|\mathbf{A}\boldsymbol{\rho} - \hat{\mathbf{g}}\|$  and  $\|\mathbf{R}\boldsymbol{\rho}\|$  axes. The corner collapses onto a much tighter volume in linear space and can sometimes be more readily chosen if the corner in log space has large curvature.
5. If there are multiple corners with  $P > 0.1$ , compare more closely the residual trend in all corners, as values  $P > 0.1$  do not necessarily indicate a lack of trend in the residual. If there is a noticeable difference in the quality of the fit associated with each corner, choose the corner that does not appear to have a trend in the residual. An additional test is to look at the error bias. Here, the error bias is simply the sum of residual error divided by the error norm,  $\frac{|\sum(\mathbf{A}\boldsymbol{\rho} - \hat{\mathbf{g}})|}{\|\mathbf{A}\boldsymbol{\rho} - \hat{\mathbf{g}}\|}$ . Experience with this data suggests choosing solutions with error bias  $< 0.25$ . If multiple corners still have a bias  $< .25$  and a similar strength in trend in their residual error, choose the corner with lowest roughness. Usually the corner having lower residual error and higher roughness tends to be not biased and to have no trend in the residual.

We apply this approach to choose the appropriate corner in Fig. 3a. The distribution corresponding to the lower corner (Fig. 3d) predicts a decay that has a large error compared to the noise in the data and has a significant trend in the residual error as seen in Fig. 3e. However, the distribution corresponding to the higher corner (Fig. 3c) predicts a decay which is hardly distinguishable from the decay predicted by the unregularized solution, shown in green, after  $t = 1$ . Therefore, the distribution shown in Fig. 3c is considered the optimal, regularized solution.



## 4.2 Results of regularization

Figure 4 shows the overall variation in the inversions due to noise by summing all 60 000 multi-pool solutions from Fig. 2a–g. We also plot the solution  $\rho(\ln k)$  from the regularized inversion. The regularization provides a simpler solution that captures the variation of the rate heterogeneity associated with uncertainty of the multi-pool solutions.

More specifically, we find that this heterogeneity takes a certain form: the regularized solution presented in Fig. 3c appears lognormal, i.e. Gaussian in  $\ln k$  space. We have shown in a separate publication (Forney and Rothman, 2012) that the entire LIDET dataset is lognormal on average. This lognormal description is considerably simpler than the multi-pool approach as the rate heterogeneity and all dynamic information are now contained within just two parameters, a mean  $\mu$  and a variance  $\sigma^2$  of the order of magnitude of decay rates. This approach also emphasizes the view that the quality of litter and young soil organic matter quality is continuously distributed.

## 5 Fitting lognormal parameters to decay data

The ubiquity of the the lognormal distribution suggests that litter decay can be well described by the two lognormal parameters,  $\mu$  and  $\sigma$ . In this section, we assume a model of lognormally distributed rates and present a procedure to estimate the values of  $\mu$  and  $\sigma$  that best fit the decay data. We then compare its results to the results of the regularized inversion.

We proceed by substituting the Gaussian distribution,

$$\rho(\ln k; \mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(\ln k - \mu)^2 / 2\sigma^2}, \quad (10)$$

**BGD**

9, 3795–3828, 2012

### Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for  $\rho(\ln k)$  in Eq. (2), providing a prediction of the mass fraction  $g(t)$  remaining as a function of the parameters  $\mu$  and  $\sigma$ :

$$g(t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma} e^{-(\ln k - \mu)^2 / 2\sigma^2} e^{-kt} d \ln k. \quad (11)$$

We then identify the values of  $\mu$  and  $\sigma$  which best fit the data by solving the non-linear least squares minimization problem

$$\min_{\mu, \sigma} \sum_{i=1}^n \left( \hat{g}_i - \int_{-\infty}^{\infty} \rho(\ln k; \mu, \sigma) e^{-kt} d \ln k \right)^2, \quad (12)$$

where  $\hat{g}_i$  are the measured data points. We solve Eq. (12) using Matlab's non-linear least squares solver `nlinfit.m`. This is repeated for all 191 LIDET datasets that are appropriately described by a superposition of exponential decays (Forney and Rothman, 2012).

Figure 5 shows a comparison of the residual error associated with the three different approaches: direct inversion (Eqs. 5–7), regularized inversion Eq. (9), and the two-parameter fit Eq. (12). The root mean square error (RMSE) of the fit to the data is calculated for each dataset and the cumulative density function of the RMSE rescaled by the total number of datasets (191) is plotted; the vertical axis shows the number of datasets having an RMSE smaller than the value on the horizontal-axis. The RMSE of the unregularized inversion technique (green) is clearly smaller than that of the other two techniques, but these solutions suffer from the problems discussed in Sect. 3.3. Mean RMSE values for all 191 datasets are 0.048, 0.053, and 0.052 for the direct inversion, regularized inversion, and lognormal model, respectively. Surprisingly, solutions from the two-parameter fit (magenta) appear to have slightly smaller residual error than solutions identified from the regularization technique (blue). Thirty-two of the datasets are predicted by both the two-parameter fit and regularization method

**Inverse method for  
estimating  
respiration rates**

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to decay exponentially; the two-parameter lognormal model however predicts the decay from 126 of the remaining 159 datasets better than the distribution obtained by regularization. Therefore, the lognormal model fits the data equally or better than the regularized solution for 158/191 or 83% of the datasets. This is unexpected because the lognormal model has only two degrees of freedom. The marginally poorer fit of the regularization method is due to the emphasis of the method on smoother (wider) solutions. We proceed to investigate this effect by comparing the fitted parameters  $\sigma$  to the standard deviation of the regularized inversions  $\rho(\ln k)$ .

Figure 6 compares the fitted values  $\mu$  and  $\sigma$  to the mean  $\langle \ln k \rangle$  and standard deviation  $\sqrt{\langle (\ln k - \langle \ln k \rangle)^2 \rangle}$  associated with the regularized solutions  $\rho$ . Figure 6a shows that values of  $\mu$  and the mean of the inversion  $\rho(\ln k)$  are consistent with one another. There is a departure at high  $\langle \ln k \rangle$  because some of these inversions are bi-modal: one mode is active whereas the other mode is distributed over extremely fast rates associated with instantaneous decays. These bi-modal datasets are effectively described by an instantaneous mass loss followed by decay that proceeds with a distribution of rates. Figure 6b, on the other hand, tells another story. Values of  $\sigma$  tend to be less than the standard deviation of  $\rho(\ln k)$ , indicating that the regularized solution is wider than the best fitting lognormal distribution. This is a consequence of weighing the solution roughness during regularization. A narrower lognormal solution with smaller residual error exists, but the weight of the roughness is large enough that the regularization method chooses a slightly wider solution which fits the data almost as well. Collectively, Figs. 5 and 6b indicate that when rates are heterogeneous, the transition from unimodal to multimodal solutions near the corner of the L-curve occurs before finding the unimodal solution with the smallest residual error.

These results indicate that the regularization method is useful for identifying that general solution shapes and trends. If regularization suggests that rate distributions are lognormal, then fitting  $\mu$  and  $\sigma$  to the data identifies more precisely the specific lognormal distribution that best fits the data.

**BGD**

9, 3795–3828, 2012

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 6 Conclusions

Direct calculation of the inverse Laplace transform with a non-negativity constraint provides the best fitting multi-pool solution without specification of the number of pools a priori. However, this multi pool solution is very sensitive to small changes in the decay function  $g(t)$  (Yeramian and Claverie, 1987) as the number of pools and the rates of each pool vary widely due to the level of noise associated with litter bag data. The regularization method described here is robust to noise and commonly indicates that a lognormal distribution provides a concise representation of the rates associated with decay data. If the lognormal distribution may be assumed, a marginally better fit to the data is found by directly estimating the lognormal parameters  $\mu$  and  $\sigma$  from the decay  $g(t)$ .

Matlab codes for all numerical procedures (direct inversion, multi-pool estimation, regularized inversion, and the two-parameter fitting procedure) are provided online in the Supplement.

**Supplementary material related to this article is available online at:**

**<http://www.biogeosciences-discuss.net/9/3795/2012/bgd-9-3795-2012-supplement.zip>**

*Acknowledgements.* We thank C. Follett, Y. Friedman, A. Petroff, O. Devauchelle, and D.M. Abrams for insightful discussions. This work was supported by NSF Grant EAR-0420592 and NASA Grant NNA08CN84A. DHR thanks the Radcliffe Institute for Advanced Study for providing a one-year fellowship during which a portion of this work was performed.

## References

Adair, E. C., Parton, W. J., Del Grosso, S. J., Silver, W. L., Harmon, M. E., Hall, S. A., Burke, I. C., and Hart, S. C.: Simple three-pool model accurately describes patterns of long-term litter

**BGD**

9, 3795–3828, 2012

### Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

decomposition in diverse climates, *Glob. Change Biol.*, 14, 2636–2660, doi:10.1111/j.1365-2486.2008.01674.x, 2008. 3798, 3799, 3804

Agren, G. I. and Bosatta, E.: *Theoretical Ecosystem Ecology: Understanding Element Cycles*, 1st edn., Cambridge University Press, available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/0521646510>, 1998. 3797, 3799

Andr n, O. and Paustian, K.: Barley straw decomposition in the field: a comparison of models, *Ecology*, 68, 1190–1200, doi:10.2307/1939203, 1987. 3797, 3798

Beare, M. H., Parmelee, R. W., Hendrix, P. F., Cheng, W., Coleman, D. C., and Crossley Jr., D. A.: Microbial and faunal interactions and effects on litter nitrogen and decomposition in agroecosystems, *Ecol. Monogr.*, 62, 569–591, doi:10.2307/2937317, 1992. 3799

Benner, R. and Herndl, G. J.: *Bacterially Derived Dissolved Organic Matter in the Microbial Carbon Pump*, available at: <http://www.sciencemag.org/site/products/microbialpump>, 2011. 3797

Berg, B. and Laskowski, R.: *Litter Decomposition: A Guide to Carbon and Nutrient Turnover*, vol. 38, Elsevier, Amsterdam, London, 2006. 3797

Berg, B. and McClaugherty, C.: *Plant Litter: Decomposition, Humus Formation, Carbon Sequestration*, 2nd edn., Springer, available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/3540749225>, 2007. 3797, 3801

Berner, R. A.: *Early Diagenesis*, Princeton University Press, available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/069108260X>, 1980. 3798

Bolin, B. and Rodhe, H.: A note on the concepts of age distribution and transit time in natural reservoirs, *Tellus*, 25, 58–62, 1973. 3798

Bolker, B. M., Pacala, S. W., and Parton, W. J.: Linear analysis of soil decomposition: insights from the century model, *Ecol. Appl.*, 8, 425–439, doi:10.2307/2641082, 1998. 3799, 3801

Bosatta, E.: Theoretical analysis of decomposition of heterogeneous substrates, *Soil Biol. Biochem.*, 17, 601–610, doi:10.1016/0038-0717(85)90035-5, 1985. 3799

Bosatta, E.: The power and reactive continuum models as particular cases of the q-theory of organic matter dynamics, *Geochimica et Cosmochimica Acta*, 59, 3833–3835, doi:10.1016/0016-7037(95)00287-A, 1995. 3799

Boudreau, B. P. and Ruddick, B. R.: On a reactive continuum representation of organic matter diagenesis, *Am. J. Sci.*, 291, 507–538, 1991. 3799, 3801

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Burdige, D. J.: Geochemistry of Marine Sediments, Princeton University Press, available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/069109506X>, 2006. 3797, 3801

Carpenter, S.: Decay of heterogenous detritus: a general model, *J. Theor. Biol.*, 89, 539–547, doi:10.1016/0022-5193(81)90026-6, 1981. 3799

Cox, P. M.: Description of the TRIFFID dynamics global vegetation model, Tech. Note 24, Tech. rep., Hadley Centre, Met Office, available at: <http://www.metoffice.com/publications/HCTN/>, 2001. 3798

Currie, W. S., Harmon, M. E., Burke, I. C., Hart, S. C., Parton, W. J., and Silver, W.: Cross-biome transplants of plant litter show decomposition models extend to a broader climatic range but lose predictability at the decadal time scale, *Glob. Change Biol.*, 16, 1744–1761, doi:10.1111/j.1365-2486.2009.02086.x, 2010. 3798, 3804, 3805

Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., da Silva Dias, P. L., Wofsy, S. C., and Zhang, X.: Couplings between changes in the climate system and biogeochemistry, in: *Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, and Miller, H. L., chap. 7, 499–587, Cambridge University Press, Cambridge UK, New York, available at: [http://www.ipcc.ch/publications\\_and\\_data/publications\\_ipcc\\_fourth\\_assessment\\_report\\_wg1\\_report\\_the\\_physical\\_science\\_basis.htm](http://www.ipcc.ch/publications_and_data/publications_ipcc_fourth_assessment_report_wg1_report_the_physical_science_basis.htm), 2007. 3797

Eijsackers, H. and Zehnder, A.: Litter decomposition: a Russian matryoshka doll, *Biogeochemistry*, 11, 153–174, doi:10.1007/BF00004495, 1990. 3797, 3799

Feng, Y.: K-model – a continuous model of soil organic carbon dynamics: model parameterization and testing, *Soil Sci.*, 174, 494–507, 2009a. 3799

Feng, Y.: K-model – a continuous model of soil organic carbon dynamics: theory, *Soil Sci.*, 174, 482–493, 2009b. 3798, 3799, 3801, 3805

Forney, D. C. and Rothman, D. H.: Common structure in the heterogeneity of plant matter decay, *J. Roy. Soc. Interface*, in press, 2012. 3798, 3799, 3800, 3801, 3803, 3805, 3806, 3811, 3812

Gallegos, D. P. and Smith, D. M.: A NMR technique for the analysis of pore structure: determination of continuous pore size distributions, *J. Coll. Interface Sci.*, 122, 143–153, doi:10.1016/0021-9797(88)90297-4, 1988. 3808

---

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Gholz, H. L., Wedin, D. A., Smitherman, S. M., Harmon, M. E., and Parton, W. J.: Long-term dynamics of pine and hardwood litter in contrasting environments: toward a global model of decomposition, *Glob. Change Biol.*, 6, 751–765, doi:10.1046/j.1365-2486.2000.00349.x, 2000. 3804
- 5 Hansen, P. C.: Rank-Deficient and Discrete Ill-Posed Problems: Numerical Aspects of Linear Inversion (Monographs on Mathematical Modeling and Computation), Society for Industrial Mathematics, available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/0898714036>, 1987. 3807, 3808, 3809, 3825
- Hansen, P. C.: Regularization tools: A Matlab package for analysis and solution of discrete ill-  
10 posed problems, *Numer. Algor.*, 6, 1–35, doi:10.1007/BF02149761, 1994. 3802, 3807, 3808, 3809, 3825
- Harmon, M.: LTER Intersite Fine Litter Decomposition Experiment (LIDET), Forest Science Data Bank code TD023, Corvallis, OR, Database, available at: <http://andrewsforest.oregonstate.edu/data/abstract.cfm?dbcode=TD023>, 2007. 3804
- 15 Harmon, M. E., Silver, W. L., Fasth, B., Chen, H., Burke, I. C., Parton, W. J., Hart, S. C., and Currie, W. S.: Long-term patterns of mass loss during the decomposition of leaf and fine root litter: an intersite comparison, *Glob. Change Biol.*, 15, 1320–1338, doi:10.1111/j.1365-2486.2008.01837.x, 2009. 3798, 3804, 3805
- Hedges, J. I. and Oades, J. M.: Comparative organic geochemistries of soils and marine sediments, *Org. Geochem.*, 27, 319–361, 1997. 3797
- 20 Horner-Devine, M. C., Carney, K. M., and Bohannan, B. J. M.: An ecological perspective on bacterial biodiversity, *Proc. R. Soc. Lond. B*, 271, 113–122, 2004. 3797
- Istratov, A. A. and Vyvenko, O. F.: Exponential analysis in physical phenomena, *Rev. Sci. Instrum.*, 70, 1233–1257, doi:10.1063/1.1149581, 1999. 3808
- 25 Janssen, B. H.: A simple method for calculating decomposition and accumulation of “young” soil organic matter, *Plant Soil*, 76, 297–304, doi:10.1007/BF02205588, 1984. 3797, 3799
- Jenkinson, D. S.: Studies on the decomposition of plant material in soil. V. The effect of plant cover and soil type on the loss of carbon from <sup>14</sup>C labeled ryegrass decomposing under field conditions, *J. Soil Sci.*, 28, 424–434, doi:10.1111/j.1365-2389.1977.tb02250.x, 1977. 3798
- 30 Jenkinson, D. S., Andrew, S. P. S., Lynch, J. M., Goss, M. J., and Tinker, P. B.: The turnover of organic carbon and nitrogen in soil [and discussion], *Philos. Trans. Biol. Sci.*, 329, 361–368, doi:10.1098/rstb.1990.0177, 1990. 3798



## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Kendall, M. and Gibbons, J. D.: Rank Correlation Methods, Oxford University Press, New York, 5th edn., available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/0195208374>, 1990. 3810
- van Keulen, H.: (Tropical) soil organic matter modelling: problems and prospects, *Nutr. Cyc. Agroecosyst.*, 61, 33–39, doi:10.1023/A:1013372318868, 2001. 3798
- Kleinberg, R.: Utility of NMR T2 distributions, connection with capillary pressure, clay effect, and determination of the surface relaxivity parameter  $\rho_2$ , *Mag. Resonance Imag.*, 14, 761–767, doi:10.1016/S0730-725X(96)00161-0, 1996. 3807, 3808
- Konishi, S. and Kitagawa, G.: Information Criteria and Statistical Modeling (Springer Series in Statistics), 1st edn., Springer, Softcover Reprint of Hardcover, 2008 edn., available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/1441924566>, 2010. 3807
- Krinner, G.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, *Global Biogeochem. Cy.*, 19, GB1015, doi:10.1029/2003GB002199, 2005. 3798
- Kroeker, R. M. and Henkelman, R. M.: Analysis of biological NMR relaxation data with continuous distributions of relaxation times, *J. Mag. Resonance* (1969), 69, 218–235, doi:10.1016/0022-2364(86)90074-0, 1986. 3807, 3808
- Lamanna, R.: On the inversion of multicomponent NMR relaxation and diffusion decays in heterogeneous systems, *Concepts Magn. Reson.*, 26A, 78–90, doi:10.1002/cmra.20036, 2005. 3802, 3808
- Lee, C., Arnosti, C., and Wakeham, S.: Particulate organic matter in the sea: the composition conundrum, *Ambio*, 33, 565–575, 2004. 3797
- Lützow, M., Kogel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., and Flessa, H.: Stabilization of organic matter in temperate soils: mechanisms and their relevance under different soil conditions – a review, *Eur. J. Soil Sci.*, 57, 426–445, doi:10.1111/j.1365-2389.2006.00809.x, 2006. 3797, 3801
- Madigan, M. T., Martinko, J. M., Dunlap, P. V., and Clark, D. P.: Brock Biology of Microorganisms, 11th edn., Benjamin Cummings, available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/0132192268>, 2005. 3797
- Manzoni, S. and Porporato, A.: A theoretical analysis of nonlinearities and feedbacks in soil carbon and nitrogen cycles, *Soil Biology and Biochemistry*, 39, 1542–1556, doi:10.1016/j.soilbio.2007.01.006, 2007. 3799



## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Manzoni, S. and Porporato, A.: Soil carbon and nitrogen mineralization: Theory and models across scales, *Soil Biology and Biochemistry*, 41, 1355–1379, doi:10.1016/j.soilbio.2009.02.031, 2009. 3798, 3799
- Manzoni, S., Katul, G. G., and Porporato, A.: Analysis of soil carbon transit times and age distributions using network theories, *J. Geophys. Res.*, 114, G04025, doi:10.1029/2009JG001070, 2009. 3798, 3799
- Mathworks: mldivide.m, Matlab Help, available at: <http://www.mathworks.com/help/releases/R2009a/techdoc/ref/mldivide.html>, 2009. 3804
- Mathworks: Matlab Help: pinv.m, available at: <http://www.mathworks.com/help/releases/R2011a/techdoc/ref/pinv.html>, 2011. 3804
- Mayer, L.: Relationships between mineral surfaces and organic carbon concentrations in soils and sediments, *Chem. Geol.*, 114, 347–363, doi:10.1016/0009-2541(94)90063-9, 1994. 3797
- Medvigy, D., Wofsy, S. C., Munger, J. W., Hollinger, D. Y., and Moorcroft, P. R.: Mechanistic scaling of ecosystem function and dynamics in space and time: ecosystem demography model version 2, *J. Geophys. Res.*, 114, G01002, doi:10.1029/2008JG000812, 2009. 3798
- Middelburg, J. J.: A simple rate model for organic matter decomposition in marine sediments, *Geochim. Cosmochim. Acta*, 53, 1577–1581, doi:10.1016/0016-7037(89)90239-1, 1989. 3797
- Minderman, G.: Addition, decomposition and accumulation of organic matter in forests, *J. Ecol.*, 56, 355–362, doi:10.2307/2258238, 1968. 3797, 3798, 3801
- Moorcroft, P. R., Hurtt, G. C., and Pacala, S. W.: A method for scaling vegetation dynamics: the ecosystem demography model (ED), *Ecol. Monogr.*, 71, 557–585, doi:10.2307/3100036, 2001. 3798
- Nieder, R. and Benbi, D. K.: *Carbon and Nitrogen in the Terrestrial Environment*, 1st edn., Springer, available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/1402084323>, 2008. 3797, 3798, 3805
- Oades, J.: The retention of organic matter in soils, *Biogeochemistry*, 5, 35–70, doi:10.1007/BF02180317, 1988. 3797
- Olson, J. S.: Energy storage and the balance of producers and decomposers in ecological systems, *Ecology*, 44, 322–331, doi:10.2307/1932179, 1963. 3798

- Parton, W. J., Schimel, D. S., Cole, C. V., and Ojima, D. S.: Analysis of factors controlling soil organic matter levels in great plains grasslands, *Soil Sci. Soc. Am. J.*, 51, 1173–1179, 1987. 3799
- Parton, W. J., Scurlock, J. M. O., Ojima, D. S., Gilmanov, T. G., Scholes, R. J., Schimel, D. S.,  
Kirchner, T., Menaut, J. C., Seastedt, T., Moya, G. E., Kamnalrut, A., and Kinyamario, J. I.:  
Observations and modeling of biomass and soil organic matter dynamics for the grassland  
biome worldwide, *Global Biogeochem. Cy.*, 7, 785–809, 1993. 3799
- Paul, E. A.: *Soil Microbiology, Ecology and Biochemistry*, 3rd edn., Academic Press,  
available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/0125468075>, 2007. 3797
- Pierce, J. R.: *An Introduction to Information Theory: Symbols, Signals and Noise*, Dover Publications, 2nd revised edn., available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/0486240614>, 1980. 3809
- Press, W. H., Flannery, B. P., Teukolsky, S. A., and Vetterling, W. T.: *Numerical Recipes in C: The Art of Scientific Computing*, 2nd edn., Cambridge University Press,  
available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/0521431085>, 1992. 3808
- Rhode, H.: *Modeling Biogeochemical Cycles*, Academic Press, San Diego, 55–72, 1992. 3798
- Rothman, D. H. and Forney, D. C.: Physical model for the decay and preservation of marine organic carbon, *Science*, 316, 1325–1328, doi:10.1126/science.1138211, 2007. 3799, 3801
- Schlesinger, W. H. and Andrews, J. A.: Soil respiration and the global carbon cycle, *Biogeochemistry*, 48, 7–20, doi:10.1023/A:1006247623877, 2000. 3797
- Sitch, S., Smith, B., Prentice, I. C., Arneeth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model, *Glob. Change Biol.*, 9, 161–185, doi:10.1046/j.1365-2486.2003.00569.x, 2003. 3798
- Strang, G.: *Introduction to Applied Mathematics*, 1st edn., Wellesley-Cambridge Press,  
available at: <http://www.amazon.com/exec/obidos/redirect?tag=citeulike07-20&path=ASIN/0961408804>, 1986. 3804
- Tenney, F. G. and Waksman, S. A.: Composition of natural organic materials and their decomposition in the soil: IV. The nature and rapidity of decomposition of the various organic complexes in different plant materials, under aerobic conditions, *Soil Sci.*, 28, 55–84, 1929. 3797, 3801

---

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Trumbore, S.: Age of soil organic matter and soil respiration: radiocarbon constraints on below-ground C dynamics, *Ecol. Appl.*, 10, 399–411, 2000. 3797
- van Veen, J. A. and Paul, E. A.: Organic carbon dynamics in grassland soils. 1. Background information and computer simulation, *Can. J. Soil Sci.*, 61, 185–201, doi:10.4141/cjss81-024, 1981. 3799
- Vetter, Y. A., Deming, J. W., Jumars, P. A., and Krieger-Brockett, B. B.: A predictive model of bacterial foraging by means of freely released extracellular enzymes, *Microb. Ecol.*, 36, 75–92, doi:10.1007/s002489900095, 1998. 3797
- Wackett, L.: The metabolic pathways of biodegradation the prokaryotes, in: *The Prokaryotes*, edited by: Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.-H., and Stackebrandt, E., chap. 29, 956–968, Springer, New York, available at: [http://dx.doi.org/10.1007/0-387-30742-7\\_29](http://dx.doi.org/10.1007/0-387-30742-7_29), 2006. 3797
- Yang, H. S. and Janssen, B. H.: A mono-component model of carbon mineralization with a dynamic rate constant, *Eur. J. Soil Sci.*, 51, 517–529, doi:10.1046/j.1365-2389.2000.00319.x, 2000. 3797, 3799
- Yeremian, E. and Claverie, P.: Analysis of multiexponential functions without a hypothesis as to the number of components, *Nature*, 326, 169–174, doi:10.1038/326169a0, 1987. 3807, 3808, 3814
- Zhou, Y. and Zhuang, X.: Robust reconstruction of the rate constant distribution using the phase function method, *Biophys. J.*, 91, 4045–4053, doi:10.1529/biophysj.106.090688, 2006. 3808

Inverse method for estimating respiration rates

D. C. Forney and D. H. Rothman

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion


**Table 1.** This table shows the wide and seemingly unpredictable variation in the types of pool models that best fit 191 LIDET datasets. Each model was obtained by solving the non-negative least squares problem (5) and analyzing the resulting  $\rho$  to determine whether the pools were active (decaying exponentially), inert, or rapid, as discussed in the text. The first column lists the model type, the second column indicates kinds of pools present in that model, the third column provides the formula for the model, the fourth column gives the total number of pools in the model, the fifth column indicates the total number of parameters in the model, and the sixth column indicates how many LIDET datasets are best fit by that model. The sizes of the constant, rapid, and active pools in the models are  $c$ ,  $r$ , and  $a$ , respectively. The size of one of the pools is determined by the others since mass fractions must sum to one.

Model type	Type of pools	Formula	Pools	Para- meters	# Data- sets
1	active	$e^{-kt}$	1	1	33
2	active, inert	$(1 - c)e^{-kt} + c$	2	2	30
3	active, rapid	$(1 - r)e^{-kt}$	2	2	37
4	active, inert, rapid	$(1 - r - c)e^{-kt} + c$	3	3	25
5	2 active	$a e^{-k_1t} + (1 - a)e^{-k_2t}$	2	3	39
6	2 active, inert	$a e^{-k_1t} + (1 - a - c)e^{-k_2t} + c$	3	4	14
7	2 active, rapid	$a e^{-k_1t} + (1 - a - r)e^{-k_2t}$	3	4	10
8	2 active, inert, rapid	$a e^{-k_1t} + (1 - a - r - c)e^{-k_2t} + c$	4	5	3



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper





**Fig. 2.** Distributions  $\rho(\ln k)$  for 60 000 trials of adding random noise to the predicted decay in Fig. 1a. For each trial,  $\rho(\ln k)$  is estimated using the method discussed in Sect. 3.1. The pools in each solution are then analyzed to determine the model type as described in the text and Table 1, except for model type 9 which is a three active-pool model and type 10 which is any combination of pools not described by models 1–9. **(A–G)** shows the mean  $\langle \rho(k) \rangle$  for each model type, calculated by summing all of the observed distributions for each type and dividing by the total number of times each type was observed. **(H)** Fraction of trials that each model type was identified.

## Inverse method for estimating respiration rates

D. C. Forney and  
D. H. Rothman

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Inverse method for  
estimating  
respiration ratesD. C. Forney and  
D. H. Rothman

**Fig. 3.** Determining the optimal value of regularization parameter  $\omega$  via the “L-Curve” method (Hansen, 1994, 1987). Counterclockwise from the upper left: **(A)** The “L-curve”,  $\|R\hat{\rho}\|$  vs.  $\|A\hat{\rho} - g\|$ . **(B)** The unregularized solution, similar to the solution found at the top of the curve, where the regularization parameter  $\omega \ll 1$ . **(C)** The solution associated with the corner of the curve corresponding to  $\omega = 0.66$ . **(D)** The solution at the corner of the curve corresponding to  $\omega = 20.1$ . **(E)** The data  $g(t)$  as well as the decay predicted from the solutions in **(B,C,D)**. The corner with  $\omega = 0.66$  is chosen as the optimal solution since it predicts the data with significantly less residual error and has no trend in the residual, unlike the corner having  $\omega = 20.1$ . The optimal solution is associated with a corner of the curve.



**Fig. 4.** The average (blue) of all 60 000 noise-generated multi-pool solutions from Fig. 2 plotted alongside the regularized solution (red).  $\langle \rho(\ln k) \rangle$  is calculated by summing  $\rho$  from all 60 000 noise-generated trials and dividing the sum by 60 000.

**Inverse method for  
estimating  
respiration rates**

D. C. Forney and  
D. H. Rothman

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion







**Fig. 5.** Comparison of root-mean-square error (RMSE) associated with the three different parameter estimation procedures. For each one, the RMSE of the predicted decay is calculated for 191 LIDET data sets. The y-axis shows the number of datasets having an RMSE smaller than the value of RMSE on the x-axis. Shown are the RMSE of the unregularized multi-pool solution (green), regularized solution (blue), and two-parameter nonlinear fit (magenta).

**Inverse method for estimating respiration rates**

D. C. Forney and  
D. H. Rothman

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 6.** Comparison of  $\mu$  and  $\sigma$  estimated from the nonlinear fitting procedure to the mean  $\langle \ln k \rangle$  and standard deviation of the regularized distribution  $\rho$ . **(A)** the mean  $\langle \ln k \rangle$  of  $\rho(\ln k)$  vs.  $\mu$ . **(B)** the standard deviation of  $\rho(\ln k)$  vs.  $\sigma$ .

**Inverse method for estimating respiration rates**

D. C. Forney and  
D. H. Rothman

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion