

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Nitrate removal in a restored riparian groundwater system: functioning and importance of individual riparian zones

S. Peter^{1,2}, **R. Rechsteiner**^{1,2}, **M. F. Lehmann**³, **R. Brankatschk**¹, **T. Vogt**^{1,4},
S. Diem⁴, **B. Wehrli**^{1,2}, **K. Tockner**⁵, and **E. Durisch-Kaiser**^{1,2}

¹Institute of Biogeochemistry and Pollutant Dynamics, ETH Zurich, 8092 Zurich, Switzerland

²Eawag, Swiss Federal Institute of Aquatic Science and Technology, 6047 Kastanienbaum, Switzerland

³Institute for Environmental Geosciences, University of Basel, 4056 Basel, Switzerland

⁴Eawag, Swiss Federal Institute of Aquatic Science and Technology, 8600 Dübendorf, Switzerland

⁵IGB, Leibniz-Institute of Freshwater Ecology and Inland Fisheries, 12587 Berlin, Germany

Received: 20 April 2012 – Accepted: 21 May 2012 – Published: 11 June 2012

Correspondence to: S. Peter (speter@env.ethz.ch)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

9, 6715–6750, 2012

**Nitrate removal in
riparian groundwater**

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

For the design and the assessment of river restoration projects, it is important to know to what extent the elimination of reactive nitrogen (N) can be improved in the riparian groundwater. We investigated the effectiveness of different riparian zones, characterized by a riparian vegetation succession, on nitrate (NO_3^-) removal from infiltrating river water in a restored and a still channelized section of the River Thur, Switzerland. Functional genes of denitrification (*nirS* and *nosZ*) were relatively abundant in groundwater from willow bush and mixed forest dominated zones, where oxygen concentrations remained low compared to the main channel and other riparian zones. After flood events, a substantial decline in NO_3^- concentration ($> 50\%$) was observed in the willow bush zone, but not in the other riparian zones closer to the river. In addition, the characteristic enrichment of ^{15}N and ^{18}O in the residual NO_3^- pool (by up to 22% for $\delta^{15}\text{N}$ and up to 12% for $\delta^{18}\text{O}$) provides qualitative evidence that the willow bush and forest zones were sites of active denitrification and, to a lesser extent, NO_3^- removal by plant uptake. Particularly in the willow bush zone, during a period of water table elevation after a flooding event, substantial input of organic carbon into the groundwater occurred, thereby fostering post-flood denitrification activity that reduced NO_3^- concentration with a rate of $\sim 21 \mu\text{mol N l}^{-1} \text{d}^{-1}$. Nitrogen removal in the forest zone was not sensitive to flood pluses, and overall NO_3^- removal rates were lower ($\sim 6 \mu\text{mol l}^{-1} \text{d}^{-1}$). Hence, discharge-modulated vegetation-soil-groundwater coupling was found to be a key driver for riparian NO_3^- removal. We estimated that, despite higher rates in the fairly constrained willow bush hot spot, total NO_3^- removal from the groundwater is lower than in the extended forest area. Overall, the aquifer in the restored section was more effective and removed $\sim 20\%$ more NO_3^- than the channelized section.

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Fixed nitrogen input into aquatic systems has tremendously increased over the past several decades (Seitzinger et al., 2006), and one third of the anthropogenic NO_3^- and ammonium is stored in groundwater, soils, and vegetation (Gruber and Galloway, 2008). High N loading can cause eutrophication of rivers, lakes, and coastal zones, and a deterioration of drinking water quality (Powlson et al., 2008). River ecosystems can play an important role in the processing of nutrients (Fischer et al., 2002), in particular through the natural removal of excess NO_3^- by denitrification and other uptake and transformation processes in riparian buffer zones (Mayer et al., 2007; Verhoeven et al., 2006). However, over the past two centuries, river regulation and the removal of wetlands have diminished the ecological and biogeochemical functionality of river systems in many parts of the world (Tockner and Stanford, 2002).

The ecological functioning (e.g., the capacity to remove NO_3^-) of a riparian system strongly depends on the quality and connectivity of different functional process zones (FPZ, *sensu* Thorp et al., 2006), which are patches that differ from their surroundings in structure and function, and are functionally connected via surface and subsurface flow paths (Fisher and Weiter, 2005; Malard et al., 2002). Hydrologic connectivity includes vertical (river-groundwater), lateral (river-landscape) and longitudinal exchange processes (Boulton, 2007; Fisher and Weiter, 2005), and can affect NO_3^- elimination along subsurface flow paths (Baker and Vervier, 2004; Burt and Pinay, 2005). Hinkle et al. (2001), for example, demonstrated that denitrification depends on subsurface travel times and the rate of water exchange between the river and its hyporheic zone.

Subsurface flow paths link different zones that carry different reactants, and thus modulate the distribution of biogeochemically relevant substrates, possibly enhancing microbial reaction rates (Edwards, 1998). Heterotrophic microbial metabolism in hyporheic zones is directly limited by the availability of organic carbon (OC) (Baker et al., 1999; Findlay and Sinsabaugh, 2003; Sobczak and Findlay, 2002). Also, denitrification in NO_3^- -rich groundwater is primarily controlled by the redox conditions and

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the availability of electron donors (i.e., bioavailable OC) (Rivett et al., 2008). Organic carbon can enter the subsurface from deep-rooted vegetation, buried C-lenses, and through episodic leaching of the overlying riparian soils (Clinton et al., 2002; Gift et al., 2010; Schade et al., 2001), leading to zones with high biogeochemical activity, often referred to as biogeochemical hot spots (McClain et al., 2003). Flood events mediate the coupling of the soil and root horizon with the groundwater leading to introduction of labile OC (Peter et al., 2012) and increased NO_3^- uptake by plants (Clement et al., 2003; Hefting et al., 2004).

In this study we hypothesize that riparian zones differing in their vegetation cover will differently affect NO_3^- removal from riparian groundwater. Furthermore, the NO_3^- removal processes are expected to be modulated by hydrological connectivity, which can represent an important constraint on substrate availability (OC, NO_3^-). To test these assumptions, we investigated the main sources of N to the groundwater and estimated N budgets for different riparian zones along a cross-section of a recently restored and a still channelized section of River Thur, Switzerland. Furthermore, we studied the potential links between water table fluctuations and OC-controlled NO_3^- removal in the groundwater. Our main objectives are (i) to investigate, which riparian zones are particularly conducive to NO_3^- removal, (ii) to identify the relevant processes for NO_3^- removal, (iii) to assess the contribution and relevance of the individual zones to whole system NO_3^- removal, and (iv) to understand the role of vegetation and flood events for NO_3^- removal. Seasonal sampling campaigns were conducted using a series of 13 piezometers placed in different zones of the restored and the channelized sections, corresponding to a river-riparian succession gradient. We collected data on nutrient and OC concentrations, the dual stable isotope ratios of NO_3^- ($\delta^{15}\text{N}$, $\delta^{18}\text{O}$), and the abundance of functional genes encoding different steps in denitrification. The observations allowed assessing the functionality of a restored river corridor for nitrogen elimination.

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 Material and methods

2.1 Study site

The River Thur originates in the limestone formation of the Mount Säntis region (2500 m a.s.l.), drains a catchment area of 1700 km² on the NE Swiss Plateau, and enters the River Rhine at 345 m a.s.l. The River Thur exhibits the dynamic flow regime of an Alpine river because reservoirs and natural lakes are absent in the catchment. Maximum, mean, and minimum flow rates are 1130, 50, and 2.24 m³ s⁻¹, respectively, for the time series of 1904 to 2000 (Binderheim-Bankay et al., 2000). It was channelized in the 1890s to prevent flooding of the river valley. In the 1970s, a plan to concurrently improve the flood protection and ameliorate the ecological state of the river corridor was elaborated. Since the early 1990s, several 1–3 km long river sections were widened to allow for the formation of alternating gravel bars and to increase hydrological connectivity between the main channel and its riparian zone. Riparian succession processes were stimulated, and as a consequence habitat diversity increased (Pasquale et al., 2011).

The study sites, instrumented for an interdisciplinary research project on river restoration (Schneider et al., 2011), were located at Niederneunforn (Canton Thurgau, Switzerland) in a 2 km long restored river corridor, and 1 km upstream in a channelized section, respectively (Fig. 1a). In the restored river sections, we identified four distinct zones across a river-riparian succession gradient (Fig. 1a): (1) a bare gravel bank (gravel); (2) exposed gravel, colonized by the tall pioneer grass *Phalaris arundinacea* (grass); (3) a second flooding terrace dominated by *Salix viminalis* (willow bush); (4) the alluvial mixed ash forest (forest) located 50–60 m away from the edge of the main channel. At the non-restored site, no riparian succession is established and pasture makes up for the only FPZ. The main channel represents the main source of the infiltrating water (Vogt et al., 2010). The piezometers penetrated the aquifer down to 8 m depth (3 to 7 m screened), allowing water sample collection from the saturated zone. The saturated zone consists of a deep sandy gravel layer (5–6 m), sealed by an impermeable

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



clay layer at the bottom and covered by a poorly permeable sand and loam layer at the top. In general, the study site is characterized by a good hydraulic and hydrological connectivity between river water and riparian groundwater (Vogt et al., 2010). The groundwater below the gravel bar and the grass zone is mainly comprised of freshly infiltrated river water (~ 23 h travel time), while subsurface travel times from the river to the willow bush zone are ~ 4.5 days, and to the forest in the order of weeks (Vogt et al., 2010). In the channelized section, the hydrological connectivity is reduced, that is, ~ 3 days travel time for the river water to the first and 11 days to the last piezometer in the transect.

2.2 Hydrodynamics

Sensors with an integrated data logger for continuous measurements (15 min interval) of hydraulic head and electrical conductivity (DL/N 70, STS AG, Switzerland; error of single measurement is $\pm 0.1\%$ for head and $\pm 2\%$ for electrical conductivity) were installed in the river channel and in the piezometer of the willow bush zone. The time series of hydraulic head and electrical conductivity were recorded from June 2008 to July 2009. A three-dimensional steady-state groundwater flow model (Feflow, DHI-WASY GmbH) was setup to derive the groundwater flow paths and Darcy fluxes for the different FPZ under base flow conditions. Details of the model setup, the boundary conditions and the model calibration are described in Diem et al. (2012). In short, the modeling domain has an extension of $1200\text{ m} \times 400\text{ m}$, with the longer axis parallel to the River Thur. The model is restricted to the gravel-and-sand aquifer with a thickness of 5–6 m. The river was included as a third-type boundary condition with water levels at a discharge of $23\text{ m}^3\text{ s}^{-1}$, which were adopted from an existing hydraulic model (Pasquale et al., 2011). In the side channels (Fig. 1a), continuously measured data at water level gauges were integrated with data from periodic measurements at fixed points. The hydraulic conductivity field was calibrated using travel time information derived by time series analysis of electrical conductivity data (Vogt et al., 2010). The river-bed transfer rates (conductance) were automatically calibrated to measured groundwater levels in

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



selected piezometers. The flow paths and the average Darcy flux along the flow paths were extracted from particle backtracking results.

2.3 Sampling

Between April 2008 and June 2009, groundwater was sampled three times during flood events, once 7 days after a peak flow event, and four times during base to average flow conditions of the River Thur (Table 1, Fig. 2a). At each date, surface river water was sampled and a 12 V purge pump was used to collect water at the piezometers. Prior to collecting groundwater samples, the electric conductivity and temperature were monitored (WTW, Nova Analytics) until constant values were reached. Samples for O₂ analysis were filled bubble-free into volumetric glass flasks and immediately fixed for Winkler-analysis (Grasshoff et al., 1999). Samples for measuring concentrations of nitrous oxide (N₂O) were taken on 15 April and 23 June 2009 only. Water was filled bubble-free in serum flasks, amended with CuCl₂, and sealed gas-tight. Samples for measuring total organic carbon (TOC), total N (TN), and dissolved OC (DOC) concentrations were transferred into muffled (450 °C) glass tubes. TOC and TN samples were acidified with 2 mol l⁻¹ HCl (pH 3). All samples were stored at 4 °C in the dark for later analysis. Samples for DOC analysis were immediately filtered through pre-combusted (220 °C, 4 h) glass fiber filters (GF/F, Whatman), stored in muffled glass bottles, and analyzed within 24 h after sample collection. To analyze the suspended bacterial gene abundance, 60–1200 ml water was filtered through 0.22 µm polycarbonate filters (Millipore), and filters were stored in sterilized plastic tubes at –20 °C until further analysis.

2.4 Chemical analyses

TOC and DOC concentrations were measured on unfiltered and filtered water samples, respectively, on a Shimadzu TOC analyzer (TOC-V CPH) using high-temperature catalytic oxidation (Benner and Strom, 1993). The accuracy of the analysis was 1.5%. Concentrations of TN (dissolved and particulate inorganic and organic N), NO₃⁻, nitrite

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(NO₂⁻), and ammonium (NH₄⁺) were determined colorimetrically on a Technicon autosampler (DEV, 2004). The analytical errors were 5 % for TN, 2.5 % for NO₂⁻, and 6 % for NO₃⁻. N₂O was quantified by headspace analysis on an Agilent 6890 N gas chromatograph (Jas, Germany) with an autosampler and a GS-Carbon PLOT column. N₂O concentrations in the samples were calculated according to Weiss and Price (1980). The analytical error was 0.2–4.6 %. Dissolved O₂ concentrations were measured using the standard Winkler method (Grasshoff et al., 1999).

The stable N and O isotopic composition of dissolved NO₃⁻, providing information on major N transformation processes, was measured using the “denitrifier method” described in detail by Sigman et al. (2001) and Casciotti et al. (2002). N and O isotopic ratios were reported in the standard δ -notation relative to atmospheric N₂ for N and V-SMOW for O, respectively. Nitrate isotope measurements were standardized using the international NO₃⁻ standards IAEA-N3 and USGS-34. The two-standard calibration allows accounting for O-isotope scale compression due to instrument properties and O-isotope exchange during bacterial conversion. Oxygen exchange was additionally monitored with ¹⁸O-enriched water standards (Casciotti et al., 2002) and was never greater than 5 %. Blank contribution was generally smaller than 1 %. The reproducibility of replicate analyses was typically $\pm 0.2\text{‰}$ for $\delta^{15}\text{N}$ and $\pm 0.3\text{‰}$ for $\delta^{18}\text{O}$. Net N and O isotope fractionation during NO₃⁻ consumption in the riparian zone was approximated assuming Rayleigh distillation kinetics (Kendall and Caldwell, 1998).

2.5 Functional gene abundance

Deoxyribonucleic acid (DNA) was extracted from filters using Fast DNA Spinkit for soil (MP Biomedicals) and quantified using Sybr green I (Matsui et al., 2004). To estimate the denitrification potential of the microbial assemblage in the mobile phase along flow paths, we analyzed the abundance of the functional genes involved in the denitrification process (NO₃⁻ reductase genes *nirK* and *nirS*, and nitrous oxide reductase gene *nosZ*) by real-time PCR on a 7300 Real-Time PCR System (Applied Biosystems) with Kapa

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sybr Fast qPCR mix (Kapa Biosystems). As qPCR standards, genomic DNA of pure cultures *Pseudomonas fluorescens* C7R12 (*nirS*), *Azospirillum irakense* DSM 11586 (*nirK*), and *Pseudomonas stutzeri* (*nosZ*) were used. The individual qPCR conditions and primers used are described in detail by Brankatschk et al. (2011). To avoid PCR inhibition, dilutions of 1 : 16 of the DNA template was found to be best for optimal detection. Copy numbers were calculated using the OPC method (Brankatschk et al., 2012) and the efficiencies of the samples were derived using the LinRegPCR program (version 11, Ruijter et al., 2009). Copy numbers were normalized to the amount of extracted DNA to reduce artifacts from sample filtration and DNA extraction. Copy numbers per ng DNA yield the proportion of the microbial community carrying the specific gene. The ratio of *nosZ*/*nirS* + *nirK* was used to quantify the proportion of the denitrifying community that can carry out all steps of denitrification (including the reduction to N₂). This approach was adapted from Philippot et al. (2009). The efficiency of the PCR reaction was 81 % for *nirS*, 70 % for *nirK*, and 86 % for *nosZ*. In the qPCR assay, each template was analyzed in triplicate, and some of the environmental samples were run with replicates of DNA extraction (April 2009: willow bush; 23 June 2009: channel, grass, and forest). The deviation for these replicates was 2–27 % of copy numbers ng⁻¹ DNA for *nirS*, 2–29 % for *nirK*, and 6–33 % for *nosZ*.

2.6 Nitrogen budget calculations

N budgets were computed along flow lines for the different FPZ, respectively, assuming that the river and the soil constitute the main sources of the N compounds, and that the river is directly connected to the individual FPZ, to which river water is supplied (Fig. 1). River water enters the FPZ near the bank and is then transported with constant velocity along the flowpath to the piezometer where the final concentration was monitored (Fig. 1a). The solute concentrations at the piezometers reflect the outflow concentration of the FPZ in this model. The difference between the compound/solute flux out of the FPZ (J_{FPZ}) and the sum of fluxes into the groundwater from the channel and the soil ($J_{CHANNEL}$ and J_{SOIL}) indicate whether there is a loss or a gain of

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a N-compound in a FPZ (Fig. 1b). A negative N budget indicates a loss of N by non-conservative processes that are not accounted for in the transport model, e.g. through microbial N dissimilation or plant assimilation. Positive values indicate additional inputs also not accounted by the model, e.g., due to underestimation of the sources that are implemented in the model, or the production of dissolved inorganic N via mineralization of organic N along the flow path. Fluxes are given in mol d^{-1} and were calculated as follows:

$$J_{\text{SOIL}} = c_{\text{SOIL}} \times Q_{\text{SOIL}} \quad (1)$$

$$J_{\text{CHANNEL}} = c_{\text{CHANNEL}} \times Q_{\text{CHANNEL}} \quad (2)$$

$$J_{\text{FPZ}} = c_{\text{FPZ}} \times Q_{\text{FPZ}} \quad (3)$$

with Q_{SOIL} in ld^{-1} calculated from specific soil exfiltration rates (q_{SOIL} in $\text{lm}^{-2} \text{d}^{-1}$) and FPZ length (l) and width (w) in m ($Q_{\text{SOIL}} = q_{\text{SOIL}} \times l \times w$). All model input data are listed in the Supplement (Table S1). Huber et al. (2012) applied the COUP numerical soil water and heat model (Jansson and Moon, 2001) in combination with measured soil properties and meteorological data monitored at the site for calculation of q_{SOIL} . Q_{CHANNEL} was calculated from groundwater flux data, q_{FPZ} (Darcy flux in m d^{-1}), adopted from the three dimensional steady-state groundwater flow model developed by Diem et al. (2012) which is described above, and the length (l) and height (h) of the FPZ ($Q_{\text{CHANNEL}} = q_{\text{FPZ}} \times l \times h$). By mass balance, Q_{FPZ} equals $Q_{\text{CHANNEL}} + Q_{\text{SOIL}}$. The data set on groundwater concentrations (c_{CHANNEL} and c_{FPZ} in mol l^{-1}) used for budget calculations is based on the sampling campaigns during 2008–2009 described in this study and an additional sampling in May 2011 (not shown). For each FPZ, data from one selected piezometer was averaged for base to average discharge conditions ($< 75 \text{ m}^3 \text{ s}^{-1}$) during spring and summer periods (Fig. 1). Soil concentration data (c_{SOIL} in mol l^{-1}) are described in Huber et al. (2012). Finally, the N fluxes were standardized to $\mu\text{mol l}^{-1} \text{d}^{-1}$ considering the effective FPZ volume (FPZ volume \times porosity). The efficiency of the individual FPZ for NO_3^- removal was calculated after Dhondt et al. (2006): efficiency =

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$(J_{\text{SOIL}} + J_{\text{CHANNEL}} - J_{\text{FPZ}}) / (J_{\text{SOIL}} + J_{\text{CHANNEL}}) \times 100\%$. Monte-Carlo-Simulations were run with 10^6 iterations on Matlab (7.8.0, The MathWorks) for mean and error estimations.

3 Results

3.1 Hydrodynamics

5 At all sampling campaigns, the water level of the River Thur was higher than the ground-water head (Fig. 2a) and the river was losing water into the saturated riparian zone (Fig. 1a). The two main piezometer transects, installed in the unconfined aquifer at the restored and the channelized section, respectively, do not reflect the subsurface flow direction (Fig. 1a). As a consequence, the individual piezometers and FPZ are
10 not directly connected by the hydrological flow. With exception of the forest zone, the river water infiltrates directly into the FPZ (Fig. 1). The hydrological regime of the river strongly differed among the sampling campaigns (Table 1), and temporal water level changes in the channel reflect these variations (Fig. 2a). In the willow bush zone, the water level responded almost instantaneously to changes in the river, indicating a good
15 hydraulic connection between river and groundwater, and during peak events groundwater reached the base of the overlying loamy soil at ~ 372.8 m a.s.l. (Fig. 2a). A similar trend in water table fluctuation was described by Vogt et al. (2010) for piezometers in the grass, forest, and pasture zones. In the willow bush zone, the electrical conductivity time series showed a slightly damped and time-shifted river signal from November
20 2008 until April 2009 (Fig. 2b), reflecting solute transport from the river to the aquifer. Additionally, elevated electrical conductivity peaks, reverse to the diluted river signal, were observed during flood events in June 2008, September 2008, and May to July 2009 (Fig. 2a).

3.2 Spatio-temporal variability of NO_3^- , NO_2^- , and N_2O concentrations

The river channel was characterized by high NO_3^- concentrations (mean $139 \pm 74 \mu\text{mol l}^{-1}$), which were marginally diluted during flood events and exhibited a marked seasonality, with higher concentrations in winter than in summer (Figs. 2c, 3a, b) (http://www.naduf.ch/daten.htm). Overall, NO_3^- represented the main fraction of TN (92 \pm 15 %), and NO_3^- concentrations were negatively (but not significantly) correlated with river discharge (Pearson correlation coefficient: $r = -0.32$, significance value: $p > 0.05$). Also, the inverse correlation with temperature was relatively weak, even if significant ($r = -0.51$, $p < 0.01$). Both, NO_2^- and N_2O concentrations in the river were low (0.5 ± 0.6 and $0.02 \pm 0.005 \mu\text{mol l}^{-1}$, respectively, Fig. 3c–e). Analysis of variance (ANOVA, SPSS 17.0, IBM) could not resolve significant differences in NO_3^- concentrations among the different riparian zones ($p > 0.05$). Nevertheless, different trends for the individual zones could be discerned for different flow regimes (Fig. 3a, b). Gravel and grass zone NO_3^- concentrations correlated with NO_3^- concentrations in the river ($r = 0.99$, $p < 0.01$ and $r = 0.78$, $p < 0.05$, respectively). During flood events, NO_3^- levels at the groundwater piezometers close to the river were lower than at the more distant ones, simply because NO_3^- concentrations were lowered in river water during these events (Fig. 3a). In the willow bush zone, the NO_3^- concentrations were only weakly correlated with those in the channel ($r = 0.33$, $p < 0.05$), and during base flow, the lowest NO_3^- concentration of all FPZ was recorded (Fig. 3b). In contrast to NO_3^- , the highest NO_2^- and N_2O concentrations were measured during base flow in the willow bush zone (Fig. 3d, e). In samples from the forest, the NO_3^- concentrations were essentially invariant, with a mean value of $136 \pm 33 \mu\text{mol l}^{-1}$. The forest zone also exhibited elevated N_2O and slightly elevated NO_2^- concentrations during base/average flow, compared to the other riparian zones. The pasture zone in the channelized river section exhibited relatively invariant NO_3^- concentrations, and no elevated NO_2^- concentrations were observed.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3 Nitrogen budgets for individual FPZ

Based on budget calculations for base and average flow conditions of the river, most of the NO_3^- originated from infiltrating channel water, whereas the contribution of soil NO_3^- to the groundwater was low in the grass and willow bush zones (1 ± 1.2 and $0.5 \pm 0.6\%$, respectively), and higher in the forest zone ($10.6 \pm 9.3\%$). For all FPZ, model results revealed a net NO_3^- loss (Fig. 4a). The highest volumetric NO_3^- removal rates were found in the willow bush zone. Grass also exhibited increased rates, but with high model error, and the lowest NO_3^- removal efficiency of all zones. A small NO_2^- (0.1 – $0.5 \mu\text{mol l}^{-1} \text{d}^{-1}$) and N_2O (0.01 – $0.02 \mu\text{mol l}^{-1} \text{d}^{-1}$) production was observed in the willow bush and the forest zones. A loss of NH_4^+ was observed in all riparian zones (-0.2 to $-1.1 \mu\text{mol l}^{-1} \text{d}^{-1}$). Extrapolation to the scale of the total restored riparian aquifer revealed that the forest zone contributed most to total NO_3^- removal (Fig. 4b). For the pasture zone, the volume of the total restored aquifer was used for scaling, which allowed a direct comparison between the restored and the channelized section. NO_3^- removal rates were higher for the restored than for the channelized section (Fig. 4).

3.4 Nitrate isotope ratios

The river water, as well as the groundwater in the adjacent gravel and grass zones exhibited the lowest $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ values for NO_3^- (~ 9.5 and $\sim 2.5\%$, respectively) among the different zones (Fig. 5). The highest NO_3^- $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ values were observed in the willow bush zone. Taking into account the complete groundwater data set, a linear relationship exists between NO_3^- $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ ($R^2 = 0.87$, slope = 0.572 , $p < 0.01$). The apparent N (and O) isotope effects ε_{N} (ε_{O}) were derived by plotting the $\delta^{15}\text{N}$ ($\delta^{18}\text{O}$) against the natural logarithm of the residual NO_3^- concentration according to Mariotti et al. (1981), where the slope of the linear regression line approximates ε_{app} :
$$\delta^{15}\text{N} \text{ (or } \delta^{18}\text{O)}_{\text{reactant}} = \delta^{15}\text{N} \text{ (or } \delta^{18}\text{O)}_{\text{initial}} - \varepsilon(\ln[\text{NO}_3^-]).$$
 For the groundwater data of the restored river section, these Rayleigh graphs yielded significant linear correlations

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for June ($R^2 = 0.94$, slope: apparent enrichment factor $\varepsilon_N = -22\text{‰}$, $p < 0.01$) and September 2008 ($R^2 = 0.75$, slope: $\varepsilon_N = -13.2\text{‰}$, $p < 0.01$) and a moderately significant linear relationship for 23 June 2009 ($R^2 = 0.42$, slope: $\varepsilon_N = -6.6\text{‰}$, $p < 0.05$) (Fig. 6). Apparent enrichment factors for ^{18}O were -14.9 , -10.4 , and -4‰ for June and September 2008, and 24 June 2009, respectively. The data for the linear regression models passed tests for normality (Shapiro-Wilk-test) and for constant variance.

3.5 Functional gene abundance

The amount of DNA extracted from water samples was generally lower in April 2009 than in June 2009, with highest concentrations in the river water in both seasons (data not shown). The *nirK*, *nirS*, and *nosZ* genes involved in denitrification were abundant throughout all the riparian aquifer, with the *nirK* gene generally showing the highest contribution (Fig. 7). The *nirS* and *nosZ* genes were significantly correlated ($r = 0.97$, $p < 0.01$) for both sampling campaigns. River water and groundwater of the grass zone generally exhibited the lowest relative abundance of *nirS* and *nosZ* genes. High abundances of *nirS* and *nosZ* genes, relative to the other zones, were observed in the willow bush zone. In the forest zone, *nirS* and *nosZ* genes were also elevated when compared to the river water samples. The *nosZ*/*(nirS + nirK)* ratio, which indicates the proportion of the denitrifying community that can carry out complete denitrification (including the reduction to N_2), was highest in the willow bush followed by the forest zone in both sampling campaigns (Fig. 7). The variations can primarily be attributed to changes in *nirS* and *nosZ* gene abundance. No samples were analyzed for the gravel and pasture groundwater.

3.6 Spatio-temporal variability of TOC and O_2 concentrations

In the River Thur, TOC concentrations were lower during base/average flow conditions than during high flow (except the sampling on 23 June 2009, which took place during a declining flood, and TOC concentrations were still similar to those at the beginning of

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the flood on 17 June 2009) (Fig. 8a, b). The TOC concentrations in the groundwater of the gravel and grass zones were closely correlated with those in the river ($r = 0.99$, $p < 0.01$ and $r = 0.87$, $p < 0.05$, respectively). In the willow bush zone, TOC concentrations were the highest during flood events. The most invariant TOC concentrations (at low levels) among all zones were observed in the forest and the pasture zones (Fig. 8a, b). There, the OC dynamics seem to be decoupled from OC fluctuations in proximity to the river corridor. Dissolved O_2 concentrations in the River Thur were close to saturated levels (280–420 $\mu\text{mol l}^{-1}$) (Fig. 8c, d), but within the first few meters distance from the channel, O_2 concentration in the groundwater dropped to 25 $\mu\text{mol l}^{-1}$ until the willow bush FPZ. In all locations, oxygen concentrations were highest during winter low flow (January 2009) and/or during early spring flooding (April 2008).

4 Discussion

4.1 N sources and net losses in individual riparian zones

The River Thur represents the main water supply for the riparian groundwater of all FPZ (Vogt et al., 2010), mirrored in the high hydraulic and hydrological connectivity in the riparian aquifer (Fig. 2a, b). Moreover, the channel constitutes the main N source to the groundwater as the River Thur receives high NO_3^- inputs from diffuse agricultural sources (Abbaspour et al., 2007), but inputs from the soil can also significantly contribute to the groundwater N pool ($\sim 11\%$ in the forest zone) (Huber et al., 2012).

Our data suggest that the individual riparian zones of the River Thur greatly differ in their capacity for net NO_3^- removal from infiltrating river water during base to average runoff (Fig. 3). Low concentrations of NO_3^- were detected in the willow bush zone, indicating a removal of the NO_3^- supplied by the river. From N budget calculations, we found that the willow bush zone can be considered a hot spot for NO_3^- removal during base and average flow, exhibiting highest removal rates per unit volume and highest fixed N removal efficiency (Fig. 4a). Lowest N removal efficiency was calculated for the

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



grass zone. However, in this zone, a relatively large variability in NO_3^- concentrations, in combination with small concentration differences with respect to the channel, and a relatively fast groundwater flow resulted in large errors for model-based estimates of N removal rates (Fig. 4a, Table S1). Taking the spatial extent of the different riparian zones into account, the forest contributes most to the total riparian aquifer NO_3^- removal (Fig. 4b). This supports the findings by Findlay et al. (2011) who found that large areas with comparatively low denitrification activity (cool matrix) can actually remove more NO_3^- than rather constrained hot spots of high denitrification rates. The budget calculations also revealed that the forest groundwater receive more soil NO_3^- than the other riparian zones ($\sim 11\%$). Comparing the pasture in the channelized part of the river with the restored aquifer, our data demonstrate that the pasture played a less important role in the ecosystem removal of NO_3^- (Fig. 4b). It needs to be noted that the model disregards NO_3^- loss at the river-aquifer boundary and assumes constant groundwater flow and transformation rates, which might not entirely reflect the reality. Especially for the forest zone with the longest infiltration path this may create additional model uncertainties that were not accounted for.

4.2 NO_3^- removal processes

In riparian aquifers, NO_3^- removal is mainly attributed to denitrification or assimilation by riparian vegetation (Dhondt et al., 2003), although a decrease in NO_3^- concentration can also be related to mixing with NO_3^- -depleted groundwater (Craig et al., 2010). In the willow bush and forest zones of the River Thur aquifer, the increased concentration of NO_2^- and N_2O (Fig. 3d, e), two intermediate products of denitrification, imply that denitrification is responsible for the apparent NO_3^- removal. Further support for denitrification as an important N loss term is provided by the functional gene analyses. The higher relative abundance of genes involved in denitrification in the willow bush and forest zones indicate that here the microbial denitrification potential was higher than in the other riparian zones (Fig. 7). It needs to be noted that we have analyzed

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the suspended microbial community, which only partially reflects the total subsurface community and does not include microbes in biofilms. In a strict sense, we cannot say a priori that gene evidence is representative of the in situ conditions, as microbes found in one FPZ could in fact originate from another FPZ or the river. In a companion study, however, Peter et al. (2012) found that the suspended microbial community is actively growing in the groundwater. Here determined gene abundances thus seem to reflect local, and to a lesser extent upstream environmental conditions, and may be used as an indicator for local processes. We found that in the willow bush zone, the relative contribution of the *nosZ* gene to the total number of genes involved in denitrification was highest compared to all other zones, independent of whether NO_3^- concentration was high or low. It has been shown that the *nosZ* abundance positively correlate with potential denitrification rates (Brankatschk et al., 2011; Hallin et al., 2009). Furthermore, it indicates the dominance of microbes that are capable of carrying out complete reduction of NO_3^- to N_2 (Fig. 7).

The patterns of ^{15}N and ^{18}O in the residual NO_3^- in groundwater helped to further discern physical mixing, plant uptake and denitrification as potential mechanisms to explain the low NO_3^- concentrations. We can rule out that mixing with water sources that are tagged with a high- $\delta^{15}\text{N}$ ($\delta^{18}\text{O}$) NO_3^- isotope signature contributed to the observed elevation in the NO_3^- delta values in the willow bush zone. Leaching of NO_3^- from willow bush soil into the groundwater was negligible ($\sim 0.5\%$) and the soil water displayed $\delta^{15}\text{N}$ values that were only slightly (if at all) elevated over the NO_3^- $\delta^{15}\text{N}$ in the River Thur and markedly lower than the observed $\delta^{15}\text{N}$ values for the willow bush zone ($9.5 \pm 2.9\%$, personal communication by Benjamin Huber). Furthermore, simple mixing with old NO_3^- -depleted groundwater is unlikely, given a very short water residence time of only 4.5 days for this zone (Vogt et al., 2010). Plotting the $\delta^{15}\text{N}$ of NO_3^- versus $1/\text{NO}_3^-$ did not reveal any evidence for mixing (plot not shown). In fact, the logarithmic correlation between NO_3^- $\delta^{15}\text{N}$ and NO_3^- concentration provides a strong argument against conservative mixing and for biological removal (Mariotti et al., 1981)

Nitrate removal in riparian groundwater

S. Peter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(Fig. 6). The calculated average $\Delta\delta^{18}\text{O}:\Delta\delta^{15}\text{N}$ enrichment ratio was 0.57. Similar values were reported for denitrification from another riparian zone (0.59; Cey et al., 1999) and other freshwater systems (0.6; Lehmann et al., 2003).

The relative contribution of denitrification and plant assimilation to the community N isotope fractionation can be approximated if the end member N isotope effects for denitrification and assimilation are known, respectively (Dhondt et al., 2003; Lund et al., 1999). Adopting the enrichment factors reported by Dhondt et al. (2003) for plant uptake ($\sim -4\text{‰}$) and denitrification in groundwater ($\sim -23\text{‰}$), and based on the average N isotope enrichment factor of -14‰ determined in this study, we can calculate that on average 62 % the observed NO_3^- deficiency can be attributed to denitrification, while the remaining 37 % was due to plant uptake. Variation in the community N isotope effect for NO_3^- removal, suggests, however, that this partitioning varied significantly. Using the same approach, we estimate that denitrification contributed 98 % in June 2008, 68 % in September 2008, and 21 % in June 2009 to the NO_3^- removal. Thus, in this riparian aquifer, NO_3^- removal seems to primarily take place through denitrification and to a lesser extent through plant uptake. Interestingly, both processes seem to be constrained to the willow bush and, to a lesser extent, to the alluvial forest zone during base and average flow conditions.

4.3 Vegetation-soil-groundwater coupling: the role of habitats and flood events

At the restored Thur site, gravel bars and grass strips close to the main channel were mainly influenced by OC introduced with infiltrating river water (Fig. 8) and exhibited no apparent denitrification activity. We assume that the subsurface transfer times were too short and OC supply insufficient for the development of denitrifying conditions (Malard et al., 2002). In the willow bush zone, where flood events result in enhanced OC availability (Fig. 8) (Peter et al., 2012), a denitrification hot spot can be formed. The alluvial forest and the pasture in the channelized river section exhibit intermediate TOC concentrations with smallest temporal concentration changes among all groundwater

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



samples and denitrification rates are lower (Fig. 4a). Hence, in the Thur aquifer, as a limiting substrate, OC appears as the main factor that controls microbial efficiency in general and NO_3^- removal in particular.

The NO_3^- removal hot spot willow bush was densely populated with *Salix viminalis*. These pioneer species are closely associated with enhanced heterotrophic growth in the rhizosphere (Lalke-Porczyk et al., 2009) through high below-ground carbon production (de Neergaard et al., 2002) and known for their high NO_3^- removal capacity (Aronsson and Perttu, 2001; Elowson, 1999). In a companion study at this field site, the OC composition in the willow bush zone groundwater could clearly be related to bioavailable soil- and vegetation-derived OC (Peter et al., 2011). Therefore, we assume that willow plants are a considerable source of bioavailable OC for the saturated subsurface zone. Here, it may directly stimulate microbial respiration, leading to the development of sub-oxic conditions, and potentially initiate denitrification. Indeed, a recent study by Clinton et al. (2010) suggests that differences in OC bioavailability in a riparian subsurface flow is related to the type of vegetation cover. Similarly, Schade et al. (2001) observed a direct effect of vegetation cover on groundwater denitrification and proposed that the organic substrates that drive denitrification originated from plant production.

Variations in water table can foster interactions of the groundwater with the root zone of the riparian vegetation. Usually roots do not grow into the saturated zone, but in riparian systems, roots may be submerged during high water periods (Dosskey et al., 2010). Flooding raises the water table by 1–2 m towards the top-soil horizon of the willow bush zone (Fig. 2a) and may stimulate NO_3^- removal through denitrification by connecting the NO_3^- -rich groundwater with the bioavailable OC pool of the overlaying soil and root zone. In the willow bush zone, the electrical conductivity measurements indicated high ionic content in the water during periods of high runoff and low conductivity in the river (Fig. 2b and Vogt et al., 2010). Conductivity peaks can be explained by the leaching of high-conductivity and, at the same time, OC-rich soil water (Huber et al., 2012) into the groundwater. Introduction of bioavailable OC stimulates microbial CO_2

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



production, resulting in an increase in the electrical conductivity through ion dissolution. These flood-related conductivity peaks were mainly observed during the vegetation period (Fig. 2). Therefore, we assume that it is the root-derived OC that stimulates microbial activity and ultimately leads to an increased conductivity signal in the groundwater during flood events. This strong river-groundwater and soil-groundwater coupling at the willow bush zone facilitates substrate input to the subsurface, and may ultimately set the optimal conditions for the development of a N turnover hot spot in the post-flooding period.

Low NO_3^- concentrations and enrichment of the heavier NO_3^- isotopologues in the willow bush zone was observed always during the vegetation period (Fig. 5). In the winter/early spring period (April 2008 and January and April 2009), the groundwater was characterized by low temperatures (0.5–10 °C) and high O_2 concentrations (Fig. 8b), indicating reduced microbial activity, consistent with no significant ^{15}N (^{18}O) enrichment in the NO_3^- pool. Also the N isotopic data suggest that in the Thur riparian aquifer little denitrification and plant uptake occurred during flooding (Figs. 3 and 5), despite substantial NO_3^- supply to the groundwater and intense contact of the groundwater with the root zone (Fig. 2a) providing bioavailable organic substrate. Flood events increase the groundwater flow velocity and can substantially decrease the actual contact time with the roots of the riparian vegetation and with the microbial “denitrifier” community, leading to diminished NO_3^- removal. One week post flooding, NO_3^- and TOC concentrations were substantially reduced in the willow bush zone (Figs. 3 and 8) and the residual NO_3^- was enriched in the heavy N and O isotopes (Fig. 5), suggesting intense post-flood NO_3^- removal through denitrification, triggered by the OC spike during the flood event, and, to a lesser degree, plant NO_3^- uptake by *Salix viminalis*.

5 Concluding remarks and implications for restoration of riparian zones

In a dynamic restored river system like at the River Thur, spatial heterogeneity translates into different FPZ with contrasting NO_3^- removal potential. Both, hot spots and

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cold matrices contribute to the enhanced NO_3^- removal capacity through denitrification and plant uptake, compared to the channelized riparian site. The rates of NO_3^- removal of $\sim 21 \mu\text{mol N l}^{-1} \text{d}^{-1}$ were similar to values reported in previous studies (Baker and Vervier, 2004; Mengis et al., 1999). The functioning of the willow bush hot spot as efficient NO_3^- filter is governed by hydraulic and biogeochemical conditions: periodic flooding transports soil- and root-derived organic matter into the groundwater and strong hydraulic connectivity warrants NO_3^- supply. The biogeochemical N-uptake by intense plant growth (Clement et al., 2003) probably also contributed to NO_3^- elimination. Denitrification not only removes NO_3^- from the system but can also be a considerable source of the greenhouse gas N_2O , especially in systems that are prone to frequent perturbations, for example due to sudden flood events. The latter may episodically “ventilate” the subsurface, thereby inhibiting N_2O reduction (Naqvi et al., 2000; Verhoeven et al., 2006; Codispoti et al., 2001). However, the generally low N_2O concentrations in the groundwater (Fig. 3e) translate into low production rates (Sect. 3.3) compared to rates from other systems (e.g., Kim et al., 2009; Mengis et al., 1996).

This study emphasizes that successful river restoration should consider the multi-dimensional linkages of alluvial groundwater systems. We recommend enhancing the environmental heterogeneity by configuring distinct riparian zones, or by allowing the occurrence of a natural geomorphic and vegetation succession. For example, we demonstrate here the value of a pioneer plant-dominated zone with regards to recharge of the soil OC inventory and the stimulation of NO_3^- removal in the water-saturated zone. Flood events are necessary to fully exploit the soil-groundwater connection and to develop optimal conditions for a denitrification hot spot. Overall, this study illustrates that the interactions between ecological and hydrological processes and riparian morphology should be taken into consideration in future restoration measures.

BGD

9, 6715–6750, 2012

Nitrate removal in riparian groundwater

S. Peter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Supplementary material related to this article is available online at:
[http://www.biogeosciences-discuss.net/9/6715/2012/
bgd-9-6715-2012-supplement.pdf](http://www.biogeosciences-discuss.net/9/6715/2012/bgd-9-6715-2012-supplement.pdf).

Acknowledgement. We are grateful to Marco Baumann and Andreas Scholtis from the Agency for the Environment, Canton Thurgau, for their continuous support and the excellent collaboration within this research project. Alessandro Brovelli (EPFL), Jörg Luster and Benjamin Huber (WSL) helped to set up the budget calculations. We would like to thank Rafael Blättler, Eytan Gur, Kate Gassner, Ruth Stierli, Beat Müller, Stefan Koetzsch, Mark Rollog, and Torsten Diem for their assistance with field experiments or laboratory analyses. We are grateful to Mario Schirmer (Eawag) and Joseph Zeyer (ETH Zurich) for supporting this work. This work is part of the project REstored COrridor Dynamics (RECORD) and supported by the Competence Center Environment and Sustainability (CCES) of the ETH Domain.

References

- Abbaspour, K. C., Yang, J., Maximov, I., Siber, R., Bogner, K., Mieleitner, J., Zobrist, J., and Srinivasan, R.: Modelling hydrology and water quality in the pre-alpine/alpine Thur watershed using SWAT, *J. Hydrol.*, 333, 413–430, 2007.
- Aronsson, P. and Perttu, K.: Willow vegetation filters for wastewater treatment and soil remediation combined with biomass production, *For. Chron.*, 77, 293–299, 2001.
- Baker, M., Dahm, C., and Valett, H.: Acetate retention and metabolism in the hyporheic zone of a mountain stream, *Limnol. Oceanogr.*, 44, 1530–1539, 1999.
- Baker, M. A. and Vervier, P.: Hydrological variability, organic matter supply and denitrification in the Garonne River ecosystem, *Freshw. Biol.*, 49, 181–190, 2004.
- Benner, R. and Strom, M.: A critical-evaluation of the analytical blank associated with DOC measurements by high-temperature catalytic-oxidation, *Mar. Chem.*, 41, 153–160, 1993.
- Binderheim-Bankay, E., Jakob, A., and Liechti, P.: NADUF – Messresultate 1977–1998, Federal Office for the Environment (FOEN), Bern, 241 pp., 2000.
- Boulton, A. J.: Hyporheic rehabilitation in rivers: restoring vertical connectivity, *Freshw. Biol.*, 52, 632–650, doi:10.1111/j.1365-2427.2006.01710.x, 2007.

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Nitrate removal in
riparian groundwater**S. Peter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Brankatschk, R., Töwe, S., Kleineidam, K., Schloter, M., and Zeyer, J.: Abundances and potential activities of nitrogen cycling microbial communities along a chronosequence of a glacier forefield, *ISME J.*, 5, 1025–1037, doi:10.1038/ismej.2010.184, 2011.
- 5 Brankatschk, R., Bodenhausen, N., Zeyer, J., and Bürgmann, H.: Efficiency of real-time qPCR depends on the template: a simple absolute quantification method correcting for qPCR efficiency variations in microbial community samples, *Appl. Environ. Microbiol.*, 78, 4481–4489, doi:10.1128/AEM.07878-11, 2012.
- Burt, T. and Pinay, G.: Linking hydrology and biogeochemistry in complex landscapes, *Progr. Phys. Geogr.*, 29, 297, 2005.
- 10 Casciotti, K. L., Sigman, D. M., Hastings, M. G., Bohlke, J. K., and Hilkert, A.: Measurement of the oxygen isotopic composition of nitrate in seawater and freshwater using the denitrifier method, *Anal. Chem.*, 74, 4905–4912, doi:10.1021/Ac020113w, 2002.
- Cey, E., Rudolph, D., Aravena, R., and Parkin, G.: Role of the riparian zone in controlling the distribution and fate of agricultural nitrogen near a small stream in Southern Ontario, *J. Contam. Hydrol.*, 37, 45–67, 1999.
- 15 Clement, J. C., Holmes, R. M., Peterson, B. J., and Pinay, G.: Isotopic investigation of denitrification in a riparian ecosystem in Western France, *J. Appl. Ecol.*, 40, 1035–1048, 2003.
- Clinton, S. M., Edwards, R. T., and Naiman, R. J.: Forest-river interactions: Influence on hyporheic dissolved organic carbon concentrations in a floodplain terrace, *J. Amer. Water Resour. Assoc.*, 38, 619–631, 2002.
- 20 Clinton, S. M., Edwards, R. T., and Findlay, S. E. G.: Exoenzyme activities as indicators of dissolved organic matter composition in the hyporheic zone of a floodplain river, *Freshw. Biol.*, 55, 1603–1615, doi:10.1111/j.1365-2427.2009.02383.x, 2010.
- Codispoti, L., Brandes, J., Christensen, J., Devol, A., Naqvi, S., Paerl, H., and Yoshinari, T.: The oceanic fixed nitrogen and nitrous oxide budgets: moving targets as we enter the anthropocene?, *Sci. Mar.*, 65, 85–105, 2001.
- 25 Craig, L., Bahr, J., and Roden, E.: Localized zones of denitrification in a floodplain aquifer in Southern Wisconsin, USA, *Hydrogeol. J.*, 18, 1867–1879, 2010.
- DEV: Deutsches Einheitsverfahren zur Wasseruntersuchung (DEW), Wiley-VCH Verlag GmbH, Weimheim, Germany, 2004.
- 30 Dhondt, K., Boeckx, P., Van Cleemput, O., and Hofman, G.: Quantifying nitrate retention processes in a riparian buffer zone using the natural abundance of N-15 in NO₃⁻, *Rapid Commun. Mass Sp.*, 17, 2597–2604, doi:10.1002/Rcm.1226, 2003.

**Nitrate removal in
riparian groundwater**

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Dhondt, K., Boeckx, P., Verhoest, N. E. C., Hofman, G., and Van Cleemput, O.: Assessment of temporal and spatial variation of nitrate removal in riparian zones, *Environ. Monit. Assess.*, 116, 197–215, doi:10.1007/s10661-006-7403-1, 2006.
- 5 Diem, S., Renard, P., and Schirmer, M.: New methods to assign water levels of dynamic rivers to groundwater models, in preparation, 2012.
- Dosskey, M. G., Vidon, P., Gurwick, N. P., Allan, C. J., Duval, T. P., and Lowrance, R.: The role of riparian vegetation in protecting and improving chemical water quality in streams, *J. Am. Water Resour. Assoc.*, 46, 261–277, doi:10.1111/j.1752-1688.2010.00419.x, 2010.
- 10 Edwards, R.: The hyporheic zone, in: *River Ecology and Management: Lessons From the Pacific Coastal Ecoregion*, edited by: Naiman, R. J. and Bilby, R. E., Springer, New York, 399–429, 1998.
- Elowson, S.: Willow as a vegetation filter for cleaning of polluted drainage water from agricultural land, *Biomass. Bioenerg.*, 16, 281–290, 1999.
- Findlay, S. and Sinsabaugh, R. L.: *Aquatic Ecosystems: Interactivity of Dissolved Organic Matter*, Academic Press, San Diego, 2003.
- 15 Findlay, S. E. G., Mulholland, P. J., Hamilton, S. K., Tank, J. L., Bernot, M. J., Burgin, A. J., Crenshaw, C. L., Dodds, W. K., Grimm, N. B., McDowell, W. H., Potter, J. D., and Sobota, D. J.: Cross-stream comparison of substrate-specific denitrification potential, *Biogeochemistry*, 104, 381–392, doi:10.1007/s10533-010-9512-8, 2011.
- 20 Fischer, H., Sachse, A., Steinberg, C. E. W., and Pusch, M.: Differential retention and utilization of dissolved organic carbon by bacteria in river sediments, *Limnol. Oceanogr.*, 47, 1702–1711, 2002.
- Fisher, S. G. and Weiter, J. R.: Flowpaths as integrators of heterogeneity in streams and landscapes, in: *Ecosystem Function in Heterogeneous Landscapes*, edited by: Lovett, G. M., Turner, M. G., Jones, C. G., and Weathers, K. C., Springer Science, New York, 311–328, 2005.
- 25 Gift, D. M., Groffman, P. M., Kaushal, S. S., and Mayer, P. M.: Denitrification potential, root biomass, and organic matter in degraded and restored urban riparian zones, *Restor. Ecol.*, 18, 113–120, doi:10.1111/j.1526-100X.2008.00438.x, 2010.
- 30 Grasshoff, K., Ehrhardt, M., Kremling, K., and Anderson, L.: *Methods of seawater analysis*, edited by: Frontmatter, Wiley-VCH Verlag GmbH, Weinheim, Germany, 1999.
- Gruber, N. and Galloway, J.: An Earth-system perspective of the global nitrogen cycle, *Nature*, 451, 293–296, 2008.

Nitrate removal in riparian groundwater

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Hallin, S., Jones, C., Schloter, M., and Philippot, L.: Relationship between N-cycling communities and ecosystem functioning in a 50-year-old fertilization experiment, *ISME J.*, 3, 597–605, 2009.
- Hefting, M., Clement, J., Dowrick, D., Cosandey, A., Bernal, S., Cimpian, C., Tatur, A., Burt, T., and Pinay, G.: Water table elevation controls on soil nitrogen cycling in riparian wetlands along a European climatic gradient, *Biogeochemistry*, 67, 113–134, 2004.
- Hinkle, S., Duff, J., Triska, F., Laenen, A., Gates, E., Bencala, K., Wentz, D., and Silva, S.: Linking hyporheic flow and nitrogen cycling near the Willamette River – a large river in Oregon, USA, *J. Hydrol.*, 244, 157–180, 2001.
- Huber, B., Luster, J., Bernasconi, S. M., Shrestha, J., and Graf Pannatier, E.: Nitrate leaching from short-hydroperiod floodplain soils, *Biogeosciences Discuss.*, 9, 5659–5694, doi:10.5194/bgd-9-5659-2012, 2012.
- Jansson, P. E. and Moon, D. S.: A coupled model of water, heat and mass transfer using object orientation to improve flexibility and functionality, *Environ. Model. Softw.*, 16, 37–46, 2001.
- Kendall, C. and Caldwell, E.: Fundamentals of isotope geochemistry, in: *Isotope Tracers in Catchment Hydrology*, edited by: Kendall, C. and McDonnell, J. J., Elsevier, Amsterdam, 51–86, 1998.
- Kim, D.-G., Isenhardt, T. M., Parkin, T. B., Schultz, R. C., and Loynachan, T. E.: Nitrate and dissolved nitrous oxide in groundwater within cropped fields and riparian buffers, *Biogeosciences Discuss.*, 6, 651–685, doi:10.5194/bgd-6-651-2009, 2009.
- Lalke-Porczyk, E., Brzezinska, M., and Donderski, W.: Influence of the root system of the common osier (*Salix viminalis* L.) on abundance of heterotrophic bacteria in the willow sewage treatment system, *Curr. Microbiol.*, 58, 571–577, 2009.
- Lehmann, M. F., Reichert, P., Bernasconi, S. M., Barbieri, A., and McKenzie, J.: Modelling nitrogen and oxygen isotope fractionation during denitrification in a lacustrine redox-transition zone, *Geochim. Cosmochim. Acta*, 67, 2529–2542, 2003.
- Lund, L., Horne, A., and Williams, A.: Estimating denitrification in a large constructed wetland using stable nitrogen isotope ratios, *Ecol. Engin.*, 14, 67–76, 1999.
- Malard, F., Tockner, K., Dole-Olivier, M. J., and Ward, J. V.: A landscape perspective of surface-subsurface hydrological exchanges in river corridors, *Freshw. Biol.*, 47, 621–640, 2002.
- Mariotti, A., Germon, J., Hubert, P., Kaiser, P., Letolle, R., Tardieux, A., and Tardieux, P.: Experimental determination of nitrogen kinetic isotope fractionation: some principles; illustration for the denitrification and nitrification processes, *Plant Soil*, 62, 413–430, 1981.

**Nitrate removal in
riparian groundwater**

S. Peter et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Mayer, P. M., Reynolds, S. K., Marshall, D., McCutchen, M. D., and Canfield, T. J.: Meta-analysis of nitrogen removal in riparian buffers, *J. Environ. Qual.*, 36, 1172–1180, doi:10.2134/Jeq2006.0462, 2007.
- 5 McClain, M. E., Boyer, E. W., Dent, C. L., Gergel, S. E., Grimm, N. B., Groffman, P. M., Hart, S. C., Harvey, J. W., Johnston, C. A., Mayorga, E., McDowell, W. H., and Pinay, G.: Biogeochemical hot spots and hot moments at the interface of terrestrial and aquatic ecosystems, *Ecosystems*, 6, 301–312, 2003.
- Mengis, M., Gachter, R., and Wehrl, B.: Nitrous oxide emissions to the atmosphere from an artificially oxygenated lake, *Limnol. Oceanogr.*, 41, 548–553, 1996.
- 10 Mengis, M., Schiff, S. L., Harris, M., English, M. C., Aravena, R., Elgood, R. J., and MacLean, A.: Multiple geochemical and isotopic approaches for assessing ground water NO_3^- elimination in a riparian zone, *Ground Water*, 37, 448–457, 1999.
- Naqvi, S., Jayakumar, D., Narvekar, P., Naik, H., Sarma, V., D'Souza, W., Joseph, S., and George, M.: Increased marine production of N_2O due to intensifying anoxia on the Indian continental shelf, *Nature*, 408, 346–349, 2000.
- 15 de Neergaard, A., Porter, J. R., and Gorissen, A.: Distribution of assimilated carbon in plants and rhizosphere soil of basket willow (*Salix viminalis* L.), *Plant Soil*, 245, 307–314, 2002.
- Pasquale, N., Perona, P., Schneider, P., Shrestha, J., Wombacher, A., and Burlando, P.: Modern comprehensive approach to monitor the morphodynamic evolution of a restored river corridor, *Hydrol. Earth Syst. Sci.*, 15, 1197–1212, doi:10.5194/hess-15-1197-2011, 2011.
- 20 Peter, S., Koetzsch, S., Traber, J., Bernasconi, S. M., Wehrl, B., and Durisch-Kaiser, E.: Intensified organic carbon dynamics in the groundwater of a restored riparian zone, *Freshwater Biology*, in press, 2012.
- Philippot, L., Cuhel, J., Saby, N., Chèneby, D., Chronakova, A., Bru, D., Arrouays, D., Martin-Laurent, F., and Simek, M.: Mapping field-scale spatial patterns of size and activity of the denitrifier community, *Environ. Microbiol.*, 11, 1518–1526, 2009.
- 25 Powlson, D. S., Addisott, T. M., Benjamin, N., Cassman, K. G., de Kok, T. M., van Grinsven, H., L'hirondel, J. L., Avery, A. A., and van Kessel, C.: When does nitrate become a risk for humans?, *J. Environ. Qual.*, 37, 291–295, Doi 10.2134/Jeq2007.0177, 2008.
- 30 Rivett, M. O., Buss, S. R., Morgan, P., Smith, J. W. N., and Bemment, C. D.: Nitrate attenuation in groundwater: a review of biogeochemical controlling processes, *Water Res.*, 42, 4215–4232, doi:10.1016/j.watres.2008.07.020, 2008.

**Nitrate removal in
riparian groundwater**S. Peter et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Ruijter, J. M., Ramakers, C., Hoogaars, W. M. H., Karlen, Y., Bakker, O., van den Hoff, M. J. B., and Moorman, A. F. M.: Amplification efficiency: linking baseline and bias in the analysis of quantitative PCR data, *Nucl. Acids Res.*, 37, 1–12, doi:10.1093/nar/gkp045, 2009.
- Schade, J. D., Fisher, S. G., Grimm, N. B., and Seddon, J. A.: The influence of a riparian shrub on nitrogen cycling in a Sonoran Desert stream, *Ecology*, 82, 3363–3376, 2001.
- Schneider, P., Vogt, T., Schirmer, M., Doetsch, J., Linde, N., Pasquale, N., Perona, P., and Cirpka, O. A.: Towards improved instrumentation for assessing river-groundwater interactions in a restored river corridor, *Hydrol. Earth Syst. Sci.*, 15, 2531–2549, doi:10.5194/hess-15-2531-2011, 2011.
- Seitzinger, S., Harrison, J. A., Bohlke, J. K., Bouwman, A. F., Lowrance, R., Peterson, B., Tobias, C., and Van Drecht, G.: Denitrification across landscapes and waterscapes: a synthesis, *Ecol. Appl.*, 16, 2064–2090, 2006.
- Sigman, D. M., Casciotti, K. L., Andreani, M., Barford, C., Galanter, M., and Bohlke, J. K.: A bacterial method for the nitrogen isotopic analysis of nitrate in seawater and freshwater, *Anal. Chem.*, 73, 4145–4153, 2001.
- Sobczak, W. V. and Findlay, S.: Variation in bioavailability of dissolved organic carbon among stream hyporheic flowpaths, *Ecology*, 83, 3194–3209, 2002.
- Thorp, J. H., Thoms, M. C., and Delong, M. D.: The riverine ecosystem synthesis: biocomplexity in river networks across space and time, *River Res. Appl.*, 22, 123–147, 2006.
- Tockner, K. and Stanford, J. A.: Riverine flood plains: present state and future trends, *Environ. Conserv.*, 29, 308–330, 2002.
- Verhoeven, J. T. A., Arheimer, B., Yin, C. Q., and Hefting, M. M.: Regional and global concerns over wetlands and water quality, *Trend. Ecol. Evol.*, 21, 96–103, doi:10.1016/j.tree.2005.11.015, 2006.
- Vogt, T., Hoehn, E., Schneider, P., Freund, A., Schirmer, M., and Cirpka, O.: Fluctuations of electrical conductivity as a natural tracer for bank filtration in a losing stream, *Adv. Water Resourc.*, 33, 1296–1308, doi:10.1016/j.advwatres.2010.02.007, 2010.

Nitrate removal in riparian groundwater

S. Peter et al.

Table 1. Temperature, mean discharge, flow regime of the River Thur, and the number (*n*) of samples taken or piezometers sampled in each riparian zone during the sampling campaigns.

Sampling date	Water temp. (°C)	Discharge (m ³ s ⁻¹)	Flow regime	Channel (<i>n</i>)	Gravel (<i>n</i>)	Grass (<i>n</i>)	Willow bush (<i>n</i>)	Forest (<i>n</i>)	Pasture (<i>n</i>)
23 Apr 2008	8.0	347	High	4	Flooded	Flooded	1	7	7
17 Jun 2008	14.4	35.6	Average	4	0	1	1	5	5
15 Jul 2008 ³	16.8	242	High	3	Flooded	1	1	5	4
2 Sep 2008	17.8	16.5	Base	1	2	2	1	4	0
6 Jan 2009 ³	0.5	14.2	Base	1	1	2	1	4	0
15 Apr 2009 ^{1,2}	9.6	62.8	Average	2	2	3	2	4	4
17 Jun 2009	15.1	100	High [†]	1	1	2	1	3	0
23 Jun 2009 ^{1,2}	11.5	201	High ^{**}	1	Flooded	2	2	4	0

[†] Onset of a flood.

^{**} Declining flood after one week of highly variable discharge conditions.

¹ Analyzed for N₂O concentration.

² Analyzed for functional gene abundance.

³ Not analyzed for stable NO₃⁻ isotopes.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







Fig. 1. (a) Field site at Niederneunforn, Switzerland, with the installed piezometers in the restored and channelized section of the River Thur and the different FPZ indicated. The modeled groundwater flow field (Diem et al., 2012) is illustrated by isopotential lines (equidistance 10 cm) and by selected flow paths. To ease a comparison with previous studies e.g., Vogt et al. (2010) and Schneider et al. (2011), the link to the original piezometer names is given here for the individual FPZ, beginning with the piezometer closest to the main channel: gravel: R050 and R060; grass: R042, R051, and R052; willow bush: R041 and R058; forest: R017, R023, R024, R019, R025, R026, and R021; pasture: R001, R004, R005, R009, R011, R012, and R014. **(b)** Schematic illustration of the N groundwater budget. River channel and the soil are the two main sources for the groundwater N pool, and the individual FPZ are directly connected to the channel.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nitrate removal in
riparian groundwater

S. Peter et al.




Fig. 2. Variations in water level (a) and electrical conductivity (EC) (b) in the channel and in the piezometer in willow bush (R041) from May 2008 until the end of June 2009 and bi-weekly NO_3^- concentration in the River Thur (c) (<http://www.naduf.ch/daten.htm>). The gray area indicates the position of the overlaying soil structure of the aquifer. Vertical bars designate the different sampling dates. Water level fluctuations in willow bush followed water level variations in the channel, reflecting high hydraulic connectivity. Electrical conductivity generally showed a slightly dampened and time-shifted signal in willow bush compared to the channel. During flood events in the periods June to October 2008 and May to July 2009, the electrical conductivity signal in willow bush was reverse to that of the channel.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nitrate removal in riparian groundwater

S. Peter et al.




Fig. 3. NO_3^- , NO_2^- and dissolved N_2O concentrations in different riparian zones during high discharge (a, c) and base to average discharge of the River Thur (b, d, e). Values from piezometers located in the same riparian zone were averaged, and error bars represent standard deviations ($n > 2$) or average deviations ($n = 2$).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nitrate removal in riparian groundwater

S. Peter et al.




Fig. 4. (a) Average NO_3^- removal rates and efficiencies with standard deviations (\pm) derived by Monte-Carlo-Simulations within a specific FPZ during base/average flow conditions of the river during spring to summer. The negative rate values designate loss from the groundwater, most probably due to dissimilative consumption by denitrification or due to plant uptake. (b) Nitrate loss rates (\pm standard deviations from Monte-Carlo-Simulations) for the individual FPZ are normalized to the whole restored river section at base/average flow conditions of the river, and during the spring to summer period. For better comparison, we used the same aquifer volume for the channelized pasture area as for the total restored aquifer.




Fig. 5. Stable N (a) and O (b) isotope ratios ($\delta^{15}\text{N}$, $\delta^{18}\text{O}$) of residual NO_3^- in different riparian zones from all sampling campaigns. In June and September 2008, as well as in June 2009 one week after a flooding event, an enrichment of ^{15}N and ^{18}O in the NO_3^- was observed in willow bush. Values from piezometers located in the same zone were averaged and error bars represent standard deviations ($n > 2$) or average deviations ($n = 2$).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. Rayleigh plots for $\delta^{15}\text{N}$ from June and September 2008 and 23 June 2009 indicate that 42–94 % of the covariance between $\ln[\text{NO}_3^-]$ and $\delta^{15}\text{N}$ can be explained by biological fractionation. The slope of the regression lines indicates the apparent (or community) enrichment factors of NO_3^- removal.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 8. TOC and O₂ concentrations in different riparian zones during high discharge (**a, c**) and base to average discharge of the River Thur (**b, d**). Values from piezometers located in the same riparian zone were averaged and error bars represent standard deviations ($n > 2$) or average deviations ($n = 2$).