

**Degradation of
planktonic and
bacterial lipids in
sinking particles**

J.-F. Rontani et al.

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Intense photooxidative degradation of planktonic and bacterial lipids in sinking particles collected with sediment traps across the Canadian Beaufort Shelf (Arctic Ocean)

J.-F. Rontani¹, B. Charriere¹, A. Forest², S. Heussner³, F. Vaultier¹, M. Petit¹, N. Delsaut³, L. Fortier², and R. Sempéré¹

¹Aix-Marseille Univ., Mediterranean Institute of Oceanography (MIO), UMR7294, CNRS/INSU, UMR235, IRD, 13288, Marseille, Cedex 09; Université du Sud Toulon-Var (MIO), 83957, La Garde cedex, France

²Université Laval, Takuvik CNRS-Laval U. Joint Laboratory, 1045 av. de la Médecine, Québec, Québec G1V 0A6, Canada

³Centre de Formation et de Recherche sur l'Environnement Méditerranéen (CEFREM, UMR5110 CNRS), Bât. U, Université de Perpignan, Via Domitia (UPVD), 66860 Perpignan, France

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Received: 31 May 2012 – Accepted: 15 June 2012 – Published: 26 June 2012

Correspondence to: J.-F. Rontani (jean-francois.rontani@univ-amu.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

9, 7743–7781, 2012

**Degradation of
planktonic and
bacterial lipids in
sinking particles**

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The lipid content of seven samples of sinking particles collected with sediment traps moored at ~100 m depth in summer and fall across the Canadian Beaufort Shelf (Arctic Ocean) was investigated. Our main goal was to quantify and characterize the biotic and abiotic degradation processes that acted on sinking material during these periods. Diatoms, which dominated the phytoplanktonic assemblage in every trap sample, appeared to be remarkably sensitive to Type II (i.e. involving singlet oxygen) photodegradation processes in summer, but seemed to be relatively unaffected by biotic degradation at the same time. Hence, the relative recalcitrance of phytodetritus towards biodegradation processes during the Arctic midnight sun period was attributed to the strong photodegradation state of heterotrophic bacteria, which likely resulted from the efficient transfer of singlet oxygen from photodegraded phytoplanktonic cells to attached bacteria. In addition, the detection in trap samples of photoproducts specific to wax ester components found in herbivorous copepods demonstrated that zooplanktonic faecal material exported out of the euphotic zone in summer were as well affected by Type II photodegradation processes. By contrast, sinking particles collected during the autumn were not influenced by any light-driven stress. Further chemical analyses showed that photodegraded sinking particles contained an important amount of intact hydroperoxides, which could then induce a strong oxidative stress in underlying sediments.

1 Introduction

Continental shelves of the Arctic Ocean receive considerable amount of terrestrial matter from river runoff mixed with important autochthonous production from microalgal photosynthesis during spring-summer (Rachold et al., 2004; Wassmann et al., 2004). The ongoing trend of declining sea ice extent and thickness in the Arctic Ocean appears to induce a steady increase in pelagic primary production (Arrigo et al., 2008),

BGD

9, 7743–7781, 2012

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

whereas permafrost thawing combined with enhanced river discharge are currently increasing the seaward flux of terrigenous material (Frey and McClelland, 2009). In turn, the annual lengthening of the ice-free period and the rise in river run-off could lead to an increase in particulate matter export that could modify the biogeochemistry and trophic balance of Arctic ecosystems through the coastal-marine realm (Vallière et al., 2008; Wassmann and Reigstad, 2011). Therefore, it is thus essential to understand how abiotic (autoxidation and photooxidation) and biotic (bacterial degradation) processes affect the dynamics of sinking fluxes of particulate organic matter in such environments.

Particles in the water column exist in a continuum of sizes (McCave, 1984), with two classes usually operationally recognized (Bacon et al., 1985; Wakeham and Lee, 1989): (i) suspended particles ($\leq 10^2 \mu\text{m}$ diameter) sinking very slowly through the water column and constituting most of the standing stock of particulate matter in the ocean and (ii) sinking particles ($\geq 10^2 \mu\text{m}$ diameter) (including zooplankton fecal pellets and marine snow aggregates) numerically less abundant but responsible for most of the downward flux of material from the upper ocean to the sea floor. Suspended particles are typically collected by filtration, whereas sinking material is commonly collected using sediment traps. It may be noted that a continual exchange exists between these two particle classes owed to the complex suite of aggregation and disaggregation processes that occur in the water column (Wakeham and Lee, 1989; Hill, 1998). The sum of these processes affects particle settling velocity, residence time, and thus the efficiency of organic matter remineralization.

In a companion paper (Rontani et al., 2012), we examined the lipid content of suspended particulate matter (SPM) samples collected in August 2009 in the Mackenzie River and in surface waters of the adjacent Beaufort Sea. Lipid biomarkers, although representing a very minor fraction of the total organic matter (OM), convey important information on the source (terrigenous, marine or bacterial) and degradation state of OM, which is commonly more diagnostic than that provided by bulk parameters (Saliot et al., 2002). Using specific lipid degradation products that have been proposed for

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

sediment and 2.1×10^6 tons yr^{-1} of particulate organic carbon respectively, Macdonald et al., 1998) and the fourth largest in terms of freshwater discharge (3.3×10^{11} $\text{m}^3 \text{yr}^{-1}$, Milliman and Meade, 1983; Brunskill, 1986; Macdonald et al., 1998). The Mackenzie River supplies about 95–99 % of the sediment to the Beaufort Shelf, with coastal erosion and other rivers (Hill et al., 1991; Rachold et al., 2004). The main river load occurs between end of May and the end of August with considerable inter-annual variance (O'Brien et al., 2006). Primary productivity over the Mackenzie Delta/Beaufort Shelf (3.3×10^6 tons yr^{-1} of particulate organic carbon) is mainly due to phytoplanktonic blooms during late spring and summer (Macdonald et al., 1998). Production by ice algae accounts for less than 10 % of the marine production in this area (Horner and Schrader, 1982).

2.2 Sample collection

Time-series sediment traps (Technicap PPS 4/3; 24 cups; cylindrico-conical shape; collecting area: 0.125 m^2) were deployed on 4 mooring lines located in the Amundsen Gulf (CA16, CA05) and on the Mackenzie Shelf (CA10, G09) in the Beaufort Sea (Fig. 1; Table 1). Baffled lids covered the opening of the sediment traps to reduce internal turbulence. Before deployment, sediment traps were thoroughly rinsed with freshwater and seawater following the JGOFS protocol (Knap et al., 1996). Sample cups were filled with filtered seawater (GFF $0.7 \mu\text{m}$) adjusted to 35 PSU with NaCl. Formalin was added to preserve the material collected (5 % v/v , sodium borate buffered).

2.3 Sample treatment

After retrieval, sample cups were checked for salinity and put aside 24 h to allow particles to settle down. Swimmers were removed from the samples then quantitative splitting into several fractions was completed using a McLane Wet Samples Divider or peristaltic pump. Sub-samples for the determination of particulate organic carbon (POC) and lipids and their degradation products were filtered in triplicates

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

through pre-weighted Whatman glass fiber filters (GFF 0.7 μm , 25 mm, combusted 4 h at 450 $^{\circ}\text{C}$). For POC analysis, filters were dried for 12 h at 60 $^{\circ}\text{C}$ and weighed again for dry weight. After exposure for 12 h to concentrated HCl fumes to remove inorganic carbon fraction, the samples were analyzed with a Perkin Elmer CHNS 2600 Series II.

5 Total and POC fluxes were expressed as daily fluxes ($\text{mg C m}^{-2} \text{d}^{-1}$) (Heussner et al., 1990; Lalande et al., 2009). The collected samples were processed in the laboratory according to the method described by Heussner et al. (1990). The total sample was divided into several aliquots to obtain different subsamples for analyzing total mass flux, TOC, lipids and their degradation products. Subsamples were filtered through a
10 precombusted quartz fiber filter (Whatman GF/F, 0.7 μm) under low vacuum.

2.3.1 Lipid extraction

All solvents employed in this study were glass distilled analytical grade. Each filter was extracted four times with $\text{CHCl}_3\text{-MeOH-H}_2\text{O}$ (1 : 2 : 0.8, $v/v/v$) using ultrasonication (separation of particles and solvents by centrifugation at 4000 G for 10 min). To initiate
15 phase separation after ultrasonication, CHCl_3 and purified H_2O were added to the combined extracts to give a final volume ratio of 1 : 1 : 0.9 ($v/v/v$). The upper aqueous phase was extracted twice with CHCl_3 and the combined CHCl_3 extracts were dried over anhydrous Na_2SO_4 , filtered and the solvent removed via rotary evaporation.

2.3.2 Hydroperoxide-reduction

20 NaBH_4 -reduction of the lipid extracts was carried out to reduce labile hydroperoxides resulting from photooxidation and autoxidation to alcohols that are amenable to gas chromatography-electron impact mass spectrometry (GC-EIMS). The filters were put in methanol (15 ml) and hydroperoxides were reduced to the corresponding alcohols by excess NaBH_4 (70 mg) (30 min at room temperature). During this treatment, ketones
25 are also reduced and the possibility of some ester cleavage cannot be excluded.

2.3.3 Alkaline hydrolysis

Metal ions can promote autoxidation during hot saponification procedures, so the prior reduction of hydroperoxides with NaBH₄ allowed us to avoid such autoxidative artifacts during the alkaline hydrolysis. After NaBH₄ reduction, 15 ml of water and 1.7 g of potassium hydroxide were added and the mixture was directly saponified by refluxing for 2 h. After cooling, the content of the flask was acidified with HCl (pH 1) and subsequently extracted three times with dichloromethane. The combined dichloromethane extracts were dried over anhydrous Na₂SO₄, filtered and concentrated to give the total lipid fraction.

2.3.4 Derivatization

After solvent evaporation, residues were taken up in 300 µl of a mixture of pyridine and N,O-bis(trimethylsilyl)trifluoroacetamide (BSTFA; Supelco) (2 : 1, v : v) and silylated for 1 h at 50 °C to convert OH-containing compounds to their TMSi-ether derivatives. After evaporation to dryness under a stream of N₂, the derivatized residues were taken up in a mixture of ethyl acetate and BSTFA (to avoid desilylation of fatty acids) for analysis using GC-EIMS. It should be noted that under these conditions steran-3β,5α,6β-triols were only silylated at the 3 and 6 positions and thus need to be analyzed with great care (Rontani et al., 2012).

2.3.5 Osmium tetroxide oxidation

Double bond positions of monounsaturated fatty acids were determined unambiguously after OsO₄ treatment. A fraction of total lipid extracts and OsO₄ (1 : 2, w : w) were added to a pyridine-dioxane mixture (1 : 8, v / v, 5 ml) and incubated for 1 hour at room temperature. Then, 6 ml of Na₂SO₃ suspension (16 % Na₂SO₃ in water-methanol, 8.5 : 2.5, v / v) was added and the mixture was again incubated for 1.5 h. The resulting mixture was acidified (pH 3) with HCl and extracted three times with DCM (5 ml). The

BGD

9, 7743–7781, 2012

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

combined DCM extracts were subsequently dried over anhydrous Na₂SO₄, filtered and concentrated.

2.4 Gas chromatography-electron impact mass spectrometry (GC-EIMS)

Compounds were identified by comparison of retention times and mass spectra with those of standards and quantified (calibration with external standards) by GC-EIMS. For low concentrations, or in the case of co-elutions, quantification was achieved using selected ion monitoring (SIM). GC-EIMS analyses were carried out with an Agilent 6890 gas chromatograph connected to an Agilent 5973 Inert mass spectrometer. The following conditions were employed: 30 m × 0.25 mm (i.d.) fused silica column coated with HP-1-MS (Agilent; 0.25 μm film thickness); oven temperature programmed in three sequential steps: (i) 70 °C to 130 °C at 20 °C min⁻¹; (ii) 130 °C to 250 °C at 5 °C min⁻¹; and (iii) 250 °C to 300 °C at 3 °C min⁻¹; carrier gas (He) maintained at 0.69 bar until the end of the temperature program and then programmed from 0.69 bar to 1.49 bar at 0.04 bar min⁻¹; injector (on column with retention gap) temperature 50 °C; electron energy 70 eV; source temperature 190 °C; cycle time 1.99 and 8.3 cycles s⁻¹ in SCAN and SIM modes, respectively.

2.5 Lipid degradation products employed for photooxidation, autoxidation and biodegradation estimation

5α(H)-stan-3β-ols, Δ⁴-stera-3β,6α/β-diols and 3β,5α,6β-steratriols (deriving from cholest-5-en-3β-ol (cholesterol), 24-ethylcholest-5-en-3β-ol (sitosterol), 24-methylcholesta-5,22E-dien-3β-ol (brassicasterol) and 24-methylcholesta-5,24(28)-dien-3β-ol (24-methylenecholesterol)) were used to estimate biodegradation, photooxidation and autoxidation state of the different components of the samples, respectively (Fig. 2) (Rontani et al., 2009, 2011; Christodoulou et al., 2009). Abiotic oxidation of the non-specific Δ⁹ monounsaturated fatty acids (oleic and palmitoleic acids) affords 8-*E*, 8-*Z*, 9-*E*, 10-*E*, 11-*E* and 11-*Z* hydroxyacids (Frankel, 1998), which were employed

BGD

9, 7743–7781, 2012

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

to estimate the importance of autoxidation and photooxidation of bulk organic matter (Fig. 2) (Marchand and Rontani, 2001; Rontani et al, 2011). In contrast, abiotic oxidation of *cis*-vaccenic acid (a typical biomarker for Gram-negative bacteria, Sicre et al., 1988, Keweloh and Heipieper, 1996) produces 10-*E*, 10-*Z*, 11-*E*, 12-*E*, 13-*E* and 13-*Z* hydroxyacids, which were useful to estimate autoxidation and photooxidation state of bacteria (Fig. 2) (Rontani et al., 2003; Christodoulou et al., 2010).

2.6 Quantification of hydroperoxides and their ketonic and alcoholic degradation products

A different treatment was employed to quantify hydroperoxides and their ketonic and alcoholic degradation products. The residue obtained after extraction was dissolved in 4 ml of dichloromethane and separated in two equal subsamples. After evaporation of the solvent, degradation products were obtained for the first subsample after acetylation (inducing complete conversion of hydroperoxides to the corresponding ketones, Mihara and Tateba, 1986) and saponification and for the second after reduction with NaBD₄ and saponification. Comparison of the amounts of alcohols present after acetylation and after NaBD₄ reduction made it possible to estimate the proportion of hydroperoxides and alcohols present in the samples, while after NaBD₄-reduction deuterium labelling allowed to estimate the proportion of ketones really present in the samples (Marchand and Rontani, 2003).

Acetylation was carried out in 300 µl of a mixture of pyridine and acetic anhydride (2 : 1, *v/v*), allowed to react at 50 °C overnight and then evaporated to dryness under nitrogen.

BGD

9, 7743–7781, 2012

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results and discussion

3.1 Classical lipid biomarker analyses

3.1.1 Sterols

In the different samples investigated the major sterols were cholesterol (10–31 %) and sitosterol (28–45 %) (Table 2). Although a substantial amount of cholesterol may be derived from diatoms or Prymnesiophycean algae (Volkman, 1986), its dominance suggests an important contribution of zooplanktonic faecal material to these samples. Indeed, it is well known that zooplankton convert much of the sterols produced by algae into cholesterol (Volkman et al., 1980; Prahl et al., 1984). Sitosterol is commonly associated with terrestrial higher plants inputs (Lütjohann, 2004), but some microalgae (and notably diatoms) are now known to produce this sterol (Volkman, 1986, 2003). The other sterols, found in lower range in this study, include 24-nor-cholesta-5,22*E*-dien-3 β -ol, 27-nor-24-methylcholesta-5,22*E*-dien-3 β -ol, cholesta-5,22*E*-dien-3 β -ol, brassicasterol, 24-methylenecholesterol, 24-methylcholest-5-en-3 β -ol (campesterol), 24-ethylcholesta-5,22*E*-dien-3 β -ol (stigmasterol) and 24-ethylcholesta-5,24(28)*E*-dien-3 β -ol (fucosterol) (Table 2). 24-nor-cholesta-5,22-dien-3 β -ol, has been previously identified in a phytoplankton sample composed principally of diatoms (Boutry et al., 1971). The important contribution of diatoms to these samples is also well supported by the presence of brassicasterol and 24-methylenecholesterol, which are major constituents of several diatom species (Lee et al., 1980). However, it may be noted that brassicasterol is also present in some dinoflagellates and in many haptophytes (Volkman, 1986, 2003). If the lack of dinosterol, which is widely accepted as a specific marker of dinoflagellates (Mansour et al., 1999), allowed excluding the presence of such organisms, a significant contribution of haptophytes to the samples cannot be totally excluded. Indeed, it was previously observed that larger haptophytes and diatoms co-dominated in near-surface assemblages of the Beaufort Sea in summer (Hill et al., 2005).

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1.2 Isoprenoids

Dihydrophytol, pristane, 3,7,11,15-tetramethylhexadecanoic (phytanic), 2,6,10,14-tetramethylpentadecanoic (pristanic) and 4,8,12-trimethyltridecanoic (4,8,12-TMTD) acids could be detected in most of the total lipid extracts analysed (Table 3). The presence of relatively high proportions of these degradation products of the chlorophyll phytyl side-chain well supports the contribution of zooplanktonic faecal pellets to the sampling material. Indeed, pelagic crustaceans decrease the abundance of the chlorophyll phytyl side-chain when feeding herbivorously (Prahl et al., 1984; Harvey et al., 1987; Bradshaw et al., 1990). Any phytol remaining in the faeces of pelagic zooplankton after herbivorous feeding will be quickly removed on subsequent reprocessing of the faecal material (Bradshaw and Eglinton, 1993). Several phytol degradation products have been thus identified in zooplanktonic faecal pellets during feeding experiments, including pristane, isomeric pristenes, isomeric phytadienes, dihydrophytol and phytanic, pristanic, 4,8,12-TMTD and isomeric phytenic acids (for a review see Rontani and Volkman, 2003).

3.1.3 Fatty acids and *n*-alkan-1-ols

Total lipid extracts of the different samples exhibited a distribution of even-carbon number dominated fatty acids ranging from C₁₄ to C₂₄ (Table 4), suggesting the presence of a material dominated by marine organisms (plankton and bacteria). Long-chain (>C₂₄) saturated fatty acids, which are characteristic of epicuticular waxes of terrestrial higher plants (Kolattukudy, 1976; Gagosian et al., 1987), could not be detected. The three dominant monounsaturated fatty acids appeared to be hexadec-9*cis*-enoic (palmitoleic), octadec-9*cis*-enoic (oleic) and octadec-11*cis*-enoic (vaccenic) acids (Table 3). Palmitoleic and oleic acids have numerous possible biological origins (plants, fungi, yeasts, bacteria, animals or algae) (Harwood and Russell, 1984), while vaccenic acid is a typical biomarker for Gram-negative bacteria (Sicre et al., 1988; Keweloh and Heipieper, 1996). Small amounts of the very unusual octadec-13-enoic acid could be

BGD

9, 7743–7781, 2012

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

“trapped” in fecal pellets (Najdek et al., 1994) produced by the large herbivorous copepods *Calanus hyperboreus* and *C. glacialis* that undergo diapause.

3.2 Lipid degradation product analyses

3.2.1 Chlorophyll

5 Although the visible light-dependent degradation rate of the chlorophyll tetrapyrrole ring is three to four times higher than for its phytyl side-chain (Cuny et al., 1999), no specific and stable tetrapyrrole photodegradation products could be identified in the literature. Type II photosensitized oxidation (i.e. involving $^1\text{O}_2$) of the phytyl side-chain, however, leads notably to the production of 3-methylidene-7,11,15-trimethylhexadecan-1,2-diol (phytyldiol) (Rontani et al., 1994). Phytyldiol is ubiquitous in the marine environment and constitutes a stable and specific tracer for photodegradation of chlorophyll phytyl side-chain (Rontani et al., 1996; Cuny and Rontani, 1999). Further, the molar ratio phytyldiol:phytol (Chlorophyll Phytyl side-chain Photodegradation Index, CPPI) was proposed to estimate the extent of chlorophyll photodegraded in natural marine sam-
10 ples (Cuny et al., 2002).

15 CPPI ranges from 36 to 121 in the case of the samples collected in summer (Table 3). These values are particularly high when compared with CPPI previously measured in particulate matter collected in summer in the Equatorial Pacific (Rontani et al., 2011) and in the north-western Mediterranean Sea (Cuny et al., 2002) (ranging from 1 to 8 and from 1 to 24, respectively). This attests to the exceptional efficiency of photoox-
20 idation processes in the Arctic Ocean region in summer, most probably because of the midnight sun that persists for 3 months (May–July) at 70° N. On the basis of these very high CPPI values, it could be estimated that during this period chlorophyll was practically entirely photodegraded in sinking particles (Table 3). In the sample CA10
25 A1 collected in October, a strong photodegradation state of chlorophyll (>94 %) was also observed (Table 3).

BGD

9, 7743–7781, 2012

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.2.2 Monounsaturated fatty acids and *n*-alkan-1-ols

Due to the lack of specificity of palmitoleic and oleic acids, their oxidation products have been used to assess abiotic degradation of bulk OM. The results obtained are summarized in Figs. 2–4. Photooxidation percentages of these two acids appeared to be very high (values ranging from 45 to 270% relative to the residual parent compound) in summer, but not in fall (values < 20%). These results, which are in good agreement with the total photodegradation of chlorophyll observed in summer (Table 3), confirm that during this period, photooxidation processes act very intensely on sinking particles of the Beaufort Sea. Autoxidation (free radical oxidation) processes also contributed to the degradation of these two fatty acids (10–30%), but to a lesser extent than light-driven degradation (Figs. 3–5). The strong spatial variability in the photooxidation stress (Figs. 3–5) could likely be attributed to differences in water clarity at the different sampling stations that typically increases from a shelf-edge location (CA05), to a mid-slope area (CA16), up to a basin-close environment (G09) (Table 1). Alternatively, visual inspection of sediment trap samples revealed that particles collected at G09 were generally finer and less aggregated than at CA16 and CA05, with the latter being obviously affected by the sinking of large diatom colonies (A. Forest, personal observation). Organic matter contained in large aggregates could then be relatively more protected against photooxidation than in fine particles that might offer a high surface-to-volume ratio.

Oxidation products of vaccenic acid allowed us to estimate photo- and autoxidation state of heterotrophic bacteria associated to sinking particles. These bacteria were also strongly photodegraded in summer (photodegradation percentage ranging from 45 to 260%) (Figs. 3–5) and weakly in fall (photodegradation percentage < 10%). During the summer period, transfer of singlet oxygen from senescent phytoplanktonic cells to bacteria (Rontani et al., 2003; Christodoulou et al., 2010) seems thus to have been especially efficient. Vaccenic acid also appeared to be affected by autoxidation but less intensively (Figs. 3–5). Indeed, reaction of singlet oxygen with unsaturated components

BGD

9, 7743–7781, 2012

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of the outer lipopolysaccharide membrane of Gram negative bacteria (the dominant bacteria in the ocean) leads to the formation of reactive secondary products, such as peroxy radicals, which may in turn accentuate cell death (Dahl et al., 1989). The intense oxidative stress resulting from singlet oxygen damages in bacteria should limit their growth (and thus biodegradation processes) during the settling of sinking particles.

Important amounts of oxidation products of C_{20:1Δ11} and C_{22:1Δ11} alkan-1-ols (values ranging from 10 to 800% of the residual parent compound), which are specific components of zooplankton wax esters (Lee and Nevenzel, 1979; Albers et al., 1996), could be also detected in the different samples (Figs. 3–5). The major part of these compounds results from the involvement of Type II (i.e. involving singlet oxygen) photoprocesses (Fig. 6). It is important to note that these oxidation products disappeared when the alkaline hydrolysis step was avoided during the treatment. These results clearly showed that photooxidation processes acted directly on wax esters and not on the corresponding *n*-alkan-1-ol after enzymatic hydrolysis. The high efficiency of Type II photooxidation processes in such micro-environment may be attributed to: (i) the high concentration of wax esters in the droplets trapped in faecal pellets (as discussed above) favoring the likelihood of interaction between singlet oxygen (produced from chlorophyll and phaeopigments contained in the pellets) and their double bonds and (ii) the apolar character of these droplets. Indeed, the lifetime of singlet oxygen in apolar environments is longer, and its potential diffusion distance greater, than under polar conditions (Suwa et al., 1977). To our knowledge, this is the first in situ demonstration of photodegradation of zooplanktonic faecal material.

Allylic hydroperoxides resulting from photo- and autoxidation of monounsaturated fatty acids may undergo: (i) heterolytic cleavage catalyzed by protons (Frimer, 1979) leading to the formation of ω-oxocarboxylic acids and other volatile products and (ii) homolytic cleavage induced by transition metal ions (Pokorny, 1987; Schaich, 2005) or UVR (since hydroperoxides absorb in the UVR range; Horspool and Armesto, 1992). Homolytic cleavage of hydroperoxyacids would lead to the formation of alkoxy radicals,

BGD

9, 7743–7781, 2012

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are generally lower for Δ^5 -sterols than for chlorophyll phytol side-chain and monounsaturated fatty acids (Rontani et al., 1998), possibly due to steric hindrance during the attack of the sterol Δ^5 double bond by $^1\text{O}_2$ (Beutner et al., 2000). Photodegradation processes acted more intensively on 24-methylenecholesterol (mainly arising from diatoms) (photodegradation percentage ranging from 40 to 81 % relative to the parent sterol) than on brassicasterol (arising from diatoms and/or Prymnesiophytes) (values ranging from 18 to 33 %) (Fig. 8). These differences suggest a higher efficiency of photodegradation processes in diatoms than in Prymnesiophytes. The similarity observed between the overall behaviors for brassicasterol and sitosterol (Fig. 8) with respect to degradation well supports a major contribution of Prymnesiophytes to sitosterol. Interestingly, the sitosterol and campesterol contained in suspended particles collected in this zone (mainly arising from terrestrial higher plants) were strongly autoxidized (Rontani et al., 2012). The lack of sitosterol autoxidation products in all the total lipid extracts obtained from sediment trap samples thus confirms that terrestrial higher plant material does not contribute significantly to this sterol in sinking particles. A release of sub-Arctic terrestrial POM in two different pools was recently proposed (Vonk et al., 2010a, b). A pool composed of mineral-bound POC derived from erosion, which has short initial residence times in the surface water and quickly settle to the sea floor and another pool composed of higher plants debris mainly contributing to suspended particulate matter. These two pools probably settle too quickly or too slowly, respectively, to contribute significantly to the material collected by the different traps deployed.

Low proportions of 5α -stanols corresponding to 24-methylenecholesterol and brassicasterol could be observed in the different samples (values ranging from 3 to 15 % of the corresponding sterol) (Fig. 8). These values, which are very close to these generally considered as typical of healthy phytoplanktonic cells (5–10 %, Wakeham et al., 1997), suggest that biodegradation processes acted only very weakly on phytoplanktonic material. The lack of the corresponding ster-4-en-3-ones, which are classical bacterial metabolites of Δ^5 -sterols (de Leeuw and Baas, 1986; Wakeham, 1989) often detected in sinking particles (Bayona et al., 1989; Christodoulou et al., 2009), well supports this

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

assumption. This apparently good resistance of phytodetritus against bacterial degradation might result from the inhibition of bacterial growth when $^1\text{O}_2$ generated by the photolysis of senescent phytoplanktonic cells in the euphotic zone is efficiently transferred to attached bacteria, as it was observed in the previous section.

As in the case of monounsaturated fatty acids, the specific proportion of hydroperoxysterols and their alcoholic and ketonic degradation products have been also determined. The results obtained are summarized in Fig. 7b. Hydroperoxysterols appear to be more stable than hydroperoxyacids in sinking particles (Fig. 7a); significant amounts of these compounds should thus reach the sediments. This observation is in good agreement with the previous detection of intact hydroperoxysterols in recent surficial sediments (Rontani and Marchand, 2000). While ratio $\Delta^4\text{-}6\alpha/6\beta\text{-hydroperoxides} / \Delta^5\text{-}7\alpha/7\beta\text{-hydroperoxysterols}$ produced during irradiation of senescent phytoplanktonic cells ranged from 0.30 to 0.35 (Rontani et al., 1997), very high values (ranging from 0.4 to 2.55) of the ratio cholest-4-en- $3\beta,6\alpha/6\beta\text{-diols} / \text{cholest-5-en-}3\beta,7\alpha/7\beta\text{-diols}$ were previously measured (after NaBH_4 -reduction) in sediment trap samples collected in Mediterranean Sea (Christodoulou et al., 2009). These high values were attributed to a faster degradation of $\Delta^5\text{-}7\text{-hydroperoxysterols}$ than $\Delta^4\text{-}6\text{-hydroperoxysterols}$ during settling through the water column (Christodoulou et al., 2009). According to the theoretical stability of the alkyl radicals formed during β -scission of the corresponding alkoxy radicals, the following order of stability was tentatively proposed by Christodoulou et al. (2009): $\Delta^4\text{-}6\text{-hydroperoxysterols} > \Delta^5\text{-}7\text{-hydroperoxysterols} > \Delta^6\text{-}5\text{-hydroperoxysterols}$. The results obtained in the present study (Fig. 7b) well support this assumption.

4 Conclusions

Lipids and their degradation products were quantified in seven samples of sinking particles collected with sediment traps in summer and fall across the Canadian Beaufort Shelf. These samples were dominated by diatoms and zooplanktonic faecal pellets.

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Terrestrial higher plants resulting from Mackenzie River discharge did not contribute significantly to the sinking material. During the summer period, Type II (i.e. involving singlet oxygen) photooxidation processes acted strongly on senescent phytoplanktonic cells, heterotrophic bacteria and zooplanktonic faecal material. Diatoms, which dominated the phytoplanktonic assemblage, appeared to be remarkably sensitive to photodegradation. Singlet oxygen transfer from phytodetritus to attached bacteria was particularly efficient inducing strong oxidative damages in these heterotrophic organisms. The presence of high amounts of photoproducts of $C_{20:1\Delta11}$ and $C_{22:1\Delta11}$ alcohols, which are specific components of wax esters found in herbivorous copepods (Lee and Nevenzel, 1979; Albers et al., 1996), allowed to demonstrate for the first time the high efficiency of Type II photodegradation processes in zooplanktonic faecal material.

In contrast, phytoplanktonic cells seemed to be relatively preserved towards biodegradation processes in sinking POM. As proposed by Rontani et al. (2011), there is a synergy between senescing phytoplanktonic cells and attached bacteria and between photooxidation and biodegradation. Photolysis of chlorophyll in senescing algal cells produces singlet oxygen, which if transferred from algal cells to attached bacteria may inhibit bacterial growth and reduce the extent of heterotrophic degradation. Therefore, it seems that there is a direct link between the photooxidation state of lipids of senescent phytoplanktonic cells in particles and their resistance towards biotic degradation. However, this does not exclude that the flux of labile dissolved organic matter produced through the photo-cleavage of phytodetritus might sustain an active community of free-living bacteria around sinking particles.

Finally, we demonstrated that strongly photodegraded sinking particles contained an important amount of intact hydroperoxides. After sedimentation, these compounds should induce a strong oxidative stress in surface sediments, which could be at the origin of the formation of epoxyacids and monounsaturated fatty acids with a *trans* double bonds previously detected in unusual high proportion in this zone (Rontani et al., 2012).

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Boutry, J., Alcaide, A., and Barbier, M.: De la présence d'un stérol en C₂₆ d'un plancton marin végétal, C. R. Acad. Sci., Paris, 272, 1022–1023, 1971.
- Bradshaw, S. A. and Eglinton, G.: Marine invertebrate feeding and the sedimentary lipid record, in: Organic Geochemistry, edited by: Engel, M. H. and Macko, S. A., Plenum Press, New York, 225–235, 1993.
- Bradshaw, S. A., O'Hara, S. C. M., Corner, E. D. S., and Eglinton, G.: Changes in lipids during simulated herbivorous feeding by the marine crustacean *Neomysis integer*, J. Mar. Biol. Assoc. UK, 70, 225–243, 1990.
- Brunskill, G. J.: Environmental features of the Mackenzie system, in: The ecology of river systems, edited by: Davis, B. R. and Walker, W. F., Dordrecht, The Netherlands, Dr. W. Junk Publications, 435–471, 1986.
- Christodoulou, S., Marty, J.-C., Miquel, J.-C., Volkman, J. K., and Rontani, J.-F.: Use of lipids and their degradation products as biomarkers for carbon cycling in the northwestern Mediterranean Sea, Mar. Chem., 113, 25–40, 2009.
- Christodoulou, S., Joux, F., Marty, J.-C., Sempéré, R., and Rontani, J.-F.: Comparative study of UV and visible light induced degradation of lipids in non-axenic senescent cells of *Emiliania huxleyi*, Mar. Chem., 119, 139–152, 2010.
- Cuny, P. and Rontani, J.-F.: On the widespread occurrence of 3-methylidene-7,11,15-trimethylhexadecan-1,2-diol in the marine environment: a specific isoprenoid marker of chlorophyll photodegradation, Mar. Chem., 65, 155–165, 1999.
- Cuny, P., Romano, J.-C., Beker, B., and Rontani, J.-F.: Comparison of the photo degradation rates of chlorophyll chlorin ring and phytol side chain in phytodetritus: is the phytyldiol versus phytol ratio (CPPI) a new biogeochemical index?, J. Exp. Mar. Biol. Ecol., 237, 271–290, 1999.
- Cuny, P., Marty, J.-C., Chiaverini, J., Vescovali, I., Raphel, D., and Rontani, J.-F.: One-year seasonal survey of the chlorophyll photodegradation process in the Northwestern Mediterranean sea, Deep-Sea Res., 49, 1987–2005, 2002.
- Dahl, T. A., Midden, W. R., and Hartman, P. E.: Comparison of killing of Gram-negative and Gram-positive bacteria by pure singlet oxygen, J. Bacteriol., 171, 2188–2194, 1989.
- De Leeuw, J. W. and Baas, M.: Early stage diagenesis of steroids, in: Biological Markers in the Sedimentary Record, edited by: Johns, R. B., Elsevier, 101–123, 1986.
- Forest, A., Stemmann, L., Picheral, M., Burdorf, L., Robert, D., Fortier, L., and Babin, M.: Size distribution of particles and zooplankton across the shelf-basin system in southeast Beaufort

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Sea: combined results from an Underwater Vision Profiler and vertical net tows, *Biogeosciences*, 9, 1301–1320, doi:10.5194/bg-9-1301-2012, 2012.

Frankel, E. N.: *Lipid Oxidation*, the Oily Press, Dundee, 1–303, 1998.

Frey, K. E. and McClelland, J. W.: Impacts of permafrost degradation on Arctic river biogeochemistry, *Hydrol. Process.*, 23, 169–182, 2009.

Frimer, A. A.: The reaction of singlet oxygen with olefins: the question of mechanism, *Chem. Rev.*, 79, 359–387, 1979.

Gagosian, R. B., Peltzer, E. T., and Merrill, J. T.: Long-range transport of terrestrially derived lipids in aerosols from the south Pacific, *Nature*, 325, 800–803, 1987.

Graeve, M., Kattner, G., and Hagen, W.: Diet-induced changes in the fatty acid composition of Arctic herbivorous copepods, *J. Exp. Mar. Biol. Ecol.*, 182, 97–110, 1994.

Harvey, H. R., Eglinton, G., O'Hara, S. C. M., and Corner, E. D. S.: Biotransformation and assimilation of dietary lipids by *Calanus* feeding on a dinoflagellate, *Geochim. Cosmochim. Ac.*, 51, 3031–3040, 1987.

Harwood, J. L. and Russell, N. L.: *Lipids in Plants and Microbes*, George Allen and Unwin, London, 1–162, 1984.

Heussner, S., Ratti, C., and Carbonne, J.: The PPS 3 time series sediment traps and the trap sample processing techniques used during the ECOMARGE experiment, *Cont. Shelf Res.*, 10, 943–958, 1990.

Hill, P. S.: Controls on floc size in the sea, *Oceanogr.*, 11, 13–18, 1998.

Hill, P. R., Blasco, S. M., Harper, J. R., and Fissel, D. B.: Sedimentation on the Canadian Beaufort Shelf, *Cont. Shelf Res.*, 11, 821–842, 1991.

Hill, V., Cota, G., and Stockwell, D.: Spring and summer phytoplankton communities in the Chukchi and eastern Beaufort Seas, *Deep-Sea Res.*, 52, 3369–3385, 2005.

Horner, R. A. and Schrader, G. C.: Relative contributions of ice algae, phytoplankton and benthic microalgae to primary production in nearshore regions of the Beaufort Sea, Arctic, 35, 485–503, 1982.

Horspool, W. and Armesto, D.: *Organic Photochemistry: A Comprehensive Treatment*, Ellis Horwood, Chichester, UK, 1–521, 1992.

Kawamura, K. and Gagosian, R. B.: Implication of ω -oxocarboxylic acids in the remote marine atmosphere for photo-oxidation of unsaturated fatty acids, *Nature*, 325, 330–332, 1987.

Keweloh, H. and Heipieper, H. J.: Trans-unsaturated fatty acids in bacteria, *Lipids*, 31, 129–137, 1996.

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Knap, A., Michaels, A., Close, A., Ducklow, H., and Dickson, A.: Protocols for the Joint Global Ocean Flux Study (JGOFS) core measurements, JGOFS Report, 19, 155–162, 1996.
- Kolattukudy, P. E.: Lipid polymers and associated phenols, their chemistry, biosynthesis, and role in pathogenesis, *Recent Adv. Phytochem.*, 77, 185–246, 1977.
- 5 Lalande, C., Bélanger, S., and Fortier, L.: Impact of a decreasing sea ice cover on the vertical export of particulate organic carbon in the northern Laptev Sea, Siberian Arctic Ocean, *Geophys. Res. Lett.*, 36, L21604, doi:10.1029/2009GL040570, 2009.
- Lee, R. F. and Nevenzel, J. C.: Wax esters in the marine environment: origin and composition of the wax from the Bute Inlet, British Columbia, *J. Fish. Res. Board Can.*, 36, 1519–1523, 1979.
- 10 Lee, C., Gagosian, R. B., and Farrington, J. W.: Geochemistry of sterols in sediments from Black Sea and the southwest African shelf and slope, *Org. Geochem.*, 2, 103–113, 1980.
- Lee, R. F., Hagen, W., and Kattner, G.: Lipid storage in marine zooplankton, *Mar. Ecol. Prog. Ser.*, 307, 273–306, 2006.
- 15 Lütjohann, D.: Sterol autoxidation: from phytosterols to oxyphytosterols, *Brit. J. Nutr.*, 91, 3–4, 2004.
- McCave, I. N.: Size spectra and aggregation of suspended particles in the ocean, *Deep-Sea Res.*, 31, 329–352, 1984.
- Macdonald, R. W., Solomon, S. M., Cranston, R. E., Welch, H. E., Yunker, M. B., and Gobeil, C.: A sediment and organic carbon budget for the Canadian Beaufort Shelf, *Mar. Geol.*, 144, 255–273, 1998.
- 20 Mansour, M. P., Volkman, J. K., Jackson, A. E., and Blackburn, S. I.: The fatty acid and sterol composition of five marine dinoflagellates, *J. Phycol.*, 35, 710–720, 1999.
- Marchand, D. and Rontani, J.-F.: Characterisation of photooxidation and autoxidation products of phytoplanktonic monounsaturated fatty acids in particulate matter and recent sediments, *Org. Geochem.*, 32, 287–304, 2001.
- 25 Marchand, D. and Rontani, J.-F.: Visible light-induced oxidation of lipid components of purple sulphur bacteria: a significant process in microbial mats, *Org. Geochem.*, 34, 61–79, 2003.
- Mihara, S. and Tateba, H.: Photosensitized oxygenation reactions of phytol and its derivatives, *J. Org. Chem.*, 51, 1142–1144, 1986.
- 30 Milliman, J. D. and Meade, R. H.: World-wide delivery of river sediment to the oceans, *J. Geol.*, 91, 1–21, 1983.

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Najdek, M., Puskaric, S., and Bochdansky, A.: Contribution of zooplankton lipids to the flux of organic matter in the northern Adriatic Sea, *Mar. Ecol.-Prog. Ser.*, 111, 241–249, 1994.
- O'Brien, M. C., Macdonald, R. W., Melling, H., and Iseki, K.: Particle fluxes and geochemistry on the Canadian Beaufort Shelf: implications for sediment transport and deposition, *Cont. Shelf Res.*, 26, 41–81, 2006.
- Pokorny, J.: Major factors affecting the autoxidation of lipids, in: *Autoxidation of unsaturated lipids*, edited by: Chan, H. W.-S., Academic Press, 141–206, 1987.
- Prahl, F. G., Eglinton, G., Corner, E. D. S., O'Hara, S. C. M., and Forsberg, T. E. V.: Changes in plant lipids during passage through the gut of *Calanus*, *J. Mar. Biol. Assoc. UK*, 64, 317–334, 1984.
- Rachold, V., Eicken, H., Gordeev, V. V., Grigoriev, M. N., Hubberten, H. W., Lisitzin, A. P., Shevchenko, V. P., and Schirrmeister, L.: Modern terrigenous organic carbon input to the Arctic Ocean, in: *The Organic Carbon Cycle in the Arctic Ocean*, edited by: Stein, R. S. and Macdonald, R. W., Springer, New York, 33–55, 2004.
- Rontani, J.-F. and Marchand, D.: D⁵-Stenol photoproducts of phytoplanktonic origin: a potential source of hydroperoxides in marine sediments?, *Org. Geochem.*, 31, 169–180, 2000.
- Rontani, J.-F. and Volkman, J. K.: Phytol degradation products as biogeochemical tracers in aquatic environments, *Org. Geochem.*, 34, 1–35, 2003.
- Rontani, J.-F., Grossi, V., Faure, R., and Aubert, C.: “Bound” 3-methylidene-7,11,15-trimethylhexadecan-1,2-diol: a new isoprenoid marker for the photodegradation of chlorophyll-a in seawater, *Org. Geochem.*, 21, 135–142, 1994.
- Rontani, J.-F., Raphel, D., and Cuny, P.: Early diagenesis of the intact and photooxidized chlorophyll phytol chain in a recent temperate sediment, *Org. Geochem.*, 24, 825–832, 1996.
- Rontani, J.-F., Cuny, P., and Aubert, C.: Rates and mechanism of light-dependent degradation of sterols in senescent cells of phytoplankton, *J. Photochem. Photobiol. A: Chem.*, 111, 139–144, 1997.
- Rontani, J.-F., Cuny, P., and Grossi, V.: Identification of a pool of lipid photoproducts in senescent phytoplanktonic cells, *Org. Geochem.*, 29, 1215–1225, 1998.
- Rontani, J.-F., Koblizek, M., Beker, B., Bonin, P., and Kolber, Z.: On the origin of *cis*-vaccenic photodegradation products in the marine environment, *Lipids*, 38, 1085–1092, 2003.
- Rontani, J.-F., Zabeti, N., and Wakeham, S. G.: The fate of marine lipids: Biotic vs. abiotic degradation of particulate sterols and alkenones in the Northwestern Mediterranean Sea, *Mar. Chem.*, 113, 9–18, 2009.

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Rontani, J.-F., Zabeti, N., and Wakeham, S. G.: Degradation of particulate organic matter in the equatorial Pacific Ocean: biotic or abiotic?, *Limnol. Oceanogr.*, 56, 333–349, 2011.
- Rontani, J.-F., Charriere, B., Petit, M., Vaultier, F., Heipieper, H. J., Link, H., Chaillou, G., and Sempéré, R.: Degradation state of organic matter in surface sediments from the Beaufort Shelf: a lipid approach, *Biogeosciences Discuss.*, 9, 3881–3916, doi:10.5194/bgd-9-3881-2012, 2012.
- Rontani, J.-F., Charriere, B., Sempéré, R., Doxaran, D., Vaultier, F., Vonk, J. E., and Volkman, J. K.: Lipid biomarker evidence for extensive biotic and abiotic degradation of terrestrial organic matter in Arctic coastal waters, *Geochim. Cosmochim. Ac.*, submitted, 2012.
- Salot, A., Parrish, C. C., Sadouni, N., Bouloubassi, I., Fillaux, J., and Cauwet, G.: Transport and fate of Danube Delta terrestrial organic matter in the Northwest Black Sea mixing zone, *Mar. Chem.*, 79, 243–259, 2002.
- Schaich, K. M.: Lipid oxidation: theoretical aspects, in: *Bailey's Industrial Oil and Fat Products*, Sixth edition, edited by: Shahidi, F., John Wiley & Sons, 269–355, 2005.
- Sicre, M.-A., Paillasseur, J. L., Marty, J.-C., and Salot, A.: Characterization of seawater samples using chemometric methods applied to biomarker fatty acids, *Org. Geochem.*, 12, 281–288, 1988.
- Suwa, K., Kimura, T., and Schaap, A. P.: Reactivity of singlet molecular oxygen with cholesterol in a phospholipidic membrane matrix: a model for oxidative damage of membranes, *Biochem. Bioph. Res. Co.*, 75, 785–792, 1977.
- Vallières, C., Retamal, L., Ramlal, P., Osburn, C. L., and Vincent, W. F.: Bacterial production and microbial food web structure in a large Arctic river and the coastal Arctic Ocean, *J. Mar. Syst.*, 74, 756–773, doi:10.1016/j.jmarsys.2007.12.002, 2008.
- Volkman, J. K.: A review of sterol markers for marine and terrigenous organic matter, *Org. Geochem.*, 9, 83–99, 1986.
- Volkman, J. K.: Sterols in microorganisms, *Appl. Microbiol. Biotechnol.*, 60, 495–506, 2003.
- Volkman, J. K., Corner, E. D. S., and Eglinton, G.: Transformations of biolipids in the marine food web and in underlying bottom sediments, *Colloq. Int. Cent. Nat. Rech. Sci.*, 293, 185–197, 1980.
- Vonk, J. E., van Dongen, B. E., and Gustafsson, O.: Selective preservation of old organic carbon fluviually released from sub-Arctic soils, *Geophys. Res. Lett.*, 37, L11605, doi:10.1029/2010gl042909, 2010a.

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Vonk, J. E., Sánchez-García, L., Semiletov, I., Dudarev, O., Eglinton, T., Andersson, A., and Gustafsson, Ö.: Molecular and radiocarbon constraints on sources and degradation of terrestrial organic carbon along the Kolyma paleoriver transect, East Siberian Sea, *Biogeochemistry*, 7, 3153–3166, doi:10.5194/bg-7-3153-2010, 2010b.
- 5 Wakeham, S. G.: Reduction of stenols to stanols in particulate matter at oxic-anoxic boundaries in seawater, *Nature*, 342, 787–790, 1989.
- Wakeham, S. G. and Lee, C.: Organic geochemistry of particulate matter in the ocean: The role of particles in oceanic sedimentary cycles, *Org. Geochem.*, 14, 83–96, 1989.
- Wakeham, S. G., Hedges, J. I., Lee, C., Peterson, M. L., and Hernes, P. J.: Compositions and
 10 transport of lipid biomarkers through the water column and surficial sediments of the EqPac Ocean, *Deep-Sea Res.*, 44, 2131–2162, 1997.
- Ward, P. F. V., Scott, T. W., and Dawson, R. M. C.: The hydrogenation of unsaturated fatty acids in the ovine digestive tract, *Biochem. J.*, 92, 60–68, 1964.
- Wassmann, P. and Reigstad, M.: Future Arctic Ocean seasonal ice zones and implications
 15 for pelagic-benthic coupling, *Oceanography*, 24, 220–231, doi:10.5670/oceanog.2011.74, 2011.
- Wassmann, P., Bauerfeind, E., Fortier, M., Fukuchi, M., Hargrave, B., Honjo, S., Moran, B., Noji, T., Nöthig, E. M, and Peinert, R.: Particulate organic carbon flux to the sea floor, in: *The Organic Carbon Cycle in the Arctic Ocean*, edited by: Stein, R. and MacDonald, R., Springer-Verlag, New York, 101–138, 2004.
- 20

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Table 1. Mooring location and deployment information.

Mooring	Location	Latitude	Longitude	Deployment period	Water depth (m)	Trap depth (m)
CA05 A22	Amundsen Gulf	71.31245	127.58237	2–7 August 2009	204	97.3
CA05 A23	Amundsen Gulf	71.31245	127.58237	8–15 August 2009	204	97.3
CA05 A24	Amundsen Gulf	71.31245	127.58237	16–31 August 2009	204	97.3
CA16 A23	Amundsen Gulf	71.78677	126.49695	8–15 August 2009	314	110.2
G09 A2	Mackenzie Shelf	71.00254	135.47930	31 July–14 August 2009	702	100.5
G09 A3	Mackenzie Shelf	71.00254	135.47930	15–31 August 2009	702	100.5
CA10 A1	Mackenzie Shelf	69.95512	138.67373	7–31 October 2003	250	105

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Table 3. Chlorophyll phytyl side-chain degradation products detected in the different samples and CPPI-based estimates of chlorophyll photodegradation.

Mooring	4,8,12-TMTD acid (%) [*]	Pristanic acid (%) [*]	Phytanic acid (%) [*]	Dihydrophytol (%) [*]	Pristane (%) [*]	CPPI photodegradation %	Chlorophyll
CA05-A22	18.8	7.9	69.0	8.1	0.4	40.0 ± 2.5	99.8 ± 0.1
CA05-A23	41.6	19.4	108.9	5.8	1.9	53.3 ± 0.4	100
CA05-A24	66.1	39.2	72.7	11.5	3.1	35.6 ± 3.7	99.6 ± 0.2
CA16-A23	55.0	27.4	87.3	15.9	2.4	104.1 ± 0.1	100
G09-A2	37.2	27.9	218.9	5.4	–	120.9 ± 41.7	100
G09-A3	37.6	10.1	110.5	6.7	14.2	36.7 ± 4.9	99.6 ± 0.2
CA10-A1	–	–	120.0	–	–	22.7 ± 9.9	94.3 ± 5.1

^{*} Relative to phytol.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4. Flux ($\mu\text{g m}^{-2} \text{d}^{-1}$) of fatty acids and *n*-alkan-1-ols.

Compounds	CA05-A22	CA05-A23	CA05-A24	CA16-A23	CA10-A1	G09-A2	G09-A3
Fatty acids							
C _{14:0}	227.2	802.0	317.9	523.6	16.5	275.3	404.2
C _{15:0}	35.4	99.2	86.3	160.3	19.1	32.8	85.3
C _{16:1Δ9} (Palmitoleic)	465.3	1066.5	306.8	570.9	81.8	457.2	507.7
C _{16:0}	971.0	2907.3	1024.7	4572.9	506.6	1581.8	1443.4
C _{17:1}	–	21.8	38.5	–	11.4	–	–
C _{17:0}	14.1	39.5	36.3	95.3	15.0	23.2	–
C _{18:1Δ9} (Oleic)	170.4	438.2	248.3	442.8	183.1	181.6	353.1
C _{18:1Δ11} (Vaccenic)	73.4	310.5	114.3	183.8	55.1	82.2	167.8
C _{18:1Δ13}	12.0	88.1	34.7	62.0	15.6	18.2	41.7
C _{18:0}	368.6	992.1	172.5	1999.0	131.0	451.1	260.6
C _{20:1Δ11}	34.0	123.3	79.8	217.5	68.9	–	–
C _{20:0}	6.0	17.4	10.5	49.8	6.7	12.0	9.2
C _{22:1Δ11}	5.7	23.7	16.1	28.6	3.8	5.6	21.7
C _{22:0}	4.7	12.4	4.2	31.4	3.2	6.9	6.6
C _{24:1Δ11}	3.3	37.7	18.3	14.3	2.0	5.7	26.3
C _{24:0}	2.7	16.4	5.8	41.3	4.9	7.2	5.7
<i>n</i>-Alkan-1-ols							
C _{14:0}	5.3	105.2	73.9	94.3	15.9	24.6	236.2
C _{16:1}	7.8	81.3	46.1	61.5	5.2	29.1	109.8
C _{16:0}	58.9	357.1	273.8	408.1	32.1	122.6	471.0
C _{18:0}	25.1	117.4	46.0	257.3	23.1	62.0	81.7
C _{20:1Δ11}	122.0	204.6	186.2	111.6	66.0	20.4	130.7
C _{22:1Δ11}	605.9	500.4	217.5	120.9	43.8	20.7	213.7

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. Map of the studied area with locations of the different stations investigated.

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.




Fig. 2. Formulae, vernacular names and potential applications of the main (non exhaustive list) lipid tracers of degradation processes employed in the present work. ¹ Quantified after NaBH₄-reduction of hydroperoxides to the corresponding alcohols and subsequent silylation.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.





Fig. 3. Percentages of photooxidation and autoxidation products (relative to the residual parent compound) of monounsaturated fatty acids and alkan-1-ols observed in G09-A2 (**A**) and G09-A3 (**B**) samples.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Percentages of photooxidation and autoxidation products (relative to the residual parent compound) of monounsaturated fatty acids and alkan-1-ols observed in CA05-A22 **(A)**, CA05-A23 **(B)** and CA05-A24 **(C)** samples.




A**B**

Fig. 5. Percentages of photooxidation and autoxidation products (relative to the residual parent compound) of monounsaturated fatty acids and alkan-1-ols observed in CA16-A23 (**A**) and CA10-A1 (**B**) samples.



R = Polyunsaturated alkyl chain

R' = CH₃- or CH₃-(CH₂)₂-

R'' = Highly photooxidized polyunsaturated alkyl chain

Fig. 6. Type II (i.e. involving singlet oxygen) photooxidation of C_{20:1Δ11} and C_{22:1Δ11} alkan-1-ols.

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.




Fig. 7. Relative percentages of intact hydroperoxides and their ketonic and alcoholic degradation products measured in the case of monounsaturated fatty acids and alkan-1-ols **(A)** and cholesterol **(B)** oxidation products.

Degradation of planktonic and bacterial lipids in sinking particles

J.-F. Rontani et al.




Fig. 8. Percentages of photooxidation and autoxidation products (relative to the residual parent compound) of sitosterol, cholesterol, brassicasterol and 24-methylenecholesterol in the different samples investigated.