

Interactive
Comment

Interactive comment on “Production of oceanic nitrous oxide by ammonia-oxidizing archaea” by C. R. Loescher et al.

C. R. Loescher et al.

cloescher@ifm-geomar.de

Received and published: 22 May 2012

Biogeosciences Discuss., 9, C770–C771, 2012 www.biogeosciences-discuss.net/9/C770/2012/ © Author(s) 2012. This work is distributed under the Creative Commons Attribute 3.0 License. Biogeosciences Discussions Interactive comment on “Production of oceanic nitrous oxide by ammonia-oxidizing archaea” by C. R. Loescher et al.

Anonymous Referee #3 Received and published: 24 April 2012

Referee 3 This manuscript describes a series of field observations, manipulations and culture experiments which are performed to investigate which group of organisms, ammonia oxidizing archaea or bacteria are the dominant producers of nitrous oxide in the

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



marine environment. This study is very timely, and following a small number of edits is entirely appropriate for publication in Biogeosciences.

We thank the anonymous reviewer #3 for the comments on our manuscript and moreover for considering our study as timely, well presented, and appropriate for publication in Biogeosciences after careful revision. We considered the comments during revision and believe we have thereby further improved the manuscript.

Q1: The manuscript is generally well presented, though I would suggest that the position of the methods section is wrong and should be re-positioned after the introduction. The present arrangement means that one is consistently looking forward to find out what, where and why something has been done.

R1: We agree with the reviewer and shifted the methods section following the introduction.

Q2: I would also suggest that the current title is not entirely representative and should be altered to reflect the coastal and shelf seas component of this study.

R2: The expression 'oceanic' represents for us a combination of coastal, shelf and open ocean regions. We thus consider the chosen title as adequate as our study includes oceanic regions reaching from the coast (with water depth below 150m in the ETNA and 100m in the ETSP) over the shelf region out to the open ocean (with water depth over 1000m in the ETNA and ETSP), thus we think the title is appropriate and like to keep the title.

Q3: In Section 2 – vertical distribution. . . I find that this discussion is not particularly easy to follow and a better approach might be to separate out the description of the two ocean areas.

R3: We presented the results and discussion section in a combined way as we considered it most focused. We chose this structure to have the possibility to directly compare the two different OMZs with regard to the strongly differing N₂O and O₂ conditions. If

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

separating the description of the two areas, this paragraph would contain too many repetitive statements. Thus we like to keep the combined comparative description.

Q4: Further to this the interrogation of the relationship between numbers of amoA genes and N₂O should be more rigorous than a simple comparison of two contour profiles, which do not actually match up as well as is described. A correlative relationship does not prove a direct link, but some statistical investigation should be performed here.

R4: We agree with the referee. Regression analysis have been applied on the overall dataset of N₂O and number of amoA genes in the ETNA resulted in a Pearson correlation coefficient of $r = 0.63$ indicating a linear correlation of the two variables. See also response to reviewer 2 question 1.

This information has now been included into the revised manuscript (Vertical distribution. . .): ‘A comparable pattern of the distribution of archaeal amoA genes and N₂O was observed in the water column of the ETNA (Fig. 3) suggesting a correlation between AOA abundance and N₂O accumulation (Pearson correlation coefficient $r = 0.63$; statistical significance is indicated) in the layers with low O₂ (Fig. 4). . .

Q5: The description of “certain depths at some stations” is very vague and this should be tightened up, I can not tell from this whether the “key genes” for denitrification and anammox were determined in the Pacific study. The lack of a relationship between deltaN₂O and AOU in the Pacific study merits further discussion.

R5: We agree that the expression ‘certain depth at some stations’ is too vague and rearranged the sentence:

‘A co-occurrence of N₂O and archaeal amoA genes was detected at certain depths, e.g. at profile V at 100m water depth (Fig.1) in the ETSP, but was not a general feature possibly resulting from additional N₂O production via other processes such as denitrification, nitrifier-denitrification or anammox (Kartal et al., 2007) at present suboxic

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

conditions (see Fig. 1).‘

Key genes for denitrification and anammox were assayed in the Atlantic and Pacific studies; however, they were only detectable in the ETSP. The respective sentence was rearranged to clarify, that the respective genes were present and that most likely mixed processes contribute to N₂O formation in this area. The contribution of e.g. denitrification to N₂O production may explain the lack of relationship between delta N₂O and AOU in the ETSP:

‘The presence of key genes of anammox and denitrification assayed and predominantly detected at coastal stations of the ETSP but also present in large parts of the area off Peru points further to an active contribution of mixed processes to N₂O production in the ETSP (the complete dataset of the ETSP can be seen in Löscher (Löscher, 2011)). N₂O production by mixed processes may explain the lack of correlation between Δ N₂O and AOU as well as NO₃⁻ in the ETSP.’

Q6: In Section 4 – Potential importance . . . I do not understand the statement: . . . AOA might dominate the production of N₂O and the balance between reduced and oxidised nitrogen species in the ocean, gradually.

Authors comment: We changed this sentence in the revised manuscript in order to make it clearer and better understandable:

‘Regarding the on-going decrease in dissolved O₂ concentrations in tropical ocean areas (Stramma et al., 2010), we hypothesize that activities of cluster B affiliated AOA might dominate the production of N₂O and the balance between reduced and oxidized nitrogen species in the ocean, as those organisms are likely more adapted to low O₂ concentration.’ Q7: Section 7 is much too short and lacking in detail. Description of the methods should be more involved as should the discussion. It would seem that culture conditions are likely to affect the mechanism by which N₂O is produced, though these are not described.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Authors comment: We agree that the description of the methods used is too short and details are missing thus we changed the section with regard to the methods and discussion and added some more details in the revised manuscript:

'AOB can produce N₂O from NH₂OH during nitrification or from NO₂⁻ during nitrifier-denitrification (Kool et al., 2010; Shaw et al., 2006). In AOA however, the pathway of ammonia oxidation is yet not understood. So far, no equivalent to the hydroxylamine-oxidoreductase, which catalyses the oxidation of NH₂OH to NO₂⁻ during nitrification, has been identified (Könneke et al., 2005; Martens-Habbenha et al., 2009) which means/indicates that AOA likely use a different pathway than AOB do, when producing N₂O. The detection of the nitrite reductase gene *nirK* in the sequenced genomes of cultured *Thaumarchaeota* (Walker et al., 2010) led to the theory that AOA might produce N₂O by nitrifier-denitrification, which might particularly impact at low O₂ concentrations. To identify the origin of N₂O formation isotopomeric studies were performed with *N. maritimus* pure cultures. Using the lowest O₂ concentration of the three chosen (112 μM), a 15N site preference (SPN₂O) in N₂O of 34 ± 12 ‰ was detected, consistent with results from AOA enrichments (Santoro et al., 2011), which is in agreement with the SPN₂O of ~33‰ typically found in AOB cultures performing ammonia oxidation (Sutka et al., 2006) (for comparison: nitrifier-denitrification of AOB results in a SPN₂O of about 0‰. Thus, our dataset points towards a production of N₂O via the oxidation of NH₄⁺ to NO₂⁻, potentially via an unknown intermediate as we were not able to detect NH₂OH in *N. maritimus* cultures using the method described in Schweiger et al. (Schweiger et al., 2007). However, taking δ¹⁸O data into account, Santoro et al. suggested a reduction of NO₂⁻ to N₂O (Santoro et al., 2011), as we have not performed O₂ isotopomeric studies, we cannot exclude N₂O production via nitrifier-denitrification, particularly, when O₂ becomes limiting as previously described for the Arabian Sea (Nicholls et al., 2007) where O₂ concentrations drop far more than in our experiments (lowest O₂ concentration ~112 μM).'

We additionally added more specific details concerning the methods description to the

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

methods section.

Interactive comment on Biogeosciences Discuss., 9, 2095, 2012.

BGD

9, C1371–C1376, 2012

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C1376

