Biogeosciences Discuss., 9, C1464–C1465, 2012 www.biogeosciences-discuss.net/9/C1464/2012/ © Author(s) 2012. This work is distributed under the Creative Commons Attribute 3.0 License.



BGD

9, C1464–C1465, 2012

Interactive Comment

Interactive comment on "Distributions of the carbonate system properties, anthropogenic CO₂, and acidification during the 2008 BOUM cruise (Mediterranean Sea)" *by* F. Touratier et al.

V. VSS Sarma

vvsarma@yahoo.com

Received and published: 25 May 2012

I would like to post a comment on this manuscript by appreciating the efforts of authors on evaluating the influence of anthropogenic CO2 and acidification in the Mediterranean Sea since preindustrial era. They tested the impact of anthropogenic CO2 influence in currently under sampled area (i.e., Med Sea). The authors have used quite accurate and valuable data set of CO2 observations from the Med Sea to estimate anthropogenic CO2 (Cant) and then evaluate its influence on pH variations in deep waters. Quantifying the influence of Cant on acidification in the marine system is one of the most challenging issues for marine biogeochemists. Earlier studies have





been shown that Mediterranean Sea is the most acidified marine system among others, and several reports mentioned that deep waters were not yet acidified. This study shown that significant acidification in the deep waters of Med Sea due to contamination of deep waters with Cant and is quite interesting for the carbon community. In order to quantify Cant, the authors have used the TrOCA method which was developed by the same authors more than a decade ago and has been tested since then at different locations including Med Sea and Atlantic Ocean and made intercomparison exercise with other existing methods. These exercises have been proved that TrOCA method can able to estimate Cant accurately with reference to many other existing methods. Despite variability with reference to space in different methods, the integrated Cant by TrOCA for the entire Atlantic Ocean was guite comparable with other methods indicating that this method can able to estimate Cant accurately. Keeping the data limitation in mind, as no CFC data available, authors have suggested that only two methods, TrOCA and CIPSL method can be used to estimate Cant to this data. Though CIPSL method yielded high total inventory during inter-comparison exercise, however it would still useful to estimate using this method as well and show the range of acidification in the deep waters of Med Sea. I appreciate authors to bring new interesting issue on ocean acidification of deep waters due to anthropogenic CO2 penetration in some parts of the world.

Interactive comment on Biogeosciences Discuss., 9, 2709, 2012.

BGD

9, C1464–C1465, 2012

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

