

Interactive
Comment

Interactive comment on “Nitrate leaching from short-hydroperiod floodplain soils” by B. Huber et al.

Anonymous Referee #1

Received and published: 23 July 2012

It is good to see work is being done on the nitrogen balance in riparian soils. In my opinion the study could be improved by increasing the depth of sampling. The question is, "Has the study reached the actual movement of nitrate into the ground water?"

In a study we were involved in it was possible to assess some of the nitrate inputs. Like you, we were not able to measure all of the nitrate leaving the system. In our study, the periods of flooding and dryness were better known. Some of the nitrate was stored in the soil beyond the reach of plants. This stored nitrate we were able to measure. The soils were in inset alluvial fan positions in a semi-desert environment. Snow floods the soils in these positions in the spring when the snow melts. These soils are known to be Holocene in age and store as much as 17,600 kg of nitrate nitrogen, or 1.8 kg of nitrate per year. Some nitrate would leave the system through the runoff in excess of

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

soil infiltration. So the loss to the system will be greater than the 1.8 kg per year that is stored (Nettleton, W.D. and Peterson, F.F., Landform, soil, and plant relationships to nitrate accumulation, Central Nevada, Geoderma, 2010.).

Hopefully your work will encourage others to investigate the nitrate balance in other riparian areas.

Interactive comment on Biogeosciences Discuss., 9, 5659, 2012.

BGD

9, C2655–C2656, 2012

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C2656

